

## 3D FLOW TRANSITION BEHIND A HEATED CYLINDER

Maosheng Ren, Camilo C.M. Rindt & Anton A. van Steenhoven

Department of Mechanical Engineering, Eindhoven University of Technology, P.O.Box 513, Eindhoven 5600 MB, The Netherlands

**Summary:** In the present study the 3D transition behind a **heated** cylinder is investigated at low Reynolds numbers:  $Re = \mathcal{O}(100)$ . For a Richardson number  $Ri \geq 1.0$ , the 3D transition manifests itself in the form of escaping thermal plumes in the far wake. From a numerical simulation at  $Re = 85$  and  $Ri = 1.0$ , it is observed that the formation of the plumes is associated with pairs of streamwise-vortices occurring close behind the cylinder.

### INTRODUCTION

Over the last ten years, a lot of attention has been paid to the 3D transition of the wake flow behind a cylinder. Experimental, numerical and theoretical studies significantly contributed to the understanding of the dynamics. From a fundamental point of view, studies on the instability mechanisms resulting in the formation of 3D vortical structures, can contribute to a better understanding of laminar-turbulent transition and may lead to ways to actively control the features of the flow.

For the unheated case, the wake flow is characterized by the Reynolds number. For a Reynolds number around 180 – 190, a 3D transition occurs with a spanwise wavelength of 3-4 cylinder diameters. For a Reynolds number around 230 – 260, 3D structures occur with a spanwise wavelength of 1 cylinder diameter.

The effect of a heat input on the behavior of the shed coherent structures received very less attention until the recent studies by Kieft *et al.* [1]. He investigated the wake structure behind a heated horizontal cylinder in cross-flow with water as the working fluid. For low Reynolds numbers, Kieft *et al.* [1] found that by heating the cylinder, the vortices in the upper and lower rows acquire different strengths due to baroclinic vorticity production. This strength difference increases for increasing Richardson numbers. It was observed that for small Richardson numbers the shed vortices move slightly downwards. A more extensive literature review and a summary of earlier results on the wake behavior behind a heated cylinder of our research group is given in van Steenhoven and Rindt [2].

### THERMAL PLUMES IN THE FAR WAKE

In the present study the 3D transition of the wake flow behind a **heated** cylinder is investigated at a Reynolds number  $Re = \mathcal{O}(100)$ . In Figure 1 (a), for  $Ri = 0$ , the wake flow manifests itself as the von Kármán vortex street. If one increases the heat input, for  $Ri \geq 1$  a 3D transition manifests itself in the form of escaping thermal plumes [3], see Figure 1 (b). The spanwise distance of the plumes is around 2 cylinder diameters, as shown in Figure 1 (c).




Figure 1: Visualization of the wake flow at  $Re = 117$  and  $Ri = 0$ ,  $Ri = 1$ .

## STREAMWISE VORTICITY IN THE NEAR WAKE

To understand the flow structures observed from the experiments, Spectral Element calculations are carried out for  $Re = 85$  and  $Ri = 1.0$ . Numerical simulations provide more detailed insight into the occurrence of the 3D transition.

In Figure 2, iso-vorticity surface  $\omega_x = \pm 0.4$  (vorticity component in streamwise direction) and iso-temperature surface  $\Theta = 0.2$  are presented. In Figure 2 (a), the flow is from left to right. It is observed that pairs of streamwise vortices appear close behind the cylinder. The streamwise-vortices-pair has a spanwise distance of around 2 cylinder diameters. In Figure 2 (b), the flow is pointing towards the observer. The fluid at the center region of the streamwise-vortices-pair has a high temperature (Ren *et al.* [4]). Due to the non-uniform temperature distribution in spanwise direction, streamwise vorticity  $\omega_x$  is generated due to the baroclinic vorticity torque  $-\partial_z \Theta \times \mathbf{g}$ .




Figure 2: Snap-shot of calculated iso-vorticity  $\omega_x = \pm 0.4$  surfaces and iso-temperature  $\Theta = 0.2$  surface at  $Ri = 1.0$ ,  $Re = 85$ . The spanwise distance of the streamwise-vortices-pair is around  $2d$ ,  $d$  is the cylinder diameters.

Furthermore, we observe the preliminary stage of plume-like structures in the upper vortices, as indicated in Figure 2. From the numerical calculations, the formation of the plume-like structures is associated with the streamwise-vortices-pair 'curling' around the upper vortices.

### AIM AND PLAN

The evolution of the streamwise-vortices-pair and the formation of the escaping plumes will be studied as function of the Richardson number. From a fundamental point of view, a detailed study on the origin of the streamwise vorticity in the near wake and the thermal plumes in the far wake will contribute to a better understanding of 3D transition behind heated bluff bodies. This will be the goal of the current research.

### References

- [1] R. N. Kieft, C. C. M. Rindt, A. A. van Steenhoven, and G. J. F. van Heijst. On the wake structure behind a heated horizontal cylinder in cross-flow. *J. Fluid Mech.*, 486:189, 2003.
- [2] A. A. van Steenhoven and C. C. M. Rindt. Flow transition behind a heated cylinder. *Int. J. Heat Fluid Flow*, 24:322, 2003.
- [3] W. J. P. M. Maas, C. C. M. Rindt, and A. A. van Steenhoven. The influence of heat on the 3d-transition of the von kármán-vortex street. *Int. J. Heat Mass Transfer*, 46:3069, 2003.
- [4] M. Ren, C. C. M. Rindt, and A. A. van Steenhoven. Buoyancy induced 3d-transition behind a heated cylinder. *Proc. Third International Symposium on Turbulence and Shear Flow Phenomena*, page 717, 2003.