

# 2

## Sequences

CHAPTER

### §7 Limits of Sequences

A *sequence* is a function whose domain is a set that has the form  $\{n \in \mathbb{Z} : n \geq m\}$ ;  $m$  is usually 1 or 0. Thus a sequence is a function that has a specified value for each integer  $n \geq m$ . It is customary to denote a sequence by a letter such as  $s$  and to denote its value at  $n$  as  $s_n$  rather than  $s(n)$ . It is often convenient to write the sequence as  $(s_n)_{n=m}^{\infty}$  or  $(s_m, s_{m+1}, s_{m+2}, \dots)$ . If  $m = 1$  we may write  $(s_n)_{n \in \mathbb{N}}$  or of course  $(s_1, s_2, s_3, \dots)$ . Sometimes we will write  $(s_n)$  when the domain is understood or when the results under discussion do not depend on the specific value of  $m$ . In this chapter, we will be interested in sequences whose range values are real numbers, i.e., each  $s_n$  represents a real number.

#### Example 1

- (a) Consider the sequence  $(s_n)_{n \in \mathbb{N}}$  where  $s_n = \frac{1}{n^2}$ . This is the sequence  $(1, \frac{1}{4}, \frac{1}{9}, \frac{1}{16}, \frac{1}{25}, \dots)$ . Formally, of course, this is the function with domain  $\mathbb{N}$  whose value at each  $n$  is  $\frac{1}{n^2}$ . The set of values is  $\{1, \frac{1}{4}, \frac{1}{9}, \frac{1}{16}, \frac{1}{25}, \dots\}$ .
- (b) Consider the sequence given by  $a_n = (-1)^n$  for  $n \geq 0$ , i.e.,  $(a_n)_{n=0}^{\infty}$  where  $a_n = (-1)^n$ . Note that the first term of the se-

quence is  $a_0 = 1$  and the sequence is  $(1, -1, 1, -1, 1, -1, 1, \dots)$ . Formally, this is a function whose domain is  $\{0, 1, 2, \dots\}$  and whose *set* of values is  $\{-1, 1\}$ .

It is important to distinguish between a sequence and its set of values, since the validity of many results in this book depends on whether we are working with a sequence or a set. We will always use parentheses  $( )$  to signify a sequence and braces  $\{ \}$  to signify a set. The sequence given by  $a_n = (-1)^n$  has an infinite number of terms even though their values are repeated over and over. On the other hand, the *set*  $\{(-1)^n : n = 0, 1, 2, \dots\}$  is exactly the set  $\{-1, 1\}$  consisting of two numbers.

- (c) Consider the sequence  $\cos(\frac{n\pi}{3})$ ,  $n \in \mathbb{N}$ . The first term of this sequence is  $\cos(\frac{\pi}{3}) = \cos 60^\circ = \frac{1}{2}$  and the sequence looks like

$$(\frac{1}{2}, -\frac{1}{2}, -1, -\frac{1}{2}, \frac{1}{2}, 1, \frac{1}{2}, -\frac{1}{2}, -1, -\frac{1}{2}, \frac{1}{2}, 1, \frac{1}{2}, -\frac{1}{2}, -1, \dots).$$

The set of values is  $\{\cos(\frac{n\pi}{3}) : n \in \mathbb{N}\} = \{\frac{1}{2}, -\frac{1}{2}, -1, 1\}$ .

- (d) If  $a_n = n^{1/n}$ ,  $n \in \mathbb{N}$ , the sequence is  $(1, \sqrt{2}, 3^{1/3}, 4^{1/4}, \dots)$ . If we approximate values to four decimal places, the sequence looks like

$$(1, 1.4142, 1.4422, 1.4142, 1.3797, 1.3480, 1.3205, 1.2968, \dots).$$

It turns out that  $a_{100}$  is approximately 1.0471 and that  $a_{1000}$  is approximately 1.0069.

- (e) Consider the sequence  $b_n = (1 + \frac{1}{n})^n$ ,  $n \in \mathbb{N}$ . This is the sequence  $(2, (\frac{3}{2})^2, (\frac{4}{3})^3, (\frac{5}{4})^4, \dots)$ . If we approximate the values to four decimal places, we obtain

$$(2, 2.25, 2.3704, 2.4414, 2.4883, 2.5216, 2.5465, 2.5658, \dots).$$

Also  $b_{100}$  is approximately 2.7048 and  $b_{1000}$  is approximately 2.7169.

The “limit” of a sequence  $(s_n)$  is a real number that the values  $s_n$  are close to for large values of  $n$ . For instance, the values of the sequence in Example 1(a) are close to 0 for large  $n$  and the values of the sequence in Example 1(d) appear to be close to 1 for large  $n$ . The sequence  $(a_n)$  given by  $a_n = (-1)^n$  requires some thought. We might say that 1 is a limit because in fact  $a_n = 1$  for the large values of  $n$  that are even. On the other hand,  $a_n = -1$  [which is quite a distance

from 1] for other large values of  $n$ . We need a concise definition in order to decide whether 1 is a limit of  $a_n = (-1)^n$ . It turns out that our definition will require the values to be close to the limit value for *all* large  $n$ , so 1 will *not* be a limit of the sequence  $a_n = (-1)^n$ .

### 7.1 Definition.

A sequence  $(s_n)$  of real numbers is said to *converge* to the real number  $s$  provided that

$$\text{for each } \epsilon > 0 \text{ there exists a number } N \text{ such that} \quad (1) \\ n > N \text{ implies } |s_n - s| < \epsilon.$$

If  $(s_n)$  converges to  $s$ , we will write  $\lim_{n \rightarrow \infty} s_n = s$ , or  $s_n \rightarrow s$ . The number  $s$  is called the *limit* of the sequence  $(s_n)$ . A sequence that does not converge to some real number is said to *diverge*.

Several comments are in order. First, in view of the Archimedean property, the number  $N$  in Definition 7.1 can be taken to be a natural number if we wish. Second, the symbol  $\epsilon$  [lower case Greek epsilon] in this definition represents a positive number, not some new exotic number. However, it is traditional in mathematics to use  $\epsilon$  and  $\delta$  [lower case Greek delta] in situations where the interesting or challenging values are the small positive values. Third, condition (1) is an infinite number of statements, one for each positive value of  $\epsilon$ . The condition states that to each  $\epsilon > 0$  there corresponds a number  $N$  with a certain property, namely  $n > N$  implies  $|s_n - s| < \epsilon$ . The value  $N$  depends on the value  $\epsilon$ , and normally  $N$  must be large if  $\epsilon$  is small. We illustrate these remarks in the next example.

### Example 2

Consider the sequence  $s_n = \frac{3n+1}{7n-4}$ . If we write  $s_n$  as  $\frac{3+\frac{1}{n}}{7-\frac{4}{n}}$  and note that  $\frac{1}{n}$  and  $\frac{4}{n}$  are very small for large  $n$ , it seems reasonable to conclude that  $\lim s_n = \frac{3}{7}$ . In fact, this reasoning will be completely valid after we have the limit theorems in §9:

$$\lim s_n = \lim \left[ \frac{3 + \frac{1}{n}}{7 - \frac{4}{n}} \right] = \frac{\lim 3 + \lim(\frac{1}{n})}{\lim 7 - 4 \lim(\frac{1}{n})} = \frac{3 + 0}{7 - 4 \cdot 0} = \frac{3}{7}.$$

However, for now we are interested in analyzing exactly what we mean by  $\lim s_n = \frac{3}{7}$ . By Definition 7.1,  $\lim s_n = \frac{3}{7}$  means that

$$\begin{aligned} &\text{for each } \epsilon > 0 \text{ there exists a number } N \text{ such that} \\ &n > N \text{ implies } \left| \frac{3n+1}{7n-4} - \frac{3}{7} \right| < \epsilon. \end{aligned} \tag{1}$$

As  $\epsilon$  varies,  $N$  varies. In Example 2 of the next section we will show that, for this particular sequence,  $N$  can be taken to be  $\frac{19}{49\epsilon} + \frac{4}{7}$ . Using this observation and a calculator, we find that for  $\epsilon$  equal to 1, 0.1, 0.01, 0.001 and 0.000001, respectively,  $N$  can be taken to be approximately 0.96, 4.45, 39.35, 388.33 and 387,755.67, respectively. Since we are interested only in integer values of  $n$ , we may as well drop the fractional part of  $N$ . Then we see that five of the infinitely many statements given by (1) are:

$$n > 0 \quad \text{implies} \quad \left| \frac{3n+1}{7n-4} - \frac{3}{7} \right| < 1; \tag{2}$$

$$n > 4 \quad \text{implies} \quad \left| \frac{3n+1}{7n-4} - \frac{3}{7} \right| < 0.1; \tag{3}$$

$$n > 39 \quad \text{implies} \quad \left| \frac{3n+1}{7n-4} - \frac{3}{7} \right| < 0.01; \tag{4}$$

$$n > 388 \quad \text{implies} \quad \left| \frac{3n+1}{7n-4} - \frac{3}{7} \right| < 0.001; \tag{5}$$

$$n > 387,755 \quad \text{implies} \quad \left| \frac{3n+1}{7n-4} - \frac{3}{7} \right| < 0.000001. \tag{6}$$

Table 7.1 partially confirms assertions (2) through (6). We could go on and on with these numerical illustrations, but it should be clear that we need a more theoretical approach if we are going to *prove* results about limits.

Table 7.1

$n$	$s_n = \frac{3n+1}{7n-4}$ approximately	$ s_n - \frac{3}{7} $ approximately
1	1.3333	.9047
2	0.7000	.2714
3	0.5882	.1597
4	0.5417	.1131
5	0.5161	.0876
6	0.5000	.0714
40	0.4384	.0098
400	0.4295	.0010

**Example 3**

We return to the examples in Example 1.

- (a)  $\lim \frac{1}{n^2} = 0$ . This will be proved in Example 1 of the next section.
- (b) The sequence  $(a_n)$  where  $a_n = (-1)^n$  does not converge. Thus the expression “ $\lim a_n$ ” is meaningless in this case. We will discuss this example again in Example 4 of the next section.
- (c) The sequence  $\cos(\frac{n\pi}{3})$  does not converge. See Exercise 8.7.
- (d) The sequence  $n^{1/n}$  appears to converge to 1. We will prove  $\lim n^{1/n} = 1$  in 9.7(c).
- (e) The sequence  $(b_n)$  where  $b_n = (1 + \frac{1}{n})^n$  converges to the number  $e$  that should be familiar from calculus. The limit  $\lim b_n$  and the number  $e$  will be discussed further in the optional §37. Recall that  $e$  is approximately 2.7182818.

We conclude this section by showing that limits are unique. That is, if  $\lim s_n = s$  and  $\lim s_n = t$ , then we must have  $s = t$ . In short, the values  $s_n$  cannot be getting arbitrarily close to different values for large  $n$ . To prove this, consider  $\epsilon > 0$ . By the definition of limit there must exist  $N_1$  so that

$$n > N_1 \quad \text{implies} \quad |s_n - s| < \frac{\epsilon}{2}$$

and there must exist  $N_2$  so that

$$n > N_2 \quad \text{implies} \quad |s_n - t| < \frac{\epsilon}{2}.$$

For  $n > \max\{N_1, N_2\}$ , the Triangle Inequality 3.7 shows that

$$|s - t| = |(s - s_n) + (s_n - t)| \leq |s - s_n| + |s_n - t| \leq \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon.$$

This shows that  $|s - t| < \epsilon$  for all  $\epsilon > 0$ . It follows that  $|s - t| = 0$ , hence  $s = t$ .

## Exercises

7.1. Write out the first five terms of the following sequences.

(a)  $s_n = \frac{1}{3n+1}$   
(c)  $c_n = \frac{n}{3^n}$

(b)  $b_n = \frac{3n+1}{4n-1}$   
(d)  $\sin(\frac{n\pi}{4})$

7.2. For each sequence in Exercise 7.1, determine whether it converges. If it converges, give its limit. No proofs are required.

7.3. For each sequence below, determine whether it converges and, if it converges, give its limit. No proofs are required.

(a)  $a_n = \frac{n}{n+1}$

(b)  $b_n = \frac{n^2+3}{n^2-3}$

(c)  $c_n = 2^{-n}$

(d)  $t_n = 1 + \frac{2}{n}$

(e)  $x_n = 73 + (-1)^n$

(f)  $s_n = (2)^{1/n}$

(g)  $y_n = n!$

(h)  $d_n = (-1)^n n$

(i)  $\frac{(-1)^n}{n}$

(j)  $\frac{7n^3+8n}{2n^3-31}$

(k)  $\frac{9n^2-18}{6n+18}$

(l)  $\sin(\frac{n\pi}{2})$

(m)  $\sin(n\pi)$

(n)  $\sin(\frac{2n\pi}{3})$

(o)  $\frac{1}{n} \sin n$

(p)  $\frac{2^{n+1}+5}{2^n-7}$

(q)  $\frac{3^n}{n!}$

(r)  $(1 + \frac{1}{n})^2$

(s)  $\frac{4n^2+3}{3n^2-2}$

(t)  $\frac{6n+4}{9n^2+7}$

7.4. Give examples of

(a) a sequence  $(x_n)$  of irrational numbers having a limit  $\lim x_n$  that is a rational number.

(b) a sequence  $(r_n)$  of rational numbers having a limit  $\lim r_n$  that is an irrational number.

7.5. Determine the following limits. No proofs are required, but show any relevant algebra.

(a)  $\lim s_n$  where  $s_n = \sqrt{n^2 + 1} - n$ ,

(b)  $\lim(\sqrt{n^2 + n} - n)$ ,

(c)  $\lim(\sqrt{4n^2 + n} - 2n).$

*Hint for (a):* First show that  $s_n = \frac{1}{\sqrt{n^2+1}+n}.$

## §8 A Discussion about Proofs

In this section we give several examples of proofs using the definition of the limit of a sequence. With a little study and practice, students should be able to do proofs of this sort themselves. We will sometimes refer to a proof as a *formal proof* to emphasize that it is a rigorous mathematical proof.

### Example 1

Prove that  $\lim \frac{1}{n^2} = 0.$

*Discussion.* Our task is to consider an arbitrary  $\epsilon > 0$  and show that there exists a number  $N$  [which will depend on  $\epsilon$ ] such that  $n > N$  implies  $|\frac{1}{n^2} - 0| < \epsilon.$  So we expect our formal proof to begin with “Let  $\epsilon > 0$ ” and to end with something like “Hence  $n > N$  implies  $|\frac{1}{n^2} - 0| < \epsilon.$ ” In between the proof should specify an  $N$  and then verify that  $N$  has the desired property, namely that  $n > N$  does indeed imply  $|\frac{1}{n^2} - 0| < \epsilon.$

As is often the case with trigonometric identities, we will initially work backward from our desired conclusion, but in the formal proof we will have to be sure that our steps are reversible. In the present example, we want  $|\frac{1}{n^2} - 0| < \epsilon$  and we want to know how big  $n$  must be. So we will operate on this inequality algebraically and try to “solve” for  $n.$  Thus we want  $\frac{1}{n^2} < \epsilon.$  By multiplying both sides by  $n^2$  and dividing both sides by  $\epsilon,$  we find that we want  $\frac{1}{\epsilon} < n^2$  or  $\frac{1}{\sqrt{\epsilon}} < n.$  If our steps are reversible, we see that  $n > \frac{1}{\sqrt{\epsilon}}$  implies  $|\frac{1}{n^2} - 0| < \epsilon.$  This suggests that we put  $N = \frac{1}{\sqrt{\epsilon}}.$

### Formal Proof

Let  $\epsilon > 0.$  Let  $N = \frac{1}{\sqrt{\epsilon}}.$  Then  $n > N$  implies  $n > \frac{1}{\sqrt{\epsilon}}$  which implies  $n^2 > \frac{1}{\epsilon}$  and hence  $\epsilon > \frac{1}{n^2}.$  Thus  $n > N$  implies  $|\frac{1}{n^2} - 0| < \epsilon.$  This proves that  $\lim \frac{1}{n^2} = 0.$  ■

**Example 2**

Prove that  $\lim_{n \rightarrow \infty} \frac{3n+1}{7n-4} = \frac{3}{7}$ .

*Discussion.* For each  $\epsilon > 0$ , we need to decide how big  $n$  must be to guarantee that  $|\frac{3n+1}{7n-4} - \frac{3}{7}| < \epsilon$ . Thus we want

$$\left| \frac{21n + 7 - 21n + 12}{7(7n - 4)} \right| < \epsilon \quad \text{or} \quad \left| \frac{19}{7(7n - 4)} \right| < \epsilon.$$

Since  $7n - 4 > 0$ , we can drop the absolute value and manipulate the inequality further to “solve” for  $n$ :

$$\frac{19}{7\epsilon} < 7n - 4 \quad \text{or} \quad \frac{19}{7\epsilon} + 4 < 7n \quad \text{or} \quad \frac{19}{49\epsilon} + \frac{4}{7} < n.$$

Our steps are reversible, so we will put  $N = \frac{19}{49\epsilon} + \frac{4}{7}$ . Incidentally, we could have chosen  $N$  to be any number larger than  $\frac{19}{49\epsilon} + \frac{4}{7}$ .

**Formal Proof**

Let  $\epsilon > 0$  and let  $N = \frac{19}{49\epsilon} + \frac{4}{7}$ . Then  $n > N$  implies  $n > \frac{19}{49\epsilon} + \frac{4}{7}$ , hence  $7n > \frac{19}{7\epsilon} + 4$ , hence  $7n - 4 > \frac{19}{7\epsilon}$ , hence  $\frac{19}{7(7n-4)} < \epsilon$ , and hence  $|\frac{3n+1}{7n-4} - \frac{3}{7}| < \epsilon$ . This proves  $\lim_{n \rightarrow \infty} \frac{3n+1}{7n-4} = \frac{3}{7}$ . ■

**Example 3**

Prove that  $\lim_{n \rightarrow \infty} \frac{4n^3+3n}{n^3-6} = 4$ .

*Discussion.* For each  $\epsilon > 0$ , we need to determine how large  $n$  must be to imply

$$\left| \frac{4n^3 + 3n}{n^3 - 6} - 4 \right| < \epsilon \quad \text{or} \quad \left| \frac{3n + 24}{n^3 - 6} \right| < \epsilon.$$

By considering  $n > 1$ , we may drop the absolute values; thus we need to find how big  $n$  must be to give  $\frac{3n+24}{n^3-6} < \epsilon$ . This time it would be very difficult to “solve” for or isolate  $n$ . Recall that we need to find some  $N$  such that  $n > N$  implies  $\frac{3n+24}{n^3-6} < \epsilon$ , but we do not need to find the least such  $N$ . So we will simplify matters by making estimates. The idea is that  $\frac{3n+24}{n^3-6}$  is bounded by some constant times  $\frac{n}{n^3} = \frac{1}{n^2}$  for sufficiently large  $n$ . To find such a bound we will find an upper bound for the numerator and a lower bound for the denominator. For example, since  $3n + 24 \leq 27n$ , it suffices for us to get  $\frac{27n}{n^3-6} < \epsilon$ . To make the denominator smaller and yet a constant multiple of  $n^3$ , we note that  $n^3 - 6 \geq \frac{n^3}{2}$  provided  $n$  is sufficiently large; in fact, all

we need is  $\frac{n^3}{2} \geq 6$  or  $n^3 \geq 12$  or  $n > 2$ . So it suffices to get  $\frac{27n}{n^{3/2}} < \epsilon$  or  $\frac{54}{n^2} < \epsilon$  or  $n > \sqrt{\frac{54}{\epsilon}}$ , provided that  $n > 2$ .

**Formal Proof**

Let  $\epsilon > 0$  and let  $N = \max\{2, \sqrt{\frac{54}{\epsilon}}\}$ . Then  $n > N$  implies  $n > \sqrt{\frac{54}{\epsilon}}$ , hence  $\frac{54}{n^2} < \epsilon$ , hence  $\frac{27n}{n^{3/2}} < \epsilon$ . Since  $n > 2$ , we have  $\frac{n^3}{2} \leq n^3 - 6$  and also  $27n \geq 3n + 24$ . Thus  $n > N$  implies

$$\frac{3n + 24}{n^3 - 6} \leq \frac{27n}{\frac{1}{2}n^3} = \frac{54}{n^2} < \epsilon,$$

and hence

$$\left| \frac{4n^3 + 3n}{n^3 - 6} - 4 \right| < \epsilon,$$

as desired. ■

Example 3 illustrates that direct proofs of even rather simple limits can get complicated. With the limit theorems of §9 we would just write

$$\lim \left[ \frac{4n^3 + 3n}{n^3 - 6} \right] = \lim \left[ \frac{4 + \frac{3}{n^2}}{1 - \frac{6}{n^3}} \right] = \frac{\lim 4 + 3 \cdot \lim(\frac{1}{n^2})}{\lim 1 - 6 \cdot \lim(\frac{1}{n^3})} = 4.$$

**Example 4**

Show that the sequence  $a_n = (-1)^n$  does not converge.

*Discussion.* We will assume that  $\lim(-1)^n = a$  and obtain a contradiction. No matter what  $a$  is, either 1 or  $-1$  will have distance at least 1 from  $a$ . Thus the inequality  $|(-1)^n - a| < 1$  will not hold for all large  $n$ .

**Formal Proof**

Assume that  $\lim(-1)^n = a$  for some  $a \in \mathbb{R}$ . Letting  $\epsilon = 1$  in the definition of the limit, we see that there exists  $N$  such that

$$n > N \quad \text{implies} \quad |(-1)^n - a| < 1.$$

By considering both an even and an odd  $n > N$ , we see that

$$|1 - a| < 1 \quad \text{and} \quad |-1 - a| < 1.$$

Now by the Triangle Inequality 3.7

$$2 = |1 - (-1)| = |1 - a + a - (-1)| \leq |1 - a| + |a - (-1)| < 1 + 1 = 2.$$

This absurdity shows that our assumption that  $\lim(-1)^n = a$  must be wrong, so the sequence  $(-1)^n$  does not converge. ■

### Example 5

Let  $(s_n)$  be a sequence of nonnegative real numbers and suppose that  $s = \lim s_n$ . Note that  $s \geq 0$ ; see Exercise 8.9(a). Prove that  $\lim \sqrt{s_n} = \sqrt{s}$ .

*Discussion.* We must consider  $\epsilon > 0$  and show that there exists  $N$  such that

$$n > N \quad \text{implies} \quad |\sqrt{s_n} - \sqrt{s}| < \epsilon.$$

This time we cannot expect to obtain  $N$  explicitly in terms of  $\epsilon$  because of the general nature of the problem. But we can hope to show such  $N$  exists. The trick here is to violate our training in algebra and “irrationalize the denominator”:

$$\sqrt{s_n} - \sqrt{s} = \frac{(\sqrt{s_n} - \sqrt{s})(\sqrt{s_n} + \sqrt{s})}{\sqrt{s_n} + \sqrt{s}} = \frac{s_n - s}{\sqrt{s_n} + \sqrt{s}}.$$

Since  $s_n \rightarrow s$  we will be able to make the numerator small [for large  $n$ ]. Unfortunately, if  $s = 0$  the denominator will also be small. So we consider two cases. If  $s > 0$ , the denominator is bounded below by  $\sqrt{s}$  and our trick will work:

$$|\sqrt{s_n} - \sqrt{s}| \leq \frac{|s_n - s|}{\sqrt{s}},$$

so we will select  $N$  so that  $|s_n - s| < \sqrt{s}\epsilon$  for  $n > N$ . Note that  $N$  exists, since we can apply the definition of limit to  $\sqrt{s}\epsilon$  just as well as to  $\epsilon$ . For  $s = 0$ , it can be shown directly that  $\lim s_n = 0$  implies  $\lim \sqrt{s_n} = 0$ ; the trick of “irrationalizing the denominator” is not needed in this case.

### Formal Proof

Case I:  $s > 0$ . Let  $\epsilon > 0$ . Since  $\lim s_n = s$ , there exists  $N$  such that

$$n > N \quad \text{implies} \quad |s_n - s| < \sqrt{s}\epsilon.$$

Now  $n > N$  implies

$$|\sqrt{s_n} - \sqrt{s}| = \frac{|s_n - s|}{\sqrt{s_n} + \sqrt{s}} \leq \frac{|s_n - s|}{\sqrt{s}} < \frac{\sqrt{s}\epsilon}{\sqrt{s}} = \epsilon.$$

Case II:  $s = 0$ . This case is left to Exercise 8.3. ■

### Example 6

Let  $(s_n)$  be a convergent sequence of real numbers such that  $s_n \neq 0$  for all  $n \in \mathbb{N}$  and  $\lim s_n = s \neq 0$ . Prove that  $\inf\{|s_n| : n \in \mathbb{N}\} > 0$ .

*Discussion.* The idea is that “most” of the terms  $s_n$  are close to  $s$  and hence not close to 0. More explicitly, “most” of the terms  $s_n$  are within  $\frac{1}{2}|s|$  of  $s$ , hence most  $s_n$  satisfy  $|s_n| \geq \frac{1}{2}|s|$ . This seems clear from Figure 8.1, but a formal proof will use the triangle inequality.

### Formal Proof

Let  $\epsilon = \frac{1}{2}|s| > 0$ . Since  $\lim s_n = s$ , there exists  $N$  in  $\mathbb{N}$  so that

$$n > N \quad \text{implies} \quad |s_n - s| < \frac{|s|}{2}.$$

Now

$$n > N \quad \text{implies} \quad |s_n| \geq \frac{|s|}{2}, \tag{1}$$

since otherwise the triangle inequality would imply

$$|s| = |s - s_n + s_n| \leq |s - s_n| + |s_n| < \frac{|s|}{2} + \frac{|s|}{2} = |s|$$

which is absurd. If we set

$$m = \min \left\{ \frac{|s|}{2}, |s_1|, |s_2|, \dots, |s_N| \right\},$$

FIGURE 8.1

then we clearly have  $m > 0$  and  $|s_n| \geq m$  for all  $n \in \mathbb{N}$  in view of (1). Thus  $\inf\{|s_n| : n \in \mathbb{N}\} \geq m > 0$ , as desired. ■

Formal proofs are required in the following exercises.

## Exercises

8.1. Prove the following:

$$\begin{array}{ll} \text{(a)} \lim \frac{(-1)^n}{n} = 0 & \text{(b)} \lim \frac{1}{n^{1/3}} = 0 \\ \text{(c)} \lim \frac{2n-1}{3n+2} = \frac{2}{3} & \text{(d)} \lim \frac{n+6}{n^2-6} = 0 \end{array}$$

8.2. Determine the limits of the following sequences, and then prove your claims.

$$\begin{array}{ll} \text{(a)} a_n = \frac{n}{n^2+1} & \text{(b)} b_n = \frac{7n-19}{3n+7} \\ \text{(c)} c_n = \frac{4n+3}{7n-5} & \text{(d)} d_n = \frac{2n+4}{5n+2} \\ \text{(e)} s_n = \frac{1}{n} \sin n \end{array}$$

8.3. Let  $(s_n)$  be a sequence of nonnegative real numbers, and suppose that  $\lim s_n = 0$ . Prove that  $\lim \sqrt{s_n} = 0$ . This will complete the proof for Example 5.

8.4. Let  $(t_n)$  be a bounded sequence, i.e., there exists  $M$  such that  $|t_n| \leq M$  for all  $n$ , and let  $(s_n)$  be a sequence such that  $\lim s_n = 0$ . Prove that  $\lim(s_n t_n) = 0$ .

8.5. (a) Consider three sequences  $(a_n)$ ,  $(b_n)$  and  $(s_n)$  such that  $a_n \leq s_n \leq b_n$  for all  $n$  and  $\lim a_n = \lim b_n = s$ . Prove that  $\lim s_n = s$ .

(b) Suppose that  $(s_n)$  and  $(t_n)$  are sequences such that  $|s_n| \leq t_n$  for all  $n$  and  $\lim t_n = 0$ . Prove that  $\lim s_n = 0$ .

8.6. Let  $(s_n)$  be a sequence in  $\mathbb{R}$ .

(a) Prove that  $\lim s_n = 0$  if and only if  $\lim |s_n| = 0$ .

(b) Observe that if  $s_n = (-1)^n$ , then  $\lim |s_n|$  exists, but  $\lim s_n$  does not exist.

8.7. Show that the following sequences do not converge.

$$\begin{array}{ll} \text{(a)} \cos\left(\frac{n\pi}{3}\right) & \text{(b)} s_n = (-1)^n n \\ \text{(c)} \sin\left(\frac{n\pi}{3}\right) \end{array}$$

8.8. Prove the following [see Exercise 7.5]:

$$\begin{array}{ll} \text{(a)} \lim[\sqrt{n^2+1} - n] = 0 & \text{(b)} \lim[\sqrt{n^2+n} - n] = \frac{1}{2} \\ \text{(c)} \lim[\sqrt{4n^2+n} - 2n] = \frac{1}{4} \end{array}$$

**8.9.** Let  $(s_n)$  be a sequence that converges.

- (a) Show that if  $s_n \geq a$  for all but finitely many  $n$ , then  $\lim s_n \geq a$ .
- (b) Show that if  $s_n \leq b$  for all but finitely many  $n$ , then  $\lim s_n \leq b$ .
- (c) Conclude that if all but finitely many  $s_n$  belong to  $[a, b]$ , then  $\lim s_n$  belongs to  $[a, b]$ .

**8.10.** Let  $(s_n)$  be a convergent sequence, and suppose that  $\lim s_n > a$ . Prove that there exists a number  $N$  such that  $n > N$  implies  $s_n > a$ .

## §9 Limit Theorems for Sequences

In this section we prove some basic results that are probably already familiar to the reader. First we prove that convergent sequences are bounded. A sequence  $(s_n)$  of real numbers is said to be *bounded* if the set  $\{s_n : n \in \mathbb{N}\}$  is a bounded set, i.e., if there exists a constant  $M$  such that  $|s_n| \leq M$  for all  $n$ .

### 9.1 Theorem.

*Convergent sequences are bounded.*

#### Proof

Let  $(s_n)$  be a convergent sequence, and let  $s = \lim s_n$ . Applying Definition 7.1 with  $\epsilon = 1$  we obtain  $N$  in  $\mathbb{N}$  so that

$$n > N \quad \text{implies} \quad |s_n - s| < 1.$$

From the triangle inequality we see that  $n > N$  implies  $|s_n| < |s| + 1$ . Define  $M = \max\{|s| + 1, |s_1|, |s_2|, \dots, |s_N|\}$ . Then we have  $|s_n| \leq M$  for all  $n \in \mathbb{N}$ , so  $(s_n)$  is a bounded sequence. ■

In the proof of Theorem 9.1 we only needed to use property 7.1(1) for a single value of  $\epsilon$ . Our choice of  $\epsilon = 1$  was quite arbitrary.

### 9.2 Theorem.

*If the sequence  $(s_n)$  converges to  $s$  and  $k \in \mathbb{R}$ , then the sequence  $(ks_n)$  converges to  $ks$ . That is,  $\lim(ks_n) = k \lim s_n$ .*

**Proof**

We assume  $k \neq 0$ , since this result is trivial for  $k = 0$ . Let  $\epsilon > 0$  and note that we need to show that  $|ks_n - ks| < \epsilon$  for large  $n$ . Since  $\lim s_n = s$ , there exists  $N$  such that

$$n > N \quad \text{implies} \quad |s_n - s| < \frac{\epsilon}{|k|}.$$

Then

$$n > N \quad \text{implies} \quad |ks_n - ks| < \epsilon. \quad \blacksquare$$

**9.3 Theorem.**

*If  $(s_n)$  converges to  $s$  and  $(t_n)$  converges to  $t$ , then  $(s_n + t_n)$  converges to  $s + t$ . That is,*

$$\lim(s_n + t_n) = \lim s_n + \lim t_n.$$

**Proof**

Let  $\epsilon > 0$ ; we need to show that

$$|s_n + t_n - (s + t)| < \epsilon \quad \text{for large } n.$$

We note that  $|s_n + t_n - (s + t)| \leq |s_n - s| + |t_n - t|$ . Since  $\lim s_n = s$ , there exists  $N_1$  such that

$$n > N_1 \quad \text{implies} \quad |s_n - s| < \frac{\epsilon}{2}.$$

Likewise, there exists  $N_2$  such that

$$n > N_2 \quad \text{implies} \quad |t_n - t| < \frac{\epsilon}{2}.$$

Let  $N = \max\{N_1, N_2\}$ . Then clearly

$$n > N \quad \text{implies} \quad |s_n + t_n - (s + t)| \leq |s_n - s| + |t_n - t| < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon. \quad \blacksquare$$

**9.4 Theorem.**

*If  $(s_n)$  converges to  $s$  and  $(t_n)$  converges to  $t$ , then  $(s_n t_n)$  converges to  $st$ . That is,*

$$\lim(s_n t_n) = (\lim s_n)(\lim t_n).$$

*Discussion.* The trick here is to look at the inequality

$$\begin{aligned} |s_n t_n - st| &= |s_n t_n - s_n t + s_n t - st| \\ &\leq |s_n t_n - s_n t| + |s_n t - st| = |s_n| \cdot |t_n - t| + |t| \cdot |s_n - s|. \end{aligned}$$

For large  $n$ ,  $|t_n - t|$  and  $|s_n - s|$  are small and  $|t|$  is, of course, constant. Fortunately, Theorem 9.1 shows that  $|s_n|$  is bounded, so we will be able to show that  $|s_n t_n - st|$  is small.

**Proof**

Let  $\epsilon > 0$ . By Theorem 9.1 there is a constant  $M > 0$  such that  $|s_n| \leq M$  for all  $n$ . Since  $\lim t_n = t$  there exists  $N_1$  such that

$$n > N_1 \quad \text{implies} \quad |t_n - t| < \frac{\epsilon}{2M}.$$

Also, since  $\lim s_n = s$  there exists  $N_2$  such that

$$n > N_2 \quad \text{implies} \quad |s_n - s| < \frac{\epsilon}{2(|t| + 1)}.$$

[We used  $\frac{\epsilon}{2(|t|+1)}$  instead of  $\frac{\epsilon}{2|t|}$ , since  $t$  could be 0.] Now if  $N = \max\{N_1, N_2\}$ , then  $n > N$  implies

$$\begin{aligned} |s_n t_n - st| &\leq |s_n| \cdot |t_n - t| + |t| \cdot |s_n - s| \\ &\leq M \cdot \frac{\epsilon}{2M} + |t| \cdot \frac{\epsilon}{2(|t| + 1)} < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon. \end{aligned}$$

■

To handle quotients of sequences, we first deal with reciprocals.

**9.5 Lemma.**

If  $(s_n)$  converges to  $s$ , if  $s_n \neq 0$  for all  $n$ , and if  $s \neq 0$ , then  $(1/s_n)$  converges to  $1/s$ .

*Discussion.* We begin by considering the equality

$$\left| \frac{1}{s_n} - \frac{1}{s} \right| = \left| \frac{s - s_n}{s_n s} \right|.$$

For large  $n$ , the numerator is small. The only possible difficulty would be if the denominator were also small for large  $n$ . This difficulty is solved in Example 6 of §8 where it is proved that  $m =$

$\inf\{|s_n| : n \in \mathbb{N}\} > 0$ . Thus

$$\left| \frac{1}{s_n} - \frac{1}{s} \right| \leq \frac{|s - s_n|}{m|s|},$$

and it is clear how our proof should proceed.

**Proof**

Let  $\epsilon > 0$ . By Example 6 of §8, there exists  $m > 0$  such that  $|s_n| \geq m$  for all  $n$ . Since  $\lim s_n = s$  there exists  $N$  such that

$$n > N \quad \text{implies} \quad |s - s_n| < \epsilon \cdot m|s|.$$

Then  $n > N$  implies

$$\left| \frac{1}{s_n} - \frac{1}{s} \right| = \frac{|s - s_n|}{|s_n s|} \leq \frac{|s - s_n|}{m|s|} < \epsilon. \quad \blacksquare$$

**9.6 Theorem.**

Suppose that  $(s_n)$  converges to  $s$  and  $(t_n)$  converges to  $t$ . If  $s \neq 0$  and  $s_n \neq 0$  for all  $n$ , then  $(t_n/s_n)$  converges to  $t/s$ .

**Proof**

By Lemma 9.5  $(1/s_n)$  converges to  $1/s$ , so

$$\lim \frac{t_n}{s_n} = \lim \frac{1}{s_n} \cdot t_n = \frac{1}{s} \cdot t = \frac{t}{s}$$

by Theorem 9.4.  $\blacksquare$

The preceding limit theorems and a few standard examples allow one to easily calculate many limits.

**9.7 Basic Examples.**

- (a)  $\lim_{n \rightarrow \infty} (\frac{1}{n^p}) = 0$  for  $p > 0$ .
- (b)  $\lim_{n \rightarrow \infty} a^n = 0$  if  $|a| < 1$ .
- (c)  $\lim(n^{1/n}) = 1$ .
- (d)  $\lim_{n \rightarrow \infty} (a^{1/n}) = 1$  for  $a > 0$ .

**Proof**

- (a) Let  $\epsilon > 0$  and let  $N = (\frac{1}{\epsilon})^{1/p}$ . Then  $n > N$  implies  $n^p > \frac{1}{\epsilon}$  and hence  $\epsilon > \frac{1}{n^p}$ . Since  $\frac{1}{n^p} > 0$ , this shows that  $n > N$  implies

$|\frac{1}{n^p} - 0| < \epsilon$ . [The meaning of  $n^p$  when  $p$  is not an integer will be discussed in §37.]

- (b) We may suppose that  $a \neq 0$ , because  $\lim_{n \rightarrow \infty} a^n = 0$  is obvious for  $a = 0$ . Since  $|a| < 1$ , we can write  $|a| = \frac{1}{1+b}$  where  $b > 0$ . By the binomial theorem [Exercise 1.12],  $(1+b^n) \geq 1+nb > nb$ , so

$$|a^n - 0| = |a^n| = \frac{1}{(1+b)^n} < \frac{1}{nb}.$$

Now consider  $\epsilon > 0$  and let  $N = \frac{1}{\epsilon b}$ . Then  $n > N$  implies  $n > \frac{1}{\epsilon b}$  and hence  $|a^n - 0| < \frac{1}{nb} < \epsilon$ .

- (c) Let  $s_n = (n^{1/n}) - 1$  and note that  $s_n \geq 0$  for all  $n$ . By Theorem 9.3 it suffices to show that  $\lim s_n = 0$ . Since  $1 + s_n = (n^{1/n})$ , we have  $n = (1 + s_n)^n$ . For  $n \geq 2$  we use the binomial expansion of  $(1 + s_n)^n$  to conclude

$$n = (1 + s_n)^n \geq 1 + ns_n + \frac{1}{2}n(n-1)s_n^2 > \frac{1}{2}n(n-1)s_n^2.$$

Thus  $n > \frac{1}{2}n(n-1)s_n^2$ , so  $s_n^2 < \frac{2}{n-1}$ . Consequently, we have  $s_n < \sqrt{\frac{2}{n-1}}$  for  $n \geq 2$ . A standard argument now shows that  $\lim s_n = 0$ ; see Exercise 9.7.

- (d) First suppose  $a \geq 1$ . Then for  $n \geq a$  we have  $1 \leq a^{1/n} \leq n^{1/n}$ . Since  $\lim n^{1/n} = 1$ , it follows easily that  $\lim(a^{1/n}) = 1$ ; compare Exercise 8.5(a). Suppose that  $0 < a < 1$ . Then  $\frac{1}{a} > 1$ , so  $\lim(\frac{1}{a})^{1/n} = 1$  from above. Lemma 9.5 now shows that  $\lim(a^{1/n}) = 1$ . ■

### Example 1

Prove that  $\lim s_n = \frac{1}{4}$ , where

$$s_n = \frac{n^3 + 6n^2 + 7}{4n^3 + 3n - 4}.$$

### Solution

We have

$$s_n = \frac{1 + \frac{6}{n} + \frac{7}{n^3}}{4 + \frac{3}{n^2} - \frac{4}{n^3}}.$$

By 9.7(a) we have  $\lim \frac{1}{n} = 0$  and  $\lim \frac{1}{n^3} = 0$ . Hence by Theorems 9.3 and 9.2 we have

$$\lim \left( 1 + \frac{6}{n} + \frac{7}{n^3} \right) = \lim(1) + 6 \cdot \lim \left( \frac{1}{n} \right) + 7 \cdot \lim \left( \frac{1}{n^3} \right) = 1.$$

Similarly, we have

$$\lim \left( 4 + \frac{3}{n^2} - \frac{4}{n^3} \right) = 4.$$

Hence Theorem 9.6 implies that  $\lim s_n = \frac{1}{4}$ . □

**Example 2**

Find  $\lim \frac{n-5}{n^2+7}$ .

**Solution**

Let  $s_n = \frac{n-5}{n^2+7}$ . We can write  $s_n$  as  $\frac{1-\frac{5}{n}}{n+\frac{7}{n}}$ , but then the denominator does not converge. So we write

$$s_n = \frac{\frac{1}{n} - \frac{5}{n^2}}{1 + \frac{7}{n^2}}.$$

Now  $\lim(\frac{1}{n} - \frac{5}{n^2}) = 0$  by 9.7(a) and Theorems 9.3 and 9.2. Likewise  $\lim(1 + \frac{7}{n^2}) = 1$ , so Theorem 9.6 tells us that  $\lim s_n = \frac{0}{1} = 0$ . □

**Example 3**

Find  $\lim \frac{n^2+3}{n+1}$ .

**Solution**

We can write  $\frac{n^2+3}{n+1}$  as

$$\frac{n + \frac{3}{n}}{1 + \frac{1}{n}} \quad \text{or} \quad \frac{1 + \frac{3}{n^2}}{\frac{1}{n} + \frac{1}{n^2}}.$$

Both fractions lead to problems: either the numerator does not converge or else the denominator converges to 0. It turns out that  $\frac{n^2+3}{n+1}$  does not converge and the symbol  $\lim \frac{n^2+3}{n+1}$  is undefined, at least for the present; see Example 6. The reader may have the urge to use the symbol  $+\infty$  here. Our next task is to make such use of the symbol  $+\infty$  legitimate. For a sequence  $(s_n)$ ,  $\lim s_n = +\infty$  will signify that the terms  $s_n$  are eventually all large. Here is the concise definition. □

**9.8 Definition.**

For a sequence  $(s_n)$ , we write  $\lim s_n = +\infty$  provided

for each  $M > 0$  there is a number  $N$  such that  
 $n > N$  implies  $s_n > M$ .

In this case we say that the sequence *diverges to*  $+\infty$ .

Similarly, we write  $\lim s_n = -\infty$  provided

for each  $M < 0$  there is a number  $N$  such that  
 $n > N$  implies  $s_n < M$ .

Henceforth we will say that  $(s_n)$  has a *limit* or that the *limit exists* provided  $(s_n)$  converges or diverges to  $+\infty$  or diverges to  $-\infty$ . In the definition of  $\lim s_n = +\infty$  the challenging values of  $M$  are large positive numbers: the larger  $M$  is the larger  $N$  will need to be. In the definition of  $\lim s_n = -\infty$  the challenging values of  $M$  are “large” negative numbers like  $-10,000,000,000$ .

**Example 4**

We have  $\lim n^2 = +\infty$ ,  $\lim(-n) = -\infty$ ,  $\lim 2^n = +\infty$  and  $\lim(\sqrt{n} + 7) = +\infty$ . Of course, many sequences do not have limits  $+\infty$  or  $-\infty$  even if they are unbounded. For example, the sequences defined by  $s_n = (-1)^n n$  and  $t_n = n \cos^2(\frac{n\pi}{2})$  are unbounded, but they do not diverge to  $+\infty$  or  $-\infty$ , so the expressions  $\lim[(-1)^n n]$  and  $\lim[n \cos^2(\frac{n\pi}{2})]$  are meaningless. Note that  $t_n = n$  when  $n$  is even and  $t_n = 0$  when  $n$  is odd.

The strategy for proofs involving infinite limits is very much the same as for finite limits. We give some examples.

**Example 5**

Give a formal proof that  $\lim(\sqrt{n} + 7) = +\infty$ .

*Discussion.* We must consider an arbitrary  $M > 0$  and show that there exists  $N$  [which will depend on  $M$ ] such that

$$n > N \quad \text{implies} \quad \sqrt{n} + 7 > M.$$

To see how big  $N$  must be we “solve” for  $n$  in the inequality  $\sqrt{n} + 7 > M$ . This inequality holds provided  $\sqrt{n} > M - 7$  or  $n > (M - 7)^2$ . Thus we will take  $N = (M - 7)^2$ .

**Formal Proof**

Let  $M > 0$  and let  $N = (M - 7)^2$ . Then  $n > N$  implies  $n > (M - 7)^2$ , hence  $\sqrt{n} > M - 7$ , hence  $\sqrt{n} + 7 > M$ . This shows that  $\lim(\sqrt{n} + 7) = +\infty$ . ■

**Example 6**

Give a formal proof that  $\lim \frac{n^2+3}{n+1} = +\infty$ ; see Example 3.

*Discussion.* Consider  $M > 0$ . We need to determine how large  $n$  must be to guarantee that  $\frac{n^2+3}{n+1} > M$ . The idea is to bound the fraction  $\frac{n^2+3}{n+1}$  below by some multiple of  $\frac{n^2}{n} = n$ ; compare Example 3 of §8. Since  $n^2 + 3 > n^2$  and  $n + 1 \leq 2n$ , we have  $\frac{n^2+3}{n+1} > \frac{n^2}{2n} = \frac{1}{2}n$ , and it suffices to arrange for  $\frac{1}{2}n > M$ .

**Formal Proof**

Let  $M > 0$  and let  $N = 2M$ . Then  $n > N$  implies  $\frac{1}{2}n > M$ , which implies

$$\frac{n^2 + 3}{n + 1} > \frac{n^2}{2n} = \frac{1}{2}n > M.$$

Hence  $\lim \frac{n^2+3}{n+1} = +\infty$ . ■

The limit in Example 6 would be easier to handle if we could apply a limit theorem. But the limit theorems 9.2–9.6 do not apply.

*WARNING.* Do not attempt to apply the limit theorems 9.2–9.6 to infinite limits. Use Theorem 9.9 or 9.10 below or Exercises 9.9–9.12.

**9.9 Theorem.**

Let  $(s_n)$  and  $(t_n)$  be sequences such that  $\lim s_n = +\infty$  and  $\lim t_n > 0$  [ $\lim t_n$  can be finite or  $+\infty$ ]. Then  $\lim s_n t_n = +\infty$ .

*Discussion.* Let  $M > 0$ . We need to show that  $s_n t_n > M$  for large  $n$ . We have  $\lim s_n = +\infty$ , and we need to be sure that the  $t_n$ 's are bounded away from 0 for large  $n$ . We will choose a real number  $m$

so that  $0 < m < \lim t_n$  and observe that  $t_n > m$  for large  $n$ . Then all we need is  $s_n > \frac{M}{m}$  for large  $n$ .

**Proof**

Let  $M > 0$ . Select a real number  $m$  so that  $0 < m < \lim t_n$ . Whether  $\lim t_n = +\infty$  or not, it is clear that there exists  $N_1$  such that

$$n > N_1 \quad \text{implies} \quad t_n > m;$$

see Exercise 8.10. Since  $\lim s_n = +\infty$ , there exists  $N_2$  so that

$$n > N_2 \quad \text{implies} \quad s_n > \frac{M}{m}.$$

Put  $N = \max\{N_1, N_2\}$ . Then  $n > N$  implies  $s_n t_n > \frac{M}{m} \cdot m = M$ . ■

**Example 7**

Use Theorem 9.9 to prove that  $\lim \frac{n^2+3}{n+1} = +\infty$ ; see Example 6.

**Solution**

We observe that  $\frac{n^2+3}{n+1} = \frac{n+\frac{3}{n}}{1+\frac{1}{n}} = s_n t_n$  where  $s_n = n + \frac{3}{n}$  and  $t_n = \frac{1}{1+\frac{1}{n}}$ . It is easy to show that  $\lim s_n = +\infty$  and  $\lim t_n = 1$ . So by Theorem 9.9, we have  $\lim s_n t_n = +\infty$ . □

Here is another useful theorem.

**9.10 Theorem.**

*For a sequence  $(s_n)$  of positive real numbers, we have  $\lim s_n = +\infty$  if and only if  $\lim(\frac{1}{s_n}) = 0$ .*

**Proof**

Let  $(s_n)$  be a sequence of positive real numbers. We have to show

$$\lim s_n = +\infty \quad \text{implies} \quad \lim \left( \frac{1}{s_n} \right) = 0 \quad (1)$$

and

$$\lim \left( \frac{1}{s_n} \right) = 0 \quad \text{implies} \quad \lim s_n = +\infty. \quad (2)$$

In this case the proofs will appear very similar, but the thought processes will be quite different.

To prove (1), suppose that  $\lim s_n = +\infty$ . Let  $\epsilon > 0$  and let  $M = \frac{1}{\epsilon}$ . Since  $\lim s_n = +\infty$ , there exists  $N$  such that  $n > N$  implies  $s_n > M = \frac{1}{\epsilon}$ . Therefore  $n > N$  implies  $\epsilon > \frac{1}{s_n} > 0$ , so

$$n > N \quad \text{implies} \quad \left| \frac{1}{s_n} - 0 \right| < \epsilon.$$

That is,  $\lim(\frac{1}{s_n}) = 0$ . This proves (1).

To prove (2), we abandon the notation of the last paragraph and begin anew. Suppose that  $\lim(\frac{1}{s_n}) = 0$ . Let  $M > 0$  and let  $\epsilon = \frac{1}{M}$ . Then  $\epsilon > 0$ , so there exists  $N$  such that  $n > N$  implies  $|\frac{1}{s_n} - 0| < \epsilon = \frac{1}{M}$ . Since  $s_n > 0$ , we can write

$$n > N \quad \text{implies} \quad 0 < \frac{1}{s_n} < \frac{1}{M}$$

and hence

$$n > N \quad \text{implies} \quad M < s_n.$$

That is,  $\lim s_n = +\infty$  and (2) holds. ■

## Exercises

**9.1.** Using the limit theorems 9.2–9.6 and 9.7, prove the following. Justify all steps.

(a)  $\lim \frac{n+1}{n} = 1$                       (b)  $\lim \frac{3n+7}{6n-5} = \frac{1}{2}$

(c)  $\lim \frac{17n^5+73n^4-18n^2+3}{23n^5+13n^3} = \frac{17}{23}$

**9.2.** Suppose that  $\lim x_n = 3$ ,  $\lim y_n = 7$  and that all  $y_n$  are nonzero. Determine the following limits:

(a)  $\lim(x_n + y_n)$                       (b)  $\lim \frac{3y_n - x_n}{y_n^2}$

**9.3.** Suppose that  $\lim a_n = a$ ,  $\lim b_n = b$ , and that  $s_n = \frac{a_n^3 + 4a_n}{b_n^2 + 1}$ . Prove  $\lim s_n = \frac{a^3 + 4a}{b^2 + 1}$  carefully, using the limit theorems.

**9.4.** Let  $s_1 = 1$  and for  $n \geq 1$  let  $s_{n+1} = \sqrt{s_n + 1}$ .

(a) List the first four terms of  $(s_n)$ .

(b) It turns out that  $(s_n)$  converges. Assume this fact and prove that the limit is  $\frac{1}{2}(1 + \sqrt{5})$ .

- 9.5.** Let  $t_1 = 1$  and  $t_{n+1} = \frac{t_n^2 + 2}{2t_n}$  for  $n \geq 1$ . Assume that  $(t_n)$  converges and find the limit.
- 9.6.** Let  $x_1 = 1$  and  $x_{n+1} = 3x_n^2$  for  $n \geq 1$ .
- (a) Show that if  $a = \lim x_n$ , then  $a = \frac{1}{3}$  or  $a = 0$ .
- (b) Does  $\lim x_n$  exist? Explain.
- (c) Discuss the apparent contradiction between parts (a) and (b).
- 9.7.** Complete the proof of 9.7(c), i.e., give the standard argument needed to show that  $\lim s_n = 0$ .
- 9.8.** Give the following when they exist. Otherwise assert "NOT EXIST."
- (a)  $\lim n^3$  (b)  $\lim(-n^3)$   
(c)  $\lim(-n)^n$  (d)  $\lim(1.01)^n$   
(e)  $\lim n^n$
- 9.9.** Suppose that there exists  $N_0$  such that  $s_n \leq t_n$  for all  $n > N_0$ .
- (a) Prove that if  $\lim s_n = +\infty$ , then  $\lim t_n = +\infty$ .
- (b) Prove that if  $\lim t_n = -\infty$ , then  $\lim s_n = -\infty$ .
- (c) Prove that if  $\lim s_n$  and  $\lim t_n$  exist, then  $\lim s_n \leq \lim t_n$ .
- 9.10.** (a) Show that if  $\lim s_n = +\infty$  and  $k > 0$ , then  $\lim(ks_n) = +\infty$ .
- (b) Show that  $\lim s_n = +\infty$  if and only if  $\lim(-s_n) = -\infty$ .
- (c) Show that if  $\lim s_n = +\infty$  and  $k < 0$ , then  $\lim(ks_n) = -\infty$ .
- 9.11.** (a) Show that if  $\lim s_n = +\infty$  and  $\inf\{t_n : n \in \mathbb{N}\} > -\infty$ , then  $\lim(s_n + t_n) = +\infty$ .
- (b) Show that if  $\lim s_n = +\infty$  and  $\lim t_n > -\infty$ , then  $\lim(s_n + t_n) = +\infty$ .
- (c) Show that if  $\lim s_n = +\infty$  and if  $(t_n)$  is a bounded sequence, then  $\lim(s_n + t_n) = +\infty$ .
- 9.12.** Assume all  $s_n \neq 0$  and that the limit  $L = \lim \left| \frac{s_{n+1}}{s_n} \right|$  exists.
- (a) Show that if  $L < 1$ , then  $\lim s_n = 0$ . *Hint:* Select  $a$  so that  $L < a < 1$  and obtain  $N$  so that  $|s_{n+1}| < a|s_n|$  for  $n \geq N$ . Then show that  $|s_n| < a^{n-N}|s_N|$  for  $n > N$ .
- (b) Show that if  $L > 1$ , then  $\lim |s_n| = +\infty$ . *Hint:* Apply (a) to the sequence  $t_n = \frac{1}{|s_n|}$ ; see Theorem 9.10.

**9.13.** Show that

$$\lim_{n \rightarrow \infty} a^n = \begin{cases} 0 & \text{if } |a| < 1 \\ 1 & \text{if } a = 1 \\ +\infty & \text{if } a > 1 \\ \text{does not exist} & \text{if } a \leq -1. \end{cases}$$

**9.14.** Let  $p > 0$ . Use Exercise 9.12 to show

$$\lim_{n \rightarrow \infty} \frac{a^n}{n^p} = \begin{cases} 0 & \text{if } |a| \leq 1 \\ +\infty & \text{if } a > 1 \\ \text{does not exist} & \text{if } a < -1. \end{cases}$$

**9.15.** Show that  $\lim_{n \rightarrow \infty} \frac{a^n}{n!} = 0$  for all  $a \in \mathbb{R}$ .

**9.16.** Use Theorems 9.9, 9.10 or Exercises 9.9–9.15 to prove the following:

(a)  $\lim_{n \rightarrow \infty} \frac{n^4 + 8n}{n^2 + 9} = +\infty$

(b)  $\lim_{n \rightarrow \infty} \left[ \frac{2^n}{n^2} + (-1)^n \right] = +\infty$

(c)  $\lim_{n \rightarrow \infty} \left[ \frac{3^n}{n^3} - \frac{3^n}{n!} \right] = +\infty$

**9.17.** Give a formal proof that  $\lim_{n \rightarrow \infty} n^2 = +\infty$  using only Definition 9.8.

**9.18. (a)** Verify  $1 + a + a^2 + \cdots + a^n = \frac{1 - a^{n+1}}{1 - a}$  for  $a \neq 1$ .

(b) Find  $\lim_{n \rightarrow \infty} (1 + a + a^2 + \cdots + a^n)$  for  $|a| < 1$ .

(c) Calculate  $\lim_{n \rightarrow \infty} (1 + \frac{1}{3} + \frac{1}{9} + \frac{1}{27} + \cdots + \frac{1}{3^n})$ .

(d) What is  $\lim_{n \rightarrow \infty} (1 + a + a^2 + \cdots + a^n)$  for  $a \geq 1$ ?

## §10 Monotone Sequences and Cauchy Sequences

In this section we obtain two theorems [Theorems 10.2 and 10.11] that will allow us to conclude that certain sequences converge *without* knowing the limit in advance. These theorems are important because in practice the limits are not usually known in advance.

### 10.1 Definition.

A sequence  $(s_n)$  of real numbers is called a *nondecreasing sequence* if  $s_n \leq s_{n+1}$  for all  $n$ , and  $(s_n)$  is called a *nonincreasing sequence*

if  $s_n \geq s_{n+1}$  for all  $n$ . Note that if  $(s_n)$  is nondecreasing, then  $s_n \leq s_m$  whenever  $n < m$ . A sequence that is nondecreasing or nonincreasing will be called a *monotone sequence* or a *monotonic sequence*.

**Example 1**

The sequences defined by  $a_n = 1 - \frac{1}{n}$ ,  $b_n = n^3$  and  $c_n = (1 + \frac{1}{n})^n$  are nondecreasing sequences, although this is not obvious for the sequence  $(c_n)$ . The sequence  $d_n = \frac{1}{n^2}$  is nonincreasing. The sequences  $s_n = (-1)^n$ ,  $t_n = \cos(\frac{n\pi}{3})$ ,  $u_n = (-1)^n n$  and  $v_n = \frac{(-1)^n}{n}$  are not monotonic sequences. Also  $x_n = n^{1/n}$  is not monotonic, as can be seen by examining the first four values; see Example 1(d) in §7.

Of the sequences above,  $(a_n)$ ,  $(c_n)$ ,  $(d_n)$ ,  $(s_n)$ ,  $(t_n)$ ,  $(v_n)$  and  $(x_n)$  are bounded sequences. The remaining sequences,  $(b_n)$  and  $(u_n)$ , are unbounded sequences.

**10.2 Theorem.**

*All bounded monotone sequences converge.*

**Proof**

Let  $(s_n)$  be a bounded nondecreasing sequence. Let  $S$  denote the set  $\{s_n : n \in \mathbb{N}\}$ , and let  $u = \sup S$ . Since  $S$  is bounded,  $u$  represents a real number. We show that  $\lim s_n = u$ . Let  $\epsilon > 0$ . Since  $u - \epsilon$  is not an upper bound for  $S$ , there exists  $N$  such that  $s_N > u - \epsilon$ . Since  $(s_n)$  is nondecreasing, we have  $s_N \leq s_n$  for all  $n \geq N$ . Of course,  $s_n \leq u$  for all  $n$ , so  $n > N$  implies  $u - \epsilon < s_n \leq u$ , which implies  $|s_n - u| < \epsilon$ . This shows that  $\lim s_n = u$ .

The proof for bounded nonincreasing sequences is left to Exercise 10.2. ■

Note that the Completeness Axiom 4.4 is a vital ingredient in the proof of Theorem 10.2.

**10.3 Discussion of Decimals.**

We have not given much attention to the notion that real numbers are simply decimal expansions. This notion is substantially correct, but there are subtleties to be faced. For example, different decimal expansions can represent the same real number. The somewhat more

abstract developments of the set  $\mathbb{R}$  of real numbers discussed in §6 turn out to be more satisfactory.

We restrict our attention to nonnegative decimal expansions and nonnegative real numbers. From our point of view, every nonnegative decimal expansion is shorthand for the limit of a bounded nondecreasing sequence of real numbers. Suppose we are given a decimal expansion  $k.d_1d_2d_3d_4\cdots$  where  $k$  is a nonnegative integer and each  $d_j$  belongs to  $\{0, 1, 2, 3, 4, 5, 6, 7, 8, 9\}$ . Let

$$s_n = k + \frac{d_1}{10} + \frac{d_2}{10^2} + \cdots + \frac{d_n}{10^n}. \quad (1)$$

Then  $(s_n)$  is a nondecreasing sequence of real numbers, and  $(s_n)$  is bounded [by  $k + 1$ , in fact]. So by Theorem 10.2,  $(s_n)$  converges to a real number that we traditionally write as  $k.d_1d_2d_3d_4\cdots$ . For example,  $3.3333\cdots$  represents

$$\lim_{n \rightarrow \infty} \left( 3 + \frac{3}{10} + \frac{3}{10^2} + \cdots + \frac{3}{10^n} \right).$$

To calculate this limit, we borrow the following fact about geometric series from Example 1 in §14:

$$\lim_{n \rightarrow \infty} a(1 + r + r^2 + \cdots + r^n) = \frac{a}{1 - r} \quad \text{for } |r| < 1; \quad (2)$$

see also Exercise 9.18. In our case,  $a = 3$  and  $r = \frac{1}{10}$ , so  $3.3333\cdots$  represents  $\frac{3}{1 - \frac{1}{10}} = \frac{10}{3}$ , as expected. Similarly,  $0.9999\cdots$  represents

$$\lim_{n \rightarrow \infty} \left( \frac{9}{10} + \frac{9}{10^2} + \cdots + \frac{9}{10^n} \right) = \frac{\frac{9}{10}}{1 - \frac{1}{10}} = 1.$$

Thus  $0.9999\cdots$  and  $1.0000\cdots$  are different decimal expansions that represent the same real number!

The converse of the preceding discussion also holds. That is, every nonnegative real number  $x$  has at least one decimal expansion. This will be proved, along with some related results, in the optional §16.

Unbounded monotone sequences also have limits.

**10.4 Theorem.**

- (i) If  $(s_n)$  is an unbounded nondecreasing sequence, then  $\lim s_n = +\infty$ .
- (ii) If  $(s_n)$  is an unbounded nonincreasing sequence, then  $\lim s_n = -\infty$ .

**Proof**

(i) Let  $(s_n)$  be an unbounded nondecreasing sequence. Let  $M > 0$ . Since the set  $\{s_n : n \in \mathbb{N}\}$  is unbounded and it is bounded below by  $s_1$ , it must be unbounded above. Hence for some  $N$  in  $\mathbb{N}$  we have  $s_N > M$ . Clearly  $n > N$  implies  $s_n \geq s_N > M$ , so  $\lim s_n = +\infty$ .

(ii) The proof is similar and is left to Exercise 10.5. ■

**10.5 Corollary.**

If  $(s_n)$  is a monotone sequence, then the sequence either converges, diverges to  $+\infty$ , or diverges to  $-\infty$ . Thus  $\lim s_n$  is always meaningful for monotone sequences.

**Proof**

Apply Theorems 10.2 and 10.4. ■

Let  $(s_n)$  be a bounded sequence in  $\mathbb{R}$ ; it may or may not converge. It is apparent from the definition of limit in 7.1 that the limiting behavior of  $(s_n)$  depends only on sets of the form  $\{s_n : n > N\}$ . For example, if  $\lim s_n$  exists, clearly it must lie in the interval  $[u_N, v_N]$  where

$$u_N = \inf\{s_n : n > N\} \quad \text{and} \quad v_N = \sup\{s_n : n > N\};$$

see Exercise 8.9. As  $N$  increases, the sets  $\{s_n : n > N\}$  get smaller, so we have

$$u_1 \leq u_2 \leq u_3 \leq \cdots \quad \text{and} \quad v_1 \geq v_2 \geq v_3 \geq \cdots;$$

see Exercise 4.7(a). By Theorem 10.2 the limits  $u = \lim_{N \rightarrow \infty} u_N$  and  $v = \lim_{N \rightarrow \infty} v_N$  both exist, and  $u \leq v$  since  $u_N \leq v_N$  for all  $N$ . If  $\lim s_n$  exists then, as noted above,  $u_N \leq \lim s_n \leq v_N$  for all  $N$ , so we must have  $u \leq \lim s_n \leq v$ . The numbers  $u$  and  $v$  are useful whether  $\lim s_n$  exists or not and are denoted  $\liminf s_n$  and  $\limsup s_n$ , respectively.

**10.6 Definition.**

Let  $(s_n)$  be a sequence in  $\mathbb{R}$ . We define

$$\limsup s_n = \lim_{N \rightarrow \infty} \sup \{s_n : n > N\} \quad (1)$$

and

$$\liminf s_n = \lim_{N \rightarrow \infty} \inf \{s_n : n > N\}. \quad (2)$$

Note that in this definition we do not restrict  $(s_n)$  to be bounded. However, we adopt the following conventions. If  $(s_n)$  is not bounded above,  $\sup\{s_n : n > N\} = +\infty$  for all  $N$  and we decree  $\limsup s_n = +\infty$ . Likewise, if  $(s_n)$  is not bounded below,  $\inf\{s_n : n > N\} = -\infty$  for all  $N$  and we decree  $\liminf s_n = -\infty$ .

We emphasize that  $\limsup s_n$  need not equal  $\sup\{s_n : n \in \mathbb{N}\}$ , but that  $\limsup s_n \leq \sup\{s_n : n \in \mathbb{N}\}$ . Some of the values  $s_n$  may be much larger than  $\limsup s_n$ ;  $\limsup s_n$  is the largest value that *infinitely many*  $s_n$ 's can get close to. Similar remarks apply to  $\liminf s_n$ . These remarks will be clarified in Theorem 11.7 and §12, where we will give a thorough treatment of  $\liminf$ 's and  $\limsup$ 's. For now, we need a theorem that shows  $(s_n)$  has a limit if and only if  $\liminf s_n = \limsup s_n$ .

**10.7 Theorem.**

Let  $(s_n)$  be a sequence in  $\mathbb{R}$ .

- (i) If  $\lim s_n$  is defined [as a real number,  $+\infty$  or  $-\infty$ ], then  $\liminf s_n = \lim s_n = \limsup s_n$ .
- (ii) If  $\liminf s_n = \limsup s_n$ , then  $\lim s_n$  is defined and  $\lim s_n = \liminf s_n = \limsup s_n$ .

**Proof**

We use the notation  $u_N = \inf\{s_n : n > N\}$ ,  $v_N = \sup\{s_n : n > N\}$ ,  $u = \lim u_N = \liminf s_n$  and  $v = \lim v_N = \limsup s_n$ .

- (i) Suppose  $\lim s_n = +\infty$ . Let  $M$  be a positive real number. Then there is a natural number  $N$  so that

$$n > N \quad \text{implies} \quad s_n > M.$$

Then  $u_N = \inf\{s_n : n > N\} \geq M$ . It follows that  $m > N$  implies  $u_m \geq M$ . In other words, the sequence  $(u_N)$  satisfies

the condition defining  $\lim u_N = +\infty$ , i.e.,  $\liminf s_n = +\infty$ . Likewise  $\limsup s_n = +\infty$ .

The case  $\lim s_n = -\infty$  is handled in a similar manner.

Now suppose that  $\lim s_n = s$  where  $s$  is a real number. Consider  $\epsilon > 0$ . There exists a natural number  $N$  such that  $|s_n - s| < \epsilon$  for  $n > N$ . Thus  $s_n < s + \epsilon$  for  $n > N$ , so

$$v_N = \sup\{s_n : n > N\} \leq s + \epsilon.$$

Also,  $m > N$  implies  $v_m \leq s + \epsilon$ , so  $\limsup s_n = \lim v_m \leq s + \epsilon$ . Since  $\limsup s_n \leq s + \epsilon$  for all  $\epsilon > 0$ , no matter how small, we conclude that  $\limsup s_n \leq s = \lim s_n$ . A similar argument shows that  $\lim s_n \leq \liminf s_n$ . Since  $\liminf s_n \leq \limsup s_n$ , we infer that all three numbers are equal:

$$\liminf s_n = \lim s_n = \limsup s_n.$$

- (ii) If  $\liminf s_n = \limsup s_n = +\infty$  it is easy to show that  $\lim s_n = +\infty$ . And if  $\liminf s_n = \limsup s_n = -\infty$  it is easy to show that  $\lim s_n = -\infty$ . We leave these two special cases to the reader.

Suppose, finally, that  $\liminf s_n = \limsup s_n = s$  where  $s$  is a real number. We need to prove that  $\lim s_n = s$ . Let  $\epsilon > 0$ . Since  $s = \lim v_N$  there exists a natural number  $N_0$  such that

$$|s - \sup\{s_n : n > N_0\}| < \epsilon.$$

Thus  $\sup\{s_n : n > N_0\} < s + \epsilon$ , so

$$s_n < s + \epsilon \quad \text{for all } n > N_0. \quad (1)$$

Similarly, since  $s = \lim u_N$  there exists  $N_1$  such that

$$|s - \inf\{s_n : n > N_1\}| < \epsilon,$$

hence  $\inf\{s_n : n > N_1\} > s - \epsilon$ , hence

$$s_n > s - \epsilon \quad \text{for all } n > N_1. \quad (2)$$

From (1) and (2) we conclude

$$s - \epsilon < s_n < s + \epsilon \quad \text{for } n > \max\{N_0, N_1\},$$

equivalently

$$|s_n - s| < \epsilon \quad \text{for } n > \max\{N_0, N_1\}.$$

This proves that  $\lim s_n = s$  as desired. ■

If  $(s_n)$  converges, then  $\liminf s_n = \limsup s_n$  by the theorem just proved, so for large  $N$  the numbers  $\sup\{s_n : n > N\}$  and  $\inf\{s_n : n > N\}$  must be close together. This implies that all the numbers in the set  $\{s_n : n > N\}$  must be close to each other. This leads us to a concept of great theoretical importance that will be used throughout the book.

### 10.8 Definition.

A sequence  $(s_n)$  of real numbers is called a *Cauchy sequence* if

$$\begin{aligned} &\text{for each } \epsilon > 0 \text{ there exists a number } N \text{ such that} \\ &m, n > N \text{ implies } |s_n - s_m| < \epsilon. \end{aligned} \tag{1}$$

Compare this definition with Definition 7.1.

### 10.9 Lemma.

*Convergent sequences are Cauchy sequences.*

#### Proof

Suppose that  $\lim s_n = s$ . The idea is that, since the terms  $s_n$  are close to  $s$  for large  $n$ , they also must be close to each other; indeed

$$|s_n - s_m| = |s_n - s + s - s_m| \leq |s_n - s| + |s - s_m|.$$

To be precise, let  $\epsilon > 0$ . Then there exists  $N$  such that

$$n > N \quad \text{implies} \quad |s_n - s| < \frac{\epsilon}{2}.$$

Clearly we may also write

$$m > N \quad \text{implies} \quad |s_m - s| < \frac{\epsilon}{2},$$

so

$$m, n > N \quad \text{implies} \quad |s_n - s_m| \leq |s_n - s| + |s - s_m| < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon.$$

Thus  $(s_n)$  is a Cauchy sequence. ■

### 10.10 Lemma.

*Cauchy sequences are bounded.*

**Proof**

The proof is similar to that of Theorem 9.1. Applying Definition 10.8 with  $\epsilon = 1$  we obtain  $N$  in  $\mathbb{N}$  so that

$$m, n > N \quad \text{implies} \quad |s_n - s_m| < 1.$$

In particular,  $|s_n - s_{N+1}| < 1$  for  $n > N$ , so  $|s_n| < |s_{N+1}| + 1$  for  $n > N$ . If  $M = \max\{|s_{N+1}| + 1, |s_1|, |s_2|, \dots, |s_N|\}$ , then  $|s_n| \leq M$  for all  $n \in \mathbb{N}$ . ■

The importance of the next theorem is the following consequence: To verify that a sequence converges it suffices to check that it is a Cauchy sequence, a property that does not involve the limit itself.

**10.11 Theorem.**

*A sequence is a convergent sequence if and only if it is a Cauchy sequence.*

**Proof**

The expression “if and only if” indicates that we have two assertions to verify: (i) convergent sequences are Cauchy sequences, and (ii) Cauchy sequences are convergent sequences. We already verified (i) in Lemma 10.9. To check (ii), consider a Cauchy sequence  $(s_n)$  and note that  $(s_n)$  is bounded by Lemma 10.10. By Theorem 10.7 we need only show

$$\liminf s_n = \limsup s_n. \tag{1}$$

Let  $\epsilon > 0$ . Since  $(s_n)$  is a Cauchy sequence, there exists  $N$  so that

$$m, n > N \quad \text{implies} \quad |s_n - s_m| < \epsilon.$$

In particular,  $s_n < s_m + \epsilon$  for all  $m, n > N$ . This shows that  $s_m + \epsilon$  is an upper bound for  $\{s_n : n > N\}$ , so  $v_N = \sup\{s_n : n > N\} \leq s_m + \epsilon$  for  $m > N$ . This, in turn, shows that  $v_N - \epsilon$  is a lower bound for  $\{s_m : m > N\}$ , so  $v_N - \epsilon \leq \inf\{s_m : m > N\} = u_N$ . Thus

$$\limsup s_n \leq v_N \leq u_N + \epsilon \leq \liminf s_n + \epsilon.$$

Since this holds for all  $\epsilon > 0$ , we have  $\limsup s_n \leq \liminf s_n$ . The opposite inequality always holds, so we have established (1). ■

The proof of Theorem 10.11 uses Theorem 10.7, and Theorem 10.7 relies implicitly on the Completeness Axiom 4.4, since without the completeness axiom it is not clear that  $\liminf s_n$  and  $\limsup s_n$  are meaningful. The completeness axiom assures us that the expressions  $\sup\{s_n : n > N\}$  and  $\inf\{s_n : n > N\}$  in Definition 10.6 are meaningful, and Theorem 10.2 [which itself relies on the completeness axiom] assures us that the limits in Definition 10.6 also are meaningful.

Exercises on  $\limsup$ 's and  $\liminf$ 's appear in §§11 and 12.

## Exercises

**10.1.** Which of the following sequences are nondecreasing? nonincreasing? bounded?

(a)  $\frac{1}{n}$

(b)  $\frac{(-1)^n}{n^2}$

(c)  $n^5$

(d)  $\sin(\frac{n\pi}{7})$

(e)  $(-2)^n$

(f)  $\frac{n}{3^n}$

**10.2.** Prove Theorem 10.2 for bounded nonincreasing sequences.

**10.3.** For a decimal expansion  $k.d_1d_2d_3d_4\cdots$ , let  $(s_n)$  be defined as in 10.3. Prove that  $s_n < k+1$  for all  $n \in \mathbb{N}$ . *Hint:*  $\frac{9}{10} + \frac{9}{10^2} + \cdots + \frac{9}{10^n} = 1 - \frac{1}{10^n}$  for all  $n$ .

**10.4.** Discuss why Theorems 10.2 and 10.11 would fail if we restricted our world of numbers to the set  $\mathbb{Q}$  of rational numbers.

**10.5.** Prove Theorem 10.4(ii).

**10.6. (a)** Let  $(s_n)$  be a sequence such that

$$|s_{n+1} - s_n| < 2^{-n} \quad \text{for all } n \in \mathbb{N}.$$

Prove that  $(s_n)$  is a Cauchy sequence and hence a convergent sequence.

**(b)** Is the result in (a) true if we only assume that  $|s_{n+1} - s_n| < \frac{1}{n}$  for all  $n \in \mathbb{N}$ ?

**10.7.** Let  $S$  be a bounded nonempty subset of  $\mathbb{R}$  and suppose  $\sup S \notin S$ . Prove that there is a nondecreasing sequence  $(s_n)$  of points in  $S$  such that  $\lim s_n = \sup S$ .

- 10.8.** Let  $(s_n)$  be a nondecreasing sequence of positive numbers and define  $\sigma_n = \frac{1}{n}(s_1 + s_2 + \cdots + s_n)$ . Prove that  $(\sigma_n)$  is a nondecreasing sequence.
- 10.9.** Let  $s_1 = 1$  and  $s_{n+1} = (\frac{n}{n+1})s_n^2$  for  $n \geq 1$ .
- (a) Find  $s_2, s_3$  and  $s_4$ .
  - (b) Show that  $\lim s_n$  exists.
  - (c) Prove that  $\lim s_n = 0$ .
- 10.10.** Let  $s_1 = 1$  and  $s_{n+1} = \frac{1}{3}(s_n + 1)$  for  $n \geq 1$ .
- (a) Find  $s_2, s_3$  and  $s_4$ .
  - (b) Use induction to show that  $s_n > \frac{1}{2}$  for all  $n$ .
  - (c) Show that  $(s_n)$  is a nonincreasing sequence.
  - (d) Show that  $\lim s_n$  exists and find  $\lim s_n$ .
- 10.11.** Let  $t_1 = 1$  and  $t_{n+1} = [1 - \frac{1}{4n^2}] \cdot t_n$  for  $n \geq 1$ .
- (a) Show that  $\lim t_n$  exists.
  - (b) What do you think  $\lim t_n$  is?
- 10.12.** Let  $t_1 = 1$  and  $t_{n+1} = [1 - \frac{1}{(n+1)^2}] \cdot t_n$  for  $n \geq 1$ .
- (a) Show that  $\lim t_n$  exists.
  - (b) What do you think  $\lim t_n$  is?
  - (c) Use induction to show that  $t_n = \frac{n+1}{2n}$ .
  - (d) Repeat part (b).

## §11 Subsequences

### 11.1 Definition.

Suppose that  $(s_n)_{n \in \mathbb{N}}$  is a sequence. A *subsequence* of this sequence is a sequence of the form  $(t_k)_{k \in \mathbb{N}}$  where for each  $k$  there is a positive integer  $n_k$  such that

$$n_1 < n_2 < \cdots < n_k < n_{k+1} < \cdots \quad (1)$$

and

$$t_k = s_{n_k}. \quad (2)$$

Thus  $(t_k)$  is just a selection of some [possibly all] of the  $s_n$ 's taken in order.

Here are some alternative ways to approach this concept. Note that (1) defines an infinite subset of  $\mathbb{N}$ , namely  $\{n_1, n_2, n_3, \dots\}$ . Conversely, every infinite subset of  $\mathbb{N}$  can be described by (1). Thus a subsequence of  $(s_n)$  is a sequence obtained by selecting, in order, an infinite subset of the terms.

For a more concise definition, recall that we can view the sequence  $(s_n)_{n \in \mathbb{N}}$  as a function  $s$  with domain  $\mathbb{N}$ ; see §7. For the subset  $\{n_1, n_2, n_3, \dots\}$ , there is a natural function  $\sigma$  [lower case Greek sigma] given by  $\sigma(k) = n_k$  for  $k \in \mathbb{N}$ . The function  $\sigma$  “selects” an infinite subset of  $\mathbb{N}$ , in order. The subsequence of  $s$  corresponding to  $\sigma$  is simply the composite function  $t = s \circ \sigma$ . That is,

$$t_k = t(k) = s \circ \sigma(k) = s(\sigma(k)) = s(n_k) = s_{n_k} \quad \text{for } k \in \mathbb{N}. \quad (3)$$

Thus a sequence  $t$  is a subsequence of a sequence  $s$  if and only if  $t = s \circ \sigma$  for some increasing function  $\sigma$  mapping  $\mathbb{N}$  into  $\mathbb{N}$ . We will usually suppress the notation  $\sigma$  and often suppress the notation  $t$  also. Thus the phrase “a subsequence  $(s_{n_k})$  of  $(s_n)$ ” will refer to the subsequence defined by (1) and (2) or by (3), depending upon your point of view.

### Example 1

Let  $(s_n)$  be the sequence defined by  $s_n = n^2(-1)^n$ . The positive terms of this sequence comprise a subsequence. In this case, the sequence  $(s_n)$  is

$$(-1, 4, -9, 16, -25, 36, -49, 64, \dots)$$

and the subsequence is

$$(4, 16, 36, 64, 100, 144, \dots).$$

More precisely, the subsequence is  $(s_{n_k})_{k \in \mathbb{N}}$  where  $n_k = 2k$  so that  $s_{n_k} = (2k)^2(-1)^{2k} = 4k^2$ . The selection function  $\sigma$  is given by  $\sigma(k) = 2k$ .

**Example 2**

Consider the sequence  $a_n = \sin(\frac{n\pi}{3})$  and its subsequence  $(a_{n_k})$  of nonnegative terms. The sequence  $(a_n)_{n \in \mathbb{N}}$  is

$$(\frac{1}{2}\sqrt{3}, \frac{1}{2}\sqrt{3}, 0, -\frac{1}{2}\sqrt{3}, -\frac{1}{2}\sqrt{3}, 0, \frac{1}{2}\sqrt{3}, \frac{1}{2}\sqrt{3}, 0, -\frac{1}{2}\sqrt{3}, -\frac{1}{2}\sqrt{3}, 0, \dots)$$

and the desired subsequence is

$$(\frac{1}{2}\sqrt{3}, \frac{1}{2}\sqrt{3}, 0, 0, \frac{1}{2}\sqrt{3}, \frac{1}{2}\sqrt{3}, 0, 0, \dots).$$

It is evident that  $n_1 = 1$ ,  $n_2 = 2$ ,  $n_3 = 3$ ,  $n_4 = 6$ ,  $n_5 = 7$ ,  $n_6 = 8$ ,  $n_7 = 9$ ,  $n_8 = 12$ ,  $n_9 = 13$ , etc. We could obtain a general formula for  $n_k$ , but the project does not seem worth the effort.

**Example 3**

It can be shown that the set  $\mathbb{Q}$  of rational numbers can be listed as a sequence  $(r_n)$ , though it is tedious to specify an exact formula. Figure 11.1 suggests such a listing [with repetitions] where  $r_1 = 0$ ,  $r_2 = 1$ ,  $r_3 = \frac{1}{2}$ ,  $r_4 = -\frac{1}{2}$ ,  $r_5 = -1$ ,  $r_6 = -2$ ,  $r_7 = -1$ , etc. Readers familiar with some set theory will recognize this assertion as the fact that “ $\mathbb{Q}$  is countable.” This sequence has an amazing property: given any real number  $a$  there exists a subsequence  $(r_{n_k})$  of  $(r_n)$  that converges to  $a$ , i.e.,  $\lim_{k \rightarrow \infty} r_{n_k} = a$ . To see this, we will show how to define or construct step-by-step a subsequence  $(r_{n_k})$  that satisfies

$$|r_{n_k} - a| < \frac{1}{k} \quad \text{for } k \in \mathbb{N}. \quad (1)$$

FIGURE 11.1

Specifically, we will assume  $n_1, n_2, \dots, n_k$  have been selected satisfying (1) and show how to select  $n_{k+1}$ . It is fairly evident that this will give us an infinite sequence  $(n_k)_{k \in \mathbb{N}}$  and hence a subsequence  $(r_{n_k})$  of  $(r_n)$  satisfying (1). To make this fully rigorous would require a technical lemma concerning step-by-step constructions whose proof depends in the end on Peano's axiom N5. For this reason, a construction of this sort is called an "inductive definition" or "definition by induction."

We now indicate the construction discussed above. Select  $n_1$  so that  $|r_{n_1} - a| < 1$ ; this is possible by the Denseness of  $\mathbb{Q}$  4.7. Suppose that  $n_1, n_2, \dots, n_k$  have been selected so that

$$n_1 < n_2 < \dots < n_k \quad (2)$$

and

$$|r_{n_j} - a| < \frac{1}{j} \quad \text{for } j = 1, 2, \dots, k. \quad (3)$$

Since there are infinitely many rational numbers in the interval  $(a - \frac{1}{k+1}, a + \frac{1}{k+1})$  by Exercise 4.11, there must exist an  $n_{k+1} > n_k$  such that  $r_{n_{k+1}}$  belongs to this interval. Then  $|r_{n_{k+1}} - a| < \frac{1}{k+1}$  and hence (2) and (3) hold for  $k+1$  in place of  $k$ . The procedure defines  $(n_k)_{k \in \mathbb{N}}$  by induction. Since (3) holds, (1) holds and we conclude that  $\lim_{k \rightarrow \infty} r_{n_k} = a$ .

#### Example 4

Suppose that  $(s_n)$  is a sequence of positive numbers such that  $\inf\{s_n : n \in \mathbb{N}\} = 0$ . The sequence  $(s_n)$  need not converge or even be bounded, but it has a subsequence that converges monotonically to 0. We will again give an inductive construction. Since  $\inf\{s_n : n \in \mathbb{N}\} = 0$ , there exists  $n_1 \in \mathbb{N}$  such that  $s_{n_1} < 1$ . Suppose that  $n_1, n_2, \dots, n_k$  have been selected so that

$$n_1 < n_2 < \dots < n_k \quad (1)$$

and

$$s_{n_{j+1}} < \min \left\{ s_{n_j}, \frac{1}{j+1} \right\} \quad \text{for } j = 1, 2, \dots, k-1. \quad (2)$$

Note that we are requiring  $s_{n_{j+1}} < s_{n_j}$  so that our subsequence will be monotonic, and we are requiring  $s_{n_{j+1}} < \frac{1}{j+1}$  to guarantee that

it will converge to 0. Since  $\min\{s_n : 1 \leq n \leq n_k\} > 0$ , it follows that  $\inf\{s_n : n > n_k\} = 0$ . Thus there exists  $n_{k+1} > n_k$  such that  $s_{n_{k+1}} < \min\{s_{n_k}, \frac{1}{k+1}\}$ . Hence (1) and (2) hold for  $k+1$  in place of  $k$ , and the construction continues by induction. As noted above, (2) shows that  $(s_{n_k})$  converges monotonically to 0.

The next theorem is almost obvious.

### 11.2 Theorem.

*If the sequence  $(s_n)$  converges, then every subsequence converges to the same limit.*

#### Proof

Let  $(s_{n_k})$  denote a subsequence of  $(s_n)$ . Note that  $n_k \geq k$  for all  $k$ . This is easy to prove by induction; in fact,  $n_1 \geq 1$  and  $n_k \geq k$  implies  $n_{k+1} > n_k \geq k$  and hence  $n_{k+1} \geq k+1$ .

Let  $s = \lim s_n$  and let  $\epsilon > 0$ . There exists  $N$  so that  $n > N$  implies  $|s_n - s| < \epsilon$ . Now  $k > N$  implies  $n_k > N$ , which implies  $|s_{n_k} - s| < \epsilon$ . Thus

$$\lim_{k \rightarrow \infty} s_{n_k} = s. \quad \blacksquare$$

Our immediate goal is to prove the Bolzano-Weierstrass theorem which asserts that every bounded sequence has a convergent subsequence. First we prove a theorem about monotonic subsequences.

### 11.3 Theorem.

*Every sequence  $(s_n)$  has a monotonic subsequence.*

#### Proof

Let's say that the  $n$ -th term is *dominant* if it is greater than every term which follows it:

$$s_m < s_n \quad \text{for all } m > n. \quad (1)$$

*Case 1.* Suppose that there are infinitely many dominant terms, and let  $(s_{n_k})$  be any subsequence consisting solely of dominant terms. Then  $s_{n_{k+1}} < s_{n_k}$  for all  $k$  by (1), so  $(s_{n_k})$  is a decreasing sequence.

*Case 2.* Suppose that there are only finitely many dominant terms. Select  $n_1$  so that  $s_{n_1}$  is beyond all the dominant terms of the

sequence. Then

$$\text{given } N \geq n_1 \text{ there exists } m > N \text{ such that } s_m \geq s_N. \quad (2)$$

Applying (2) with  $N = n_1$  we select  $n_2 > n_1$  such that  $s_{n_2} \geq s_{n_1}$ . Suppose that  $n_1, n_2, \dots, n_k$  have been selected so that

$$n_1 < n_2 < \dots < n_k \quad (3)$$

and

$$s_{n_1} \leq s_{n_2} \leq \dots \leq s_{n_k}. \quad (4)$$

Applying (2) with  $N = n_k$  we select  $n_{k+1} > n_k$  such that  $s_{n_{k+1}} \geq s_{n_k}$ . Then (3) and (4) hold with  $k+1$  in place of  $k$ , the procedure continues by induction, and we obtain a nondecreasing subsequence  $(s_{n_k})$ . ■

The elegant proof in Theorem 11.3 was brought to our attention by David M. Bloom and is based on a solution in D. J. Newman's beautiful book *A Problem Seminar*, Springer-Verlag, New York-Berlin-Heidelberg: 1982.

#### 11.4 Corollary.

*Let  $(s_n)$  be any sequence. There exists a monotonic subsequence whose limit is  $\limsup s_n$ , and there exists a monotonic subsequence whose limit is  $\liminf s_n$ .*

##### Proof

For  $N$  in  $\mathbb{N}$ , let  $v_N = \sup\{s_n : n > N\}$ . Then  $v_1 \geq v_2 \geq v_3 \geq \dots$  and  $v = \lim_N v_N = \limsup s_n$ ; see Definition 10.6. Our task is to show that there is a monotonic subsequence of  $(s_n)$  that converges to  $v$ . If  $v = -\infty$ , then  $\lim s_n = -\infty$  by Theorem 10.7, the sequence  $(s_n)$  itself converges to  $\limsup s_n$ , and so a monotonic subsequence of  $(s_n)$  converges to  $\limsup s_n$  by Theorem 11.3. Henceforth we assume that  $v \neq -\infty$ .

First look at Case 1 of the previous proof. Then  $s_{n_k} = \sup\{s_n : n \geq n_k\} = v_{n_k-1}$ , so  $\lim_{k \rightarrow \infty} s_{n_k} = \lim_N v_N = \limsup s_n$ , as desired.

Suppose now that there are only finitely many dominant terms, and let  $s_{m_0}$  be the last dominant term. We claim

$$\sup\{s_n : n > N\} = v_N = v \quad \text{for } N > m_0. \quad (2.1)$$

Otherwise, since  $(v_N)$  is a nonincreasing sequence, there is an  $N > m_0$  so that  $v_{N+1} < v_N$ . So  $s_{N+1}$  must be bigger than  $v_{N+1} = \sup\{s_n :$

$n \geq N + 2\}$ , but this implies that  $s_{N+1}$  is dominant, contrary to the choice of  $m_0$ .

If infinitely many  $s_n$  equal  $v$ , simply select a subsequence of  $(s_n)$  consisting of terms equal to  $v$ . Otherwise, there exists  $n_1 \geq m_0$  so that

$$s_n < v \quad \text{for all } n \geq n_1. \quad (2.2)$$

Select a sequence  $(t_N)$  that increases to  $v$ . If  $v$  is finite,  $t_N = v - \frac{1}{N}$  will do if  $v$  is finite, and  $t_N = N$  will do if  $v = +\infty$ . The desired subsequence is obtained by induction, the first term being  $s_{n_1}$ . Note that, by (2), we have  $s_{n_1} < v$ . Assume that  $n_1 < n_2 < \cdots < n_k$  have been selected so that

$$s_{n_1} < s_{n_2} < \cdots < s_{n_k} < v, \quad (2.3)$$

$$\text{and } t_k < s_{n_k} \quad \text{for } k \geq 2. \quad (2.4)$$

Using (1), we select  $n_{k+1} > n_k$  so that  $s_{n_{k+1}} > s_{n_k}$  and  $s_{n_{k+1}} > t_{k+1}$ . By (2), we also have  $s_{n_{k+1}} < v$ . The procedure continues by induction, and by (3) we obtain an increasing subsequence of  $(s_n)$ . Also, by (4), we have  $t_k < s_{n_k} < v$  for all  $k$ , so  $v = \lim_{k \rightarrow \infty} t_k \leq \lim_{k \rightarrow \infty} s_{n_k} \leq v$ , and the subsequence  $(s_{n_k})$  converges to  $v$  as desired.

The assertion about  $\liminf s_n$  has a similar proof, but it also can be derived from the first assertion; see Exercise 11.8. This revised proof is based on correspondence with Ray Hoobler, City College, CUNY. ■

### 11.5 Bolzano-Weierstrass Theorem.

*Every bounded sequence has a convergent subsequence.*

#### Proof

If  $(s_n)$  is a bounded sequence, it has a monotonic subsequence by Theorem 11.3. The subsequence converges by Theorem 10.2. ■

The Bolzano-Weierstrass theorem is very important and will be used at critical points in Chapter 3. Our proof, based on Theorem 11.3, is somewhat nonstandard for reasons we now discuss. Many of the notions introduced in this chapter make equally good sense in more general settings. For example, the ideas of convergent sequence, Cauchy sequence and bounded sequence all make sense for a sequence  $(s_n)$  where each  $s_n$  belongs to the plane. But

the idea of a monotonic sequence does not carry over. It turns out that the Bolzano-Weierstrass theorem also holds in the plane and in many other settings [see Theorem 13.5], but clearly it would no longer be appropriate to prove it directly from an analogue of Theorem 11.3. Since the Bolzano-Weierstrass Theorem 11.5 generalizes to settings where Theorem 11.3 makes little sense, in applications we will emphasize 11.5 rather than 11.3.

We need one more notion, and then we will be able to tie our various concepts together in Theorem 11.7.

### 11.6 Definition.

Let  $(s_n)$  be a sequence in  $\mathbb{R}$ . A *subsequential limit* is any real number or symbol  $+\infty$  or  $-\infty$  that is the limit of some subsequence of  $(s_n)$ .

When a sequence has a limit  $s$ , then all subsequences have limit  $s$ , so  $\{s\}$  is the set of subsequential limits. The interesting case is when the original sequence does not have a limit. We return to some of the examples discussed after Definition 11.1.

### Example 5

Consider  $(s_n)$  where  $s_n = n^2(-1)^n$ . The subsequence of even terms diverges to  $+\infty$ , and the subsequence of odd terms diverges to  $-\infty$ . All subsequences that have a limit diverge to  $+\infty$  or  $-\infty$ , so that  $\{-\infty, +\infty\}$  is exactly the set of subsequential limits of  $(s_n)$ .

### Example 6

Consider the sequence  $a_n = \sin(\frac{n\pi}{3})$ . This sequence takes each of the values  $\frac{1}{2}\sqrt{3}$ ,  $0$  and  $-\frac{1}{2}\sqrt{3}$  an infinite number of times. The only convergent subsequences are constant from some term on, and  $\{-\frac{1}{2}\sqrt{3}, 0, \frac{1}{2}\sqrt{3}\}$  is the set of subsequential limits of  $(a_n)$ . If  $n_k = 3k$ , then  $a_{n_k} = 0$  for all  $k \in \mathbb{N}$  and obviously  $\lim_{k \rightarrow \infty} a_{n_k} = 0$ . If  $n_k = 6k + 1$ , then  $a_{n_k} = \frac{1}{2}\sqrt{3}$  for all  $k$  and  $\lim_{k \rightarrow \infty} a_{n_k} = \frac{1}{2}\sqrt{3}$ . And if  $n_k = 6k + 4$ , then  $\lim_{k \rightarrow \infty} a_{n_k} = -\frac{1}{2}\sqrt{3}$ .

### Example 7

Let  $(r_n)$  be a list of all rational numbers. It was shown in Example 3 that every real number is a subsequential limit of  $(r_n)$ . Also,  $+\infty$

and  $-\infty$  are subsequential limits; see Exercise 11.7. Consequently,  $\mathbb{R} \cup \{-\infty, +\infty\}$  is the set of subsequential limits of  $(r_n)$ .

**Example 8**

Let  $b_n = n[1 + (-1)^n]$  for  $n \in \mathbb{N}$ . Then  $b_n = 2n$  for even  $n$  and  $b_n = 0$  for odd  $n$ . Thus  $\{0, +\infty\}$  is the set of subsequential limits of  $(b_n)$ .

**11.7 Theorem.**

Let  $(s_n)$  be any sequence in  $\mathbb{R}$ , and let  $S$  denote the set of subsequential limits of  $(s_n)$ .

- (i)  $S$  is nonempty.
- (ii)  $\sup S = \limsup s_n$  and  $\inf S = \liminf s_n$ .
- (iii)  $\lim s_n$  exists if and only if  $S$  has exactly one element, namely  $\lim s_n$ .

**Proof**

(i) is an immediate consequence of Corollary 11.4.

To prove (ii), consider any limit  $t$  of a subsequence  $(s_{n_k})$  of  $(s_n)$ . By Theorem 10.7 we have  $t = \liminf s_{n_k} = \limsup s_{n_k}$ . Since  $\{s_{n_k} : k > N\} \subseteq \{s_n : n > N\}$  for each  $N \in \mathbb{N}$ , we have

$$\liminf s_n \leq \liminf s_{n_k} = t = \limsup s_{n_k} \leq \limsup s_n.$$

This inequality holds for all  $t$  in  $S$ ; therefore

$$\liminf s_n \leq \inf S \leq \sup S \leq \limsup s_n.$$

Corollary 11.4 shows that  $\liminf s_n$  and  $\limsup s_n$  both belong to  $S$ . Therefore (ii) holds.

Assertion (iii) is simply a reformulation of Theorem 10.7. ■

Theorem 11.7 and Corollary 11.4 show that  $\limsup s_n$  is exactly the largest subsequential limit of  $(s_n)$ , and  $\liminf s_n$  is exactly the smallest subsequential limit of  $(s_n)$ . This makes it easy to calculate  $\limsup$ 's and  $\liminf$ 's.

We return to the examples given before Theorem 11.7.

**Example 9**

If  $s_n = n^2(-1)^n$ , then  $S = \{-\infty, +\infty\}$  as noted in Example 5. Therefore  $\limsup s_n = \sup S = +\infty$  and  $\liminf s_n = \inf S = -\infty$ .

**Example 10**

If  $a_n = \sin(\frac{n\pi}{3})$ , then  $S = \{-\frac{1}{2}\sqrt{3}, 0, \frac{1}{2}\sqrt{3}\}$  as observed in Example 6. Hence  $\limsup a_n = \sup S = \frac{1}{2}\sqrt{3}$  and  $\liminf a_n = \inf S = -\frac{1}{2}\sqrt{3}$ .

**Example 11**

If  $(r_n)$  denotes a list of all rational numbers, then the set  $\mathbb{R} \cup \{-\infty, +\infty\}$  is the set of subsequential limits of  $(r_n)$ . Consequently we have  $\limsup r_n = +\infty$  and  $\liminf r_n = -\infty$ .

**Example 12**

If  $b_n = n[1 + (-1)^n]$ , then  $\limsup b_n = +\infty$  and  $\liminf b_n = 0$ ; see Example 8.

The next result shows that the set  $S$  of subsequential limits always contains all limits of sequences *from*  $S$ . Such sets are called *closed sets*. Sets of this sort will be discussed further in the optional §13.

**11.8 Theorem.**

*Let  $S$  denote the set of subsequential limits of a sequence  $(s_n)$ . Suppose  $(t_n)$  is a sequence in  $S \cap \mathbb{R}$  and that  $t = \lim t_n$ . Then  $t$  belongs to  $S$ .*

**Proof**

Since a subsequence of  $(s_n)$  converges to  $t_1$ , there exists  $n_1$  such that  $|s_{n_1} - t_1| < 1$ . Assume that  $n_1, n_2, \dots, n_k$  have been selected so that

$$n_1 < n_2 < \dots < n_k \quad (1)$$

and

$$|s_{n_j} - t_j| < \frac{1}{j} \quad \text{for } j = 1, 2, \dots, k. \quad (2)$$

Since a subsequence of  $(s_n)$  converges to  $t_{k+1}$ , there exists  $n_{k+1} > n_k$  such that  $|s_{n_{k+1}} - t_{k+1}| < \frac{1}{k+1}$ . Thus (1) and (2) hold for  $k+1$ .

For the rest of the proof we need to consider cases. Suppose first that  $t \in \mathbb{R}$ , i.e., that  $t$  is not  $+\infty$  or  $-\infty$ . Since

$$|s_{n_k} - t| \leq |s_{n_k} - t_k| + |t_k - t| < \frac{1}{k} + |t_k - t| \quad (3)$$

for all  $k \in \mathbb{N}$ , it follows easily that  $\lim_{k \rightarrow \infty} s_{n_k} = t$ , so  $t$  belongs to  $S$ . [To check that  $\lim_{k \rightarrow \infty} s_{n_k} = t$ , consider  $\epsilon > 0$ . There exists  $N$  so that  $k > N$  implies  $|t_k - t| < \frac{\epsilon}{2}$ . If  $k > \max\{N, \frac{2}{\epsilon}\}$ , then  $\frac{1}{k} < \frac{\epsilon}{2}$  and  $|t_k - t| < \frac{\epsilon}{2}$ , so  $|s_{n_k} - t| < \epsilon$  by (3).]

Suppose next that  $t = +\infty$ . From (2) we have

$$s_{n_k} > t_k - \frac{1}{k} \quad \text{for } k \in \mathbb{N}. \quad (4)$$

Since  $\lim t_k = +\infty$  it follows easily that  $\lim_{k \rightarrow \infty} s_{n_k} = +\infty$ . Therefore  $t = +\infty$  belongs to  $S$ . The case  $t = -\infty$  is handled in a similar way. ■

## Exercises

**11.1.** Let  $a_n = 3 + 2(-1)^n$  for  $n \in \mathbb{N}$ .

- (a) List the first eight terms of the sequence  $(a_n)$ .
- (b) Give a subsequence that is constant [takes a single value]. Specify the selection function  $\sigma$ .

**11.2.** Consider the sequences defined as follows:

$$a_n = (-1)^n, \quad b_n = \frac{1}{n}, \quad c_n = n^2, \quad d_n = \frac{6n+4}{7n-3}.$$

- (a) For each sequence, give an example of a monotone subsequence.
- (b) For each sequence, give its set of subsequential limits.
- (c) For each sequence, give its  $\limsup$  and  $\liminf$ .
- (d) Which of the sequences converges? diverges to  $+\infty$ ? diverges to  $-\infty$ ?
- (e) Which of the sequences is bounded?

**11.3.** Repeat Exercise 11.2 for the sequences:

$$s_n = \cos\left(\frac{n\pi}{3}\right), \quad t_n = \frac{3}{4n+1}, \quad u_n = \left(-\frac{1}{2}\right)^n, \quad v_n = (-1)^n + \frac{1}{n}.$$

**11.4.** Repeat Exercise 11.2 for the sequences:

$$w_n = (-2)^n, \quad x_n = 5^{(-1)^n}, \quad y_n = 1 + (-1)^n, \quad z_n = n \cos\left(\frac{n\pi}{4}\right).$$

**11.5.** Let  $(q_n)$  be an enumeration of all the rationals in the interval  $(0, 1]$ .

- (a) Give the set of subsequential limits for  $(q_n)$ .
- (b) Give the values of  $\limsup q_n$  and  $\liminf q_n$ .

- 11.6.** Show that every subsequence of a subsequence of a given sequence is itself a subsequence of the given sequence. *Hint:* Define subsequences as in (3) of Definition 11.1.
- 11.7.** Let  $(r_n)$  be an enumeration of the set  $\mathbb{Q}$  of all rational numbers. Show that there exists a subsequence  $(r_{n_k})$  such that  $\lim_{k \rightarrow \infty} r_{n_k} = +\infty$ .
- 11.8. (a)** Use Definition 10.6 and Exercise 5.4 to prove that  $\liminf s_n = -\limsup(-s_n)$ .
- (b)** Let  $(t_k)$  be a monotonic subsequence of  $(-s_n)$  converging to  $\limsup(-s_n)$ . Show that  $(-t_k)$  is a monotonic subsequence of  $(s_n)$  converging to  $\liminf s_n$ . Observe that this completes the proof of Corollary 11.4.
- 11.9. (a)** Show that the closed interval  $[a, b]$  is a closed set.
- (b)** Is there a sequence  $(s_n)$  such that  $(0, 1)$  is its set of subsequential limits?
- 11.10.** Let  $(s_n)$  be the sequence of numbers in Figure 11.2 listed in the indicated order.
- (a)** Find the set  $S$  of subsequential limits of  $(s_n)$ .
- (b)** Determine  $\limsup s_n$  and  $\liminf s_n$ .

**FIGURE 11.2**

## §12 lim sup's and lim inf's

Let  $(s_n)$  be any sequence of real numbers, and let  $S$  be the set of subsequential limits of  $(s_n)$ . Recall that

$$\limsup s_n = \lim_{N \rightarrow \infty} \sup \{s_n : n > N\} = \sup S \quad (*)$$

and

$$\liminf s_n = \lim_{N \rightarrow \infty} \inf \{s_n : n > N\} = \inf S. \quad (**)$$

The first equalities in  $(*)$  and  $(**)$  are the definitions given in 10.6, and the second equalities are proved in Theorem 11.7. This section is designed to increase the students' familiarity with these concepts. Most of the material is given in the exercises. We illustrate the techniques by proving some results that will be needed later in the text.

### 12.1 Theorem.

*If  $(s_n)$  converges to a positive real number  $s$  and  $(t_n)$  is any sequence, then*

$$\limsup s_n t_n = s \cdot \limsup t_n.$$

*Here we allow the conventions:  $s \cdot (+\infty) = +\infty$  and  $s \cdot (-\infty) = -\infty$  for  $s > 0$ .*

### Proof

We first show

$$\limsup s_n t_n \geq s \cdot \limsup t_n. \quad (1)$$

We have three cases. Let  $\beta = \limsup t_n$ .

*Case 1.* Suppose  $\beta$  is finite.

By Corollary 11.4, there exists a subsequence  $(t_{n_k})$  of  $(t_n)$  such that  $\lim_{k \rightarrow \infty} t_{n_k} = \beta$ . We also have  $\lim_{k \rightarrow \infty} s_{n_k} = s$  [by Theorem 11.2], so  $\lim_{k \rightarrow \infty} s_{n_k} t_{n_k} = s\beta$ . Thus  $(s_{n_k} t_{n_k})$  is a subsequence of  $(s_n t_n)$  that converges to  $s\beta$ , and therefore  $s\beta \leq \limsup s_n t_n$ . [Recall that  $\limsup s_n t_n$  is the largest possible limit of a subsequence of  $(s_n t_n)$ .] Thus (1) holds.

Case 2. Suppose  $\beta = +\infty$ .

There exists a subsequence  $(t_{n_k})$  of  $(t_n)$  such that  $\lim_{k \rightarrow \infty} t_{n_k} = +\infty$ . Since  $\lim_{k \rightarrow \infty} s_{n_k} = s > 0$ , Theorem 9.9 shows that  $\lim_{k \rightarrow \infty} s_{n_k} t_{n_k} = +\infty$ . Hence  $\limsup s_n t_n = +\infty$ , so (1) clearly holds.

Case 3. Suppose  $\beta = -\infty$ .

Since  $s > 0$ , the right-hand side of (1) is equal to  $s \cdot (-\infty) = -\infty$ . Hence (1) is obvious in this case.

We have now established (1) in all cases. For the reversed inequality, we resort to a little trick. First note that we may ignore the first few terms of  $(s_n)$  and assume that all  $s_n \neq 0$ . Then we can write  $\lim_{s_n} \frac{1}{s_n} = \frac{1}{s}$  by Lemma 9.5. Now we apply (1) with  $s_n$  replaced by  $\frac{1}{s_n}$  and  $t_n$  replaced by  $s_n t_n$ :

$$\limsup t_n = \limsup \left( \frac{1}{s_n} \right) (s_n t_n) \geq \left( \frac{1}{s} \right) \limsup s_n t_n,$$

i.e.,

$$\limsup s_n t_n \leq s \cdot \limsup t_n.$$

This inequality and (1) prove the theorem. ■

The next theorem will be useful in dealing with infinite series; see the proof of the Ratio Test 14.8.

### 12.2 Theorem.

Let  $(s_n)$  be any sequence of nonzero real numbers. Then we have

$$\liminf \left| \frac{s_{n+1}}{s_n} \right| \leq \liminf |s_n|^{1/n} \leq \limsup |s_n|^{1/n} \leq \limsup \left| \frac{s_{n+1}}{s_n} \right|.$$

#### Proof

The middle inequality is obvious. The first and third inequalities have similar proofs. We will prove the third inequality and leave the first inequality to Exercise 12.11.

Let  $\alpha = \limsup |s_n|^{1/n}$  and  $L = \limsup \left| \frac{s_{n+1}}{s_n} \right|$ . We need to prove that  $\alpha \leq L$ . This is obvious if  $L = +\infty$ , so we assume  $L < +\infty$ . To prove  $\alpha \leq L$  it suffices to show

$$\alpha \leq L_1 \quad \text{for any } L_1 > L. \tag{1}$$

Since

$$L = \limsup \left| \frac{s_{n+1}}{s_n} \right| = \lim_{N \rightarrow \infty} \sup \left\{ \left| \frac{s_{n+1}}{s_n} \right| : n > N \right\} < L_1,$$

there exists a natural number  $N$  such that

$$\sup \left\{ \left| \frac{s_{n+1}}{s_n} \right| : n \geq N \right\} < L_1.$$

Thus

$$\left| \frac{s_{n+1}}{s_n} \right| < L_1 \quad \text{for } n \geq N. \quad (2)$$

Now for  $n > N$  we can write

$$|s_n| = \left| \frac{s_n}{s_{n-1}} \right| \cdot \left| \frac{s_{n-1}}{s_{n-2}} \right| \cdots \left| \frac{s_{N+1}}{s_N} \right| \cdot |s_N|.$$

There are  $n - N$  fractions here, so applying (2) we see that

$$|s_n| < L_1^{n-N} |s_N| \quad \text{for } n > N.$$

Since  $L_1$  and  $N$  are fixed in this argument,  $a = L_1^{-N} |s_N|$  is a positive constant and we may write

$$|s_n| < L_1^n a \quad \text{for } n > N.$$

Therefore we have

$$|s_n|^{1/n} < L_1 a^{1/n} \quad \text{for } n > N.$$

Since  $\lim_{n \rightarrow \infty} a^{1/n} = 1$  by Example 9.7(d), we conclude that  $\alpha = \limsup |s_n|^{1/n} \leq L_1$ ; see Exercise 12.1. Consequently (1) holds as desired. ■

### 12.3 Corollary.

If  $\lim \left| \frac{s_{n+1}}{s_n} \right|$  exists [and equals  $L$ ], then  $\lim |s_n|^{1/n}$  exists [and equals  $L$ ].

#### Proof

If  $\lim \left| \frac{s_{n+1}}{s_n} \right| = L$ , then all four values in Theorem 12.2 must equal  $L$ . Hence  $\lim |s_n|^{1/n} = L$ ; see Theorem 10.7. ■

**Exercises**

**12.1.** Let  $(s_n)$  and  $(t_n)$  be sequences and suppose that there exists  $N_0$  such that  $s_n \leq t_n$  for all  $n > N_0$ . Show that  $\liminf s_n \leq \liminf t_n$  and  $\limsup s_n \leq \limsup t_n$ . *Hint:* Use Definition 10.6 and Exercise 9.9(c).

**12.2.** Prove that  $\limsup |s_n| = 0$  if and only if  $\lim s_n = 0$ .

**12.3.** Let  $(s_n)$  and  $(t_n)$  be the following sequences that repeat in cycles of four:

$$(s_n) = (0, 1, 2, 1, 0, 1, 2, 1, 0, 1, 2, 1, 0, 1, 2, 1, 0, \dots)$$

$$(t_n) = (2, 1, 1, 0, 2, 1, 1, 0, 2, 1, 1, 0, 2, 1, 1, 0, 2, \dots)$$

Find

- |  |                                   |
|--|-----------------------------------|
| <b>(a)</b> $\liminf s_n + \liminf t_n$ , | <b>(b)</b> $\liminf(s_n + t_n)$ , |
| <b>(c)</b> $\liminf s_n + \limsup t_n$ , | <b>(d)</b> $\limsup(s_n + t_n)$ , |
| <b>(e)</b> $\limsup s_n + \limsup t_n$ , | <b>(f)</b> $\liminf(s_n t_n)$ ,   |
| <b>(g)</b> $\limsup(s_n t_n)$            |                                   |

**12.4.** Show that  $\limsup(s_n + t_n) \leq \limsup s_n + \limsup t_n$  for bounded sequences  $(s_n)$  and  $(t_n)$ . *Hint:* First show

$$\sup\{s_n + t_n : n > N\} \leq \sup\{s_n : n > N\} + \sup\{t_n : n > N\}.$$

Then apply Exercise 9.9(c).

**12.5.** Use Exercises 11.8(a) and 12.4 to prove

$$\liminf(s_n + t_n) \geq \liminf s_n + \liminf t_n$$

for bounded sequences  $(s_n)$  and  $(t_n)$ .

**12.6.** Let  $(s_n)$  be a bounded sequence, and let  $k$  be a nonnegative real number.

- (a)** Prove that  $\limsup(ks_n) = k \cdot \limsup s_n$ .
- (b)** Do the same for  $\liminf$ . *Hint:* Use Exercise 11.8(a).
- (c)** What happens in (a) and (b) if  $k < 0$ ?

**12.7.** Prove that if  $\limsup s_n = +\infty$  and  $k > 0$ , then  $\limsup(ks_n) = +\infty$ .

**12.8.** Let  $(s_n)$  and  $(t_n)$  be bounded sequences of nonnegative numbers. Prove that  $\limsup s_n t_n \leq (\limsup s_n)(\limsup t_n)$ .

**12.9.** **(a)** Prove that if  $\lim s_n = +\infty$  and  $\liminf t_n > 0$ , then  $\lim s_n t_n = +\infty$ .

(b) Prove that if  $\limsup s_n = +\infty$  and  $\liminf t_n > 0$ , then  $\limsup s_n t_n = +\infty$ .

(c) Observe that Exercise 12.7 is the special case of (b) where  $t_n = k$  for all  $n \in \mathbb{N}$ .

12.10. Prove that  $(s_n)$  is bounded if and only if  $\limsup |s_n| < +\infty$ .

12.11. Prove the first inequality in Theorem 12.2.

12.12. Let  $(s_n)$  be a sequence of nonnegative numbers, and for each  $n$  define  $\sigma_n = \frac{1}{n}(s_1 + s_2 + \cdots + s_n)$ .

(a) Show that

$$\liminf s_n \leq \liminf \sigma_n \leq \limsup \sigma_n \leq \limsup s_n.$$

*Hint:* For the last inequality, show first that  $M > N$  implies

$$\sup\{\sigma_n : n > M\} \leq \frac{1}{M}(s_1 + s_2 + \cdots + s_N) + \sup\{s_n : n > N\}.$$

(b) Show that if  $\lim s_n$  exists, then  $\lim \sigma_n$  exists and  $\lim \sigma_n = \lim s_n$ .

12.13. Let  $(s_n)$  be a bounded sequence in  $\mathbb{R}$ . Let  $A$  be the set of  $a \in \mathbb{R}$  such that  $\{n \in \mathbb{N} : s_n < a\}$  is finite, i.e., all but finitely many  $s_n$  are  $\geq a$ . Let  $B$  be the set of  $b \in \mathbb{R}$  such that  $\{n \in \mathbb{N} : s_n > b\}$  is finite. Prove that  $\sup A = \liminf s_n$  and  $\inf B = \limsup s_n$ .

12.14. Calculate (a)  $\lim(n!)^{1/n}$ , (b)  $\lim \frac{1}{n}(n!)^{1/n}$ .

## §13 \* Some Topological Concepts in Metric Spaces

In this book we are restricting our attention to analysis on  $\mathbb{R}$ . Accordingly, we have taken full advantage of the order properties of  $\mathbb{R}$  and studied such important notions as  $\limsup$ 's and  $\liminf$ 's. In §3 we briefly introduced a distance function on  $\mathbb{R}$ . Most of our analysis could have been based on the notion of distance, in which case it becomes easy and natural to work in a more general setting. For example, analysis on the  $k$ -dimensional Euclidean spaces  $\mathbb{R}^k$  is important, but these spaces do not have the useful natural ordering that  $\mathbb{R}$  has, unless of course  $k = 1$ .

**13.1 Definition.**

Let  $S$  be a set, and suppose  $d$  is a function defined for all pairs  $(x, y)$  of elements from  $S$  satisfying

**D1.**  $d(x, x) = 0$  for all  $x \in S$  and  $d(x, y) > 0$  for distinct  $x, y$  in  $S$ .

**D2.**  $d(x, y) = d(y, x)$  for all  $x, y \in S$ .

**D3.**  $d(x, z) \leq d(x, y) + d(y, z)$  for all  $x, y, z \in S$ .

Such a function  $d$  is called a *distance function* or a *metric* on  $S$ . A *metric space*  $S$  is a set  $S$  together with a metric on it. Properly speaking, the metric space is the pair  $(S, d)$  since a set  $S$  may well have more than one metric on it; see Exercise 13.1.

**Example 1**

As in Definition 3.4, let  $\text{dist}(a, b) = |a - b|$  for  $a, b \in \mathbb{R}$ . Then  $\text{dist}$  is a metric on  $\mathbb{R}$ . Note that Corollary 3.6 gives D3 in this case. As remarked there, the inequality

$$\text{dist}(a, c) \leq \text{dist}(a, b) + \text{dist}(b, c)$$

is called the triangle inequality. In fact, for any metric  $d$ , property D3 is called the *triangle inequality*.

**Example 2**

The space of all  $k$ -tuples

$$\mathbf{x} = (x_1, x_2, \dots, x_k) \quad \text{where} \quad x_j \in \mathbb{R} \quad \text{for} \quad j = 1, 2, \dots, k,$$

is called *k-dimensional Euclidean space* and written  $\mathbb{R}^k$ . As noted in Exercise 13.1,  $\mathbb{R}^k$  has several metrics on it. The most familiar metric is the one that gives the ordinary distance in the plane  $\mathbb{R}^2$  or in 3-space  $\mathbb{R}^3$ :

$$d(\mathbf{x}, \mathbf{y}) = \left[ \sum_{j=1}^k (x_j - y_j)^2 \right]^{1/2}.$$

[The summation notation  $\sum$  is explained in 14.1.] Obviously this function  $d$  satisfies properties D1 and D2. The triangle inequality D3 is not so obvious. For a proof, see for example [33], §6.1, or [36], 1.37.

### 13.2 Definition.

A sequence  $(s_n)$  in a metric space  $(S, d)$  *converges to*  $s$  in  $S$  if  $\lim_{n \rightarrow \infty} d(s_n, s) = 0$ . A sequence  $(s_n)$  in  $S$  is a *Cauchy sequence* if for each  $\epsilon > 0$  there exists an  $N$  such that

$$m, n > N \quad \text{implies} \quad d(s_m, s_n) < \epsilon.$$

The metric space  $(S, d)$  is said to be *complete* if every Cauchy sequence in  $S$  converges to some element in  $S$ .

Since the Completeness Axiom 4.4 deals with least upper bounds, the word “complete” now appears to have two meanings. However, these two uses of the term are very closely related and both reflect the property that the space is complete, i.e., has no gaps. Theorem 10.11 asserts that the metric space  $(\mathbb{R}, \text{dist})$  is a complete metric space, and the proof uses the Completeness Axiom 4.4. We could just as well have taken as an axiom the completeness of  $(\mathbb{R}, \text{dist})$  as a metric space and proved the least upper bound property in 4.4 as a theorem. We did not do so because the concept of least upper bound in  $\mathbb{R}$  seems to us more fundamental than the concept of Cauchy sequence.

We will prove that  $\mathbb{R}^k$  is complete. But we have a notational problem, since we like subscripts for sequences and for coordinates of points in  $\mathbb{R}^k$ . When there is a conflict, we will write  $(\mathbf{x}^{(n)})$  for a sequence instead of  $(\mathbf{x}_n)$ . In this case,

$$\mathbf{x}^{(n)} = (x_1^{(n)}, x_2^{(n)}, \dots, x_k^{(n)}).$$

Unless otherwise specified, *the metric in  $\mathbb{R}^k$  is always as given in Example 2.*

### 13.3 Lemma.

A sequence  $(\mathbf{x}^{(n)})$  in  $\mathbb{R}^k$  converges if and only if for each  $j = 1, 2, \dots, k$ , the sequence  $(x_j^{(n)})$  converges in  $\mathbb{R}$ . A sequence  $(\mathbf{x}^{(n)})$  in  $\mathbb{R}^k$  is a Cauchy sequence if and only if each sequence  $(x_j^{(n)})$  is a Cauchy sequence in  $\mathbb{R}$ .

#### Proof

The proof of the first assertion is left to Exercise 13.2. For the second assertion, we first observe for  $\mathbf{x}, \mathbf{y}$  in  $\mathbb{R}^k$  and  $j = 1, 2, \dots, k$ ,

$$|x_j - y_j| \leq d(\mathbf{x}, \mathbf{y}) \leq \sqrt{k} \max\{|x_j - y_j| : j = 1, 2, \dots, k\}. \quad (1)$$

Suppose  $(\mathbf{x}^{(n)})$  is a Cauchy sequence in  $\mathbb{R}^k$ , and consider fixed  $j$ . If  $\epsilon > 0$ , there exists  $N$  such that

$$m, n > N \quad \text{implies} \quad d(\mathbf{x}^{(m)}, \mathbf{x}^{(n)}) < \epsilon.$$

From (1) we see that

$$m, n > N \quad \text{implies} \quad |x_j^{(m)} - x_j^{(n)}| < \epsilon,$$

so  $(x_j^{(n)})$  is a Cauchy sequence in  $\mathbb{R}$ .

Now suppose each sequence  $(x_j^{(n)})$  is a Cauchy sequence in  $\mathbb{R}$ . Let  $\epsilon > 0$ . For  $j = 1, 2, \dots, k$ , there exist  $N_j$  such that

$$m, n > N_j \quad \text{implies} \quad |x_j^{(m)} - x_j^{(n)}| < \frac{\epsilon}{\sqrt{k}}.$$

If  $N = \max\{N_1, N_2, \dots, N_k\}$ , then by (1)

$$m, n > N \quad \text{implies} \quad d(\mathbf{x}^{(m)}, \mathbf{x}^{(n)}) < \epsilon,$$

i.e.,  $(\mathbf{x}^{(n)})$  is a Cauchy sequence in  $\mathbb{R}^k$ . ■

### 13.4 Theorem.

*Euclidean  $k$ -space  $\mathbb{R}^k$  is complete.*

#### Proof

Consider a Cauchy sequence  $(\mathbf{x}^{(n)})$  in  $\mathbb{R}^k$ . By Lemma 13.3,  $(x_j^{(n)})$  is a Cauchy sequence in  $\mathbb{R}$  for each  $j$ . Hence by Theorem 10.11,  $(x_j^{(n)})$  converges to some real number  $x_j$ . By Lemma 13.3 again, the sequence  $(\mathbf{x}^{(n)})$  converges, in fact to  $\mathbf{x} = (x_1, x_2, \dots, x_k)$ . ■

We now can prove the Bolzano-Weierstrass theorem for  $\mathbb{R}^k$ ; compare Theorem 11.5. A set  $S$  in  $\mathbb{R}^k$  is *bounded* if there exists  $M > 0$  such that

$$\max\{|x_j| : j = 1, 2, \dots, k\} \leq M \quad \text{for all} \quad \mathbf{x} \in S.$$

### 13.5 Bolzano-Weierstrass Theorem.

*Every bounded sequence in  $\mathbb{R}^k$  has a convergent subsequence.*

#### Proof

Let  $(\mathbf{x}^{(n)})$  be a bounded sequence in  $\mathbb{R}^k$ . Then each sequence  $(x_j^{(n)})$  is bounded in  $\mathbb{R}$ . By Theorem 11.5, we may replace  $(\mathbf{x}^{(n)})$  by a subsequence such that  $(x_1^{(n)})$  converges. By the same theorem, we may

replace  $(\mathbf{x}^{(n)})$  by a subsequence of the subsequence such that  $(x_2^{(n)})$  converges. Of course,  $(x_1^{(n)})$  still converges by Theorem 11.2. Repeating this argument  $k$  times, we obtain a sequence  $(\mathbf{x}^{(n)})$  so that each sequence  $(x_j^{(n)})$  converges,  $j = 1, 2, \dots, k$ . This sequence represents a subsequence of the original sequence, and it converges in  $\mathbb{R}^k$  by Lemma 13.3. ■

### 13.6 Definition.

Let  $(S, d)$  be a metric space. Let  $E$  be a subset of  $S$ . An element  $s_0 \in E$  is *interior to  $E$*  if for some  $r > 0$  we have

$$\{s \in S : d(s, s_0) < r\} \subseteq E.$$

We write  $E^\circ$  for the set of points in  $E$  that are interior to  $E$ . The set  $E$  is *open in  $S$*  if every point in  $E$  is interior to  $E$ , i.e., if  $E = E^\circ$ .

### 13.7 Discussion.

One can show [Exercise 13.4]

- (i)  $S$  is open in  $S$  [trivial].
- (ii) The empty set  $\emptyset$  is open in  $S$  [trivial].
- (iii) The union of *any* collection of open sets is open.
- (iv) The intersection of *finitely many* open sets is again an open set.

Our study of  $\mathbb{R}^k$  and the exercises suggest that metric spaces are fairly general and useful objects. When one is interested in convergence of certain objects [such as points or functions], there is often a metric that assists in the study of the convergence. But sometimes no metric will work and yet there is still some sort of convergence notion. Frequently the appropriate vehicle is what is called a *topology*. This is a set  $S$  for which certain subsets are decreed to be *open sets*. In general, all that is required is that the family of open sets satisfies (i)–(iv) above. In particular, the open sets defined by a metric form a topology. We will not pursue this abstract theory. However, because of this abstract theory, concepts that can be defined in terms of open sets [see Definitions 13.8, 13.11 and 22.1] are called *topological*, hence the title of this section.

**13.8 Definition.**

Let  $(S, d)$  be a metric space. A subset  $E$  of  $S$  is *closed* if its complement  $S \setminus E$  is an open set. In other words,  $E$  is closed if  $E = S \setminus U$  where  $U$  is an open set.

Because of (iii) in Discussion 13.7, the intersection of *any* collection of closed sets is closed [Exercise 13.5]. The *closure*  $E^-$  of a set  $E$  is the intersection of all closed sets containing  $E$ . The *boundary* of  $E$  is the set  $E^- \setminus E^\circ$ ; points in this set are called *boundary points* of  $E$ .

To get a feel for these notions, we state some easy facts and leave the proofs as exercises.

**13.9 Proposition.**

Let  $E$  be a subset of a metric space  $(S, d)$ .

- (a) The set  $E$  is closed if and only if  $E = E^-$ .
- (b) The set  $E$  is closed if and only if it contains the limit of every convergent sequence of points in  $E$ .
- (c) An element is in  $E^-$  if and only if it is the limit of some sequence of points in  $E$ .
- (d) A point is in the boundary of  $E$  if and only if it belongs to the closure of both  $E$  and its complement.

**Example 3**

In  $\mathbb{R}$ , open intervals  $(a, b)$  are open sets. Closed intervals  $[a, b]$  are closed sets. The interior of  $[a, b]$  is  $(a, b)$ . The boundary of both  $(a, b)$  and  $[a, b]$  is the two-element set  $\{a, b\}$ .

Every open set in  $\mathbb{R}$  is the union of a disjoint sequence of open intervals [Exercise 13.7]. A closed set in  $\mathbb{R}$  need not be the union of a disjoint sequence of closed intervals and points; such a set appears in Example 5.

**Example 4**

In  $\mathbb{R}^k$ , open balls  $\{\mathbf{x} : d(\mathbf{x}, \mathbf{x}_0) < r\}$  are open sets, and closed balls  $\{\mathbf{x} : d(\mathbf{x}, \mathbf{x}_0) \leq r\}$  are closed sets. The boundary of each of these sets is  $\{\mathbf{x} : d(\mathbf{x}, \mathbf{x}_0) = r\}$ . In the plane  $\mathbb{R}^2$ , the sets

$$\{(x_1, x_2) : x_1 > 0\} \quad \text{and} \quad \{(x_1, x_2) : x_1 > 0 \text{ and } x_2 > 0\}$$

are open. If  $>$  is replaced by  $\geq$ , we obtain closed sets. Many sets are neither open nor closed, for example

$$\{(x_1, x_2) : x_1 > 0 \text{ and } x_2 \geq 0\}.$$

### 13.10 Theorem.

Let  $(F_n)$  be a decreasing sequence [i.e.,  $F_1 \supseteq F_2 \supseteq \cdots$ ] of closed bounded nonempty sets in  $\mathbb{R}^k$ . Then  $F = \bigcap_{n=1}^{\infty} F_n$  is also closed, bounded and nonempty.

#### Proof

Clearly  $F$  is closed and bounded. It is the nonemptiness that needs proving! For each  $n$ , select an element  $\mathbf{x}_n$  in  $F_n$ . By the Bolzano-Weierstrass theorem 13.5, a subsequence  $(\mathbf{x}_{n_m})_{m=1}^{\infty}$  of  $(\mathbf{x}_n)$  converges to some element  $\mathbf{x}_0$  in  $\mathbb{R}^k$ . To show  $\mathbf{x}_0 \in F$ , it suffices to show  $\mathbf{x}_0 \in F_{n_0}$  with  $n_0$  fixed. If  $m \geq n_0$ , then  $n_m \geq n_0$ , so  $\mathbf{x}_{n_m} \in F_{n_m} \subseteq F_{n_0}$ . Hence the sequence  $(\mathbf{x}_{n_m})_{m=n_0}^{\infty}$  consists of points in  $F_{n_0}$  and converges to  $\mathbf{x}_0$ . Thus  $\mathbf{x}_0$  belongs to  $F_{n_0}$  by (b) of Proposition 13.9. ■

### Example 5

Here is a famous nonempty closed set in  $\mathbb{R}$  called the *Cantor set*. Pictorially,  $F = \bigcap_{n=1}^{\infty} F_n$  where  $F_n$  are sketched in Figure 13.1. The Cantor set has some remarkable properties. The sum of the lengths of the intervals comprising  $F_n$  is  $(\frac{2}{3})^{n-1}$  and this tends to 0 as  $n \rightarrow \infty$ . Yet the intersection  $F$  is so large that it cannot be written as a sequence; in set-theoretic terms it is “uncountable.” The interior of

FIGURE 13.1

$F$  is the empty set, so  $F$  equals its boundary. For more details, see [36], 2.44, or [23], 6.62.

### 13.11 Definition.

Let  $(S, d)$  be a metric space. A family  $\mathcal{U}$  of open sets is said to be an *open cover* for a set  $E$  if each point of  $E$  belongs to at least one set in  $\mathcal{U}$ , i.e.,

$$E \subseteq \bigcup \{U : U \in \mathcal{U}\}.$$

A *subcover* of  $\mathcal{U}$  is any subfamily of  $\mathcal{U}$  that also covers  $E$ . A cover or subcover is *finite* if it contains only finitely many sets; the sets themselves may be infinite.

A set  $E$  is *compact* if every open cover of  $E$  has a finite subcover of  $E$ .

This rather abstract definition is very important in advanced analysis; see, for example, [22]. In  $\mathbb{R}^k$ , compact sets are nicely characterized, as follows.

### 13.12 Heine-Borel Theorem.

*A subset  $E$  of  $\mathbb{R}^k$  is compact if and only if it is closed and bounded.*

#### Proof

Suppose that  $E$  is compact. For each  $m \in \mathbb{N}$ , let  $U_m$  consist of all  $\mathbf{x}$  in  $\mathbb{R}^k$  such that

$$\max\{|\mathbf{x}_j| : j = 1, 2, \dots, k\} < m.$$

The family  $\mathcal{U} = \{U_m : m \in \mathbb{N}\}$  is an open cover of  $E$  [it covers  $\mathbb{R}^k$ !], so a finite subfamily of  $\mathcal{U}$  covers  $E$ . If  $U_{m_0}$  is the largest member of the subfamily, then  $E \subseteq U_{m_0}$ . It follows that  $E$  is bounded. To show that  $E$  is closed, consider any point  $\mathbf{x}_0$  in  $\mathbb{R}^k \setminus E$ . For  $m \in \mathbb{N}$ , let

$$V_m = \left\{ \mathbf{x} \in \mathbb{R}^k : d(\mathbf{x}, \mathbf{x}_0) > \frac{1}{m} \right\}.$$

Then each  $V_m$  is open in  $\mathbb{R}^k$  and  $\mathcal{V} = \{V_m : m \in \mathbb{N}\}$  covers  $E$  since  $\bigcup_{m=1}^{\infty} V_m = \mathbb{R}^k \setminus \{\mathbf{x}_0\}$ . Since  $E$  can be covered by finitely many  $V_m$ , for some  $m_0$  we have

$$E \subseteq \left\{ \mathbf{x} \in \mathbb{R}^k : d(\mathbf{x}, \mathbf{x}_0) > \frac{1}{m_0} \right\}.$$

Thus  $\{\mathbf{x} \in \mathbb{R}^k : d(\mathbf{x}, \mathbf{x}_0) < \frac{1}{m_0}\} \subseteq \mathbb{R}^k \setminus E$ , so that  $\mathbf{x}_0$  is interior to  $\mathbb{R}^k \setminus E$ . Since  $\mathbf{x}_0$  in  $\mathbb{R}^k \setminus E$  was arbitrary,  $\mathbb{R}^k \setminus E$  is an open set. Hence  $E$  is a closed set.

Now suppose that  $E$  is closed and bounded. Since  $E$  is bounded,  $E$  is a subset of some set  $F$  having the form

$$F = \{\mathbf{x} \in \mathbb{R}^k : |x_j| \leq m \text{ for } j = 1, 2, \dots, k\}.$$

As noted in Exercise 13.12, it suffices to prove that  $F$  is compact. We do so in the next proposition after some preparation. ■

The set  $F$  in the last proof is a  $k$ -cell because it has the following form. There exist closed intervals  $[a_1, b_1], [a_2, b_2], \dots, [a_k, b_k]$  so that

$$F = \{\mathbf{x} \in \mathbb{R}^k : x_j \in [a_j, b_j] \text{ for } j = 1, 2, \dots, k\}.$$

The *diameter* of  $F$  is

$$\delta = \left[ \sum_{j=1}^k (b_j - a_j)^2 \right]^{1/2};$$

that is,  $\delta = \sup\{d(\mathbf{x}, \mathbf{y}) : \mathbf{x}, \mathbf{y} \in F\}$ . Using midpoints  $c_j = \frac{1}{2}(a_j + b_j)$  of  $[a_j, b_j]$ , we see that  $F$  is a union of  $2^k$   $k$ -cells each having diameter  $\frac{\delta}{2}$ . If this remark is not clear, consider first the cases  $k = 2$  and  $k = 3$ .

### 13.13 Proposition.

*Every  $k$ -cell  $F$  in  $\mathbb{R}^k$  is compact.*

#### Proof

Assume  $F$  is not compact. Then there exists an open cover  $\mathcal{U}$  of  $F$ , no finite subfamily of which covers  $F$ . Let  $\delta$  denote the diameter of  $F$ . As noted above,  $F$  is a union of  $2^k$   $k$ -cells having diameter  $\frac{\delta}{2}$ . At least one of these  $2^k$   $k$ -cells, which we denote by  $F_1$ , cannot be covered by finitely many sets from  $\mathcal{U}$ . Likewise,  $F_1$  contains a  $k$ -cell  $F_2$  of diameter  $\frac{\delta}{4}$  which cannot be covered by finitely many sets from  $\mathcal{U}$ . Continuing in this fashion, we obtain a sequence  $(F_n)$  of  $k$ -cells such that

$$F_1 \supseteq F_2 \supseteq F_3 \supseteq \dots; \tag{1}$$

$$F_n \text{ has diameter } \delta \cdot 2^{-n}; \tag{2}$$

$$F_n \text{ cannot be covered by finitely many sets from } \mathcal{U}. \tag{3}$$

By Theorem 13.10, the intersection  $\cap_{n=1}^{\infty} F_n$  contains a point  $\mathbf{x}_0$ . This point belongs to some set  $U_0$  in  $\mathcal{U}$ . Since  $U_0$  is open, there exists  $r > 0$  so that

$$\{\mathbf{x} \in \mathbb{R}^k : d(\mathbf{x}, \mathbf{x}_0) < r\} \subseteq U_0.$$

It follows that  $F_n \subseteq U_0$  provided  $\delta \cdot 2^{-n} < r$ , but this contradicts (3) in a dramatic way. ■

Since  $\mathbb{R} = \mathbb{R}^1$ , the preceding results apply to  $\mathbb{R}$ .

## Exercises

**13.1.** For points  $\mathbf{x}, \mathbf{y}$  in  $\mathbb{R}^k$ , let

$$d_1(\mathbf{x}, \mathbf{y}) = \max\{|x_j - y_j| : j = 1, 2, \dots, k\}$$

and

$$d_2(\mathbf{x}, \mathbf{y}) = \sum_{j=1}^k |x_j - y_j|.$$

(a) Show that  $d_1$  and  $d_2$  are metrics for  $\mathbb{R}^k$ .

(b) Show that  $d_1$  and  $d_2$  are complete.

**13.2.** (a) Prove (1) in Lemma 13.3.

(b) Prove the first assertion in Lemma 13.3.

**13.3.** Let  $B$  be the set of all bounded sequences  $\mathbf{x} = (x_1, x_2, \dots)$ , and define  $d(\mathbf{x}, \mathbf{y}) = \sup\{|x_j - y_j| : j = 1, 2, \dots\}$ .

(a) Show that  $d$  is a metric for  $B$ .

(b) Does  $d^*(\mathbf{x}, \mathbf{y}) = \sum_{j=1}^{\infty} |x_j - y_j|$  define a metric for  $B$ ?

**13.4.** Prove (iii) and (iv) in Discussion 13.7.

**13.5.** (a) Verify one of DeMorgan's Laws for sets:

$$\bigcap \{S \setminus U : U \in \mathcal{U}\} = S \setminus \bigcup \{U : U \in \mathcal{U}\}.$$

(b) Show that the intersection of any collection of closed sets is a closed set.

**13.6.** Prove Proposition 13.9.

- 13.7.** Show that every open set in  $\mathbb{R}$  is the disjoint union of a finite or infinite sequence of open intervals.
- 13.8.** (a) Verify the assertions in Example 3.  
 (b) Verify the assertions in Example 4.
- 13.9.** Find the closures of the following sets:  
 (a)  $\{\frac{1}{n} : n \in \mathbb{N}\}$ ,  
 (b)  $\mathbb{Q}$ , the set of rational numbers,  
 (c)  $\{r \in \mathbb{Q} : r^2 < 2\}$ .
- 13.10.** Show that the interior of each of the following sets is the empty set.  
 (a)  $\{\frac{1}{n} : n \in \mathbb{N}\}$ ,  
 (b)  $\mathbb{Q}$ , the set of rational numbers,  
 (c) the Cantor set in Example 5.
- 13.11.** Let  $E$  be a subset of  $\mathbb{R}^k$ . Show that  $E$  is compact if and only if every sequence in  $E$  has a subsequence that converges to a point in  $E$ .
- 13.12.** Let  $(S, d)$  be any metric space.  
 (a) Show that if  $E$  is a closed subset of a compact set  $F$ , then  $E$  is also compact.  
 (b) Show that the finite union of compact sets in  $S$  is compact.
- 13.13.** Let  $E$  be a compact nonempty subset of  $\mathbb{R}$ . Show that  $\sup E$  and  $\inf E$  belong to  $E$ .
- 13.14.** Let  $E$  be a compact nonempty subset of  $\mathbb{R}^k$ , and let  $\delta = \sup\{d(\mathbf{x}, \mathbf{y}) : \mathbf{x}, \mathbf{y} \in E\}$ . Show that  $E$  contains points  $\mathbf{x}_0, \mathbf{y}_0$  such that  $d(\mathbf{x}_0, \mathbf{y}_0) = \delta$ .
- 13.15.** Let  $(B, d)$  be as in Exercise 13.3, and let  $F$  consist of all  $\mathbf{x} \in B$  such that  $\sup\{|\mathbf{x}_j| : j = 1, 2, \dots\} \leq 1$ .  
 (a) Show that  $F$  is closed and bounded. [A set  $F$  in a metric space  $(S, d)$  is *bounded* if there exist  $s_0 \in S$  and  $r > 0$  such that  $F \subseteq \{s \in S : d(s, s_0) \leq r\}$ .]  
 (b) Show that  $F$  is not compact. *Hint:* For each  $\mathbf{x}$  in  $F$ , let  $U(\mathbf{x}) = \{\mathbf{y} \in B : d(\mathbf{y}, \mathbf{x}) < 1\}$ , and consider the cover  $\mathcal{U}$  of  $F$  consisting of all  $U(\mathbf{x})$ . For each  $n \in \mathbb{N}$ , let  $\mathbf{x}^{(n)}$  be defined so that  $x_n^{(n)} = -1$  and  $x_j^{(n)} = 1$  for  $j \neq n$ . Show that distinct  $\mathbf{x}^{(n)}$  cannot belong to the same member of  $\mathcal{U}$ .

## §14 Series

Our thorough treatment of sequences now allows us to quickly obtain the basic properties of infinite series.

### 14.1 Summation Notation.

The notation  $\sum_{k=m}^n a_k$  is shorthand for the sum  $a_m + a_{m+1} + \cdots + a_n$ . The symbol “ $\sum$ ” instructs us to sum and the decorations “ $k = m$ ” and “ $n$ ” tell us to sum the summands obtained by successively substituting  $m, m+1, \dots, n$  for  $k$ . For example,  $\sum_{k=2}^5 \frac{1}{k^2+k}$  is shorthand for

$$\frac{1}{2^2+2} + \frac{1}{3^2+3} + \frac{1}{4^2+4} + \frac{1}{5^2+5} = \frac{1}{6} + \frac{1}{12} + \frac{1}{20} + \frac{1}{30}$$

and  $\sum_{k=0}^n 2^{-k}$  is shorthand for  $1 + 1/2 + 1/4 + \cdots + 1/2^n$ .

The symbol  $\sum_{n=m}^{\infty} a_n$  is shorthand for  $a_m + a_{m+1} + a_{m+2} + \cdots$ , although we have not yet assigned meaning to such an infinite sum. We now do so.

### 14.2 Infinite Series.

To assign meaning to  $\sum_{n=m}^{\infty} a_n$ , we consider the sequences  $(s_n)_{n=m}^{\infty}$  of *partial sums*:

$$s_n = a_m + a_{m+1} + \cdots + a_n = \sum_{k=m}^n a_k.$$

The infinite series  $\sum_{n=m}^{\infty} a_n$  is said to *converge* provided the sequence  $(s_n)$  of partial sums converges to a real number  $S$ , in which case we define  $\sum_{n=m}^{\infty} a_n = S$ . Thus

$$\sum_{n=m}^{\infty} a_n = S \quad \text{means} \quad \lim s_n = S \quad \text{or} \quad \lim_{n \rightarrow \infty} \left( \sum_{k=m}^n a_k \right) = S.$$

A series that does not converge is said to *diverge*. We say that  $\sum_{n=m}^{\infty} a_n$  *diverges to  $+\infty$*  and we write  $\sum_{n=m}^{\infty} a_n = +\infty$  provided  $\lim s_n = +\infty$ ; a similar remark applies to  $-\infty$ . The symbol  $\sum_{n=m}^{\infty} a_n$  has no meaning unless the series converges or diverges to  $+\infty$  or  $-\infty$ . Often we will be concerned with properties of infinite series but not their exact values or precisely where the summation begins, in which case we may write  $\sum a_n$  rather than  $\sum_{n=m}^{\infty} a_n$ .

If the terms  $a_n$  of an infinite series  $\sum a_n$  are all nonnegative, then the partial sums  $(s_n)$  form a nondecreasing sequence, so Theorems 10.2 and 10.4 show that  $\sum a_n$  either converges or diverges to  $+\infty$ . In particular,  $\sum |a_n|$  is meaningful for any sequence  $(a_n)$  whatever. The series  $\sum a_n$  is said to *converge absolutely* or to be *absolutely convergent* if  $\sum |a_n|$  converges. Absolutely convergent series are convergent, as we shall see in 14.7.

### Example 1

A series of the form  $\sum_{n=0}^{\infty} ar^n$  for constants  $a$  and  $r$  is called a *geometric series*. These are the easiest series to sum. For  $r \neq 1$ , the partial sums  $s_n$  are given by

$$\sum_{k=0}^n ar^k = a \frac{1 - r^{n+1}}{1 - r}. \quad (1)$$

This identity can be verified by mathematical induction or by multiplying both sides by  $1 - r$ , in which case the right hand side equals  $a - ar^{n+1}$  and the left side becomes

$$\begin{aligned} (1 - r) \sum_{k=0}^n ar^k &= \sum_{k=0}^n ar^k - \sum_{k=0}^n ar^{k+1} \\ &= a + ar + ar^2 + \cdots + ar^n \\ &\quad - (ar + ar^2 + \cdots + ar^n + ar^{n+1}) \\ &= a - ar^{n+1}. \end{aligned}$$

For  $|r| < 1$ , we have  $\lim_{n \rightarrow \infty} r^{n+1} = 0$  by Example 7(b) in §9, so from (1) we have  $\lim_{n \rightarrow \infty} s_n = \frac{a}{1-r}$ . This proves

$$\sum_{n=0}^{\infty} ar^n = \frac{a}{1-r} \quad \text{if } |r| < 1. \quad (2)$$

If  $a \neq 0$  and  $|r| \geq 1$ , then the sequence  $(ar^n)$  does not converge to 0, so the series  $\sum ar^n$  diverges by Corollary 14.5 below.

### Example 2

Formula (2) of Example 1 and the next result are very important and both should be used whenever possible, even though we will not prove (1) below until the next section. Consider a fixed positive

real number  $p$ . Then

$$\sum_{n=1}^{\infty} \frac{1}{n^p} \text{ converges if and only if } p > 1. \quad (1)$$

In particular, for  $p \leq 1$ , we can write  $\sum 1/n^p = +\infty$ . The exact value of the series for  $p > 1$  is not easy to determine. Here are some remarkable formulas that can be shown by techniques [Fourier series or complex variables, to name two possibilities] that will not be covered in this text.

$$\sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{\pi^2}{6} = 1.6449 \dots, \quad (2)$$

$$\sum_{n=1}^{\infty} \frac{1}{n^4} = \frac{\pi^4}{90} = 1.0823 \dots. \quad (3)$$

Similar formulas hold for  $\sum_{n=1}^{\infty} \frac{1}{n^p}$  when  $p$  is any even integer, but no such elegant formulas are known for  $p$  odd. In particular, no such formula is known for  $\sum_{n=1}^{\infty} \frac{1}{n^3}$  though of course this series converges and can be approximated as closely as desired.

It is worth emphasizing that it is often easier to prove that a limit exists or that a series converges than to determine its exact value. In the next section we will show without much difficulty that  $\sum \frac{1}{n^p}$  converges for all  $p > 1$ , but it is a lot harder to show that the sum is  $\frac{\pi^2}{6}$  when  $p = 2$  and no one knows exactly what the sum is for  $p = 3$ .

### 14.3 Definition.

We say that a series  $\sum a_n$  satisfies the *Cauchy criterion* if its sequence  $(s_n)$  of partial sums is a Cauchy sequence [see Definition 10.8]:

$$\begin{aligned} &\text{for each } \epsilon > 0 \text{ there exists a number } N \text{ such that} \\ &m, n > N \text{ implies } |s_n - s_m| < \epsilon. \end{aligned} \quad (1)$$

Nothing is lost in this definition if we impose the restriction  $n > m$ . Moreover, it is only a notational matter to work with  $m - 1$  where  $m \leq n$  instead of  $m$  where  $m < n$ . Therefore (1) is equivalent to

$$\begin{aligned} &\text{for each } \epsilon > 0 \text{ there exists a number } N \text{ such that} \\ &n \geq m > N \text{ implies } |s_n - s_{m-1}| < \epsilon. \end{aligned} \quad (2)$$

Since  $s_n - s_{m-1} = \sum_{k=m}^n a_k$ , condition (2) can be rewritten

$$\text{for each } \epsilon > 0 \text{ there exists a number } N \text{ such that} \\ n \geq m > N \text{ implies } \left| \sum_{k=m}^n a_k \right| < \epsilon. \quad (3)$$

We will usually use version (3) of the *Cauchy criterion*. Theorem 10.11 implies the following.

#### 14.4 Theorem.

*A series converges if and only if it satisfies the Cauchy criterion.*

#### 14.5 Corollary.

*If a series  $\sum a_n$  converges, then  $\lim a_n = 0$ .*

#### Proof

Since the series converges, (3) in Definition 14.3 holds. In particular, (3) in 14.3 holds for  $n = m$ ; i.e., for each  $\epsilon > 0$  there exists a number  $N$  such that  $n > N$  implies  $|a_n| < \epsilon$ . Thus  $\lim a_n = 0$ . ■

The converse of Corollary 14.5 does not hold as the example  $\sum 1/n = +\infty$  shows.

We next give several tests to assist us in determining whether a series converges. The first test is elementary but useful.

#### 14.6 Comparison Test.

*Let  $\sum a_n$  be a series where  $a_n \geq 0$  for all  $n$ .*

- (i) *If  $\sum a_n$  converges and  $|b_n| \leq a_n$  for all  $n$ , then  $\sum b_n$  converges.*
- (ii) *If  $\sum a_n = +\infty$  and  $b_n \geq a_n$  for all  $n$ , then  $\sum b_n = +\infty$ .*

#### Proof

- (i) For  $n \geq m$  we have

$$\left| \sum_{k=m}^n b_k \right| \leq \sum_{k=m}^n |b_k| \leq \sum_{k=m}^n a_k;$$

the first inequality follows from the triangle inequality [Exercise 3.6(b)]. Since  $\sum a_n$  converges, it satisfies the Cauchy criterion 14.3(3). It follows from (1) that  $\sum b_n$  also satisfies the Cauchy criterion, and hence  $\sum b_n$  converges.

- (ii) Let  $(s_n)$  and  $(t_n)$  be the sequences of partial sums for  $\sum a_n$  and  $\sum b_n$ , respectively. Since  $b_n \geq a_n$  for all  $n$ , we obviously have  $t_n \geq s_n$  for all  $n$ . Since  $\lim s_n = +\infty$ , we conclude that  $\lim t_n = +\infty$ , i.e.,  $\sum b_n = +\infty$ . ■

**14.7 Corollary.**

*Absolutely convergent series are convergent.*

**Proof**

Suppose that  $\sum b_n$  is absolutely convergent. This means that  $\sum a_n$  converges where  $a_n = |b_n|$  for all  $n$ . Then  $|b_n| \leq a_n$  trivially, so  $\sum b_n$  converges by 14.6(i). ■

We next state the Ratio Test which is popular because it is often easy to use. But it has defects: It isn't as general as the Root Test. An important result concerning the radius of convergence of a power series uses the Root Test. Finally, the Ratio Test is worthless if some of the  $a_n$ 's equal 0. To review  $\limsup$ 's and  $\liminf$ 's, see 10.6, 10.7, 11.7 and §12.

**14.8 Ratio Test.**

*A series  $\sum a_n$  of nonzero terms*

- (i) *converges absolutely if  $\limsup |a_{n+1}/a_n| < 1$ ,*
- (ii) *diverges if  $\liminf |a_{n+1}/a_n| > 1$ .*
- (iii) *Otherwise  $\liminf |a_{n+1}/a_n| \leq 1 \leq \limsup |a_{n+1}/a_n|$  and the test gives no information.*

We give the proof after the proof of the Root Test.

Remember that if  $\lim |a_{n+1}/a_n|$  exists, then it is equal to both  $\limsup |a_{n+1}/a_n|$  and  $\liminf |a_{n+1}/a_n|$  and hence the Ratio Test will give information unless, or course, the limit  $\lim |a_{n+1}/a_n|$  equals 1.

**14.9 Root Test.**

*Let  $\sum a_n$  be a series and let  $\alpha = \limsup |a_n|^{1/n}$ . The series  $\sum a_n$*

- (i) *converges absolutely if  $\alpha < 1$ ,*
- (ii) *diverges if  $\alpha > 1$ .*
- (iii) *Otherwise  $\alpha = 1$  and the test gives no information.*

**Proof**

- (i) Suppose  $\alpha < 1$ , and select  $\epsilon > 0$  so that  $\alpha + \epsilon < 1$ . Then by Definition 10.6 there is a natural number  $N$  such that

$$\alpha - \epsilon < \sup\{|a_n|^{1/n} : n > N\} < \alpha + \epsilon.$$

In particular, we have  $|a_n|^{1/n} < \alpha + \epsilon$  for  $n > N$ , so

$$|a_n| < (\alpha + \epsilon)^n \quad \text{for } n > N.$$

Since  $0 < \alpha + \epsilon < 1$ , the geometric series  $\sum_{n=N+1}^{\infty} (\alpha + \epsilon)^n$  converges, and the Comparison Test shows that the series  $\sum_{n=N+1}^{\infty} a_n$  also converges. Then clearly  $\sum a_n$  converges; see Exercise 14.9.

- (ii) If  $\alpha > 1$ , then by Corollary 11.4 a subsequence of  $|a_n|^{1/n}$  has limit  $\alpha > 1$ . It follows that  $|a_n| > 1$  for infinitely many choices of  $n$ . In particular, the sequence  $(a_n)$  cannot possibly converge to 0, so the series  $\sum a_n$  cannot converge by Corollary 14.5.
- (iii) For each of the series  $\sum \frac{1}{n}$  and  $\sum \frac{1}{n^2}$ ,  $\alpha$  turns out to equal 1 as can be seen by applying 9.7(c). Since  $\sum \frac{1}{n}$  diverges and  $\sum \frac{1}{n^2}$  converges, the equality  $\alpha = 1$  does not guarantee either convergence or divergence of the series. ■

**Proof of the Ratio Test**

Let  $\alpha = \limsup |a_n|^{1/n}$ . By Theorem 12.2 we have

$$\liminf \left| \frac{a_{n+1}}{a_n} \right| \leq \alpha \leq \limsup \left| \frac{a_{n+1}}{a_n} \right|. \quad (1)$$

If  $\limsup |a_{n+1}/a_n| < 1$ , then  $\alpha < 1$  and the series converges by the Root Test. If  $\liminf |a_{n+1}/a_n| > 1$ , then  $\alpha > 1$  and the series diverges by the Root Test. Assertion 14.8(iii) is verified by again examining the series  $\sum 1/n$  and  $\sum 1/n^2$ . ■

Inequality (1) in the proof of the Ratio Test shows that the Root Test is superior to the Ratio Test in the following sense: Whenever the Root Test gives no information [i.e.,  $\alpha = 1$ ] the Ratio Test will surely also give no information. On the other hand, Example 8 below gives a series for which the Ratio Test gives no information but which converges by the Root Test. Nevertheless, the tests usually fail together as the next remark shows.

**14.10 Remark.**

If the terms  $a_n$  are nonzero and if  $\lim |a_{n+1}/a_n| = 1$ , then  $\alpha = \limsup |a_n|^{1/n} = 1$  by Corollary 12.3, so neither the Ratio Test nor the Root Test gives information concerning the convergence of  $\sum a_n$ .

We have three tests for convergence of a series [Comparison, Ratio, Root] and we will obtain two more in the next section. There is no clearcut strategy advising us which test to try first. However, if the form of a given series  $\sum a_n$  does not suggest a particular strategy, and if the ratios  $a_{n+1}/a_n$  are easy to calculate, one may as well try the Ratio Test first.

**Example 3**

Consider the series

$$\sum_{n=2}^{\infty} \left(-\frac{1}{3}\right)^n = \frac{1}{9} - \frac{1}{27} + \frac{1}{81} - \frac{1}{243} + \cdots \quad (1)$$

This is a geometric series and has the form  $\sum_{n=0}^{\infty} ar^n$  if we write it as  $(1/9) \sum_{n=0}^{\infty} (-1/3)^n$ . Here  $a = 1/9$  and  $r = -1/3$ , so by (2) of Example 1 the sum is  $(1/9)/[1 - (-1/3)] = 1/12$ .

The series (1) can also be shown to converge by the Comparison Test, since  $\sum 1/3^n$  converges by the Ratio Test or by the Root Test. In fact, if  $a_n = (-1/3)^n$ , then  $\lim |a_{n+1}/a_n| = \limsup |a_n|^{1/n} = 1/3$ . Of course, none of these tests will give us the exact value of the series (1).

**Example 4**

Consider the series

$$\sum \frac{n}{n^2 + 3}. \quad (1)$$

If  $a_n = \frac{n}{n^2 + 3}$ , then

$$\frac{a_{n+1}}{a_n} = \frac{n+1}{(n+1)^2 + 3} \cdot \frac{n^2 + 3}{n} = \frac{n+1}{n} \cdot \frac{n^2 + 3}{n^2 + 2n + 4},$$

so  $\lim |a_{n+1}/a_n| = 1$ . As noted in 14.10, neither the Ratio Test nor the Root Test gives any information in this case. Before trying the Comparison Test we need to decide whether we *believe* the series

converges or not. Since  $a_n$  is approximately  $1/n$  for large  $n$  and since  $\sum(1/n)$  diverges, we expect the series (1) to diverge. Now

$$\frac{n}{n^2 + 3} \geq \frac{n}{n^2 + 3n^2} = \frac{n}{4n^2} = \frac{1}{4n}.$$

Since  $\sum(1/n)$  diverges,  $\sum(1/4n)$  also diverges [its partial sums are  $s_n/4$  where  $s_n = \sum_{k=1}^n (1/k)$ ], so (1) diverges by the Comparison Test.

### Example 5

Consider the series

$$\sum \frac{1}{n^2 + 1}. \quad (1)$$

As the reader should check, neither the Ratio Test nor the Root Test gives any information. The  $n$ th term is approximately  $\frac{1}{n^2}$  and in fact  $\frac{1}{n^2+1} \leq \frac{1}{n^2}$ . Since  $\sum \frac{1}{n^2}$  converges, the series (1) converges by the Comparison Test.

### Example 6

Consider the series

$$\sum \frac{n}{3^n}. \quad (1)$$

If  $a_n = n/3^n$ , then  $a_{n+1}/a_n = (n+1)/(3n)$ , so  $\lim |a_{n+1}/a_n| = 1/3$ . Hence the series (1) converges by the Ratio Test. In this case, applying the Root Test is not much more difficult provided we recall  $\lim n^{1/n} = 1$ . It is also possible to show that (1) converges by comparing it with a suitable geometric series.

### Example 7

Consider the series

$$\sum a_n \text{ where } a_n = \left[ \frac{2}{(-1)^n - 3} \right]^n. \quad (1)$$

The form of  $a_n$  suggests the Root Test. Since  $|a_n|^{1/n} = 1$  for even  $n$  and  $|a_n|^{1/n} = 1/2$  for odd  $n$ , we have  $\alpha = \limsup |a_n|^{1/n} = 1$ . So the Root Test gives no information and the Ratio Test cannot help either. On the other hand, if we had been alert, we would have observed that  $a_n = 1$  for even  $n$ , so  $(a_n)$  cannot converge to 0. Therefore the series (1) diverges by Corollary 14.5.

**Example 8**

Consider the series

$$\sum_{n=0}^{\infty} 2^{(-1)^n - n} = 2 + \frac{1}{4} + \frac{1}{2} + \frac{1}{16} + \frac{1}{8} + \frac{1}{64} + \cdots. \quad (1)$$

Let  $a_n = 2^{(-1)^n - n}$ . Since  $a_n \leq \frac{1}{2^{n-1}}$  for all  $n$ , we can quickly conclude that the series converges by the Comparison Test. But our real interest in this series is that it illustrates the difference between the Ratio Test and the Root Test. Since  $a_{n+1}/a_n = 1/8$  for even  $n$  and  $a_{n+1}/a_n = 2$  for odd  $n$ , we have

$$\frac{1}{8} = \liminf \left| \frac{a_{n+1}}{a_n} \right| < 1 < \limsup \left| \frac{a_{n+1}}{a_n} \right| = 2.$$

Hence the Ratio Test gives no information.

Note that  $(a_n)^{1/n} = 2^{\frac{1}{n}-1}$  for even  $n$  and  $(a_n)^{1/n} = 2^{-\frac{1}{n}-1}$  for odd  $n$ . Since  $\lim 2^{\frac{1}{n}} = \lim 2^{-\frac{1}{n}} = 1$  by Example 7(d) in §9, we conclude that  $\lim(a_n)^{1/n} = \frac{1}{2}$ . Therefore  $\alpha = \limsup(a_n)^{1/n} = \frac{1}{2} < 1$  and the series (1) converges by the Root Test.

**Example 9**

Consider the series

$$\sum \frac{(-1)^n}{\sqrt{n}}. \quad (1)$$

Since  $\lim \sqrt{n/(n+1)} = 1$ , neither the Ratio Test nor the Root Test gives any information. Since  $\sum \frac{1}{\sqrt{n}}$  diverges, we will not be able to use the Comparison Test 14.6(i) to show that (1) converges. Since the terms of the series (1) are not all nonnegative, we will not be able to use the Comparison Test 14.6(ii) to show that (1) diverges. It turns out that this series converges by the Alternating Series Test 15.3, which we have deferred to the next section.

**Exercises**

**14.1.** Determine which of the following series converge. Justify your answers.

(a)  $\sum \frac{n^4}{2^n}$

(b)  $\sum \frac{2^n}{n!}$

$$\begin{array}{ll} \text{(c)} \sum \frac{n^2}{3^n} & \text{(d)} \sum \frac{n!}{n^4+3} \\ \text{(e)} \sum \frac{\cos^2 n}{n^2} & \text{(f)} \sum_{n=2}^{\infty} \frac{1}{\log n} \end{array}$$

**14.2.** Repeat Exercise 14.1 for the following.

$$\begin{array}{ll} \text{(a)} \sum \frac{n-1}{n^2} & \text{(b)} \sum (-1)^n \\ \text{(c)} \sum \frac{3n}{n^3} & \text{(d)} \sum \frac{n^3}{3^n} \\ \text{(e)} \sum \frac{n^2}{n!} & \text{(f)} \sum \frac{1}{n^n} \\ \text{(g)} \sum \frac{n}{2^n} & \end{array}$$

**14.3.** Repeat Exercise 14.1 for the following.

$$\begin{array}{ll} \text{(a)} \sum \frac{1}{\sqrt{n!}} & \text{(b)} \sum \frac{2+\cos n}{3^n} \\ \text{(c)} \sum \frac{1}{2^n+n} & \text{(d)} \sum \left(\frac{1}{2}\right)^n \left(50 + \frac{2}{n}\right) \\ \text{(e)} \sum \sin\left(\frac{n\pi}{9}\right) & \text{(f)} \sum \frac{(100)^n}{n!} \end{array}$$

**14.4.** Repeat Exercise 14.1 for the following.

$$\begin{array}{ll} \text{(a)} \sum_{n=2}^{\infty} \frac{1}{[n+(-1)^n]^2} & \text{(b)} \sum [\sqrt{n+1} - \sqrt{n}] \\ \text{(c)} \sum \frac{n!}{n^n} & \end{array}$$

**14.5.** Suppose that  $\sum a_n = A$  and  $\sum b_n = B$  where  $A$  and  $B$  are real numbers. Use limit theorems from §9 to quickly prove the following.

- (a)  $\sum (a_n + b_n) = A + B$ .
- (b)  $\sum ka_n = kA$  for  $k \in \mathbb{R}$ .
- (c) Is  $\sum a_n b_n = AB$  a reasonable conjecture? Discuss.

**14.6.** (a) Prove that if  $\sum |a_n|$  converges and  $(b_n)$  is a bounded sequence, then  $\sum a_n b_n$  converges. *Hint:* Use Theorem 14.4.

(b) Observe that Corollary 14.7 is a special case of part (a).

**14.7.** Prove that if  $\sum a_n$  is a convergent series of nonnegative numbers and  $p > 1$ , then  $\sum a_n^p$  converges.

**14.8.** Show that if  $\sum a_n$  and  $\sum b_n$  are convergent series of nonnegative numbers, then  $\sum \sqrt{a_n b_n}$  converges. *Hint:* Show that  $\sqrt{a_n b_n} \leq a_n + b_n$  for all  $n$ .

**14.9.** The convergence of a series does not depend on any finite number of the terms, though of course the value of the limit does. More precisely, consider series  $\sum a_n$  and  $\sum b_n$  and suppose that the set  $\{n \in \mathbb{N} : a_n \neq b_n\}$  is finite. Then the series both converge or else they both diverge. Prove this. *Hint:* This is almost obvious from Theorem 14.4.

- 14.10.** Find a series  $\sum a_n$  which diverges by the Root Test but for which the Ratio Test gives no information. Compare Example 8.
- 14.11.** Let  $(a_n)$  be a sequence of nonzero real numbers such that the sequence  $(\frac{a_{n+1}}{a_n})$  of ratios is a constant sequence. Show that  $\sum a_n$  is a geometric series.
- 14.12.** Let  $(a_n)_{n \in \mathbb{N}}$  be a sequence such that  $\liminf |a_n| = 0$ . Prove that there is a subsequence  $(a_{n_k})_{k \in \mathbb{N}}$  such that  $\sum_{k=1}^{\infty} a_{n_k}$  converges.
- 14.13.** We have seen that it is often a lot harder to find the value of an infinite sum than to show that it exists. Here are some sums that can be handled.
- (a) Calculate  $\sum_{n=1}^{\infty} (\frac{2}{3})^n$  and  $\sum_{n=1}^{\infty} (-\frac{2}{3})^n$ .
- (b) Prove  $\sum_{n=1}^{\infty} \frac{1}{n(n+1)} = 1$ . *Hint:* Note that  $\sum_{k=1}^n \frac{1}{k(k+1)} = \sum_{k=1}^n [\frac{1}{k} - \frac{1}{k+1}]$ .
- (c) Prove that  $\sum_{n=1}^{\infty} \frac{n-1}{2^{n+1}} = \frac{1}{2}$ . *Hint:* Note that  $\frac{k-1}{2^{k+1}} = \frac{k}{2^k} - \frac{k+1}{2^{k+1}}$ .
- (d) Use (c) to calculate  $\sum_{n=1}^{\infty} \frac{n}{2^n}$ .
- 14.14.** Prove that  $\sum_{n=2}^{\infty} \frac{1}{n}$  diverges by comparing with the series  $\sum_{n=2}^{\infty} a_n$  where  $(a_n)$  is the sequence

$$\left( \frac{1}{2}, \frac{1}{4}, \frac{1}{4}, \frac{1}{8}, \frac{1}{8}, \frac{1}{8}, \frac{1}{8}, \frac{1}{16}, \frac{1}{16}, \frac{1}{16}, \frac{1}{16}, \frac{1}{16}, \frac{1}{16}, \frac{1}{16}, \frac{1}{16}, \frac{1}{32}, \frac{1}{32}, \dots \right).$$

## §15 Alternating Series and Integral Tests

Sometimes one can check convergence or divergence of series by comparing the partial sums with familiar integrals. We illustrate.

### Example 1

We show that  $\sum \frac{1}{n} = +\infty$ .

Consider the picture of the function  $f(x) = \frac{1}{x}$  in Figure 15.1. For  $n \geq 1$  it is evident that

$$\sum_{k=1}^n \frac{1}{k} = \text{Sum of the areas of the first } n \text{ rectangles in Figure 15.1}$$

**FIGURE 15.1**

$$\begin{aligned} &\geq \text{Area under the curve } \frac{1}{x} \text{ between } 1 \text{ and } n+1 \\ &= \int_1^{n+1} \frac{1}{x} dx = \log(n+1). \end{aligned}$$

Since  $\lim_{n \rightarrow \infty} \log(n+1) = +\infty$ , we conclude that  $\sum_{n=1}^{\infty} \frac{1}{n} = +\infty$ .

The series  $\sum \frac{1}{n}$  diverges very slowly. In Example 7 of §16, we observe that  $\sum_{n=1}^N \frac{1}{n}$  is approximately  $\log_e N + 0.5772$ . Thus for  $N = 1,000$  the sum is approximately 7.485, and for  $N = 1,000,000$  the sum is approximately 14.393.

Another proof that  $\sum \frac{1}{n}$  diverges was indicated in Exercise 14.14. However, an integral test is useful to establish the next result.

**Example 2**

We show that  $\sum \frac{1}{n^2}$  converges.

Consider the graph of  $f(x) = \frac{1}{x^2}$  in Figure 15.2. Then we have

$$\begin{aligned} \sum_{k=1}^n \frac{1}{k^2} &= \text{Sum of the areas of the first } n \text{ rectangles} \\ &\leq 1 + \int_1^n \frac{1}{x^2} dx = 2 - \frac{1}{n} < 2 \end{aligned}$$

for all  $n \geq 1$ . Thus the partial sums form an increasing sequence that is bounded above by 2. Therefore  $\sum_{n=1}^{\infty} \frac{1}{n^2}$  converges and its sum is less than or equal to 2. Actually, we have already mentioned [without proof!] that the sum is  $\frac{\pi^2}{6} = 1.6449 \dots$ .

**FIGURE 15.2**

Note that in estimating  $\sum_{k=1}^n \frac{1}{k^2}$  we did not simply write  $\sum_{k=1}^n \frac{1}{k^2} \leq \int_0^n \frac{1}{x^2} dx$ , even though this is true, because this integral is infinite. We were after a *finite* upper bound for the partial sums.

The techniques just illustrated can be used to prove the following theorem.

**15.1 Theorem.**

$\sum \frac{1}{n^p}$  converges if and only if  $p > 1$ .

**Proof**

Supply your own picture and observe that if  $p > 1$ , then

$$\sum_{k=1}^n \frac{1}{k^p} \leq 1 + \int_1^n \frac{1}{x^p} dx = 1 + \frac{1}{p-1} \left(1 - \frac{1}{n^{p-1}}\right) < 1 + \frac{1}{p-1} = \frac{p}{p-1}.$$

Consequently  $\sum_{n=1}^{\infty} \frac{1}{n^p} \leq \frac{p}{p-1} < +\infty$ .

Suppose that  $0 < p \leq 1$ . Then  $\frac{1}{n} \leq \frac{1}{n^p}$  for all  $n$ . Since  $\sum \frac{1}{n}$  diverges, we see that  $\sum \frac{1}{n^p}$  diverges by the Comparison Test. ■

**15.2 Integral Tests.**

Here are the conditions under which an integral test is advisable:

- (a) The tests in §14 do not seem to apply.
- (b) The terms of the series  $\sum a_n$  are nonnegative.
- (c) There is a nice nonincreasing function  $f$  on  $[1, \infty)$  such that  $f(n) = a_n$  for all  $n$  [ $f$  is *nonincreasing* if  $x < y$  implies  $f(x) \geq f(y)$ ].

(d) The integral of  $f$  is easy to calculate or estimate.

If  $\lim_{n \rightarrow \infty} \int_1^n f(x) dx = +\infty$ , then the series will diverge just as in Example 1. If  $\lim_{n \rightarrow \infty} \int_1^n f(x) dx < +\infty$ , then the series will converge just as in Example 2. The interested reader may formulate and prove the general result [Exercise 15.8].

The following result is a bit tricky to prove, but it enables us to conclude that series like  $\sum \frac{(-1)^n}{\sqrt{n}}$  converge even though they do not converge absolutely. See Example 9 in §14.

### 15.3 Alternating Series Theorem.

If  $a_1 \geq a_2 \geq \cdots \geq a_n \geq \cdots \geq 0$  and  $\lim a_n = 0$ , then the alternating series  $\sum (-1)^n a_n$  converges.

The series  $\sum (-1)^n a_n$  is called an *alternating series* because the signs of the terms alternate between  $+$  and  $-$ .

#### Proof

It suffices to show that the series satisfies the Cauchy criterion 14.3(3). This will follow easily from

$$n \geq m > N \quad \text{implies} \quad \left| \sum_{k=m}^n (-1)^k a_k \right| \leq a_N, \quad (1)$$

since for each  $\epsilon > 0$  there exists  $N \in \mathbb{N}$  such that  $a_N < \epsilon$ .

To prove (1), we fix  $n \geq m$  and define

$$A = a_m - a_{m+1} + a_{m+2} - a_{m+3} + \cdots \pm a_n$$

so that

$$\sum_{k=m}^n (-1)^k a_k = (-1)^m A. \quad (2)$$

If  $n - m$  is odd, the last term of  $A$  is  $-a_n$ , so

$$A = [a_m - a_{m+1}] + [a_{m+2} - a_{m+3}] + \cdots + [a_{n-1} - a_n] \geq 0$$

and also

$$A = a_m - [a_{m+1} - a_{m+2}] - [a_{m+3} - a_{m+4}] - \cdots - [a_{n-2} - a_{n-1}] - a_n \leq a_m.$$

Remember that the numbers in brackets are nonnegative, since  $(a_n)$  is nonincreasing. If  $n - m$  is even, the last term of  $A$  is  $+a_n$ , so

$$A = [a_m - a_{m+1}] + [a_{m+2} - a_{m+3}] + \cdots + [a_{n-2} - a_{n-1}] + a_n \geq 0$$

and

$$A = a_m - [a_{m+1} - a_{m+2}] - [a_{m+3} - a_{m+4}] - \cdots - [a_{n-1} - a_n] \leq a_m.$$

In either case we have  $0 \leq A \leq a_m$ . Hence from (2) we see that

$$\left| \sum_{k=m}^n (-1)^k a_k \right| = A \leq a_m.$$

Assertion (1) now follows since  $n \geq m > N$  implies

$$\left| \sum_{k=m}^n (-1)^k a_k \right| \leq a_m \leq a_N. \quad \blacksquare$$

## Exercises

**15.1.** Determine which of the following series converge. Justify your answers.

(a)  $\sum \frac{(-1)^n}{n}$

(b)  $\sum \frac{(-1)^n n!}{2^n}$

**15.2.** Repeat Exercise 15.1 for the following.

(a)  $\sum [\sin(\frac{n\pi}{6})]^n$

(b)  $\sum [\sin(\frac{n\pi}{7})]^n$

**15.3.** Show that  $\sum_{n=2}^{\infty} \frac{1}{n(\log n)^p}$  converges if and only if  $p > 1$ .

**15.4.** Determine which of the following series converge. Justify your answers.

(a)  $\sum_{n=2}^{\infty} \frac{1}{\sqrt{n} \log n}$

(b)  $\sum_{n=2}^{\infty} \frac{\log n}{n}$

(c)  $\sum_{n=4}^{\infty} \frac{1}{n(\log n)(\log \log n)}$

(d)  $\sum_{n=2}^{\infty} \frac{\log n}{n^2}$

**15.5.** Why didn't we use the Comparison Test to prove Theorem 15.1 for  $p > 1$ ?

**15.6. (a)** Give an example of a divergent series  $\sum a_n$  for which  $\sum a_n^2$  converges.

**(b)** Observe that if  $\sum a_n$  is a convergent series of nonnegative terms, then  $\sum a_n^2$  also converges. See Exercise 14.7.

- (c) Give an example of a convergent series  $\sum a_n$  for which  $\sum a_n^2$  diverges.
- 15.7. (a) Prove that if  $(a_n)$  is a nonincreasing sequence of real numbers and if  $\sum a_n$  converges, then  $\lim na_n = 0$ . *Hint:* Consider  $|a_{N+1} + a_{N+2} + \cdots + a_n|$  for suitable  $N$ .
- (b) Use (a) to give another proof that  $\sum \frac{1}{n}$  diverges.
- 15.8. Formulate and prove a general integral test as advised in 15.2.

## §16 \* Decimal Expansions of Real Numbers

We begin by recalling the brief discussion of decimals in Discussion 10.3. There we considered a decimal expansion  $k.d_1d_2d_3\cdots$  where  $k$  is a nonnegative integer and each digit  $d_j$  belongs to  $\{0, 1, 2, 3, 4, 5, 6, 7, 8, 9\}$ . This expansion represents the real number

$$k + \sum_{j=1}^{\infty} \frac{d_j}{10^j} = k + \sum_{j=1}^{\infty} d_j \cdot 10^{-j}$$

which we also can write as

$$\lim_{n \rightarrow \infty} s_n \quad \text{where} \quad s_n = k + \sum_{j=1}^n d_j \cdot 10^{-j}.$$

Thus *every such decimal expansion represents a nonnegative real number*. We will prove the converse after we formalize the process of long division. The development here is based on some suggestions by Karl Stromberg.

### 16.1 Long Division.

Let's first consider positive integers  $a$  and  $b$  where  $a < b$ . We analyze the familiar long division process which gives a decimal expansion for  $\frac{a}{b}$ . Figure 16.1 shows the first few steps where  $a = 3$  and  $b = 7$ . If we name the digits  $d_1, d_2, d_3, \dots$  and the remainders  $r_1, r_2, r_3, \dots$ , then so far  $d_1 = 4$ ,  $d_2 = 2$  and  $r_1 = 2$ ,  $r_2 = 6$ . At the next step we divide 7 into  $60 = 10 \cdot r_2$  and obtain  $60 = 7 \cdot 8 + 4$ . The quotient 8 becomes the

**FIGURE 16.1**

third digit  $d_3$ , we place the product 56 under 60, subtract and obtain a new remainder  $4 = r_3$ . That is, we are calculating the remainder obtained by dividing 60 by 7. Next we multiply the remainder  $r_3 = 4$  by 10 and repeat the process. At each stage

$$\begin{aligned} d_n &\in \{0, 1, 2, 3, 4, 5, 6, 7, 8, 9\} \\ r_n &= 10 \cdot r_{n-1} - 7 \cdot d_n \\ 0 &\leq r_n < 7. \end{aligned}$$

These results hold for  $n = 1, 2, \dots$  if we set  $r_0 = 3$ . In general, we set  $r_0 = a$  and obtain

$$d_n \in \{0, 1, 2, 3, 4, 5, 6, 7, 8, 9\} \tag{1}$$

$$r_n = 10 \cdot r_{n-1} - b \cdot d_n \tag{2}$$

$$0 \leq r_n < b. \tag{3}$$

We next show that this construction is well defined in general and that the decimal expansion represents  $\frac{a}{b}$ . In what follows we *do not need to assume that  $a$  and  $b$  are integers*;  $a$  and  $b$  will represent positive numbers. The only noticeable change in our construction will be that the “remainders”  $r_n$  will not necessarily be integers. We also do not assume that  $a < b$ , so the first step will be a little different than in our example. The first step will provide us with the integer part of  $\frac{a}{b}$ .

Let  $\mathbb{Z}^+ = \mathbb{N} \cup \{0\}$ . By the Archimedean property 4.6, we have  $a \leq nb$  for some positive integer  $n$ . Hence  $\{m \in \mathbb{Z}^+ : mb \leq a\}$  is finite. This set is also nonempty, since it contains 0, so we can

define

$$k = \max\{m \in \mathbb{Z}^+ : mb \leq a\}.$$

Thus  $kb \leq a < (k+1)b$ . Let  $r_0 = a - kb$  and note that  $0 \leq r_0 < b$ . Next define

$$d_1 = \max\{d \in \mathbb{Z}^+ : db \leq 10 \cdot r_0\}$$

and

$$r_1 = 10 \cdot r_0 - d_1 b.$$

Note that  $d_1 \leq 9$ , because  $10 \cdot b \leq 10 \cdot r_0$  would imply  $b \leq r_0$ , a contradiction. Also note that  $d_1 b \leq 10 \cdot r_0 < (d_1 + 1)b$ , so  $0 \leq r_1 = 10 \cdot r_0 - d_1 b < b$ . Thus the following holds for  $n = 1$ :

$$d_n \in \{0, 1, 2, 3, 4, 5, 6, 7, 8, 9\} \quad (1)$$

$$r_n = 10 \cdot r_{n-1} - d_n b \quad (2)$$

$$0 \leq r_n < b. \quad (3)$$

Suppose that  $d_1, d_2, \dots, d_n \in \mathbb{Z}^+$  and  $r_0, r_1, \dots, r_n$  have been defined satisfying (1)–(3). Next define

$$d_{n+1} = \max\{d \in \mathbb{Z}^+ : db \leq 10 \cdot r_n\}$$

and

$$r_{n+1} = 10 \cdot r_n - d_{n+1} b.$$

Then  $d_{n+1} \leq 9$  since  $10 \cdot b \leq 10 \cdot r_n$  would imply  $b \leq r_n$ , violating (3). Hence (1) holds for  $n+1$  and (2) is obvious for  $n+1$  by our definition of  $r_{n+1}$ . Finally  $d_{n+1} b \leq 10 \cdot r_n < (d_{n+1} + 1)b$  implies  $0 \leq r_{n+1} < b$ , so (3) holds for  $n+1$ . The construction of the sequences  $(d_n)$  and  $(r_n)$  satisfying (1)–(3) is completed by an appeal to the principle of induction.

To see that the decimal expansion  $k.d_1 d_2 d_3 \dots$  represents  $\frac{a}{b}$ , we observe that (2) implies

$$r_n \cdot 10^{-n} = r_{n-1} \cdot 10^{-n+1} - d_n \cdot 10^{-n} \cdot b$$

for  $n \geq 1$ . Transposing and changing  $n$  to  $j$ , we obtain

$$d_j \cdot 10^{-j} \cdot b = r_{j-1} \cdot 10^{-j+1} - r_j \cdot 10^{-j}$$

for  $j \geq 1$ . When we sum from  $j = 1$  to  $j = n$ , most of the terms on the right side cancel [it's called a telescoping sum]. Hence the partial sums  $s_n$  for the decimal expansion satisfy

$$s_n \cdot b = \left[ k + \sum_{j=1}^n d_j \cdot 10^{-j} \right] \cdot b = kb + r_0 - r_n \cdot 10^{-n}.$$

In view of (3), we have  $\lim_n [r_n \cdot 10^{-n}] = 0$ , so  $\lim_n s_n = k + \frac{r_0}{b}$ . Recall that  $r_0 = a - kb$ ; hence

$$\lim_{n \rightarrow \infty} s_n = k + \frac{a - kb}{b} = \frac{a}{b}.$$

Thus  $k.d_1d_2d_3\cdots$  is a decimal expansion for  $\frac{a}{b}$ .

### 16.2 Theorem.

*Every nonnegative real number  $x$  has at least one decimal expansion.*

#### Proof

Let  $a = x$  and  $b = 1$  in 16.1 above. ■

As noted in Discussion 10.3,  $1.000\cdots$  and  $.999\cdots$  are decimal expansions for the same real number. That is, the series

$$1 + \sum_{j=1}^{\infty} 0 \cdot 10^{-j} \quad \text{and} \quad \sum_{j=1}^{\infty} 9 \cdot 10^{-j}$$

have the same value, namely 1. Similarly,  $2.75000\cdots$  and  $2.74999\cdots$  are both decimal expansions for  $\frac{11}{4}$  [Exercise 16.1]. The next theorem shows that this is essentially the only way a number can have distinct decimal expansions.

### 16.3 Theorem.

*A real number  $x$  has exactly one decimal expansion or else  $x$  has two decimal expansions, one ending in a sequence of all 0's and the other ending in a sequence of all 9's.*

#### Proof

We assume  $x \geq 0$ . If  $x$  has decimal expansions  $k.000\cdots$  with  $k > 0$ , then it has one other decimal expansion, namely  $(k - 1).999\cdots$ . If  $x$  has decimal expansion  $k.d_1d_2d_3\cdots d_r000\cdots$  where  $d_r \neq 0$ , then it

has one other decimal expansion  $k.d_1d_2d_3\cdots(d_r-1)9999\cdots$ . The reader can easily check these claims [Exercise 16.2].

Now suppose that  $x$  has two distinct decimal expansions  $k.d_1d_2d_3\cdots$  and  $\ell.e_1e_2e_3\cdots$ . Suppose that  $k < \ell$ . If any  $d_j < 9$ , then by Exercise 16.3 we have

$$x < k + \sum_{j=1}^{\infty} 9 \cdot 10^{-j} = k + 1 \leq \ell \leq x,$$

a contradiction. It follows that  $x = k + 1 = \ell$  and its decimal expansions must be  $k.999\cdots$  and  $(k+1).000\cdots$ . In the remaining case, we have  $k = \ell$ . Let

$$m = \min\{j : d_j \neq e_j\}.$$

We may assume that  $d_m < e_m$ . If  $d_j < 9$  for any  $j > m$ , then by Exercise 16.3,

$$\begin{aligned} x &< k + \sum_{j=1}^m d_j \cdot 10^{-j} + \sum_{j=m+1}^{\infty} 9 \cdot 10^{-j} = k + \sum_{j=1}^m d_j \cdot 10^{-j} + 10^{-m} \\ &= k + \sum_{j=1}^{m-1} e_j \cdot 10^{-j} + d_m \cdot 10^{-m} + 10^{-m} \leq k + \sum_{j=1}^m e_j \cdot 10^{-j} \leq x, \end{aligned}$$

a contradiction. Thus  $d_j = 9$  for  $j > m$ . Likewise, if  $e_j > 0$  for any  $j > m$ , then

$$\begin{aligned} x &> k + \sum_{j=1}^m e_j \cdot 10^{-j} = k + \sum_{j=1}^{m-1} d_j \cdot 10^{-j} + e_m \cdot 10^{-m} \\ &\geq k + \sum_{j=1}^{m-1} d_j \cdot 10^{-j} + d_m \cdot 10^{-m} + 10^{-m} \\ &= k + \sum_{j=1}^m d_j \cdot 10^{-j} + \sum_{j=m+1}^{\infty} 9 \cdot 10^{-j} \geq x, \end{aligned}$$

a contradiction. So in this case,  $d_j = 9$  for  $j > m$ ,  $e_m = d_m + 1$  and  $e_j = 0$  for  $j > m$ . ■

#### 16.4 Definition.

An expression of the form

$$k.d_1d_2\cdots d_\ell \overline{d_{\ell+1}\cdots d_{\ell+r}}$$

represents the decimal expansion in which the block  $d_{\ell+1} \cdots d_{\ell+r}$  is repeated indefinitely:

$$k.d_1d_2 \cdots d_\ell d_{\ell+1} \cdots d_{\ell+r} d_{\ell+1} \cdots d_{\ell+r} d_{\ell+1} \cdots d_{\ell+r} d_{\ell+1} \cdots d_{\ell+r} \cdots$$

We call such an expansion a *repeating decimal*.

### Example 1

Every integer is a repeating decimal. For example,  $17 = 17.\overline{0} = 17.000 \cdots$ . Another simple example is

$$.\overline{8} = .888 \cdots = \sum_{j=1}^{\infty} 8 \cdot 10^{-j} = \frac{8}{10} \sum_{j=0}^{\infty} 10^{-j} = \frac{8}{10} \cdot \frac{10}{9} = \frac{8}{9}.$$

### Example 2

The expression  $3.9\overline{67}$  represents the repeating decimal  $3.9676767 \cdots$ . We evaluate this as follows:

$$\begin{aligned} 3.9\overline{67} &= 3 + 9 \cdot 10^{-1} + 6 \cdot 10^{-2} + 7 \cdot 10^{-3} + 6 \cdot 10^{-4} + 7 \cdot 10^{-5} + \cdots \\ &= 3 + 9 \cdot 10^{-1} + 67 \cdot 10^{-3} \sum_{j=0}^{\infty} (10^{-2})^j \\ &= 3 + 9 \cdot 10^{-1} + 67 \cdot 10^{-3} \left( \frac{100}{99} \right) = 3 + \frac{9}{10} + \frac{67}{990} \\ &= \frac{3928}{990} = \frac{1964}{495}. \end{aligned}$$

Thus the repeating decimal  $3.9\overline{67}$  represents the rational number  $\frac{1964}{495}$ . Any repeating decimal can be evaluated as a rational number in this way, as we'll show in the next theorem.

### Example 3

We find the decimal expansion for  $\frac{11}{7}$ . By the usual long division process in 16.1, we find

$$\frac{11}{7} = 1.571428571428571428571428571 \cdots,$$

i.e.,  $\frac{11}{7} = 1.\overline{571428}$ . To check this, observe

$$\begin{aligned} 1.\overline{571428} &= 1 + 571428 \cdot 10^{-6} \sum_{j=0}^{\infty} (10^{-6})^j = 1 + \frac{571428}{999999} \\ &= 1 + \frac{4}{7} = \frac{11}{7}. \end{aligned}$$

Many books give the next theorem as an exercise, probably to avoid the complicated notation.

### 16.5 Theorem.

*A real number  $x$  is rational if and only if its decimal expansion is repeating. [Theorem 16.3 shows that if  $x$  has two decimal expansions, they are both repeating.]*

#### Proof

First assume  $x \geq 0$  has a repeating decimal expansion  $x = k.d_1d_2 \cdots d_\ell \overline{d_{\ell+1} \cdots d_{\ell+r}}$ . Then

$$x = k + \sum_{j=1}^{\ell} d_j \cdot 10^{-j} + 10^{-\ell} y$$

where

$$y = \overline{.d_{\ell+1} \cdots d_{\ell+r}},$$

so it suffices to show such  $y$  are rational. To simplify the notation, we write

$$y = \overline{.e_1e_2 \cdots e_r}.$$

A little computation shows that

$$y = \sum_{j=1}^r e_j \cdot 10^{-j} \left[ \sum_{j=0}^{\infty} (10^{-r})^j \right] = \sum_{j=1}^r e_j \cdot 10^{-j} \frac{10^r}{10^r - 1}.$$

In fact, if we write  $e_1e_2 \cdots e_r$  for the usual *decimal*  $\sum_{j=0}^{r-1} e_j \cdot 10^{r-1-j}$  *not the product*, then  $y = \frac{e_1e_2 \cdots e_r}{10^r - 1}$ ; see Example 3.

Next consider any positive rational, say  $\frac{a}{b}$  where  $a, b \in \mathbb{N}$ . We may assume that  $a < b$ . As we saw in 16.1,  $\frac{a}{b}$  is given by the decimal

expansion  $.d_1d_2d_3\cdots$  where  $r_0 = a$ ,

$$d_n \in \{0, 1, 2, 3, 4, 5, 6, 7, 8, 9\} \quad (1)$$

$$r_n = 10 \cdot r_{n-1} - d_n b \quad (2)$$

$$0 \leq r_n < b, \quad (3)$$

for  $n \geq 1$ . Since  $a$  and  $b$  are integers, each  $r_n$  is an integer. Thus (3) can be written

$$r_n \in \{0, 1, 2, \dots, b-1\} \quad \text{for } n \geq 0. \quad (4)$$

This set has  $b$  elements, so the first  $b+1$  remainders  $r_n$  cannot all be distinct. That is, there exist integers  $m \geq 0$  and  $p > 0$  so that

$$0 \leq m < m+p \leq b \quad \text{and} \quad r_m = r_{m+p}.$$

From the construction giving (1)–(3) it is clear that given  $r_{n-1}$ , the integers  $r_n$  and  $d_n$  are uniquely determined. Thus

$$r_j = r_k \quad \text{implies} \quad r_{j+1} = r_{k+1} \quad \text{and} \quad d_{j+1} = d_{k+1}.$$

Since  $r_m = r_{m+p}$ , we conclude that  $r_{m+1} = r_{m+1+p}$  and  $d_{m+1} = d_{m+1+p}$ . A simple induction shows that the statement

$$"r_n = r_{n+p} \quad \text{and} \quad d_n = d_{n+p}"$$

holds for all integers  $n \geq m+1$ . Thus the decimal expansion of  $\frac{a}{b}$  is periodic with period  $p$  after the first  $m$  digits. That is,

$$\frac{a}{b} = .d_1d_2\cdots d_m \overline{d_{m+1}\cdots d_{m+p}}.$$

■

#### Example 4

An expansion such as

$.101001000100001000001000000100000001000000001000000000100\cdots$

must represent an irrational number, since it cannot be a repeating decimal: we've arranged for arbitrarily long blocks of 0's.

#### Example 5

We do not know the complete decimal expansions of  $\sqrt{2}$ ,  $\sqrt{3}$  and many other familiar irrational numbers, but we know that they cannot be repeating by virtue of the last theorem.

**Example 6**

We have claimed that  $\pi$  and  $e$  are irrational. These facts and many others are proved in a fascinating book by Ivan Niven [30]. Here is the proof that

$$e = \sum_{k=0}^{\infty} \frac{1}{k!}$$

is irrational. Assume that  $e = \frac{a}{b}$  where  $a, b \in \mathbb{N}$ . Then both  $b!e$  and  $b! \sum_{k=0}^b \frac{1}{k!}$  must be integers, so the difference

$$b! \sum_{k=b+1}^{\infty} \frac{1}{k!}$$

must be a positive integer. On the other hand, this last number is less than

$$\frac{1}{b+1} + \frac{1}{(b+1)^2} + \frac{1}{(b+1)^3} + \cdots = \frac{1}{b} \leq 1,$$

a contradiction.

**Example 7**

There is a famous number introduced by Euler over 200 years ago that arises in the study of the gamma function. It is known as *Euler's constant* and is defined by

$$\gamma = \lim_{n \rightarrow \infty} \left[ \sum_{k=1}^n \frac{1}{k} - \log_e n \right].$$

Even though

$$\lim_{n \rightarrow \infty} \sum_{k=1}^n \frac{1}{k} = +\infty \quad \text{and} \quad \lim_{n \rightarrow \infty} \log_e n = +\infty,$$

the limit defining  $\gamma$  exists and is finite [Exercise 16.9]. In fact,  $\gamma$  is approximately .577216. The amazing fact is that no one knows whether  $\gamma$  is rational or not. Most mathematicians believe  $\gamma$  is irrational. This is because it is “easier” for a number to be irrational, since repeating decimal expansions must be regular. The remark in Exercise 16.8 hints at another reason it is easier for a number to be irrational.

**Exercises**

- 16.1.** (a) Show that  $2.74\overline{9}$  and  $2.75\overline{0}$  are both decimal expansions for  $\frac{11}{4}$ .  
(b) Which of these expansions arises from the long division process described in 16.1?
- 16.2.** Verify the claims in the first paragraph of the proof of Theorem 16.3.
- 16.3.** Suppose that  $\sum a_n$  and  $\sum b_n$  are convergent series of nonnegative numbers. Show that if  $a_n \leq b_n$  for all  $n$  and if  $a_n < b_n$  for at least one  $n$ , then  $\sum a_n < \sum b_n$ .
- 16.4.** Write the following repeating decimals as rationals, i.e., as fractions of integers.
- |                      |                        |
|----------------------|------------------------|
| (a) $.2$             | (b) $.0\overline{2}$   |
| (c) $.\overline{02}$ | (d) $3.\overline{14}$  |
| (e) $.\overline{10}$ | (f) $.149\overline{2}$ |
- 16.5.** Find the decimal expansions of the following rational numbers.
- |                    |                    |
|--------------------|--------------------|
| (a) $\frac{1}{8}$  | (b) $\frac{1}{16}$ |
| (c) $\frac{2}{3}$  | (d) $\frac{7}{9}$  |
| (e) $\frac{6}{11}$ | (f) $\frac{22}{7}$ |
- 16.6.** Find the decimal expansions of  $\frac{1}{7}$ ,  $\frac{2}{7}$ ,  $\frac{3}{7}$ ,  $\frac{4}{7}$ ,  $\frac{5}{7}$  and  $\frac{6}{7}$ . Note the interesting pattern.
- 16.7.** Is  $.1234567891011121314151617181920212223242526 \dots$  rational?
- 16.8.** Let  $(s_n)$  be a sequence of numbers in  $(0, 1)$ . Each  $s_n$  has a decimal expansion  $.d_1^{(n)}d_2^{(n)}d_3^{(n)} \dots$ . For each  $n$ , let  $e_n = 6$  if  $d_n^{(n)} \neq 6$  and  $e_n = 7$  if  $d_n^{(n)} = 6$ . Show that  $e_1e_2e_3 \dots$  is the decimal expansion for some number  $y$  in  $(0, 1)$  and that  $y \neq s_n$  for all  $n$ . *Remark:* This shows that the elements of  $(0, 1)$  cannot be listed as a sequence. In set-theoretic parlance,  $(0, 1)$  is “uncountable.” Since the set  $\mathbb{Q} \cap (0, 1)$  can be listed as a sequence, there must be a lot of irrational numbers in  $(0, 1)$ !
- 16.9.** Let  $\gamma_n = (\sum_{k=1}^n \frac{1}{k}) - \log_e n = \sum_{k=1}^n \frac{1}{k} - \int_1^n \frac{1}{t} dt$ .
- (a) Show that  $(\gamma_n)$  is a decreasing sequence. *Hint:* Look at  $\gamma_n - \gamma_{n+1}$ .
- (b) Show that  $0 < \gamma_n \leq 1$  for all  $n$ .
- (c) Observe that  $\gamma = \lim_n \gamma_n$  exists and is finite.



<http://www.springer.com/978-0-8176-3943-3>

Harmonic Analysis and Hypergroups

Ross, K. (Ed.)

1998, VI, 249 p., Hardcover

ISBN: 978-0-8176-3943-3

A product of Birkhäuser Basel