

The Prandtl-Meyer Solution

S.P. Kiselev, E.V. Vorozhtsov, and V.M. Fomin

■ Impressum

This Mathematica-Notebook is part of the book entitled

S.P. Kiselev, E.V. Vorozhtsov, and V.M. Fomin
 Foundations of Fluid Mechanics with Applications
 Problem Solving Using *Mathematica*.
 ■ Birkhauser Boston, Basel, 1999.

■ Copyright

Foundations of Fluid Mechanics with Applications
 Problem Solving Using *Mathematica*.
 ■ Birkhauser Boston, Basel, 1999.

■ General Description

The program prog6-3.nb, which follows below, enables the user to construct the solution in the centered Prandtl-Meyer rarefaction wave arising in a supersonic flow around a dihedral corner for the case where $|\mathbf{j}_0| < |\mathbf{q}_+|$.

The formulation of this problem and the discussion of its solution can be found in Section 6.4 of the above mentioned book.

■ User's Guide

■ Step 1

Load and compile the program file beginning with the line

```
q [M1_, M_] := g1 * (ArcTan[Sqrt[g2 * (M12 - 1)]] -
```

(see the Section "Program Listing")

■ Step 2

Specify the input data by entering them in the line (see also Section "Examples of the Input Data" below)

Meyer [1.4, 4.0, 1, 10, 1, -1, 1, -0.5, 0.6, 30, 80]

Then click in this line and wait for the result of symbolic/numerical computation.
The meaning of the input parameters is explained in the Section "Parameters Used in Program prog6-3.nb".

■ Program Listing

```

    q [M1_, M_] := g1 * (ArcTan[Sqrt[g2 * (M12 - 1)]] -
ArcTan[Sqrt[g2 * (M2 - 1)]] - ArcTan[Sqrt[M12 - 1]] +
ArcTan[Sqrt[M2 - 1]]];

cl2[f_] := Hue[1.2 * (Af * f + Bf)];

fij[mm_, mm1_] := ArcTan[Sqrt[mm2 - 1], 1] - tj + q[mm1, mm];

Meyer[g_, M1r_, M10_, qmax_, surf_, xl_, xr_,
      yr_, h0_, ix_, mlint_] :=
( g2 = (g - 1) / (g + 1); g1 = 1 / Sqrt[g2];
If[surf == 1, Plot3D[q[m1, m], {m1, 1, M1r}, {m, 1, M1r},
      AspectRatio -> Automatic,
      AxesLabel -> {"M", "M1", "q"}]]];
(* The computation of the Mach number and the
   ratios P/P1 and r/r1 *)
P = {}; r = {}; Mach = {}; xx = {};
g2 = (g - 1) / 2; yr1 = yr; xr1 = xr; h1 = h0;
g4 = g / (g - 1); g5 = 1 / (g - 1); z1 = 1 + g2 M102;
dq = N[Pi / 180];
tcr = Floor[N[90 * (g1 - 1)]]; tmax = Min[qmax, tcr];
Do[qj = (1 - j) * dq;
  (* Plot[qj - q[M10, m], {m, 1, M1r}]; *)
a0 = 1; b0 = M1r; isl = Sign[qj - q[M10, a0]];
isr = Sign[qj - q[M10, b0]];
While[N[Abs[a0 - b0]] > 10-7,
c = (a0 + b0) / 2; isc = Sign[qj - q[M10, c]];
  If[isc == isr, b0 = c, a0 = c]];
AppendTo[Mach, c], {j, tmax}];
Do[z = 1 + g2 Mach[[i]]2; pi = N[(z1 / z)g4]; dd = N[(z1 / z)g5];
AppendTo[P, pi]; AppendTo[r, dd], {i, tmax}];
Print["q, deg.      M      P/P1      r/r1      a, deg."];
Do[ qj = 1 - j; mj = N[Mach[[j]], 9]; pj = N[P[[j]], 9];
  dd = N[r[[j]], 9]; AppendTo[xx, qj];
  aj = N[(180 / Pi) * ArcTan[1 / Sqrt[Mach[[j]]2 - 1]], 9];
Print[" ", qj, " ", mj, " ", pj,
      " ", dd, " ", aj], {j, tmax}];

```

```

(* --- Graphics --- *)
lc = MapThread[List, {xx, Mach}];
gr1 = ListPlot[lc, PlotJoined -> True,
    DisplayFunction -> Identity];
lc = MapThread[List, {xx, P}];
gr2 = ListPlot[lc, PlotJoined -> True,
    PlotStyle -> {Dashing[{0.02}]},
    DisplayFunction -> Identity];
lc1 = MapThread[List, {xx, r}];
gr3 = ListPlot[lc1, PlotStyle -> {PointSize[0.01]},
    DisplayFunction -> Identity];
Show[gr1, gr3, gr2, Frame -> True,
    PlotRange -> All,
    DisplayFunction -> $DisplayFunction];
(* The flowfield in the Prandtl-Meyer wave *)
If[N[Abs[xr]] < 10-7, j0 = N[-Pi/2], j0 = N[ArcTan[yr/xr]]];
If[xr < 0, j0 = N[j0 - Pi]];
(* --- The determination of M2 --- *)
a0 = 1; b0 = M1r; isl = Sign[j0 - q[M10, a0]];
isr = Sign[j0 - q[M10, b0]];
While[N[Abs[a0 - b0]] > 10-7,
c = (a0 + b0) / 2; isc = Sign[j0 - q[M10, c]];
    If[isc == isr, b0 = c, a0 = c]];
M2 = c; fa2 = N[j0 + ArcSin[1/M2]]; cf2 = N[Cos[fa2]];
Print["j0 = ", j0, "; M2 = ", M2, "; a2 = ", fa2];
p2 = N[(z1 / (1 + g2 * M22))g4];
xpl = x1; xpr = xr; ypt = N[Abs[x1]]; ypb = yr;
If[xr < 0.3 * Abs[x1], xpr = 0.5 * Abs[x1]];
jy = Floor[ix * (ypt - ypb) / (xpr - xpl)];
hx = (xpr - xpl) / (ix - 1); hy = (ypt - ypb) / (jy - 1);
Print["The color map on the original ", ix, "x", jy, " mesh"];
mij = Table[M10, {i, ix}, {j, jy}];
pij = Table[1, {i, ix}, {j, jy}];
ctal = N[Sqrt[M102 - 1]];
Do[ yj = ypb + (j - 0.5) * hy; xwj = N[xr * (yj / yr)];
    xcj = -yj * cf2; xuj = N[yj * ctal];
    Do[ xi = x1 + (i - 0.5) * hx;
If[(xi > xcj && yj < 0) || (yj > 0 && xi > xuj),
    tj = ArcTan[xi, yj]; a0 = 1; b0 = M1r;
    isl = Sign[fij[a0, M10]]; isr = Sign[fij[b0, M10]];
    While[N[Abs[a0 - b0]] > 10-4, c = (a0 + b0) / 2;

```

```

isc = Sign[fij[c, M10]]; If[isc == isr, b0 = c, a0 = c]];
mij[[i, j]] = c; pij[[i, j]] = N[(z1 / (1 + g2 * c2))g4]];
If[(xi > xwj && xi < xcj) && (yj < 0),
  mij[[i, j]] = M2; pij[[i, j]] = p2], {i, ix}], {j, jy}];
Pmax = Max[pij]; Pmin = Min[pij];
Af = h0 / (Pmin - Pmax); Bf = -Af * Pmax;
Print["Pmin = ", Pmin, ", Pmax = ", Pmax];
(* Print["P = ", pt]; *)
gr1 = DensityGraphics[Transpose[pij]];
gr2 = Show[gr1, ColorFunction -> (cl2[#]&),
  Mesh -> False,
  DisplayFunction -> Identity];

ynl = -ypb / hy; xn0 = -xpl / hx; xnr = (xr - xpl) / hx;
obj = Polygon[{{0, ynl}, {xn0, ynl}, {xnr, 0}, {0, 0}, {0, ynl}}];
wall = Graphics[{RGBColor[0.3, 0.9, 1], obj}];
gg1 = Show[gr2, wall, AspectRatio -> Automatic,
  DisplayFunction -> Identity];
(* -- Making a color map showing the correspondence
   between individual colors and the numerical
   values of P/P1 ----- *)
dy = N[jy / 10]; dx = dy; mc = {};
Do[ybj = (j - 1) * dy; ytj = ybj + dy; ycj = ybj + 0.5 dy;
  pcj = Pmin + ((j - 0.5) * dy / jy) * (Pmax - Pmin);
  lis = {{0, ybj}, {0, ytj}, {dx, ytj}, {dx, ybj}, {0, ybj}};
  boxj = Graphics[{cl2[pcj], Polygon[lis]}];
  cont = ListPlot[lis, PlotJoined -> True,
    DisplayFunction -> Identity];
  txt = Graphics[Text[pcj, {2 dx, ycj}, {-1, 0}]];
  gj = Show[txt, boxj, cont, DisplayFunction -> Identity];
  AppendTo[mc, gj], {j, 10}];
gg2 = Show[mc, AspectRatio -> Automatic,
  PlotRange -> All,
  DisplayFunction -> Identity];
Show[GraphicsArray[{gg1, gg2}]];
(* --- Interpolation on a finer mesh --- *)
tb = {};
Do[yj = ypb + (j - 1) * hy;
  Do[xi = xpl + (i - 1) * hx;
    AppendTo[tb, {xi, yj, pij[[i, j]]}], {i, ix}], {j, jy}];
Print["Calculation of the interpolating function"];
vint = Interpolation[tb, InterpolationOrder -> 1];

```

```

h1n = (xpr - xpl) / (m1int - 1);
m2int = Floor[m1int * (ypt - ypb) / (xpr - xpl)];
h2n = (ypt - ypb) / (m2int - 1);
vn = Table[0.0, {i, m1int}, {j, m2int}];
(* -- Calculation of the values of P (x,y)/P1 in
    the nodes of a fine mesh ----- *)
Do[yj = ypb + (j - 1) * h2n;
    Do[xi = xpl + (i - 1) * h1n;
        vn[[i, j]] = vint[xi, yj], {i, m1int}], {j, m2int}];
gr3 = DensityGraphics[Transpose[vn]];
gr4 = Show[gr3, ColorFunction -> (cl2[#]&),
            Mesh -> False,
            DisplayFunction -> Identity];
ynl = - ypb / h2n; xn0 = - xpl / h1n;
xnr = (xr - xpl) / h1n;
obj = Polygon[{{0, ynl}, {xn0, ynl}, {xnr, 0},
               {0, 0}, {0, ynl}}];
wall = Graphics[{RGBColor[0.3, 0.9, 1], obj}];
gg3 = Show[gr4, wall, AspectRatio -> Automatic,
            DisplayFunction -> Identity];
Print["The color map on the fine ", m1int, "x", m2int, " mesh"]
Show[GraphicsArray[{gg3, gg2}]]]; )

```

■ Parameters Used in Program prog6-3.nb

Parameter	Description
g	the ratio of the gas specific heats; $g > 0$
$M1r$	the right end of the interval $[1, M1r]$, in which the local value $M(x, y)$ of the Mach number is computed; $M1r > 1$; $M1r > 3$ for $g = 1.4$
$M10$	the freestream Mach number; $M10 \geq 1$
q_{\max}	the maximum value of the angle q , in degrees, for making the one-dimensional plots of $M(q)$, $P(q)/P_1$, $r(q)/r_1$, $a(q)$; q is a positive integer, $1 < q < q_+$
surf	if surf = 1 then the surface $q = f(M_1, M)$ is plotted; otherwise this surface is not plotted
x_l	the abscissa of the left end $(x_l, 0)$ of a dihedral corner, $x_l < 0$
x_r	the abscissa of the right end of the right facet of the dihedral corner
y_r	the ordinate of the right end of the right facet of the dihedral corner, $y_r < 0$
h_0	the number4 in the interval $0 < h_0 \leq 1$ to specify the color

corresponding to the minimum value of the function $P(x, y)/P_1$ in the spatial region under consideration


ix the number of cells of an uniform rectangular (rough) grid in the interval $x_l \leq x \leq x_r$; ix is a positive integer, $ix > 1$

m1int the number of cells of a fine mesh in the interval $x_l \leq x \leq x_r$; m1int $> ix$

■ Examples of the Input Data

■ Example 1


Meyer[1.4, 4.0, 1, 60, 1, -1, 1, -0.5, 0.6, 30, 100]



q, deg.	M	P/P1	r/r1	a, deg.
0	1.00000009	0.999999896	0.999999925	89.9757717
-1	1.08181283	0.906677246	0.932414639	67.5741964
-2	1.1326026	0.851129804	0.891244779	61.9969031
-3	1.17686996	0.804379084	0.855996478	58.1803101
-4	1.2177352	0.762678786	0.82405905	55.2047574
-5	1.25647327	0.724491496	0.79437173	52.7383063
-6	1.29376224	0.688998854	0.766375805	50.6185967
-7	1.33001676	0.655701263	0.739734253	48.7526432
-8	1.36551365	0.62426639	0.714226041	47.081676
-9	1.4004505	0.594458493	0.689696846	45.5658835
-10	1.43497446	0.566103152	0.666034672	44.1769403
-11	1.46919981	0.539066156	0.643155165	42.8938798
-12	1.50321791	0.513241232	0.620993156	41.7007055
-13	1.53710368	0.488542152	0.599497218	40.5849082

- 14	1.57092008	0.464897337	0.57862594	39.5364978
- 15	1.6047211	0.442246189	0.558345381	38.5473547
- 16	1.63855377	0.420536622	0.538627403	37.6107847
- 17	1.67245975	0.399723146	0.519448331	36.7211959
- 18	1.70647642	0.379765522	0.500788045	35.8738665
- 19	1.74063793	0.360627604	0.482629148	35.0647656
- 20	1.77497575	0.342276624	0.46495649	34.2904245
- 21	1.80951938	0.324682486	0.44775665	33.5478313
- 22	1.84429637	0.307817492	0.431017805	32.8343606
- 23	1.87933299	0.291655766	0.414729273	32.1477025
- 24	1.91465446	0.276173001	0.398881347	31.4858147
- 25	1.95028564	0.261346018	0.383464914	30.8468734
- 26	1.98624995	0.247153106	0.36847187	30.2292637
- 27	2.022571	0.233573254	0.353894368	29.6315263
- 28	2.05927184	0.220586379	0.339725101	29.0523502
- 29	2.09637555	0.208173063	0.325957051	28.4905474
- 30	2.13390502	0.196314573	0.312583536	27.9450415
- 31	2.17188314	0.184992756	0.299598122	27.4148536
- 32	2.21033314	0.174189902	0.286994486	26.8990864
- 33	2.24927828	0.163888829	0.274766526	26.3969215
- 34	2.28874233	0.154072717	0.262908181	25.9076047
- 35	2.32874945	0.144725148	0.251413488	25.4304418
- 36	2.36932448	0.135830021	0.240276481	24.96479
- 37	2.41049227	0.127371699	0.229491384	24.5100608
- 38	2.45227894	0.11933474	0.21905228	24.065702
- 39	2.49471077	0.111704123	0.208953395	23.6312062
- 40	2.53781548	0.104465015	0.199188797	23.2060948
- 41	2.58162096	0.0976029996	0.189752705	22.7899284
- 42	2.62615672	0.0911038447	0.180639171	22.3822902
- 43	2.67145279	0.0849536856	0.171842333	21.9827949
- 44	2.71754029	0.0791389419	0.163356299	21.5910817
- 45	2.76445213	0.0736462453	0.155175039	21.2068068

- 46	2.81222209	0.0684625682	0.147292573	20.8296509
- 47	2.86088505	0.0635751985	0.139702931	20.4593154
- 48	2.91047803	0.0589716428	0.13240001	20.0955142
- 49	2.9610391	0.0546397492	0.125377758	19.7379812
- 50	3.01260832	0.0505676313	0.118630052	19.386463
- 51	3.06522754	0.0467436995	0.112150755	19.0407203
- 52	3.11894056	0.0431566586	0.105933702	18.7005263
- 53	3.17379317	0.0397955173	0.099972727	18.3656671
- 54	3.22983363	0.0366495658	0.094261616	18.035938
- 55	3.28711239	0.033708409	0.0887941733	17.7111459
- 56	3.34568235	0.0309619533	0.0835641982	17.3911077
- 57	3.4055995	0.0284003915	0.0785654607	17.0756475
- 58	3.46692267	0.0260142237	0.0737917423	16.7645984
- 59	3.5297139	0.0237942504	0.069236828	16.4578007




$j_0 = -0.463648$; $M_2 = 2.00673$; $a_2 = 0.0580166$

The color map on the original 30x22 mesh

$P_{\min} = 0.0216725$, $P_{\max} = 1$

Calculation of the interpolating function

The color map on the fine 100x75 mesh




■ Example 2

The dihedral angle magnitude = $\pi/2$

Meyer[1.4, 8.0, 1, 10, 0, -1, 0.0, -0.5, 0.6, 30, 50]

q, deg.	M	P/Pl	r/r1	a, deg.
0	1.00000005	0.999999939	0.999999957	89.9814953
-1	1.08181288	0.906677188	0.932414597	67.5741897
-2	1.13260259	0.851129812	0.891244785	61.9969039
-3	1.17687003	0.804379014	0.855996426	58.1803049
-4	1.21773513	0.762678853	0.824059102	55.204762
-5	1.25647319	0.724491576	0.794371792	52.7383112
-6	1.2937622	0.688998888	0.766375832	50.6185987
-7	1.33001668	0.655701337	0.739734312	48.7526472
-8	1.36551367	0.62426637	0.714226025	47.081675
-9	1.40045054	0.594458461	0.68969682	45.565882



j0 = -1.5708; M2 = 6.81904; a2 = -1.42362

The color map on the original 30x30 mesh

Pmin = 0.000539792, Pmax = 1

Calculation of the interpolating function


The color map on the fine 50x50 mesh

■ Example 3

The dihedral angle magnitude $< \mathbf{p}/2$

Meyer[1.4, 10.0, 1, 10, 0, -1, -0.07, -0.5, 0.6, 57, 90]

q, deg.	M	P/Pl	r/r1	a, deg.
0	1.00000007	0.999999922	0.999999944	89.9790177
-1	1.08181281	0.906677271	0.932414657	67.5741993
-2	1.13260258	0.851129828	0.891244797	61.9969053
-3	1.17687003	0.804379014	0.855996426	58.1803049
-4	1.21773509	0.762678898	0.824059136	55.204765
-5	1.25647316	0.724491604	0.794371815	52.738313
-6	1.29376217	0.688998916	0.766375855	50.6186003
-7	1.33001678	0.655701243	0.739734237	48.7526421
-8	1.36551367	0.62426637	0.714226025	47.081675
-9	1.40045052	0.594458474	0.68969683	45.5658826




j0 = -1.70989; M2 = 8.6092; a2 = -1.59347

The color map on the original 57x57 mesh

Pmin = 0.000120105, Pmax = 1

Calculation of the interpolating function

The color map on the fine 90x90 mesh



■ The Structure of the Output

The following information is printed out in the process of the work of the program:

1. The surface $q = q(M_1, M)$;
2. The numerical values of the functions $M(q)$, $P(q)/P_1$, $r(q)/r_1$, $\alpha(q)$ for $q = 1, 2, \dots, q_{\max}$ degrees;
3. The picture showing the curves of the functions listed in item No. 2;
4. The computed numerical values of the angle j_0 (in radians), the Mach number M_2 , and the angle α_2 (in radians);
5. The numerical values of $\min(P(x, y)/P_1)$ and $\max(P(x, y)/P_1)$;
6. The color map of the pressure field $P(x, y)/P_1$ on a rough mesh of $ix * jy$ cells along with a colored column giving the correspondence between the individual colors and the values of P/P_1 ;
7. The color map of the pressure field $P(x, y)/P_1$ on a fine mesh of $m1int * m2int$ cells along with a colored column giving the correspondence between the individual colors and the values of P/P_1 ;

To resize an individual picture obtained by *Mathematica*, please

- (i) Click anywhere inside the cell, but not the cell bracket itself. A bounding box with small handles appears around the graphic image.
- (ii) Drag one of the handles to adjust the size and shape of the bounding box. In this way it is possible to resize the height and width of the graphic image.