

Contents

| | |
|--|----------|
| 1 Introduction | 1 |
| 2 Optics of Nonlinear Crystals | 3 |
| 2.1 Three- and Four-Wave (Three- and Four-Frequency) Interactions in Nonlinear Media | 3 |
| 2.2 Phase-Matching Conditions | 5 |
| 2.3 Optics of Uniaxial Crystals | 6 |
| 2.4 Types of Phase Matching in Uniaxial Crystals | 10 |
| 2.5 Calculation of Phase-Matching Angles in Uniaxial Crystals | 13 |
| 2.6 Reflection and Refraction of Light Waves at the Surfaces of Uniaxial Crystals | 14 |
| 2.7 Optics of Biaxial Crystals | 16 |
| 2.8 Types of Phase Matching in Biaxial Crystals | 18 |
| 2.9 Calculation of Phase-Matching Angles in Biaxial Crystals | 19 |
| 2.10 Crystal Symmetry and Effective Nonlinearity: Uniaxial Crystals | 23 |
| 2.11 Crystal Symmetry and Effective Nonlinearity: Biaxial Crystals | 25 |
| 2.12 Theory of Nonlinear Frequency-Conversion Efficiency | 32 |
| 2.13 Wave Mismatch and Phase-Matching Bandwidth | 40 |
| 2.14 Calculation of Nonlinear Frequency-Conversion Efficiency in Some Special Cases | 48 |
| 2.14.1 Plane-Wave Fixed-Field Approximation | 49 |
| 2.14.2 Fundamental Wave Depletion (“Nonlinear Regime”) | 52 |
| 2.14.3 SHG of a Divergent Fundamental Radiation Beam in the Fixed-Field Approximation | 54 |
| 2.14.4 SHG of a Divergent Fundamental Radiation Beam in the Nonlinear Regime | 55 |
| 2.14.5 Fixed-Intensity Approximation | 57 |
| 2.14.6 Frequency Conversion of Ultrashort Laser Pulses | 59 |
| 2.14.7 Frequency Conversion of Laser Beams with Limited Aperture in the Stationary Regime | 61 |
| 2.14.8 Linear Absorption | 65 |
| 2.15 Additional Comments | 65 |

| | |
|--|-----------|
| 3 Properties of Nonlinear Optical Crystals | 67 |
| 3.1 Basic Nonlinear Optical Crystals | 68 |
| 3.1.1 LiB_3O_5 , Lithium Triborate (LBO) | 68 |
| 3.1.2 KH_2PO_4 , Potassium Dihydrogen Phosphate (KDP) | 78 |
| 3.1.3 KD_2PO_4 , Deuterated Potassium Dihydrogen Phosphate (DKDP) | 85 |
| 3.1.4 $\text{NH}_4\text{H}_2\text{PO}_4$, Ammonium Dihydrogen Phosphate (ADP) | 90 |
| 3.1.5 $\beta\text{-BaB}_2\text{O}_4$, Beta-Barium Borate (BBO) | 96 |
| 3.1.6 LiIO_3 , Lithium Iodate | 103 |
| 3.1.7 KTiOPO_4 , Potassium Titanyl Phosphate (KTP) | 107 |
| 3.1.8 LiNbO_3 , Lithium Niobate | 119 |
| 3.1.9 KNbO_3 , Potassium Niobate | 126 |
| 3.1.10 AgGaS_2 , Silver Thiogallate | 132 |
| 3.1.11 ZnGeP_2 , Zinc Germanium Phosphide | 136 |
| 3.2 Frequently Used Nonlinear Optical Crystals | 142 |
| 3.2.1 $\text{KB}_5\text{O}_8 \cdot 4\text{H}_2\text{O}$, Potassium Pentaborate Tetrahydrate (KB5) | 142 |
| 3.2.2 $\text{CO}(\text{NH}_2)_2$, Urea | 146 |
| 3.2.3 CsH_2AsO_4 , Cesium Dihydrogen Arsenate (CDA) | 149 |
| 3.2.4 CsD_2AsO_4 , Deuterated Cesium Dihydrogen Arsenate (DCDA) | 152 |
| 3.2.5 KTiOAsO_4 , Potassium Titanyl Arsenate (KTA) | 156 |
| 3.2.6 $\text{MgO} : \text{LiNbO}_3$, Magnesium-Oxide-Doped Lithium Niobate | 159 |
| 3.2.7 Ag_3AsS_3 , Proustite | 162 |
| 3.2.8 GaSe , Gallium Selenide | 166 |
| 3.2.9 AgGaSe_2 , Silver Gallium Selenide | 169 |
| 3.2.10 CdSe , Cadmium Selenide | 173 |
| 3.2.11 CdGeAs_2 , Cadmium Germanium Arsenide | 176 |
| 3.3 Other Inorganic Nonlinear Optical Crystals | 179 |
| 3.3.1 $\text{KB}_5\text{O}_8 \cdot 4\text{D}_2\text{O}$, Deuterated Potassium Pentaborate Tetrahydrate (DKB5) | 179 |
| 3.3.2 CsB_3O_5 , Cesium Triborate (CBO) | 180 |
| 3.3.3 $\text{BeSO}_4 \cdot 4\text{H}_2\text{O}$, Beryllium Sulfate | 182 |
| 3.3.4 MgBaF_4 , Magnesium Barium Fluoride | 184 |
| 3.3.5 $\text{NH}_4\text{D}_2\text{PO}_4$, Deuterated Ammonium Dihydrogen Phosphate (DADP) | 186 |
| 3.3.6 RbH_2PO_4 , Rubidium Dihydrogen Phosphate (RDP) | 188 |
| 3.3.7 RbD_2PO_4 , Deuterated Rubidium Dihydrogen Phosphate (DRDP) | 192 |
| 3.3.8 KH_2AsO_4 , Potassium Dihydrogen Arsenate (KDA) | 192 |
| 3.3.9 KD_2AsO_4 , Deuterated Potassium Dihydrogen Arsenate (DKDA) | 195 |

| | | |
|--------|--|-----|
| 3.3.10 | $\text{NH}_4\text{H}_2\text{AsO}_4$, Ammonium Dihydrogen Arsenate (ADA) | 196 |
| 3.3.11 | $\text{NH}_4\text{D}_2\text{AsO}_4$, Deuterated Ammonium Dihydrogen Arsenate (DADA) | 198 |
| 3.3.12 | RbH_2AsO_4 , Rubidium Dihydrogen Arsenate (RDA) | 199 |
| 3.3.13 | RbD_2AsO_4 , Deuterated Rubidium Dihydrogen Arsenate (DRDA) | 202 |
| 3.3.14 | $\text{LiCOOH} \cdot \text{H}_2\text{O}$, Lithium Formate Monohydrate (LFM) | 204 |
| 3.3.15 | NaCOOH , Sodium Formate | 207 |
| 3.3.16 | $\text{Ba}(\text{COOH})_2$, Barium Formate | 209 |
| 3.3.17 | $\text{Sr}(\text{COOH})_2$, Strontium Formate | 210 |
| 3.3.18 | $\text{Sr}(\text{COOH})_2 \cdot 2\text{H}_2\text{O}$, Strontium Formate Dihydrate | 211 |
| 3.3.19 | LiGaO_2 , Lithium Gallium Oxide | 213 |
| 3.3.20 | $\alpha\text{-HIO}_3$, α -Iodic Acid | 214 |
| 3.3.21 | $\text{K}_2\text{La}(\text{NO}_3)_5 \cdot 2\text{H}_2\text{O}$, Potassium Lanthanum Nitrate Dihydrate (KLN) | 217 |
| 3.3.22 | CsTiOAsO_4 , Cesium Titanyl Arsenate (CTA) | 220 |
| 3.3.23 | NaNO_2 , Sodium Nitrite | 221 |
| 3.3.24 | $\text{Ba}_2\text{NaNb}_5\text{O}_{15}$, Barium Sodium Niobate (“Banana”) | 224 |
| 3.3.25 | $\text{K}_2\text{Ce}(\text{NO}_3)_5 \cdot 2\text{H}_2\text{O}$, Potassium Cerium Nitrate Dihydrate (KCN) | 227 |
| 3.3.26 | $\text{K}_3\text{Li}_2\text{Nb}_5\text{O}_{15}$, Potassium Lithium Niobate | 229 |
| 3.3.27 | HgGa_2S_4 , Mercury Thiogallate | 231 |
| 3.3.28 | HgS , Cinnibar | 233 |
| 3.3.29 | Ag_3SbS_3 , Pyraryrite | 235 |
| 3.3.30 | Se, Selenium | 236 |
| 3.3.31 | Tl_3AsS_3 , Thallium Arsenic Selenide (TAS) | 238 |
| 3.3.32 | Te, Tellurium | 240 |
| 3.4 | Other Organic Nonlinear Optical Crystals | 243 |
| 3.4.1 | $\text{C}_{12}\text{H}_{22}\text{O}_{11}$, Sucrose (Saccharose) | 243 |
| 3.4.2 | L-Arginine Phosphate Monohydrate (LAP) | 245 |
| 3.4.3 | Deuterated L-Arginine Phosphate Monohydrate (DLAP) | 247 |
| 3.4.4 | L-Pyrrolidone-2-carboxylic Acid (L-PCA) | 250 |
| 3.4.5 | $\text{CaC}_4\text{H}_4\text{O}_6 \cdot 4\text{H}_2\text{O}$, Calcium Tartrate Tetrahydrate (L-CTT) | 251 |
| 3.4.6 | $(\text{NH}_4)_2\text{C}_2\text{O}_4 \cdot \text{H}_2\text{O}$, Ammonium Oxalate (AO) | 253 |
| 3.4.7 | m-Bis(aminomethyl)benzene (BAMB) | 254 |
| 3.4.8 | 3-Methoxy-4-hydroxy-benzaldehyde (MHBA) | 256 |
| 3.4.9 | 2-Furyl Methacrylic Anhydride (FMA) | 258 |
| 3.4.10 | 3-Methyl-4-nitropyridine-1-oxide (POM) | 259 |
| 3.4.11 | Thienylchalcone (T-17) | 261 |
| 3.4.12 | 5-Nitrouracil (5NU) | 263 |
| 3.4.13 | 2-(N-Prolinol)-5-nitropyridine (PNP) | 265 |

| | | |
|----------|---|------------|
| 3.4.14 | 2-Cyclooctylamino-5-nitropyridine (COANP) | 266 |
| 3.4.15 | L-N-(5-Nitro-2-pyridyl)leucinol (NPLO) | 268 |
| 3.4.16 | C ₆ H ₄ (NO ₂) ₂ , m-Dinitrobenzene (MDNB) | 270 |
| 3.4.17 | 4-(N,N-Dimethylamino)-3-acetamidonitrobenzene (DAN) | 272 |
| 3.4.18 | Methyl-(2,4-dinitrophenyl)-aminopropanoate (MAP) | 274 |
| 3.4.19 | m-Nitroaniline (MNA) | 276 |
| 3.4.20 | N-(4-Nitrophenyl)-N-methylaminoacetonitrile (NPAN) | 278 |
| 3.4.21 | N-(4-Nitrophenyl)-L-prolinol (NPP) | 280 |
| 3.4.22 | 3-Methyl-4-methoxy-4'-nitrostilbene (MMONS) | 281 |
| 3.5 | Properties of Crystalline Quartz (α -SiO ₂) | 283 |
| 3.6 | New Developments | 286 |
| 4 | Applications of Nonlinear Crystals | 289 |
| 4.1 | Generation of Neodymium Laser Harmonics | 289 |
| 4.1.1 | Second-Harmonic Generation of Neodymium Laser Radiation in Inorganic Crystals | 289 |
| 4.1.2 | Second-Harmonic Generation of 1,064 μ m Radiation in Organic Crystals | 294 |
| 4.1.3 | Intracavity SHG | 296 |
| 4.1.4 | Third-Harmonic Generation | 298 |
| 4.1.5 | Fourth-Harmonic Generation | 301 |
| 4.1.6 | Fifth-Harmonic Generation | 301 |
| 4.1.7 | Harmonic Generation of 1,318 μ m Radiation | 304 |
| 4.2 | Harmonic Generation of High-Power Large-Aperture Neodymium Glass Laser Radiation | 306 |
| 4.2.1 | "Angle-Detuning" Scheme | 306 |
| 4.2.2 | "Polarization-Mismatch" Scheme | 306 |
| 4.2.3 | "Polarization-Bypass" Scheme | 308 |
| 4.2.4 | Comparison of Schemes | 308 |
| 4.2.5 | Experimental Results | 308 |
| 4.2.6 | "Quadrature" Scheme | 310 |
| 4.3 | Harmonic Generation for Other Laser Sources | 311 |
| 4.3.1 | Ruby Laser | 311 |
| 4.3.2 | Ti:sapphire Laser | 312 |
| 4.3.3 | Semiconductor Lasers | 312 |
| 4.3.4 | Dye Lasers | 315 |
| 4.3.5 | Gas Lasers | 320 |
| 4.3.6 | Iodine Laser | 321 |
| 4.3.7 | CO ₂ Laser | 324 |
| 4.3.8 | Other Lasers | 324 |
| 4.3.9 | Frequency Conversion of Femtosecond Pulses | 326 |

| | | |
|-------|--|------------|
| 4.4 | Sum-Frequency Generation | 327 |
| 4.4.1 | Up-Conversion to the UV Region | 328 |
| 4.4.2 | Infrared Up-Conversion | 333 |
| 4.4.3 | Up-Conversion of CO ₂ Laser Radiation to the Near IR and Visible Regions | 336 |
| 4.5 | Difference Frequency Generation | 339 |
| 4.5.1 | DFG in the Visible Region | 339 |
| 4.5.2 | DFG in the Mid IR Region | 340 |
| 4.5.3 | DFG in the Far IR Region | 344 |
| 4.6 | Optical Parametric Oscillation | 345 |
| 4.6.1 | OPO in the UV, Visible, and Near IR Spectral Regions | 345 |
| 4.6.2 | OPO in the Mid IR Region | 359 |
| 4.6.3 | Conversion of OPO Radiation to the UV Region | 360 |
| 4.7 | Stimulated Raman Scattering and Picosecond Continuum Generation in Crystals | 362 |
| | References | 367 |
| | Appendix: List of Commonly Used Laser Wavelengths | 405 |
| | Subject Index | 407 |

Handbook of Nonlinear Optical Crystals

Dmitriev, V.G.; Gurzadyan, G.G.; Nikogosyan, D.N.

1999, XVIII, 414 p., Hardcover

ISBN: 978-3-540-65394-3