

Prion Protein as Copper-Binding Protein at the Synapse

Hans A. Kretzschmar, Tobias Tings, Axel Madlung, Armin Giese, and Jochen Herms

1. Introduction

Various approaches have been taken to study the function of prion proteins. Biochemical methods were applied to search for a binding partner of PrP^C which is attached to the cell surface by a glycosylphosphatidylinositol GPI anchor (1). The glial fibrillary acidic protein was one of the first possible binding partners to be described (2) followed by Bcl-2 (3,4), molecular chaperones (5), amyloid precursor-like protein 1 (6), the 37-kDa laminin receptor (7) and a 66-kDa membrane protein which has not been characterized in more detail (8). However, it has not been possible to show any biological significance for PrP^C binding of these proteins. Based on biochemical analyses of chicken PrP^C, Harris et al. (9) hypothesized that PrP^C may play a role in the regulation of the expression of cholinergic receptors at the neuromuscular endplate.

Biochemical, morphological, and electrophysiological studies of the first PrP gene (*Prnp*) knockout mouse (Prnp^{0/0} mouse), which was generated by Büeler et al. (10), showed a regular expression of the acetylcholine receptor (11). Except for changes in its circadian rhythm (12,13) and increased sensitivity to seizures (14), this Prnp^{0/0} mouse showed no developmental or behavioral changes (10). These findings were confirmed in studies of another Prnp^{0/0} line generated by Manson et al. (15). The lack of severe defects in these two lines of Prnp^{0/0} mice was ascribed to adaptation, because PrP^C was absent throughout embryogenesis. However, transgenic mice expressing inducible PrP^C-

transgenes that were rendered PrP^C-deficient as adults by administration of doxycycline have remained healthy for more than 1.5 yr (16). A third Prnp^{0/0} mouse generated by Sakaguchi et al. (17) showed progressive ataxia and loss of Purkinje cells in mice aged more than 70 wk. Also, a fourth independently generated Prnp^{0/0} mouse (18,19) exhibits ataxia and Purkinje cell degeneration. Weissmann (20) suggested that additional deletions of intronic sequences of Prnp may play a role in this knockout line. Most recently the upregulation of a novel PrP^C-like protein, designated Doppel, whose gene is located 16 kb downstream of the mouse PrP, has been speculated to be the cause of Purkinje cell degeneration observed in two of the Prnp^{0/0} mouse lines (21). Even though the hypothesis of the interaction of prion proteins with cholinergic receptors thus could not be confirmed, the studies of Harris et al. (9) indicated that PrP^C is enriched at the neuromuscular end-plate, i.e. at synaptic endings. Indeed immunohistochemistry of PrP^C-overexpressing transgenic mice reveal a synaptic expression pattern of PrP^C (22,23). PrP^C is predominantly expressed in regions of high synaptic density, such as the inner and outer plexiform layer of the retina or the cerebellar molecular layer (Fig. 1), in contrast to earlier studies in which a predominantly somatic expression of PrP^C was described (24–26). Further evidence for a preferentially synaptic location of the prion protein in the central nervous system was shown in immunoelectron microscopic studies by Fournier et al. (27) and Salès et al. (28). Electron microscopic evidence for a synaptic location of PrP^C has proven very difficult, however. Thus, it was necessary to use embedding techniques leading to destruction of cell membranes. As a consequence, the electron microscopic evidence for PrP^C location in synaptic vesicles has been disputed. Biochemical studies showed that the prion protein is located predominantly in the synaptic plasma membrane (23) and, to a lesser extent, in the synaptic vesicle fraction. Fig. 2 shows a Western blot analysis of PrP^C expression in various synaptic fractions. The enrichment of PrP^C in the synaptic plasma membrane fraction is evident (Fig. 2A, lane 4).

2. Electrophysiological Studies

Electrophysiological studies in Prnp^{0/0} mice have been used to identify the function of PrP^C in neurons. Collinge et al. (29) were the first to describe a change in long-term potentiation (LTP), i.e., a change of synaptic transmission after repetitive stimulation in the Prnp^{0/0} mouse generated by Büeler et al. (10). This finding was confirmed in a second Prnp^{0/0} mouse generated by Manson et al. (30). However, Lledo et al. (31) did not observe LTP changes.

In addition, Collinge et al. (29) found altered kinetics of the inhibitory postsynaptic currents (IPSCs), i.e., a prolongation of the rise time of GABA_A receptor-mediated IPSCs in hippocampal neurons of Prnp^{0/0} mice. The authors argue that this may be caused by changes in the GABA_A receptor on the




Fig. 1. Synaptic expression pattern of PrP^C in PrP^C-overexpressing transgenic mice. Laser scanning confocal images of PrP^C expression in the retina and cerebellar cortex of PrP^C-overexpressing mice. Expression of PrP^C (A) and synaptophysin (B) in Tg20 retina. PrP^C is strongly expressed in the inner and outer plexiform layer, similar to synaptophysin. PrP^C expression in Tg35 (C) and Tg20 (D) cerebellar cortex. Strong PrP^C expression was observed in the molecular and granule cell layers in both transgenic mouse lines. However PrP^C expression in Purkinje cells was only observed in Tg35 (C).

postsynaptic membrane since a decrease of the amplitude of stimulated inhibitory postsynaptic currents and a shift of the reverse potential of GABA_A receptor-mediated chloride currents were also observed. Lledo et al. (31) did not confirm this finding for hippocampal neurons of the same knockout line. Also, a more detailed analysis of the kinetics of GABA_A-induced currents in outside-out patches from cerebellar Purkinje cells of Prnp^{0/0} mice did not reveal significant deviations from control cells (32). Moreover, studies on the kinetics of spontaneous inhibitory postsynaptic currents (sIPSCs) in cerebellar Purkinje cells of Prnp^{0/0} mice initially showed significant differences between the rise time of wild-type and that of Prnp^{0/0} Purkinje cells (32). Further experiments with Purkinje cells of younger animals, with a better voltage clamp (and consequently a more exact estimation of the rise time [33]) showed a significant increase in the rise time, from 1.9 ms in wild-type to 2.81 ms in Prnp^{0/0} mouse Purkinje cells (Fig. 3D; $P = 0.001$). No differences were found in the decay time (Fig. 3E). Evidence for the hypothesis that the increased rise time is caused by loss of the PrP^C was found in studies on the rise time in Prnp^{0/0} mice that were Prnp reconstituted (Fig. 3D; Tg35; [34]). The IPSC rise time in Purkinje cells of these animals corresponds to the rise time in wildtype ani-




Fig. 2. Enrichment of PrP^C in the synaptic plasma membrane fraction. Preparations of the synaptic plasma membrane fraction and synaptic vesicle fractions from synaptosomes (54). Equal amounts (100 µg/per lane) of brain homogenate and various subcellular fractions from wild-type (lane 1–4), Prnp^{0/0} (lane 6), and Tg35 (lane 7) mice were investigated in Western blots. The monoclonal antibody 3B5 (A); hybridoma supernatant 1:50) (55) was used to identify PrP^C. A polyclonal antiserum (1:2000) was used to identify the synaptic vesicle protein synaptotagmin (B) (56). The N-methyl-D-aspartate (NMDA) receptor subunit, R1, was shown using the monoclonal antibody, Akp (C); (1:2000) (55,57). Subcellular fractions are designated as follows: lane 1, WT homogenate; lane 2, WT crude synaptic vesicle fraction; lane 3, WT cytosolic synaptic fraction; lane 4, WT synaptic plasma membrane fraction; lane 5, mol w. standards; lane 5 synaptic plasma membrane fraction from Prnp^{0/0} mouse brains. An enrichment of PrP^C (A) is noted in the synaptic plasma membrane fraction of wild-type mouse (lane 4), in analogy to the subunit R1 of the NMDA receptor in lane 4 (C). In contrast to synaptotagmin, a protein that is predominantly localized to the membranes of synaptic vesicles, PrP^C is not enriched in the synaptic vesicle fraction (lane 2), although it may be found in this location in low concentration.

mals. To clarify the question of whether the increase in rise time in Prnp^{0/0} mice is caused by the loss of PrP^C expression in the presynapse or postsynapse, an additional Tg line, which expresses PrP^C only at the presynapse (Tg20) (34) was examined. In this line, rise times corresponding to the wildtype were found (Fig. 3D). Thus, it appears that the loss of the presynaptic PrP^C expression at the inhibitory synapse is responsible for the prolongation of the rise time of inhibitory postsynaptic currents in Prnp^{0/0} mice.

Independent of the findings at inhibitory synapses, Colling et al. (35) described an additional electrophysiological phenotype in Prnp^{0/0} mice, i. e. a disturbance of the late afterhyperpolarization current, I_{AHP}. This current is involved in action potential repolarization and therefore influences the frequency of action potentials. Colling et al. (35) reasoned that the disturbed I_{AHP} in Prnp^{0/0} mice is caused by a decreased conductance of calcium-activated




Fig. 3. Presynaptic PrP^C expression modulates the kinetics of inhibitory postsynaptic currents (IPSC). (A), Spontaneous IPSCs from a Purkinje cell of a 10-d-old wild-type mouse using the patch-clamp technique, as described (32) (B), Using the effect of 10 μ M bicucullin, a γ -aminobutyric acid A (GABA_A) receptor blocker, it is shown that the synaptic currents are inhibitory GABA_A receptor-mediated conductances. (C), rise time and decay time in wildtype IPSCs. During rise time, there is a linear increase of GABA_A receptor-mediated current from 10 to 90% of the maximum (gray line *a*). The decay time (τ) is calculated from the kinetics of an exponential function (gray line *b*) that shows the best fit to the actual decay of the current. (D), Rise time in WT, Prnp^{0/0}, Tg20, and Tg35. Shown is the mean of results from each of 10 measurements in Purkinje cells of 9–12d-old animals. Each point corresponds to the rise time of inhibitory postsynaptic currents of a Purkinje cell (mean of the rise time of 20 consecutive IPSCs for each cell). The mean of all measurements is shown as black line. The IPSC rise time is significantly prolonged in Prnp^{0/0} mice compared to wild-type mice ($p = 0.001$, t -test according to Welch). No significant differences were found among the rise times of wild-type, Tg20, and Tg35 cells. (E), Means of the decay time of IPSCs in wildtype, Prnp^{0/0}, Tg20 and Tg35. There are no differences among these mouse lines.

potassium channels, which may be related to a disturbed intracellular calcium homeostasis. This concept is based on findings by Whatley et al. (36) that indicated an effect of recombinant PrP^C on the intracellular calcium concentration




Fig. 4. Copper concentration in synaptosomes correlates with PrP^C expression. The copper concentrations in whole-brain homogenates and synaptosomal fractions from wild-type (open columns), Prnp^{0/0} (black columns), and Tg20 (gray columns) mice were studied by atomic absorption spectroscopy. Shown are the mean and SE of the arithmetic mean of 3–7 preparations from each of five brains of age-matched (2 ± 0.4 mo) female animals of various lines. The copper concentration related to protein concentration in whole-brain homogenates shows no significant differences among wild-type, Prnp^{0/0} and Tg20 mice, but the synaptosomal fraction shows a significant reduction of copper in Prnp^{0/0} mice compared to wildtype and Tg20 mice ($p = 0.03$; t -test).

in synaptosomes. Indeed, a study of calcium-activated potassium currents in Purkinje cells of Prnp^{0/0} mice showed a reduced amplitude of these currents (Herms et al., in preparation). Further investigations of transgenic animals which were Prnp reconstituted on the Prnp^{0/0} background (Tg35, Tg20) showed that loss of PrP^C expression in Purkinje cells is responsible for this finding (37). Thus, a reconstitution of the amplitude of calcium-activated potassium conductances was observed in a transgenic line that shows overexpression of PrP^C in all neurons (Tg35), whereas a transgenic line that overexpresses PrP^C in all neurons but Purkinje cells, showed no reconstitution of the amplitude. The subsequent microfluorometric investigation of the intracellular calcium homeostasis in Prnp^{0/0} mice confirmed that the reduction of calcium-activated potassium currents is probably caused by reduced calcium release from intracellular calcium-sensitive calcium stores (37) (Herms et al., in preparation).

3. The Role of Copper

The cause of the observed electrophysiological alterations in Prnp^{0/0} mice is not yet known. They may be related to the decreased copper concentration in synaptic membranes of Prnp^{0/0} mice (Fig. 4; [23]). The N-terminus of PrP^C has

a highly conserved octapeptide repeat sequence (PHGGGWGQ) $\times 4$ (38), whose possible copper-binding properties were first shown by Hornshaw et al. (39,40) and later by Miura et al. (41). The recombinant N-terminus of PrP^C from amino acid 23 to 98 (PrP 23–98) shows a cooperative binding of 5–6 copper ions (42). Half-maximal cooperative copper binding of PrP23–98 is in the micromolar range (5.9 μM). Further investigations, using synthetic octapeptides (43) confirmed cooperative copper binding by PrP^C.

The significant decrease of synaptosomal copper concentration in Prnp^{0/0} mouse synaptosomes (Fig. 4) may be caused by a decreased reuptake of copper released into the synaptic cleft during synaptic vesicle release, since the difference in the synaptosomal copper concentration between Prnp^{0/0} mice and wildtype mice seems to be too large to be explained solely by the loss of copper bound to PrP^C. In addition, one would then also expect differences in the copper concentration of the crude homogenate in wildtype, Tg20 and Prnp^{0/0} mice (Fig. 4). The findings may therefore be explained by a dysregulation of the copper concentration in the brains of Prnp^{0/0} mice caused by loss of PrP^C.

In addition to the decreased synaptosomal copper concentration, a number of further changes were observed that indicated a biological function of copper binding by PrP^C. Thus, significant differences between Prnp^{0/0} mice and wildtype mice were found in inhibitory synaptic transmission in the presence of copper (42). The application of copper elicited a significant reduction of the mean amplitude of spontaneous inhibitory postsynaptic GABA_A receptor-mediated currents in Purkinje cells of Prnp^{0/0} mice at a concentration of 2 μM Cu²⁺, whereas this concentration showed no effect on the IPSCs of the wildtype mice. Because it is well known that the GABA_A receptor is functionally disturbed at a concentration of copper in the range of 1 μM (44), this finding indicates that differences between Prnp^{0/0} and wildtype mice may be caused by missing copper buffering in the synaptic cleft by PrP^C.

It is difficult to verify whether the loss of PrP^C indeed leads to a reduction of the amount of copper located at the synaptic plasma membrane in intact synapses because direct synaptic measurements in vivo are not possible at present. We used an indirect approach to assess the problem of copper binding at the synapse, by studying the effect of hydrogen peroxide on inhibitory synaptic transmission (23). H₂O₂ is known to alter the probability of synaptic vesicle release by reacting with metal ions, particularly iron and copper at the presynapse, by increasing the presynaptic calcium concentration. By performing patch-clamp measurements on cerebellar slice preparations of wildtype, Prnp^{0/0} and PrP^C reconstituted transgenic mice, we observed the effect of 0.01% H₂O₂ on the frequency of spontaneous IPSCs in Purkinje cells correlate with the amount of PrP^C expressed in the presynaptic neuron (Fig. 5). This indicates that the amount of copper at the synapse may indeed be PrP^C-related.




Fig. 5. Enhancement of inhibitory synaptic activity by hydrogen peroxide is related to the amount of PrP^C at the presynaptic plasma membrane. Effect of 0.01% H₂O₂ on the frequency of inhibitory postsynaptic currents in the different mouse lines. Each point represents the mean \pm SEM sIPSC frequency in 1-min intervals normalized to the values before H₂O₂ application of wild-type ($n = 14$), Prnp^{0/0} ($n = 21$), Tg35 ($n = 15$) and Tg20 ($n = 4$) mouse Purkinje cells. The bar indicates the time during which H₂O₂ was applied. The application of H₂O₂ led to a marked enhancement of synaptic activity in wild-type mice, there is no comparable effect in Prnp^{0/0} mice. In transgenic mice that overexpress PrP^C on a Prnp^{0/0} background in all neurons (Tg35), the sIPSC frequency increase after H₂O₂ application is rescued. Also, PrP^C-reconstituted mice, which express PrP^C in cerebellar interneurons, but not in Purkinje cells (Tg20), show a rescue, indicating that the presynaptic PrP^C expression is important for the rescue of the H₂O₂ effect on IPSC frequency.

It remains to be shown whether buffering of copper released during synaptic vesicle release, which prevents or minimizes unspecific binding of copper to other proteins, is the primary function of PrP^C (**Fig. 6**). Alternatively, the binding of copper to PrP^C may primarily serve the reuptake of copper into the presynapse by endocytosis of PrP^C (45,46) or may be of structural importance for the N-terminus of PrP^C (47).

The hypothesis of a functional re-uptake of copper in the synaptic cleft by the prion protein (**Fig. 6**) may explain electrophysiological findings in Prnp^{0/0} mice, which, on first glance, seem contradictory. A slight increase of extracellular copper concentration, caused by decreased or missing copper buffering in




Fig. 6. Hypothetical model showing a possible function of copper binding by PrP^C at the synaptic plasma membrane. The prion protein is attached to the presynaptic plasma membrane (1) (23), where its N-terminal moiety (2) binds free copper that is released into the synaptic cleft with synaptic vesicle release (3) (58,59). There is an endocytotic uptake of PrP^C into the presynapse (4) (45,46) where PrP^C-bound copper is released, possibly induced by endosomal pH changes (5) (43). Thus PrP^C serves to keep the copper concentration in the presynaptic cytosol and the synaptic cleft constant despite copper losses during synaptic vesicle release (3).

the synaptic cleft in Prnp^{0/0} mice, may cause a decrease in the conductance of voltage-activated calcium channels and a change in the kinetics of the GABA_A receptor. Thus, the conductance of the GABA_A receptor and voltage-activated calcium channels, which modulate intracellular calcium homeostasis is clearly disturbed by copper concentrations of 1–10 μ M (44,48). This would explain the alteration of the intracellular calcium homeostasis in Prnp^{0/0} mice, changes in the conductance of calcium-related ion currents, and changes in GABA_A receptor-related inhibitory postsynaptic currents observed under certain conditions. Reduced LTP in Prnp^{0/0} mice may be explained by this hypothesis, as well. As shown by Doreulee et al. (49), LTP is blocked by concentrations of free copper as low as 1 μ M. Changes in the circadian rhythm observed by Tobler et al. (12,13) in Prnp^{0/0} mice could be related to a disturbed copper uptake and a decreased activity of copper-dependent enzymes, since the synthesis of melatonin, which is important in the regulation of circadian rhythms (50), is regulated by the copper-dependent enzyme monamine oxidase (51). Also, the activity of two other copper-dependent enzymes, the Cu/Zn superox-

ide dismutase and the glutathione reductase have been found to be altered in PrP^{0/0} mice (52,53).

4. Conclusion

In summary, our studies have shown that PrP^C binds copper cooperatively and with high affinity. In the brain highest concentrations of PrP^C are found at synapses. Synaptosomes of Prnp^{0/0} mice demonstrate a strong reduction of copper concentration. Copper binding by PrP^C in the synaptic cleft has a significant influence on synaptic transmission. It remains to be shown whether additional phenotypes observed in Prnp^{0/0} mice result from decreased copper binding or from a disturbance of copper distribution in the absence of PrP^C.

Acknowledgment

This work was supported by the BMBF (German Federal Ministry of Science and Technology) grant KI9461/8, the Deutsche Forschungsgemeinschaft (grant Kr. 1561/21), and Sonderforschungsbereich 406, as well as the Wilhelm-Sander-Stiftung (grant 9343008). We thank Charles Weissmann (University of Zürich) for Prnp^{0/0}, Tg35 and Tg20 mice.

References.

1. Stahl, N., Borchelt, D. A., Hsiao, K., and Prusiner, S. B. (1987) Scrapie prion protein contains a phosphatidylinositol glycolipid. *Cell* **51**, 229–240.
2. Oesch, B., Teplow, T. B., Stahl, N., Serban, D., Hood, L. E., and Prusiner, S. B. (1990) Identification of cellular proteins binding to the scrapie prion protein. *Biochemistry* **29**, 5848–5855.
3. Kurschner, C., and Morgan, J. I. (1995) The cellular prion protein (PrP) selectively binds to Bcl-2 in the yeast two-hybrid system. *Mol. Brain Res.* **30**, 165–167.
4. Kurschner, C., and Morgan, J. I. (1996) Analysis of interaction sites in homo- and heteromeric complexes containing Bcl-2 family members and the cellular prion protein. *Mol. Brain Res.* **37**, 249–258.
5. Edenhofer, F., Rieger, R., Famulok, M., Wendler, W., Weiss, S., and Winnacker, E.-L. (1996) Prion protein PrP^c interacts with molecular chaperones of the Hsp60 family. *J. Virol.* **70**, 4724–4728.
6. Yehiely, F., Bamborough, P., DaCosta, M., Perry, B. J., Thinakaran, G., Cohen, F. E., et al. (1997) Identification of candidate proteins binding to prion protein. *Neurobiol. Dis.* **3**, 339–355.
7. Rieger, R., Edenhofer, F., Lasmézas, C. I., and Weiss, S. (1998) The human 37-kDa laminin receptor precursor interacts with the prion protein in eukaryotic cells. *Nature Med.* **3**, 1383–1388.
8. Martins, V. R., Graner, E., Garcia-Abreu, J., deSouza, S. J., Mercadante, A. F., Veiga, S. S., et al. (1998) Complementary hydrophathy identifies a cellular prion protein receptor. *Nature Med.* **3**, 1376–1382.

9. Harris, D. A., Falls, D. L., Johnson, F. A., and Fischbach, G. D. (1991) A prion-like protein from chicken brain copurifies with an acetylcholine receptor-inducing activity. *Proc. Natl. Acad. Sci. USA* **88**, 7664–7668.
10. Büeler, H., Fischer, M., Lang, Y., Bluethmann, H., Lipp, H.-P., DeArmond, S. J., et al. (1992) Normal development and behaviour of mice lacking the neuronal cell-surface PrP protein. *Nature* **356**, 577–582.
11. Brenner, H. R., Herczeg, A., and Oesch, B. (1992) Normal development of nerve-muscle synapses in mice lacking the prion protein gene. *Proc. Roy. Soc. Lond. (Biol.)* **250**, 151–156.
12. Tobler, I., Gaus, S. E., Deboer, T., Achermann, P., Fischer, M., Rüllicke, T., et al. (1996) Altered circadian activity rhythms and sleep in mice devoid of prion protein. *Nature* **380**, 639–642.
13. Tobler, I., Deboer, T., and Fischer, M. (1997) Sleep and sleep regulation in normal and prion protein-deficient mice. *J. Neurosci.* **17**, 1869–1879.
14. Walz, W., Amaral, O. B., Rockenback, I. C., Roesler, R., Izquierdo, I., Cavalheiro, E. A., et al. (1999) Increased sensitivity to seizures in mice lacking cellular prion protein. *Epilepsia* **40**, 1679–1682.
15. Manson, J. C., Clarke, A. R., McBride, P. A., McConnell, I., and Hope, J. (1994) PrP gene dosage determines the timing but not the final intensity or distribution of lesions in scrapie pathology. *Neurodegeneration* **3**, 331–340.
16. Tremblay, P., Meiner, Z., Galou, M., Heinrich, C., Petromilli, C., Lisse, T., et al. (1998) Doxycycline control of prion protein transgene expression modulates prion disease in mice. *Proc. Natl. Acad. Sci. USA* **95**, 12,580–12,585.
17. Sakaguchi, S., Katamine, S., Nishida, N., Moriuchi, R., Shigematsu, K., Sugimoto, T., et al. (1996) Loss of cerebellar Purkinje cells in aged mice homozygous for a disrupted PrP gene. *Nature* **380**, 528–531.
18. Nishida, N., Tremblay, P., Sugimoto, T., Shigematsu, K., Shirabe, S., Petromilli, C., et al. (1999) A mouse prion protein transgene rescues mice deficient for the prion protein gene from Purkinje cell degeneration and demyelination. *Lab. Invest.* **79**, 689–697.
19. Moore, R. C., Hope, J., McBride, P. A., McConnell, I., Selfridge, J., Melton, D. W., et al. (1998) Mice with gene targeted prion protein alterations show that *Prnp*, *Sinc* and *Prni* are congruent. *Nature Genet.* **18**, 118–125.
20. Weissmann, C. (1996) PrP effects clarified. *Curr. Biol.* **6**, 1369
21. Moore, R. C., Lee, I. Y., Silverman, G. L., Harrison, P. M., Strome, R., Heinrich, C., et al. (1999) Ataxia in prion protein (PrP)-deficient mice is associated with upregulation of the novel PrP-like protein doppel. *J. Mol. Biol.* **292**, 797–818.
22. Chishti, M. A., Strome, R., Carlson, G. A., and Westaway, D. (1997) Syrian hamster prion protein (PrP^C) is expressed in photoreceptor cells of the adult retina. *Neurosci. Lett.* **234**, 11–14.
23. Herms, J., Tings, T., Gall, S., Madlung, A., Giese, A., Siebert, H. P. et al. (1999) Evidence of presynaptic location and function of the prion protein. *J. Neurosci.* **19**, 8866–8875.

24. DeArmond, S. J., Mobley, W. C., DeMott, D. L., Barry, R. A., Beckstead, J. H., and Prusiner, S. B. (1987) Changes in the localization of brain prion proteins during scrapie infection. *Neurology* **37**, 1271–1280.
25. Piccardo, P., Safar, J., Ceroni, M., Gajdusek, D. C., and Gibbs Jr., C.J. (1990) Immunohistochemical localization of prion protein in spongiform encephalopathies and normal brain tissue. *Neurology* **40**, 518–522.
26. Bendheim, P. E., Brown, H. R., Rudelli, R. D., Scala, L. J., Goller, N., L., Wen, G. Y., et al. (1992) Nearly ubiquitous tissue distribution of the scrapie agent precursor protein. *Neurology* **42**, 149–156.
27. Fournier, J.-G., Escaig-Haye, F., De Villemeur, T. B., and Robain, O. (1995) Ultrastructural localization of cellular prion protein (PrP_C) in synaptic boutons of normal hamster hippocampus. *C. R. Acad. Sci. Paris, Sciences de la vie/Life sciences* **318**, 339–344.
28. Salès, N., Rodolfo, K., Hässig, R., Faucheux, B., Di Giamberardino, L., and Moya, K. L. (1998) Cellular prion protein localization in rodent and primate brain. *Eur. J. Neurosci.* **10**, 2464–2471.
29. Collinge, J., Whittington, M. A., Sidle, K. C. L., Smith, C. J., Palmer, M. S., Clarke, A. R., et al. (1994) Prion protein is necessary for normal synaptic function. *Nature* **370**, 295–297.
30. Manson, J. C., Hope, J., Clarke, A. R., Johnston, A., Black, C., and MacLeod, N. (1995) PrP gene dosage and long term potentiation. *Neurodegeneration* **4**, 113–115.
31. Lledo, P. M., Tremblay, P., DeArmond, S. J., Prusiner, S. B., and Nicoll, R. A. (1996) Mice deficient for prion protein exhibit normal neuronal excitability and synaptic transmission in the hippocampus. *Proc. Natl. Acad. Sci. USA* **93**, 2403–2407.
32. Herms, J. W., Kretzschmar, H. A., Titz, S., and Keller, B. U. (1995) Patch-clamp analysis of synaptic transmission to cerebellar Purkinje cells of prion protein knockout mice. *Eur. J. Neurosci.* **7**, 2508–2512.
33. Llano, I., Marty, A., Armstrong, C. M., and Konnerth, A. (1991) Synaptic- and agonist-induced excitatory currents of Purkinje cells in rat cerebellar slices. *J. Physiol.* **434**, 183–213.
34. Fischer, M., Rülcke, T., Raeber, A., Sailer, A., Moser, M., Oesch, B., et al. (1996) Prion protein (PrP) with amino-proximal deletions restoring susceptibility of PrP knockout mice to scrapie. *EMBO J.* **15**, 1255–1264.
35. Colling, S. B., Collinge, J., and Jefferys, J. G. R. (1996) Hippocampal slices from prion protein null mice: disrupted Ca²⁺ activated K⁺ currents. *Neurosci. Lett.* **209**, 49–52.
36. Whatley, S. A., Powell, J. F., Politopoulou, G., Campbell, I. C., Brammer, M. J., and Percy, N. S. (1995) Regulation of intracellular free calcium levels by the cellular prion protein. *Neuroreport* **6**, 2333–2337.
37. Tings, T., Herms, J., Dunker, S., and Kretzschmar, H., A. (1998) Rescue of Ca²⁺-activated K⁺-currents in cerebellar Purkinje cells of transgenic mice overexpressing the prion protein on a knock-out background. *Clin. Neuropathol.* **17**, 254

38. Schätzl, H. M., Da Costa, M., Taylor, L., Cohen, F. E., and Prusiner, S. B. (1995) Prion protein gene variation among primates. *J. Mol. Biol.* **245**, 362–374.
39. Hornshaw, M. P., McDermott, J. R., and Candy, J. M. (1995) Copper binding to the N-terminal tandem repeat regions of mammalian and avian prion protein. *Biochem. Biophys. Res. Commun.* **207**, 621–629.
40. Hornshaw, M. P., McDermott, J. R., Candy, J. M., and Lakey, J. H. (1995) Copper binding to the N-terminal tandem repeat region of mammalian and avian prion protein: Structural studies using synthetic peptides. *Biochem. Biophys. Res. Commun.* **214**, 993–999.
41. Miura, T., Hori-i, A., and Takeuchi, H. (1996) Metal-dependent α -helix formation promoted by the glycine-rich octapeptide region of prion protein. *FEBS Lett.* **396**, 248–252.
42. Brown, D. R., Qin, K., Herms, J. W., Madlung, A., Manson, J., Strome, R., et al. (1997) The cellular prion protein binds copper *in vivo*. *Nature* **390**, 684–687.
43. Viles, J. H., Cohen, F. E., Prusiner, S. B., Goodin, D. B., Wright, P. E., and Dyson, H. J. (1999) Copper binding to the prion protein: Structural implications of four identical cooperative binding sites. *Proc. Natl. Acad. Sci. USA* **96**, 2042–2047.
44. Sharonova, I. N., Vorobjev, V. S., and Haas, H. L. (1998) High-affinity copper block of GABA_A-receptor-mediated currents in acutely isolated cerebellar Purkinje cells of the rat. *Eur. J. Neurosci.* **10**, 522–528.
45. Shyng, S. L., Heuser, J. E., and Harris, D. A. (1994) A glycolipid-anchored prion protein is endocytosed via clathrin-coated pits. *J. Cell Biol.* **125**, 1239–1250.
46. Pauly, P. C. and Harris, D. A. (1998) Copper stimulates endocytosis of the prion protein. *J. Biol. Chem.* **273**, 33,107–33,110.
47. Marcotte, E. M. and Eisenberg, D. (1999) Chicken prion tandem repeats form a stable, protease-resistant domain. *Biochemistry* **38**, 667–676.
48. Nam, S. C. and Hockberger, P. E. (1992) Divalent ions released from stainless steel hypodermic needles reduce neuronal calcium currents. *Pflugers Arch.* **420**, 106–108.
49. Doreulee, N., Yanovsky, Y., and Haas, H. L. (1997) Suppression of long-term potentiation in hippocampal slices by copper. *Hippocampus* **7**, 666–669.
50. McArthur, A. J., Gillette, M. U., and Prosser, R. A. (1991) Melatonin directly resets the rat suprachiasmatic circadian clock in vitro. *Brain Res.* **565**, 158–161.
51. Oxenburg, G. F., and Requintin, P. J. (1998) The effect of MAO-A inhibition and cold-immobilisation stress on N-acetylserotonin and melatonin in SHR and WKY rats. *J. Neural. Transmission Suppl.* **52**, 333–336.
52. Brown, D. R., Schulz-Schaeffer, W. J., Schmidt, B., and Kretzschmar, H. A. (1997) Prion protein-deficient cells show altered response to oxidative stress due to decreased SOD-1 activity. *Exp. Neurol.* **146**, 104–112.
53. White, A. R., Collins, S. J., Maher, F., Jobling, M. F., Stewart, L. R., Thyer, J. M., et al. (1999) Prion protein-deficient neurons reveal lower glutathione reductase activity and increased susceptibility to hydrogen peroxide toxicity. *Am. J. Pathol.* **155**, 1723–1730.

54. Huttner, W. B., Schiebler, W., Greengard, P., and De Camilli, P. (1983) Synapsin I (protein I), a nerve terminal-specific phosphoprotein. III. Its association with synaptic vesicles studied in a highly purified synaptic vesicle preparation. *J. Cell Biol.* **96**, 1374–1388.
55. Krasemann, S., Groschup, M. H., Harmeyer, S., Hunsmann, G., and Bodemer, W. (1996) Generation of monoclonal antibodies against human prion proteins in PrP^{0/0} mice. *Mol. Medicine* **2**, 725–734.
56. Brose, N., Petrenko, A. G., Südhof, T. C., and Jahn, R. (1992) Synaptotagmin: a calcium sensor on the synaptic vesicle surface. *Science* **256**, 1021–1025.
57. Brose, N., Huntley, G. W., Stern-Bach, Y., Sharma, G., Morrison, J. H., and Heinemann, S. F. (1994) Differential assembly of coexpressed glutamate receptor subunits in neurons of rat cerebral cortex. *J. Biol. Chem.* **269**, 16,780–16,784.
58. Kardos, J., Kovács, I., Hajós, F., Kálmán, M. and Simonyi, M. (1989) Nerve endings from rat brain tissue release copper upon depolarization. A possible role in regulating neuronal excitability. *Neurosci. Lett.* **103**, 139–144.
59. Hartter, D. E. and Barnea, A. (1988) Evidence for release of copper in the brain: depolarisation-induced release of newly taken-up 67 copper. *Synapse* **2**, 412–415.



<http://www.springer.com/978-0-89603-924-7>

Molecular Pathology of the Prions

Baker, H.F. (Ed.)

2001, XII, 279 p., Hardcover

ISBN: 978-0-89603-924-7

A product of Humana Press