

Preface

One of the fundamental problems in any area of mathematics is to determine the distinct variants of an object under consideration. As for complex-functional analysis, one is interested, for example, in studying the equivalent representations of the conformally invariant classes of holomorphic functions. This problem is addressed here for the holomorphic \mathcal{Q} classes.

For $p \in [0, \infty)$ and dm – the element of the two dimensional Lebesgue measure, we say that f , a holomorphic function in the unit disk \mathbf{D} , is of the class \mathcal{Q}_p provided

$$E_p(f) = \sup \left\{ \left(\int_{\mathbf{D}} |f'(z)|^2 \left(\log \left| \frac{1 - \bar{w}z}{w - z} \right| \right)^p dm(z) \right)^{1/2} : w \in \mathbf{D} \right\} < \infty.$$

It is clear that each \mathcal{Q}_p can serve as a sample of the conformally invariant classes in the sense of: $E_p(f \circ \sigma) = E_p(f)$ for all $f \in \mathcal{Q}_p$ and $\sigma \in \text{Aut}(\mathbf{D})$ – the group of all conformal automorphisms of \mathbf{D} .

The goal of this monograph is to bring the major features of \mathcal{Q}_p to light, in particular, to characterize \mathcal{Q}_p in different terms. More precisely:

Chapter 1 contains some (most) basic properties of \mathcal{Q}_p such as Möbius boundedness, image area and higher derivatives. The aim is to show that the classes \mathcal{Q}_p , $p \in (0, 1)$ are of independent interest.

Chapter 2 discusses the problem of how a \mathcal{Q}_p can be embedded into a Bloch-type space and vice versa. This problem will be solved by considering boundedness and compactness of a composition operator acting between two spaces.

Chapter 3 describes the coefficients of either Taylor or random series of functions from \mathcal{Q}_p . In particular, we will see that there is a big difference between the cases: $p \in (0, 1)$ and $p = 1$.

Chapter 4 exhibits a geometric way to understand \mathcal{Q}_p , that is, p -Carleson measure characterization of \mathcal{Q}_p . This simple but important property induces certain deep relations between \mathcal{Q}_p and the mean Lipschitz spaces, as well as the Besov spaces which are conformally invariant, too.

Chapter 5 characterizes the inner and outer functions in \mathcal{Q}_p by means of p -Carleson measure and other two conformally invariant measures: Poisson measure and hyperbolic measure.

Chapter 6 gives the boundary value behavior of a \mathcal{Q}_p -function for $p \in (0, 1)$. This allows us to study \mathcal{Q}_p via those non-holomorphic functions on the unit

circle \mathbf{T} and even on the exterior of the unit disk $\mathbf{C} \setminus \mathbf{D}$, and hence leads to a consideration of harmonic analysis.

Chapter 7 explores a list of properties of $\mathcal{Q}_p(\mathbf{T})$ (i.e. \mathcal{Q} class on \mathbf{T}). Specially, the $\mathcal{Q}_p(\mathbf{T})$ -solutions of the $\bar{\partial}$ -equation produce a decomposition of \mathcal{Q}_p through the bounded functions on \mathbf{T} . As applications, the corona theorem and interpolation theorem related to \mathcal{Q}_p are established.

Chapter 8 deals with a localization of $\mathcal{Q}_p(\mathbf{T})$ based on the dyadic partitions of all subarcs of \mathbf{T} . The results enable us to recognize \mathcal{Q}_p from mean oscillation to dyadic model, and finally to wavelet basis.

The exposition is at as elementary a level as possible, and it is intended to be accessible to graduate students with a basic knowledge of complex-functional-real analysis. The material of this monograph has been collected from a series of talks that I gave over the past six years most in Canada, China, Finland, Germany, Greece and Sweden, but also from a lecture course at University of La Laguna in the fall semester of 1999. The selection of topics is rather arbitrary, but reflects the author's preference for the analytic approach. There is no attempt to cover all recent advances (for instance, \mathcal{Q} classes of higher dimensions), and yet, it is hoped that the reader will be intrigued by this monograph and will, at some point, read the notes presented at the end of each chapter as well as the papers listed in the references, and proceed to a further research.

Here, I owe a great debt of gratitude to the many people who assisted me with this work. R. Aulaskari and M. Essén read the whole manuscript, caught a number of errors and offered many helpful suggestions. G. Dafni, P. Gauthier and K. J. Wirths read parts of the manuscript and contributed significantly, by valuable queries and comments, to the accuracy of the final version. M. Anderson, S. Axler, H. Carlsson, D. C. Chang, Y. He, S. Janson, H. Jarchow, F. Jafari, H. Kisilevsky, L. Lindahl, A. Nicolau, J. Peetre, L. Peng, F. Pérez-González, H. Proppe, S. Ruscheweyh, W. Sander, A. Siskakis, W. Smith, K. Sten, D. Stegenga, M. Wong, K. Xiong and G. Zhang made friendly advice and warm encouragement.

The following also gave aid and comfort. The Alexander von Humboldt Foundation, Germany and the Swedish Institute, Sweden supported my work on this book. The Institute of Mathematics at Technical University of Braunschweig supplied the computer facilities, and moreover its faculty member H. Weiss kindly helped me create LaTeX working directory and taught me much knowledge about computer. The Department of Mathematics and Statistics at Concordia University provided a good place to carry my writing and revising through to the end. Without their help I would not have gone ahead with publishing this book.

I am grateful to the editors of Springer-Verlag for accepting this monograph for publication in the LNM series, as well as to S. Zoeller for the efficient handling of the editing.

Finally, I want to express my special thanks to my wife, Xianli, and my son, Sa, for their understanding and support.

Montreal, July 2001

Jie Xiao



<http://www.springer.com/978-3-540-42625-7>

Holomorphic Q Classes

Xiao, J.

2001, X, 104 p., Softcover

ISBN: 978-3-540-42625-7