

2

Diversity, Sources, and Detection of Human Bacterial Pathogens in the Marine Environment

Janelle R. Thompson, Luisa A. Marcelino, and Martin F. Polz

2.1. INTRODUCTION

Disease outbreaks in marine organisms appear to be escalating worldwide (Harvell et al., 1999, 2002) and a growing number of human bacterial infections have been associated with recreational and commercial uses of marine resources (Tamplin, 2001). Whether these increases reflect better reporting or global trends is a subject of active research (reviewed in Harvell et al., 1999, 2002; Rose et al., 2001; Lipp et al., 2002); however, in light of heightened human dependence on marine environments for fisheries, aquaculture, waste disposal, and recreation, the potential for pathogen emergence from ocean ecosystems requires investigation.

A surprising number of pathogens have been reported from marine environments and the probability of their transmission to humans is correlated to factors that affect their distribution. Both indigenous and introduced pathogens can be the cause of illness acquired from marine environments and their occurrence depends on their ecology, source, and survival. To judge the risk from introduced pathogens, levels of indicator organisms are routinely monitored at coastal sites. However, methods targeting specific pathogens are increasingly used and are the only way to judge or predict risk associated with the occurrence of indigenous pathogen populations.

In this chapter, we review the recognized human pathogens that have been found in associations with marine environments (Section 2.2), the potential routes of transmission of marine pathogens to humans, including seafood consumption, seawater exposure (including marine aerosols), and marine zoonoses (Section 2.3), and we discuss the methods available to assess the public-health risks associated with marine pathogens (Sections 2.4 and 2.5).

Janelle R. Thompson, Luisa A. Marcelino, and Martin F. Polz • Department of Civil and Environmental Engineering, Massachusetts Institute of Technology, 77 Massachusetts Avenue, Cambridge, MA 02139, USA.

2.2. DIVERSITY AND ECOLOGY

Our current knowledge of the diversity and ecology of bacterial pathogens associated with marine environments stems from (i) clinical accounts of marine-acquired illnesses, (ii) disease outbreaks of known etiology in marine animals, and (iii) testing of marine environments for the presence of pathogen populations. In particular, surveys of environmental microbial communities based on 16S ribosomal RNA (rRNA) gene sequence diversity have revealed a large number of organisms closely related to human pathogens; however, the public health risk of many of these pathogen-like populations remains unknown. This is largely due to a poorly defined relationship between clinical isolates and pathogen-like populations detected in the environment because many methods used to detect environment populations do not possess high enough resolution to discriminate virulent from harmless strains.

The genetic elements encoding virulence properties are not uniformly distributed among strains within a potentially pathogenic species. For marine pathogens, this has been explored in some detail in *Vibrio* species. Environmental populations of *Vibrio* are characterized by heterogeneous distributions of multiple virulence factors, combinations of which regulate the epidemic potential (e.g. Faruque et al., 1998; Karaolis et al., 1998; Chakraborty et al., 2000). Similarly, comparisons of the genomic diversity of clinical and environmental *Vibrio vulnificus* isolates suggest that seafood-borne human infections are established by a single highly virulent strain among coexisting genetically heterogeneous populations (Jackson et al., 1997). However, what leads to the occurrence of one strain over another remains poorly understood.

Whether environmental conditions select for strains possessing human virulence factors is an area of increased research (e.g. Tamplin et al., 1996; Jackson et al., 1997; Faruque et al., 1998; Chakraborty et al., 2000). Such factors may include attachment mechanisms to organic matter, motility, secretion of lytic compounds, and the ability to grow rapidly under nutrient-replete conditions. Transfer of virulence properties between different species has been observed (Faruque et al., 1999; Boyd et al., 2000), and specific virulence factors (e.g., hemolysins, toxins, attachment pili) may be borne on mobile genetic elements. Thus, environmental interaction may confer enhanced pathogenicity on a subset of an environmental population. In general, the marine environment may be a powerful incubator for new combinations of virulence properties due to the extremely large overall population size of bacterial populations and efficient mixing timescales. These natural phenomena may be further enhanced by human activity such as increased sewage input and ballast water transport (Ruiz et al., 2000) both of which introduce microbial species across geographical barriers.

2.2.1. Pathogenic Species

The known diversity of human pathogens in the ocean continues to expand as the virulence of emerging pathogens is recognized. Pathogens associated with marine environments and their observed routes of transmission to humans are presented in Table 2.1. Of the 23 lineages currently characterized within the domain Bacteria by 16S rRNA phylogeny (Cole et al., 2003), six harbor human pathogens, and of these six, all lineages contain strains found as human and/or animal pathogens in marine environments (i.e., the Bacteroides-Flavobacterium group (Bernardet, 1998), the Spirochetes, the Gram-positive Bacteria, the Chlamydia (Johnson & Lepennec, 1995; Kent et al., 1998), the Cyanobacteria (Carmichael, 2001), and the Proteobacteria) (see also references in Table 2.1).

Table 2.1. Human-pathogenic bacteria detected in marine environments.

Genus	Species ^a	Hosts of marine disease		Observed routes of human infection ^b				Estimated infectious dose ^c	Human syndrome	References
		Marine animals								
		Humans	Seafood	Sea water	Zoonoses	Aerosols				
<i>Acinetobacter</i>	<i>A. calcoaceticus</i>	ND	X				X		Sepsis, meningitis, pneumonia (nonmarine acquired)	(Austin et al., 1979; Grimes et al., 1984; Grimes, 1991)
<i>Aeromonas</i>	<i>A. hydrophila</i>	X		X	X			10 ⁷ and 10 ⁹ cells in 2 of 57 tests (Morgan, 1985)	GI, sepsis, wound infection	(Morgan et al., 1985; Chowdhury et al., 1990; Ashbolt et al., 1995; Jones & Wilcox, 1995; Caudell & Kuhn, 1997; Fiorentini et al., 1998; Dumontet et al., 2000)
	<i>A. caviae</i>	X		X	X				Wound infection	(Ashbolt et al., 1995; Jones & Wilcox, 1995; Dumontet et al., 2000)
	<i>A. sobria</i>	X		X	X				Wound infection	(Ashbolt et al., 1995; Jones & Wilcox, 1995; Itoh et al., 1999; Dumontet et al., 2000)
<i>Brucella</i>	<i>B. maris</i>	X	X			X			Neurobrucellosis, brucellosis	(Corbel, 1997; Brew et al., 1999; Foster et al., 2002; Sohn et al., 2003)
<i>Burkholderia</i>	<i>B. pseudomallei</i> **	ND	X						Melioidosis (Nonmarine acquired)	(Hicks et al., 2000)
<i>Campylobacter</i>	<i>C. lari</i> *	X		X					GI	(Endtz et al., 1997)
	<i>C. jejuni</i> *	X		X				<500 cells (**)	GI	(Abeyta et al., 1993)
<i>Clostridium</i>	<i>C. botulinum</i> (type E)	X		X				0.1-lug toxin (*) 10 ⁵ cells (**)	Botulism, GI	(Huss, 1980; Weber et al., 1993)
(continued)										

(continued)

Table 2.1. (Continued)

Hosts of marine disease		Observed routes of human infection ^b				Estimated infectious dose ^c	Human syndrome	References
		Humans	Marine animals	Seafood	Sea water	Zoonoses	Aerosols	
Genus	Species ^a							
		X	X	X				(Feldhusen, 2000; Aschfalk & Muller, 2001)
<i>Edwardsiella</i>	<i>E. tarda</i>	X			X			(Kusuda & Kawai, 1998; Slaven et al., 2001)
<i>Enterobacter</i>	<i>E. cloacae</i>	ND						(Salas & Geesey, 1983; Grimes, 1991)
<i>Erysipelothrix</i>	<i>E. rhusiopathiae</i>	X			X	X		(Brooke & Riley, 1999; Fidalgo et al., 2000; Lehane & Rawlin, 2000)
<i>Escherichia</i>	<i>E. coli</i> *	X	X	X	X			(Kueh et al., 1992; Raidal et al., 1998; Feldhusen, 2000)
<i>Francisella</i>	<i>F. philomiragia</i>	X			X		X	(Wenger et al., 1989; Ender & Dolan, 1997)
<i>Halomonas</i>	<i>H. venusta</i>	X			X	X		(von Graevenitz et al., 2000)
<i>Klebsiella</i>	<i>K. pneumoniae</i>	X			X		X	(Kueh et al., 1992; Ritter et al., 1993; Ender & Dolan, 1997)
<i>Lactococcus</i>	<i>K. oxytoca</i>	ND		X				(Lopez-Sabater et al., 1996)
	<i>L. garvieae</i>	ND	X					(Fefer et al., 1998; Kusuda & Kawai, 1998) (Ravelo et al., 2003)
<i>Legionella</i>	<i>L. pneumophila</i>	ND					X	(Fliermans et al., 1981; Ortizroque & Hazen, 1987; Grimes, 1991)
	<i>L. bozemanii</i>	X			X		X	(Losonsky, 1991)

<i>Leptospira</i>	<i>L. interrogans</i> **	X	X	X	X	Wound, Respiratory, Leptospirosis (primarily freshwater)	(Thomas & Scott, 1997; Tryland, 2000; Levett, 2001; Arzouni et al., 2002)
	Spp.**	ND	X			Leptospirosis (Nonmarine acquired)	(Gulland et al., 1996; Levett, 2001; Colagross-Schouten et al., 2002)
<i>Listeria</i>	<i>L. monocytogenes</i> *	X		X		Flu-like symptoms	(Colburn et al., 1990; Dillon et al., 1994)
<i>Morganella</i>	<i>M. morganii</i>	ND				Histamine production	(Lopez-Sabater et al., 1996)
<i>Mycobacterium</i>	<i>M. tuberculosis</i>	ND	X	X	10 cells (**)	Tuberculosis	(Bernardelli et al., 1996; Dobos et al., 1999; Lehane & Rawlin, 2000; Montali et al., 2001)
	<i>M. bovis</i>	X	X		X	Tuberculosis	(Thompson et al., 1993; Bernardelli et al., 1996)
	<i>M. marinum</i>	X	X	X	X	Wound, "fish tank granuloma"	(Dobos et al., 1999; De la Torre et al., 2001)
	<i>M. haemophilum</i>	X		X	X	Wound from coral injury	(Saubolle et al., 1996; Dobos et al., 1999; Smith et al., 2003)
<i>Mycoplasma</i>	<i>M. phocacebrale</i>	X	X		X	Skin, "seal finger"	(Stadtlander & Madoff, 1994; Baker et al., 1998)
<i>Photobacterium</i>	<i>P. damsela</i>	X	X	X	X	Wound, sepsis	(Fraser et al., 1997; Kusuda & Kawai, 1998; Rodgers & Furones, 1998; CDC, 1999; Barber & Swygert, 2000)
<i>Plesiomonas</i>	<i>P. shigelloides</i>	X		X		GI	(Gonzalez et al., 1999; Oxley et al., 2002; Chan et al., 2003)
<i>Pseudomonas</i>	<i>P. aeruginosa</i>	X		X		Skin, wound, ear infection, "diver's hand"	(Erickson et al., 1992; Ritter et al., 1993; Ahlen et al., 2000)

(continued)

Table 2.1 (Continued)

Hosts of marine disease		Observed routes of human infection ^b				Estimated infectious dose ^c	Human syndrome	References
		Humans	Marine animals	Seafood	Sea water	Zoonoses	Aerosols	
Genus	Species ^a							
<i>Rhodococcus</i>	<i>R. equi</i>	ND	X					(Prescott, 1991; Weinstock & Brown, 2002)
<i>Salmonella</i>	<i>S. enteritidis</i> **	X		X	X	X		(Dalsgaard, 1998; Polo et al., 1999; Tryland, 2000; Aschfalk et al., 2002)
	<i>S. spp</i> **	X	X	X	X	X	10 ² CFU/g varies (*)	(Tryland, 2000; Aschfalk et al., 2002)
<i>Serratia</i>	<i>S. liquefaciens</i>	ND	X				Sepsis	(Starliper, 2001) (Grohskopf et al., 2001)
<i>Shewanella</i>	<i>S. putrefaciens</i>	X		X			Wound infection, sepsis	(Dominguez et al., 1996; Iwata et al., 1999; Leong et al., 2000; Vogel et al., 2000; Pagani et al., 2003)
	<i>S. alga</i>	X		X			Sepsis, ear infection	(Nozue et al., 1992; Holt et al., 1997; Gram et al., 1999; Iwata et al., 1999)
<i>Shigella</i>	<i>S. dysenteriae</i> *	X		X	X		GI	(Kueh et al., 1992; Feldhusen, 2000)
<i>Staphylococcus</i>	<i>S. aureus</i> *	X		X	X		GI, wound, ear, skin infections	(Charoencra & Fujioka, 1993; Thomas & Scott, 1997; Feldhusen, 2000; Tryland, 2000)
							10 ⁵ –10 ⁶ CFU/g (*) (oral)	
<i>Streptococcus</i>	<i>S. iniae</i>	X	X			X	Skin, wound infections	(Lehane & Rawlin, 2000) (Thomas & Scott, 1997) (Weinstein, 2003)

<i>Vibrio</i>	<i>V. alginolyticus</i>	X	X	X	X	X	Wound/ear infections, Sepsis	(Howard & Bennett, 1993; CDC, 1999, 2000)
	<i>V. carchariae</i>	X	X		X	X	Wound infection	(Pavia et al., 1989; Lee et al., 2002; Nicolas et al., 2002)
	<i>V. cholerae O1</i>	X	X	X	X		GI, sepsis	(CDC, 1999; Lipp et al., 2002)
	<i>V. cholerae non-O1</i>	X	X	X	X	X	GI, wound/ear infection, sepsis	(Howard & Bennett, 1993; CDC, 1999, 2000)
	<i>V. cincinnatiensis</i>	X		X	X		Sepsis	(Brayton et al., 1986)
	<i>V. fluvialis</i>	X		X	X		GI	(Howard & Bennett, 1993; CDC, 1999, 2000)
	<i>V. furnissii</i>	X		X	X		GI	(Dalsgaard et al., 1997)
	<i>V. hollisae</i>	X		X	X		GI, sepsis	(Howard & Bennett, 1993; CDC, 1999, 2000)
	<i>V. metschnikovii</i>	X	X	X	X		GI	(Buck, 1991; CDC, 2000)
	<i>V. mimicus</i>	X		X	X	X	GI, wound/ear infection	(CDC, 1999, 2000)
	<i>V. parahaemolyticus</i>	X	X	X	X		GI, wound/ear infection, sepsis	(CDC, 1999, 2000)
	<i>V. vulnificus</i>	X	X	X	X	X	GI, wound/ear infection, sepsis	(Howard & Bennett, 1993; Howard & Burgess, 1993; CDC, 1999, 2000; Johnson & Arnett, 2001)
	<i>Y. enterocolitica*</i>	X		X	X		GI	(Feldhusen, 2000)

^a Marine-indigenous species, unless otherwise indicated; * marine-contaminant from anthropogenic or natural sources, ** marine source not determined.

^b Routes of human marine-acquired disease, except for aerosol inhalation where nonmarine aerosol transmission was noted.

^c Infectious doses are indicated as reported, either as total cells or as an environmental concentration. References indicated *Feldhusen (2000) or **PHAC (2001).

A large majority of known marine pathogens belong to the gamma-Proteobacteria. Within these, the genus *Vibrio* alone contains 11 recognized human pathogens including *V. cholera*, the etiological agent of epidemic cholera, and the hazardous seafood poisoning agents *V. vulnificus* and *Vibrio parahaemolyticus*. Many more vibrios are associated with diseases in marine animals, and only a handful of the 40 or more species currently described within the genus appear to be benign. Other notable gamma-proteobacterial pathogens are members of the *Aeromonas* and *Shewanella* genera, which are also widely distributed throughout marine environment. The proportion of marine human pathogenic species within the gamma-Proteobacteria is in contrast to terrestrial environments where groups such as the alpha-Proteobacteria and the spirochetes also contain many pathogenic members. Such discrepancy could reflect differing evolutionary trajectories of marine and terrestrial communities, or could reflect preferential culturability of gamma-proteobacterial pathogens as has generally been observed for heterotrophic gamma-Proteobacteria from the marine environment (e.g. Eilers et al., 2000b).

The deeply branching lineages of the Gram-positive bacteria also contain a high diversity of recognized marine pathogens. The *Mycobacterium* group is represented with several notable human pathogens including agents of tuberculosis, skin disease, and an expanding diversity of fish and marine mammal pathogens (Saubolle et al., 1996; Kusuda & Kawai, 1998; Dobos et al., 1999; Rhodes et al., 2001). Other Gram-positive human pathogens found in associations with marine environments include members of the *Clostridia*, *Listeria*, *Rhodococcus*, *Streptococcus*, and *Mycoplasma* group (Table 2.1).

2.2.2. Environmental Associations

Marine pathogens are often found in association with the surfaces of marine animals, phytoplankton, sediments and suspended detritus. The association of pathogens with marine biota has been compared to vector-borne disease in terrestrial environments as variability in environmental conditions can affect both the vector distribution and pathogen growth (Lipp et al., 2002). For example, algal and zooplankton blooms can promote proliferation of associated bacterial communities by providing microenvironments favoring growth and by exuding nutrients into the water (Lipp et al., 2002). Associations between zooplankton and pathogenic *Vibrio* and *Aeromonas* species have been observed (Kaneko and Colwell, 1978; Colwell, 1996; Dumontet et al., 2000; Heidelberg et al., 2002a) and the dynamics of attached pathogenic *Vibrio* species and *Vibrio* mediated disease (i.e. cholera) have been correlated to seasonal algal and zooplankton blooms (Kaneko and Colwell, 1978; Colwell, 1996; Heidelberg et al., 2002a).

Association with larger marine animals also influences the abundance of pathogens in the environment through activities including bioconcentration, fecal contamination, and by creating conditions favoring growth. Marine sediments with high overlying fish abundance have been found to be enriched in *Clostridium botulinum* spores suggesting deposition (Huss, 1980) while sediments underlying farmed mussels have been observed to support an enriched presence of vibrios relative to surrounding environments, possibly due to stimulated *Vibrio* growth in an organic-enriched environment (La Rosa, 2001). Filter-feeding shellfish are effective bioconcentrators of small particles and pathogenic contaminants in marine environments. Shellfish samples have been observed to harbor marine contaminants including Enteric bacteria (Burkhardt et al., 1992), *Campylobacter* (Abeyta et al., 1993; Endtz et al., 1997) and *Listeria* species (Colburn et al., 1990) in addition to potentially pathogenic indigenous flora

including *Vibrio* species (Olafsen et al., 1993; Lipp and Rose, 1997) thus, it is not surprising that shellfish have long been recognized as a potential source for marine-acquired illness.

Active growth of certain marine pathogens may occur only in association with nutrient-rich environments such as animal guts or organic-rich sediments. Such populations, when dislodged, may occur as inactive transients in seawater and act as seed populations for inoculating new habitats (Ruby and Nealson, 1978). This life-cycle has been suggested for certain fish-associated vibrios based on their ability to grow rapidly in response to nutrient addition even after prolonged incubation in seawater under starvation conditions (Jensen et al., 2003). Gastrointestinal tracts of marine animals have been shown to harbor a wide diversity of organisms closely related to bacterial pathogens (MacFarlane et al., 1986; Oxley et al., 2002). Similarly, organisms commonly associated with sediment environments include enteric pathogens (Grimes et al., 1986), and members of the genera *Vibrio* (Watkins, 1985; Hoi et al., 1998; Dumontet et al., 2000), *Aeromonas* (Dumontet et al., 2000), *Shewanella* (Myers and Nealson, 1990), *Clostridia* (Huss, 1980) and *Listeria* (Colburn et al., 1990).

Intracellular associations of bacteria with protozoan and algal hosts have been described in natural and clinical settings and may represent an additional source of pathogens in marine environments. Colonization of amoeboid hosts has been observed for several human bacterial pathogens including *Mycobacterium* (Cirillo et al., 1997; Steinert et al., 1998), *Burkholderia* (Michel and Hauröder, 1997; Marolda et al., 1999; Landers et al., 2000) and *Legionella* species (Cianciotto and Fields, 1992; Fields, 1996). *Legionella pneumophila* can replicate inside amoebas in natural waters and it is currently held that adaptation to the intracellular environment of a protozoan host predisposed *L. pneumophila*, the agent of Legionnaires' disease, to infect mammalian cells (Cianciotto and Fields, 1992; Fields, 1996; DePaola et al., 2000; Harb et al., 2000; Swanson and Hammer, 2000). Relatively high concentrations of *L. pneumophila* have been found in fresh water and coastal systems (10^2 to 10^4 CFU per ml) (Fliermans et al., 1981; Ortizroque and Hazen, 1987; Fliermans, 1996). Survival of free-living *L. pneumophila* in seawater over several days has been demonstrated (Heller et al., 1998); however, extracellular growth in natural water has not been observed (Steinert et al., 1998; Swanson and Hammer, 2000). Whether associations of *Legionella* spp. or other marine pathogens with protozoan hosts promotes growth of these bacteria in marine environments remains to be determined.

Algal cells have been shown to harbor intracellular bacterial associations (Biegala et al., 2002) and it is currently debated whether agents of harmful algal blooms (HAB) maintain bacterial symbionts that participate in toxin production (Gallacher and Smith, 1999). Bacteria found in association with cultures of HAB algae have been reported to produce a level of toxin per cell volume that is equivalent to the production of toxin in the alga (Gallacher and Smith, 1999). In addition, autonomous toxin production by free-living bacteria has been observed under marine conditions (Michaud et al., 2002). The relative contribution to toxin production during HABs by free-living, surface associated, or intracellular bacteria is an area of active investigation (Carmichael, 2001; Vasquez et al., 2001; Smith et al., 2002) (see also Chapter 10 of this book). Overall, the role of protist and algal hosts for harboring marine pathogens in the environment remains an important but poorly understood factor to be considered in risk assessment.

2.2.3. Abiotic Factors

Environmental parameters such as salinity, temperature, nutrients, and solar radiation influence the survival and proliferation of pathogens directly by affecting their growth and death

rates and indirectly through ecosystem interactions. The survival of contaminant pathogens in marine environments has been shown to decrease with elevated sunlight (Rozen & Belkin, 2001; Fujioka & Yoneyama, 2002; Hughes, 2003), high salinity (Anderson et al., 1979; Sinton et al., 2002), and increased temperature (Faust et al., 1975). However, elevated nutrients and particle associations have been shown to promote the survival of marine contaminants (Gerba & McLeod, 1976). There is increasing evidence that many pathogens found as pollutants in marine environments can survive harsh environmental conditions for prolonged periods of time in a spore-like, “viable but nonculturable” (VBNC) state (e.g. Grimes et al., 1986; Rahman et al., 1996; Rigsbee et al., 1997; Steinert et al., 1997; Cappelier et al., 1999a, 1999b; Besnard et al., 2000; Asakura et al., 2002; Bates et al., 2002). The effects of environmental parameters on the survival of enteric bacteria are reviewed in detail in Chapter 10 of this book.

In contrast to microbial contaminants, marine-indigenous pathogens are adapted to prevalent environmental conditions and their proliferation may be triggered by specific factors. For example, warm water temperatures appear to have a positive effect on the abundance of human-invasive pathogens, which tend to have mesophilic growth optima. In temperate environments, the distribution of such pathogens is typically seasonal with peaks in both environmental abundance and human infection occurring during the warmer months. This has been demonstrated for human pathogenic *Aeromonas* spp. (Kaper et al., 1981; Burke et al., 1984), *Shewanella algae* (Gram et al., 1999) and vibrios (CDC, 1999, 2000; Heidelberg et al., 2002b; Thompson et al., 2004b), including *V. cholerae* (Jiang & Fu, 2001) *V. parahaemolyticus* (Kaneko & Colwell, 1978), and *V. vulnificus* (Wright et al., 1996). In addition, elevated sunlight can stimulate growth of marine indigenous heterotrophic bacteria by increasing nutrient availability by photochemical breakdown of complex polymers to release organic metabolites (Chrost & Faust, 1999; Tranvik & Bertilsson, 2001). Nutrient enrichment in seawater samples and sediments has been correlated to increases in the relative abundance of *Vibrio* populations (Eilers et al., 2000a; La Rosa et al., 2001). It remains to be established whether stimulated growth of opportunistic invasive pathogens, in response to nutrient enrichment, is a general feature of seawater environments.

2.3. ROUTES OF TRANSMISSION

Transmission of pathogens to humans through marine environments most frequently occurs by eating contaminated seafood, but can also follow other routes including seawater contact or exposure to marine aerosols and zoonoses. The potential for contracting human diseases through marine environments depends on several factors including the susceptibility of the human host, the degree of exposure to a pathogen population, and the virulence of the pathogenic agent. Individuals with medical conditions such as liver disease and diabetes, or who are immunocompromised, are most susceptible to infections (Howard & Bennett, 1993; Howard & Burgess, 1993); however, infections also occur in healthy individuals. The degree of host exposure to a marine pathogen varies with the route of transmission and has been correlated to both the environmental concentration of the pathogen and the duration of exposure. For the purposes of risk assessment for seafood consumption, an average amount of ingested seafood is assumed (e.g., 110 g oyster meat (Miliotis et al., 2000)) and swimming related illnesses have been correlated to time spent in the water (Corbett et al., 1993). However, no explicit models appear to have been formulated for prediction of other routes of exposure (e.g., animal

contact, or aerosol inhalation). Finally, the virulence of the pathogenic population determines the dose needed to establish human disease. In several cases, it has been observed that strains most closely resembling clinical isolates represent only a small subset of related co-occurring organisms, suggesting that infections from marine environments may frequently be initiated by small numbers of highly virulent variants (Jackson et al., 1997).

2.3.1. Seafood Consumption

The most important route of infection by marine pathogens is by consumption of contaminated seafood resulting in symptoms from self-limiting gastroenteritis (typical seafood poisoning) to invasive infections that are potentially fatal. *Vibrio* species are the most significant risk in seafood consumption and an estimated 10,000 cases of food-borne infection occurs in the United States each year (FDA, 1994; Altekruze et al., 1997). But other bacterial genera naturally found in association with fish and shellfish have also been implicated in seafood-borne diseases (e.g., *Aeromonas*, *Clostridium*, *Plesiomonas*). Fecal contamination from human sewage or animal sources is recognized as an additional important source of seafood-borne pathogens (e.g., *Campylobacter*, *Escherichia*, *Listeria*, *Salmonella*, *Shigella*, and *Yersinia*) (Feldhusen, 2000). However, in several cases a clear distinction cannot be made whether a pathogen is a fecal contaminant or a natural part of the marine community. For example, *Salmonella*, generally considered a marine contaminant, may be a natural part of marine ecosystems (Tryland, 2000; Aschfalk et al., 2002). Other genera, such as *Campylobacter*, are detected in the feces of marine birds (Endtz et al., 1997) and could be described as “endemic contaminants” since their presence can be detected in environments not polluted by humans.

Infection by ingestion generally requires relatively large doses of pathogens (e.g., 10^5 – 10^{10} cells for most gamma proteobacterial pathogens), although some highly virulent pathogens such as *Shigella* or enterohemorrhagic *Escherichia coli* can establish infections with doses as small as 10–100 cells (PHAC, 2001) (Table 2.1). Levels of marine-indigenous pathogens in fresh seafood are usually low enough to be considered safe so that only the growth of these organisms is regarded as a hazard (e.g., during periods of improper handling) (Feldhusen, 2000). For example, nonrefrigeration of oysters after harvesting can amplify the endemic *Vibrio* population 10,000-fold (Miliotis et al., 2000) resulting in levels that are deemed unsafe for human consumption (i.e., $\geq 10^4$ cells/g oyster (FDA, 1997)).

While cooking minimizes the risk of seafood-borne infection, poisoning can occur from heat-stable bacterial toxins or compounds. Scombroid (or histamine) fish poisoning is caused when bacteria containing the enzyme histadine-decarboxylase proliferate in improperly stored fish rich in the amino acid histadine (e.g., tuna, sardines, and salmon) (Burke & Tester, 2002). Bacterial transformation of histadine can produce dangerous levels of histamine, consumption of which can lead to severe allergic reactions. Several types of bacteria including *Morganella morganii* and *Klebsiella oxytoca* have been implicated in histamine production in fish (Lopez-Sabater et al., 1996). In addition, toxins produced by marine bacterial species may be concentrated by the activities of filter feeding shellfish. Although this has not been confirmed as a route of human pathogenicity in marine environments, toxin production has been observed by bacterial strains associated with HAB algae including members of the *Roseobacter* and *Alteromonas* genera, and cyanobacterial species (Gallacher & Smith, 1999; Carmichael, 2001).

2.3.2. Seawater Exposure

Pathogens can be transmitted to humans through seawater during accidental ingestion, inhalation, or by direct exposure of ears, eyes, nose, and wounded soft tissue. Although sewage contamination has long been recognized as a significant risk factor in acquiring illnesses after seawater exposure, sewage-borne pathogens are primarily viral rather than bacterial (Cabelli et al., 1982; Griffin et al., 2001). Invasive bacterial infections acquired in marine environments have primarily been attributed to marine endemic species including gamma-proteobacterial strains related to *Aeromonas*, *Halomonas*, *Pseudomonas*, *Shewanella*, and *Vibrio* (Table 2.1). In beaches with high swimmer density, human-shed *Staphylococcus* or *Streptococcus* can cause minor wound and ear infections (Charoenca & Fujioka, 1993; Thomas & Scott, 1997). Other bacterial infections that have been reported after exposure to marine or estuarine waters include leptospirosis (Thomas & Scott, 1997) and skin granulomas caused by water-borne *Mycobacterium marinum* (Dobos et al., 1999). Near-drowning experiences in marine environments bring seawater into the lungs and can result in pneumonia (Ender & Dolan, 1997; Thomas and Scott, 1997). Such infections have been reported for marine indigenous pathogens including *Legionella bozemanii*, *Francisella philomiragia*, *Klebsiella pneumonia* and several *Vibrio* and *Aeromonas* species (Ender & Dolan, 1997).

Although the range of infectious doses for wound and skin infections is not known and the degree of exposure is difficult to estimate, the danger may potentially be high. Fifty percent mortality was observed for artificially wounded rats exposed to $\sim 10^7$ CFUs of marine and clinical isolates of *Aeromonas hydrophila*, *V. parahaemolyticus*, and *V. vulnificus* (Kueh et al., 1992). In the same study, similar mortalities were observed in rats exposed to 1 ml aliquots of seawater from multiple sites, suggesting a high degree of indigenous seawater-associated virulence (Kueh et al., 1992).

2.3.3. Aerosol Exposure

The first case of Legionnaires Disease in 1976 demonstrated the importance of airborne transmission of the water-borne bacterial pathogen *Legionella pneumophila* (McDade et al., 1977). Transmission of bacterial disease by marine aerosols has not been documented but should be considered as a potential route of infection. Studies have shown that *Mycobacterium* species are enriched in aerosols from natural waters (Wendt et al., 1980; Parker et al., 1983) and additional respiratory disease agents, which have been detected in seawater, include *F. philomiragia*, *Legionella* spp., *Acinetobacter calcoaceticus*, and *K. pneumoniae* (Grimes, 1991; Ender & Dolan, 1997). In general, infectious doses for respiratory agents are small, e.g. 5–10 organisms for *Mycobacterium tuberculosis* infection. In addition, aerosols, generated in coastal environments by wave activity, can transmit algal toxins to humans (Van Dolah, 2000) and cause viruses to become airborne (Baylor et al., 1977). Thus, marine aerosols may be an unrecognized factor in the transmission of diseases from marine environments.

2.3.4. Marine Zoonoses

Zoonoses are naturally transmissible diseases from animals to humans. Warm-blooded marine mammals harbor and are afflicted by a wide variety of pathogens posing zoonotic risk to humans including *Brucella*, *Burkholderia*, *Clostridium*, *Helicobacter*, *Mycobacterium*,

Rhodococcus, and *Salmonella* species (Bernardelli et al., 1996; Harper et al., 2000; Tryland, 2000; Aschfalk & Muller, 2001; Aschfalk et al., 2002) (Table 2.1). Tuberculosis, a chronic respiratory disease caused by *Mycobacterium* species including *M. tuberculosis* and *M. bovis*, has afflicted natural and captive populations of marine mammals (Bernardelli et al., 1996; Montali et al., 2001) and transmission from seal to man has been documented (Thompson et al., 1993). Brucellosis, a systemic infection, is transmitted to humans from infected animals, meat, or dairy products in many parts of the world. Brucellosis has also been observed in a wide range of marine animals including dolphins, porpoises, whales, seals, and otters (Tryland, 2000; Foster et al., 2002). The zoonotic potential of these marine *Brucella* species has been recognized after three incidents of infection, first of a researcher handling a marine isolate (Brew et al., 1999) and then in two cases of neurobrucellosis attributed to a marine *Brucella* strain in Peru (Sohn et al., 2003).

Injuries inflicted by marine animals or sustained during their handling are especially susceptible to infection by associated microorganisms and therefore emergency treatment of bites (e.g., from sharks, moray eels) includes broad-spectrum antibiotics (Erickson et al., 1992; Howard & Burgess, 1993). Handling of fish or crabs has been associated with infection by *Erysipelothrix rhusopathiae*, a mycoplasma-like organism common on the skin of fish, which manifests as a localized swollen purple area around a wound (fish handler's disease) (Thomas & Scott, 1997). Other mycoplasma-like organisms including *Mycoplasma phocacerebrale* have been isolated from seals during pneumonia epizootics and have been implicated in development of "seal finger," a local infection of the hands in humans (Kirchhoff et al., 1989; Stadtlander & Madoff, 1994; Baker et al., 1998).

The transmission of disease between farmed and wild fish populations is one of many concerns regarding the sustainability of aquaculture practices (Garrett et al., 1997; Naylor et al., 2000). The zoonotic potential of farmed fish environments has also been recognized on several occasions. The fish pathogen, *Streptococcus inae* (Zlotkin et al., 1998; Colorni et al., 2002), caused an outbreak of infection in fish farmers in British Columbia (Weinstein et al., 1996, 1997). Additional health hazards of fish handlers include infections with *A. hydrophila*, *Edwardsiella tarda*, *E. rhusopathiae*, *M. marinum*, and *Vibrio* species (Lehane & Rawlin, 2000). In addition, several currently emerging pathogens of fish populations are closely related to human pathogens (Fryer & Mauel, 1997; Rhodes et al., 2001; Starliper, 2001). Recently, *Serratia liquefaciens* was identified as an agent of deadly systemic hospital infections in humans (Grohskopf et al., 2001) and in the same year was identified as a pathogen of farmed Atlantic salmon (Starliper, 2001).

2.4. INDICATORS FOR MARINE RISK ASSESSMENT

The quality of marine waters has been routinely monitored using detection of indicator organisms found in association with human pollution. Indicators are elements that can be efficiently monitored to approximate the risk of human exposure to a given environment. While the indicators themselves do not necessarily cause disease, their presence in an environment suggests a high probability of co-occurring pathogens. Although traditionally indicator organisms have been relied upon for water quality assessment, the use of physical and chemical proxies and direct detection of pathogen populations are showing promise as tools for future water quality management.

2.4.1. Indicators for Sewage Pollution

Sewage-associated public health risks continue to plague coastal environments worldwide. The NRDC¹ reports that 12,184 U.S. beach closings or advisories were issued in 2002 (of 2922 reporting beaches) of which 87% were attributed to poor bacterial water quality (as monitored by indicators for fecal pollution) (Dorfman, 2003). In a landmark epidemiological study, Cabelli et al. (1982) found that illness (primarily gastroenteritis and respiratory infections) associated with swimming in several marine environments increased linearly with the degree of site pollution. They further showed that levels of Gram-positive fecal enterococci and fecal coliforms were good proxies for sewage contamination. Based upon this and similar studies the current USEPA² standard for acceptably safe beaches is a monthly geometric mean of 35 enterococci per 100 ml (Dufour et al., 1986) and a median of 14 fecal coliforms per 100 ml in shellfish harvesting waters (USEPA, 1988).

The use of enterococci and fecal coliform levels as indicator organisms for marine water quality assessment has been repeatedly called into question. These indicator species have shown varying degrees of specificity for detecting sewage contamination against background environmental fluctuations from animal and environmental sources (Grant et al., 2001; Boehm et al., 2002). Boehm et al. (2002) showed that coastal enterococci levels are enriched by bird activity in adjacent estuaries. Alternative sewage-borne indicators, such as *Clostridium perfringens*, have been considered due to their stability in the marine environment (Fujioka, 1997); however, they too are found in association with marine animals (e.g., Aschfalk & Muller, 2001) and may be subject to environmental variability. In addition, their correlation to human illness has not been convincing (Dufour et al., 1986). Furthermore, exclusive reliance on fecal indicator bacteria for marine water quality assessment has been challenged due to their limited ability to predict viral contamination and the presence of marine-indigenous pathogens (Dumontet et al., 2000; Tamplin, 2001). While sewage indicators remain a useful tool for monitoring water pollution, continued efforts to establish alternative indicators for nonsewage related risks hold promise for future risk assessment.

2.4.2. Indicators for Nonsewage Related Risk

Additional factors that have been related to human risks from seawater exposure include swimmer density, eutrophication, and thermal pollution. High swimmer density at bathing beaches has been correlated to the acquisition of ear and minor skin infections from human shed bacteria. Levels of the pathogen, *Staphylococcus aureus*, have been proposed as an indicator for exposure to human-shed bacteria with levels above 100 CFU per 100 ml of seawater considered unsafe (Charoenca & Fujioka, 1993; Fujioka, 1997).

Eutrophication of coastal environments may be linked to infections by marine indigenous pathogens (e.g., Kueh et al., 1992). The relative abundance of *Vibrio* populations in seawater samples increases in response to organic nutrient enrichment, and pollution from aquaculture environments has been correlated to increased proportions of vibrios in underlying sediments (Eilers et al., 2000a; La Rosa et al., 2001). Accordingly, the prevalence of vibrios or other aerobic heterotrophs has been suggested as an indicator for nutrient enrichment in marine environments (La Rosa et al., 2001).

¹ National Resources Defense Council.

² United States Environmental Protection Agency.

That high seawater temperature bears higher risk of exposure to marine pathogens has been established in studies of shellfish (Wright et al., 1996; Motes et al., 1998; Miliotis et al., 2000), natural waters (Wright et al., 1996; Jiang & Fu, 2001; Heidelberg et al., 2002b; Louis et al., 2003; Thompson et al., 2004b), and the incidence of epidemic cholera (Colwell, 1996; Pascual et al., 2000). Remote sensing of sea surface temperature is currently being explored as a means to predict the onset of cholera outbreaks along the Indian and Bangladesh coasts (Lobitz et al., 2000).

2.5. DETECTION AND QUANTIFICATION

In this section an overview of the methods currently available to detect, identify, and enumerate marine pathogen (or indicator) populations is presented. At the center of the discussion will be methods with proven utility for targeting specific populations within environmental microbial communities. However, several techniques used to isolate and identify marine pathogens in clinical specimens will also be briefly evaluated. Methods used to identify and quantify microbial populations can be divided into three main groups: culture, immunology, and nucleic acid based. However, protocols frequently do not fall exclusively into one category but represent combinations. Because of the considerable number of published protocol and commercial kits, this overview presents the general principles that define these three main groups of methods. Where specific examples are given these have been selected because they have been (i) employed by several laboratories and/or (ii) characterized with respect to their limits of sensitivity and specificity. A summary of representative nucleic acid- and immunology-based methods for detection or quantification of marine-relevant pathogen populations is presented in Table 2.2. In a few cases, methods are described that have not yet been applied to pathogen detection but hold potential.

Methods for monitoring pathogen populations should be selected by evaluating the factors that mediate exposure of humans to the pathogen (e.g., abundance, virulence/infectious dose, route of exposure) and the constraints of the method (e.g., sensitivity, specificity, dynamic range, cost). Methods targeting pathogen populations must be sensitive enough to monitor populations at levels below the infectious dose, and specific enough to recognize the target group without generating false positives by cross-reacting with nontarget organisms. Detection requires positive identification at or above specified threshold concentrations while enumeration requires flexibility to identify a range of population levels. For clinical purposes, detection is often sufficient, while quantification of hazardous populations is preferable for analysis of environmental samples. The methods also differ greatly in speed and cost of implementation and therefore the most accurate method may not always be the most preferable when rapid decision making is required. The following sections present our attempt to take these considerations into account while evaluating the strengths and weaknesses of various methods.

2.5.1. Culture-Based Methods

Detection of pathogens via culturing requires enrichment of a target population over other environmental bacteria. This employs selective and/or differential media, which provide a 'presumptive identification' and can be followed by any number of tests (e.g. biochemical, immunological or molecular) to confirm the identity of isolates. A medium is selective if it favors the growth of a specific population of organisms and is differential if it allows distinction

Table 2.2 Representative immunological and molecular methods for detection of human pathogens in environmental samples.

Bacteria	Sample	Target	Method	Sensitivity	Reference
<i>Escherichia coli</i>	Food	Shiga-toxin gene (<i>stx</i>)	PCR following enrichment Colony hybridization	ND	(Dutta et al., 2000)
	Waste and drinking water	16S rRNA gene	Fish	10^7 – 10^8 cells/ 100 mL	(Baudart et al., 2002)
	Artificially contaminated food	NA	Real-Time PCR (BAX®kit, Dupont)	10^2 – 10^3 cells/100 mL	(Bhagwat, 2003)
	Water	β-D-galactosidase gene	Real-Time PCR following enrichment with magnetic beads	10^3 cells/100 mL	(Foulds et al., 2002)
	Waste water	<i>gcvp</i> gene	Multiplex PCR	ND	(Turner et al., 1997)
	Food outbreaks—clinical isolated strains	Genome	PFGE (PulseNet)	ND	(Breuer et al., 2001)
	Wastewater wetlands	<i>stx</i> 1, 2 genes and intimin (<i>eae</i>) gene	Real-time PCR	W/o enrichment: 3×10^4 CFU/g With enrichment: 10^4 CFU/g	(Ibekwe et al., 2002)
<i>Campylobacter</i> spp.	Poultry	16S rRNA gene	Modified RT-PCR followed by hybridization	<1 CFU/g	(Uyttendaele et al., 1995, 1996)
	Mussels and oysters—isolated strains	GTPase gene	PCR and line-blot hybridization	ND	(Van Doorn et al., 1998)
	Water, food	VSI sequence	Real-time PCR	6–15 CFU/reaction	(Yang et al., 2003)
	Food outbreaks—clinical isolated strains	Genome	PFGE (PulseNet)	ND	(Ribot et al., 2001)
<i>Listeria monocytogenes</i>	Water, and waste water	<i>flaA</i> , B genes	Modified PCR	3–15 CFU/ 100 mL	(Waage et al., 1999c)
	Food outbreaks—clinical isolated strains	Genome	PFGE (PulseNet)	ND	(Graves & Swaminathan, 2001)
	Food outbreaks—clinical isolated strains	Listeriolysin O (<i>hly</i>) gene	PCR	ND	(Jackson et al., 2000)
	Food outbreaks—clinical isolated strains	<i>hly</i> gene	Dot-blot hybridization (US-FDA established protocols)	ND	(Jackson et al., 2000)

Food	16S rRNA gene	PCR following culture enrichment	0.04–0.2 CFU/g	(Somer & Kashi, 2003)
	<i>iap</i> gene	PCR following culture enrichment or directly from the sample	W/o enrichment: 10 ⁷ CFU/g waste sample With enrichment: <10 CFU/g waste sample	(Burtscher & Wuertz, 2003)
<i>Staphylococcus aureus</i>	Water/food outbreaks—clinical isolates	Colony hybridization (US-FDA established protocols)	ND	(Jackson et al., 2000)
	Household waste samples	PCR following culture enrichment or directly from the sample	W/o enrichment: 10 ⁶ CFU/g waste sample With enrichment: <10 CFU/g waste sample	(Burtscher & Wuertz, 2003)
<i>Yersinia</i> spp.	Water/food outbreaks—clinical isolates	Colony hybridization (US-FDA established protocols)	ND	(Jackson et al., 2000)
	Raw meat	Real-time PCR	10 ⁶ CFU/g	(Vishnubhatla et al., 2001)
	Water/food outbreaks—clinical isolates	Dot-blot hybridization—(US-FDA established protocols)	ND	(Jackson et al., 2000)
	Household waste samples	PCR following enrichment or directly from the sample	w/o enrichment: 10 ⁷ CFU/g waste sample/ with enrichment: <10 FU/g waste sample	(Burtscher & Wuertz, 2003)
	Water and waste water	PCR following culture enrichment	8–17 CFU/ 100 mL	(Waage et al., 1999a)

(continued)

Table 2.2 (Continued)

Bacteria	Sample	Target	Method	Sensitivity	Reference
<i>Shigella</i> spp.	Sewage polluted seawater	Invasion plasmid antigen H (<i>ipaH</i>)	PCR	10 to 100 CFU/reaction	(Kong et al., 2002)
	Water	<i>ipaBDC</i> , <i>ipaH</i> genes and <i>stx1</i> gene	PCR	5×10^2 CFU/100 mL	(Faruque et al., 2002)
<i>Salmonella</i> spp.	Food (outbreak)	<i>invA</i> gene	Real-time PCR	ND	(Daum et al., 2002)
	Waste samples	<i>ompC</i> gene	PCR following enrichment or directly from the sample	w/o enrichment: 10^7 CFU/g waste sample/with enrichment: <10 CFU/g waste sample	(Burtischer & Wuertz, 2003)
<i>Vibrio cholerae</i>	Water and shellfish	<i>invA</i> gene	PCR	3×10^5 cells/100 ml	(Dupray et al., 1997)
	Sewage polluted seawater	<i>ipaB</i> gene	PCR	10 to 100 CFU/reaction	(Kong et al., 2002)
	Water, and waste water	ST11, 15 sequences	PCR	10 CFU/100 ml	(Waage et al., 1999b)
	Harp seals	LPS	ELISA	ND	(Aschfalk et al., 2002)
	Raw oysters	Hemolysin (<i>hlyA</i>)	Real-Time PCR	10^2 CFU/100 mL	(Lyon, 2001)
	Waste waters estuarine waters—environmental isolates	<i>ctxA</i> , <i>hlyA</i> , <i>ompU</i> , <i>stn/sto</i> , <i>tcpA</i> , <i>tcpL</i> , <i>toxR</i> , and <i>zot</i> genes	Multiplex PCR	ND	(Rivera et al., 2001)
	Water/food outbreaks—clinical isolates	Cholera enterotoxin gene (<i>ctx</i>)	ELISA (US-FDA established protocols)	ND	(Jackson et al., 2000)
	Sewage polluted seawater	<i>ctx</i> gene	PCR	10–100 CFU/reaction	(Kong et al., 2002)
	Seawater	<i>ctxA</i> , 16s–23s rRNA	PCR	ND	(Lipp et al., 2003)
	Estuarine water	O1 or O139 surface antigens	Agglutination following culture enrichment	2×10^3 CFU/mL ND	(Hasan et al., 1995; Louis et al., 2003)
	Seawater	16s–23s rRNA	Colony hybridization following enrichment	ND, but <100 CFU/100 mL was detected	(Jiang & Fu, 2001)
	Seawater where <i>V. cholera</i> is endemic (Bangladesh)	O1 or O139 LPS antigens	Immunofluorescence microscopy	$1.5\text{--}4 \times 10^4$ cells/100 mL	(Brayton et al., 1987; Hasan et al., 1995)

<i>V. parahaemolyticus</i>	Raw oysters	Thermostable direct hemolysin (<i>tdh</i>) gene	Colony hybridization following culture enrichment	10 CFU/g D	(DePaola et al., 2000; Nordstrom & DePaola, 2003)
	Raw oysters	<i>tdh</i> gene	Real-time PCR following culture enrichment	1 CFU/reaction	(Blackstone et al., 2003)
	Sewage polluted seawater	16s–23s rRNA	PCR	10–100 CFU/reaction	(Kong et al., 2002)
	Seawater	<i>toxR</i> , <i>tdh</i> , <i>trh</i> genes	Most-probable-number PCR	3 cells/100 mL	(Alam et al., 2003)
<i>V. vulnificus</i>	Seafood	<i>tdh</i> gene	PCR with enrichment	0.3 cells/g	(Hara-Kudo et al., 2003)
	Seawater, sediment, fish	Cytolysin gene	Hybridization following culture enrichment	2 CFU/100 mL	(Hoi et al., 1998)
	Oysters—isolated strains	Genome	PFGE	ND	(Jackson et al., 1997; Tamplin et al., 1996)
	Eels	LPS	ELISA	10 ⁴ –10 ⁵ cells/reaction	(Biosca et al., 1997)
	Water	Hemolysin (<i>vvhA</i>) gene	Colony hybridization	2 × 10 ³ /100 mL	(Lee et al., 2001)

NA: not applicable.

of specific properties of the target population. Formulations of media designed to isolate specific organisms have been widely published in the literature and are available through various microbiological handbooks [e.g. (Atlas, 1995)]. For example, bacteria in the genus *Vibrio* (a marine-endemic genus containing a high diversity of human and animal pathogens) can be readily isolated using thiosulfate-citrate bile salts sucrose (TCBS) media where selectivity for organisms tolerant of both intestinal and marine environments (characteristic of vibrios) is provided by the combination of bile salts and alkaline pH, respectively. The TCBS medium is also differential for the trait of sucrose fermentation because it contains an indicator dye, which responds to acid produced by sucrose fermentation during growth. TCBS media has been routinely employed in clinical settings to diagnose gastrointestinal diseases, seafood poisoning, or wound infections mediated by *Vibrio* species. For diagnosis of the diarrheal disease cholera, presumptive identification of the etiological agent *V. cholerae* as small yellow colonies on TCBS media must be confirmed with subsequent tests, as certain marine vibrios and alkaline-tolerant enteric bacteria can manifest similar morphologies on the media (Lotz, 1983). However, the specificity for the target group (e.g. vibrios) of such media can be surprisingly good. For example, >95% of seawater isolates grown on an improved formulation of TCBS (2–3% salt) (Toro et al., 1995) were *Vibrio* sp. as determined by 16S rRNA sequencing and the remainder were closely related genera (Thompson et al., 2005). Similarly, a selective media designed for presumptive identification of *V. vulnificus* by combination of antibiotic resistance, metabolism of cellobiose, and colony morphology, yielded 79% specificity for target organisms as confirmed by hybridization with DNA probes (Hoi and Dalsgaard, 2000).

Growth-based quantification of pathogen abundance has long tradition and is often referred to as direct viable counts (DVC). Abundance is either inferred from the number of colony forming units (CFUs) on culture plates or by Most Probable Number (MPN) dilutions of environmental samples. However, to ensure the accuracy of detection, representative presumptive positive strains must be corroborated by more extensive characterization with biochemical tests or molecular assays (described in the next sections). The dilution or concentration (e.g., by filtration) of samples prior to culture-based enumeration can accommodate a wide dynamic range of environmental microbial population sizes. Protocols for culture-based enumeration of marine pathogens include those for *Aeromonas* (Villari et al., 1999), *Clostridium* (Glasby and Hatheway, 1985), *Legionella* (Boulanger and Edelstein, 1995; Bartie et al., 2003), *Vibrio* sp. (Hernandezlopez et al., 1995) and *V. vulnificus* (Hoi and Dalsgaard, 2000; Cerda-Cuellar et al., 2001).

A disadvantage of culture-based detection and enumeration methods is the dependence on reproducible and quantitative growth of target pathogen populations on culture media. Indeed, the majority of natural bacteria have been shown to be inherently difficult to culture and even those that are typically easy to culture can enter stages where their culture efficiency drops dramatically. For example, certain pathogens have been shown to enter a viable but non-culturable state (VBNC) in response to shifts in environmental conditions, complicating the interpretation of population dynamics observed in culture-based studies (Grimes et al., 1986; Rahman et al., 1996; Rigsbee et al., 1997; Steinert et al., 1997; Cappelier et al., 1999a; Cappelier et al., 1999b; Besnard et al., 2000; Asakura et al., 2002; Bates et al., 2002). Thus, it is important to evaluate whether non-culturable states have been described for the target pathogens and to take these into account in the evaluation of protocols.

An additional limitation of culture-based techniques is the rate at which the target population grows to detectable levels. Several assays designed for routine monitoring of marine water

quality have been optimized for speed. For example, detection and enumeration of Fecal Enterococci using USEPA Method 1600 requires a 24 hr incubation for presumptive results, which are then verified by biochemical testing over an additional 48 hours. However, with notable exceptions, most culture-based identification schemes for specific populations are time and labor-intensive, and may require preliminary enrichment or decontamination steps that confound enumeration. For example, pathogenic *Mycobacteria* species grow relatively slowly in culture (1 to >20 weeks) and thus can easily be overgrown by faster-growing organisms. Since *Mycobacteria* are resistant to harsh conditions (i.e. alkaline and acidic treatments), washing environmental samples at high or low pH can be coupled with selective media to eliminate faster growing competitors and increase the efficiency of their isolation (Songer, 1981) (Hartmans and DeBont, 1999).

Despite some disadvantages of culture-based methods, including the variability in culturing efficiency of target populations and the labor intensive nature of microbial cultivation, significant benefits remain. Most notably, the cost of materials needed for culture-based assays is often less than for molecular methods, which can require extensive training, and highly specialized materials and equipment. In addition, cultured isolates allow subsequent investigations into the virulence and/or clinical significance of environmental pathogen populations.

2.5.2. Immunological Methods

Immunological detection has been used to identify and in some cases enumerate pathogen populations in clinical and environmental samples. These methods rely on the inherently high specificity of immune reactions and typically target pathogen-specific antigens such as cell-wall lipopolysaccharides (LPSs), membrane and flagellar proteins or toxins. Immunoassays can be categorized into three main groups: enzyme-linked immunosorbent assay (ELISA), immunofluorescent microscopy, and agglutination assays. These have been essential diagnostic tools in medicine and food quality monitoring because they are fast and accurate (for a detailed description see Schlöter et al., 1995; Rose et al., 2002).

There are several notable challenges for the implementation of immunological methods to detection of pathogens in environmental samples, which contain a large diversity of unknown bacteria. First, the sensitivity of many current methods is not high enough for detection of pathogens at low, environmentally relevant, concentrations. Second, false positive results can be generated by cross-reaction of antibodies with antigens of similar but nontargeted organisms. This is particularly problematic when polyclonal antibodies are used since these are complex mixtures of antibodies against multiple, mostly uncharacterized cell structures. However, the increased facility with which antibodies specific for single antigenic determinants (monoclonal antibodies) can be produced is improving the specificity of assays (Schlöter et al., 1995; Mitov et al., 2003). Finally, design and production of specific antibodies generally requires growth of target microorganisms, constraining the applicability of the methods to culturable populations. Despite these limitations, immunological methods have many potential applications for detection of pathogens in clinical and environmental settings.

2.5.2.1. Enzyme-Linked Immunosorbent Assay

Several ELISA assays have been developed for identification of marine-pathogen populations in human or animal clinical samples. For the indirect ELISA assay, bacteria (or

bacterial antigens) are immobilized in microtiter wells and are challenged with pathogen-specific antibodies. These antibodies can be contained in anti-sera collected from infected individuals or laboratory animals (polyclonal) or can be derived from clonal cell lines (monoclonal). In the direct ELISA assay, antibodies linked to microtiter plates are challenged with antigens (e.g., bacterial cells). In both assays, detection of positive antigen–antibody complexes is accomplished by activation of an enzyme reporter system (e.g., alkaline phosphatase, peroxidase or β -galactosidase) upon binding. This typically results in formation of colored product, which can be measured. A considerable number of ELISA assays are available for pathogen-specific antigens including the LPS of *Salmonella* (House et al., 2001), the cholera toxin antigen of *V. cholerae* (Jackson et al., 2000), the heat-labile enterotoxin of enterotoxigenic *E. coli* (Germani et al., 1994b; Koike et al., 1997), the Shiga-like toxin I of diarrhoeogenic *E. coli* (Germani et al., 1994a), and the listeriolysin O and internalin A of *Listeria monocytogenes* (Jackson et al., 2000; Boerlin et al., 2003; Palumbo et al., 2003). Many of these assays are commercially available, are routinely applied to clinical specimens or contaminated food samples, and possess high potential for automation.

The sensitivity achieved by most ELISA assays makes them useful for clinical detection of pathogens, and in some cases quantification. However, the application of the ELISA assay to environmental samples frequently requires careful evaluation and optimization due to the generally low concentration of pathogens. For example, a direct ELISA assay for *V. vulnificus* was evaluated in artificially infected eel and water samples (Biosca et al., 1997). Antibodies targeted against biotype 2 LPS yielded a detection limit of 10^4 – 10^5 cells per well, corresponding to water-borne *V. vulnificus* populations near 10^6 CFU per ml (Biosca et al., 1997). Because typical environmental concentrations do not exceed 10^3 cells per ml the assay was not adequate for *V. vulnificus* detection in natural seawater. ELISA assays have been successfully used in several studies to characterize the pathogen populations present in marine mammals. This included detection of *Salmonella* spp. LPS (Aschfalk et al., 2002) and *C. perfringens* toxin (Aschfalk & Muller, 2001) in seal populations of the Greenland Sea, and *Brucella* populations in marine mammals in North Atlantic coastal waters (Tryland et al., 1999; Foster et al., 2002).

2.5.2.2. Immunofluorescence Microscopy

Immunofluorescence has been used to identify and quantify marine pathogens in environmental, food, and clinical samples. In these assays, fluorescence-conjugated antibodies are incubated with fixed samples (e.g., cell suspensions, filter concentrated cells, or tissue sections) and positive reactions are detected by epifluorescent microscopy. Pathogen-specific antibodies can be conjugated directly to a fluorescent marker (e.g., fluorescein isothiocyanate (FITC) or Texas Red), or can be targeted by a second, fluorescently labeled antibody. When samples are prepared quantitatively, enumeration of positive reactions provides a measure of population size. Indeed, several examples highlight the sensitive detection of water- and food-borne pathogen populations. *E. coli* abundances in seawater were detected above 1 cell per ml by applying a primary polyclonal mixture followed by a secondary, FITC-conjugated antibody to filter concentrated samples (Caruso et al., 2000, 2002). Similarly, FITC-conjugated monoclonal antibodies targeting *V. cholerae* O1 or O139 detected between 10^2 and 10^4 cells per ml in filter-concentrated river and estuarine waters in Bangladesh (Brayton et al., 1987; Hasan et al., 1995). *L. pneumophila* abundance in lake water was determined over a range of 9–3000 cells per ml by a direct immunofluorescence assay with monoclonal antibodies against serogroups 1–4 following 500-fold concentration of samples by centrifugation (Fliermans et al., 1981). For

routine detection of pathogens in food and clinical samples a number of fluorescently conjugated antibodies are commercially available. For example, polyclonal *Salmonella* spp. antibody mixture, directly conjugated with Texas-Red, allowed the detection of *Salmonella* spp. in fresh and processed meats (Duffy et al., 2000). These examples illustrate that immunofluorescence holds promise for sensitive and accurate detection of pathogens in environmental samples.

2.5.2.3. Agglutination Assays

Agglutination assays are routinely used for identification of clinical isolates and have in some cases been applied to detection of environmental pathogens. The assay is based on antigen binding to antibodies that are linked to particles (e.g., latex beads). Antibody–antigen aggregates result in the formation of visible clumps that are easily observed on a microscope slide or in a liquid test tube format. Several agglutination kits are commercially available including diagnostic tests for *S. aureus* (targeting protein A and clumping factor) (Wilkerson et al., 1997) and the BengalScreen agglutination test, which has been shown to identify *V. cholerae* O139 above 2×10^3 CFU per ml in clinical and environmental samples (Hasan et al., 1995). Additional agglutination assays have been developed targeting the LPS and outer membrane proteins of *Pseudomonas anguilliseptica* (Lopez-Romalde et al., 2003), *Brucella bacteremia* (Almuneef & Memish, 2003), and *Salmonella* spp. (Jackson et al., 2000). The greatest advantage of agglutination assays is that they are relatively simple, rapid, and inexpensive yet retain the potentially high specificity of immunological methods. However, as with most immunological methods, the sensitivity needs to be carefully evaluated for environmental applications.

2.5.3. Nucleic-Acid-Based Methods

Advances in molecular biology have revolutionized clinical and environmental microbiology by facilitating the identification of emerging pathogens, the detection of environmental populations, and the discrimination between closely related pathogenic and nonpathogenic bacteria. Molecular methods allow the characterization of bacteria by genotype rather than by phenotype and thus require identification of a unique genetic signature for individual or groups of pathogenic strains. Determination of genetic signatures remains the biggest challenge and typically requires extensive sequence characterization of the pathogen and related bacteria. However, if specific signatures can be identified, molecular methods provide a powerful diagnostic tool because nucleic acids can be rapidly and sensitively measured.

Discrimination of nucleotide variation among genes, whose occurrence is specific to an organism or whose sequence differentiates organisms, is often achieved by nucleic acid hybridization; other methods rely on restriction cutting of the chromosome. Hybridization-based methods include fluorescence in situ hybridization (FISH) and filter hybridization (e.g., colony and dot-blot hybridization), and the polymerase chain reaction (PCR). The PCR couples hybridization of short DNA molecules (primers) to template molecules followed by amplification with a polymerase (see below). Molecular typing methods have used PCR (e.g., multilocus sequence typing (MLST)) or restriction cutting (e.g., pulsed field gel electrophoresis (PFGE)) for analyzing genomic signatures. The general principles of hybridization-based, PCR-based, and molecular typing methods have been reviewed in widely available protocol books (Sambrook & Russel, 2001; Persing, 2003).

Important considerations for development of hybridization-based or PCR-based pathogen detection assays are those of probe specificity and sensitivity in the choice of target genes.

Short probes (oligonucleotides) can be hybridized with the highest specificity since they can differentiate as little as a single nucleotide change between targets; however, they can only carry a limited amount of label so that their detection limit is relatively high. On the other hand, longer probes (polynucleotides) can carry multiple labels but cannot distinguish closely related sequences because mismatches up to a certain level cannot be differentiated. Thus, knowledge of sequence variation among genes in related pathogenic and nonpathogenic strains is important for judgment of specificity. For environmental pathogens this remains a challenge since it has been shown that very similar pathogenic and nonpathogenic strains can coexist (Zo et al., 2002). Furthermore, genome sequencing has demonstrated that pathogenicity has frequently arisen via transfer of genes from other bacterial groups, and even genes, which are unique to a pathogen among closely related bacteria, may have close sequence relatives in overall distantly related bacteria (Welch et al., 2002; Ivanova et al., 2003). Thus, ideally, assay development should be coupled to exploration of population genetics and dynamics of the target pathogens and related groups.

2.5.3.1. Hybridization Methods

Fluorescent in situ hybridization. FISH enables detection of specific nucleic acid sequences inside intact cells. Fixed cells are immobilized on microscope slides and permeabilized with chemical reagents. Probes, primarily oligonucleotides (<25 nucleotides long), complementary to specific regions in the cellular DNA or RNA molecules are applied to the cells under optimized incubation and wash conditions. Fluorescent labeling of the probes allows visualization of the target cells by epifluorescent microscopy. Several different labeling techniques are available and include direct labeling of the probe (e.g., FITC or cyanin dye 3) or indirect labeling of probes with enzymes (e.g., horseradish peroxidase), antibodies, or the (strept) avidin system (Moter & Gobel, 2000). For a general review on the use of FISH to detect microbial populations in natural environments see Moter & Gobel (2000).

Several publications have tested the applicability of FISH for environmental detection and enumeration of pathogens or indicators. For example, 16S rRNA targeted oligonucleotide probes have been designed to differentiate Enterobacteriaceae both as a group and as individual species (Loge et al., 1999; Baudart et al., 2002; Rompre et al., 2002). Specifically, *E. coli*, *Enterobacter cloacae*, and *Citrobacter freundii* were identified after membrane filtration followed by FISH for water quality control purposes (Loge et al., 1999; Baudart et al., 2002). This enabled detection of 10^5 – 10^7 *E. coli* cells per ml of wastewater (Baudart et al., 2002) (Table 2.2). In general, because of the reliance on microscopy, the target population has to be present at >0.1% of the total cell numbers in the community, which translates to $\sim 10^3$ cells per ml for bacterioplankton in most natural waters. However, Colwell and colleagues have combined FISH with the high throughput cell counting ability of flow cytometry and were able to detect *Vibrio* populations at abundances as low as 13 cells per ml (Heidelberg et al., 2002b).

One major problem in FISH arises from the generally low signal level per probe provided by direct labeling procedures, and the low diffusion of large molecules through the cell wall in indirect labeling procedures. This has confined routine FISH application to use of rRNA as targets since these are present in hundreds and thousands of copies in actively growing cells (DeLong et al., 1989; Amann et al., 1990). Unfortunately, the ability of the rRNAs to discriminate among closely related organisms is limited since they are highly conserved molecules and contain only relatively short, variable nucleotide stretches. For example, while it is possible to identify *E. coli* on the species level, pathogenic strains cannot be distinguished

from harmless strains. However, over the past decade several improvements have been made to increase the sensitivity of FISH by use of brighter fluorochromes, signal amplification systems coupled to reporter enzymes, and multiply labeled probes (reviewed in Pernthaler et al., 2002a, 2002b). Thus, it is possible that in the near future more variable targets, such as messenger RNA, will be among the targets for FISH. Furthermore, if flow cytometry can be routinely combined with FISH more efficient sample analysis may arise since labor-intensive microscopy may be circumvented.

Dot-blot and Colony Hybridizations. In all filter hybridizations, nucleic acids are immobilized on membranes and hybridized with specific labeled probes. Various labels are available ranging from radionucleides to biotin or digoxigenin. The latter are detected with antibodies carrying enzymes, which elicit either a color precipitation or chemiluminescent reaction. In dot-blot hybridizations, the target nucleic acids are purified either from isolates or environmental samples; in colony hybridization, filter membranes are applied directly to culture plates and cells are transferred to the membranes, lysed, and their nucleic acids hybridized. In both methods either oligonucleotides against rRNA or polynucleotides against protein-coding mRNA (or genes) can be used as probes and the same considerations of varying ability of different types of probes to discriminate strains and species apply as for all hybridizations (see above).

Although dot-blot hybridization is routinely applied to detect bacterial populations in ecological studies (Koizumi et al., 2002; Polz and Cavanaugh, 1997; Raskin et al., 1994) it has only rarely been applied to monitoring of pathogens. It was recommended for its accuracy, speed, and low cost for detection of drug resistant *M. tuberculosis* strains (Victor et al., 1999), and produced a detection limit of 10^2 cells when albacore tuna muscle extract was artificially contaminated with the pathogen *Stenotrophomonas maltophilia* (Ben-Gigirey et al., 2002). Nonetheless, for reliable detection of environmental pathogens, culture enrichment prior to hybridization has been recommended due to uncertain detection limits and possible interference of inhibitors (Straub & Chandler, 2003).

Colony hybridization is essentially an extension of culture-dependent detection of pathogens and, although the same limitations based on culturability apply, it allows rapid, sensitive, and accurate identification of strains. Probes targeting the thermostable direct hemolysin (*tdh*) and/or *tdh*-related hemolysin (*trh*) genes enabled the detection of oyster-associated pathogenic *V. parahaemolyticus* strains at low densities (usually <10 CFU per g of oyster) (Blackstone et al., 2003). Colony hybridization has also been used for the study of the seasonal dynamics of *V. cholerae* along the California coastline with a dynamic range of three orders of magnitude and 1 CFU per ml as the lowest observed abundance (Jiang & Fu, 2001). The USFDA³ has recognized the high accuracy of colony hybridization and has approved a number of gene targets specific for food- and water-borne pathogens including Listeriolysin O 11 and *msp* genes of *L. monocytogenes*, the invasive genes of *Shigella* spp., enterotoxin B of *S. aureus*, the heat-stable toxin genes of *E. coli*, and the *ail* gene and *inv* genes of *Yersinia pseudotuberculosis* and *Yersinia enterocolitica* (Jackson et al., 2000).

2.5.3.2. Polymerase Chain Reaction

PCR-based detection has revolutionized diagnostic microbiology due to the combination of sensitivity, specificity, and fast turnaround time for identification of infectious agents. The PCR represents an enzymatic copying of specific genes allowing million- to billion-fold

³ United States Food and Drug Administration.

amplification above the background of single or mixtures of genomes. Double-stranded (genomic) DNA template is denatured and the resulting single strands hybridized with one of two primers, which flank the target gene. These primers are then extended with a thermostable DNA polymerase (e.g., Taq polymerase) generating copies of complementary DNA. This overall process is repeated between 20 and 45 times (cycles) in a single test tube. The specificity of the target amplification is determined by the design of sequence-specific primers and optimization of reaction conditions.

Due to the exponential amplification of templates, the PCR has an exceptional sensitivity of, theoretically, a single target copy; however, the PCR also has high potential for misleading results due to contamination, biases, and inhibition. Contamination most often stems from previous PCR reactions carried out in the same laboratory. For example, aerosols generated by pipetting or opening of reaction tubes are a major source of false positives but use of stuffed tips, laminar flow benches, and UV treatment of tubes can minimize such problems. The PCR is also subject to a number of biases, which generally become more pronounced with increasing reaction cycles. Specifically, after initial exponential amplification, the product accumulation becomes less efficient until a maximum product concentration is reached, which is independent of the amount of starting template. Since the starting template concentration and the efficiency of the amplification are unknown in the reaction, the results of simple PCR assays should never be quantitatively interpreted unless appropriate controls are included (see quantitative PCR (QPCR) below) (von Wintzingerode et al., 1997; Polz & Cavanaugh, 1998). Finally, inhibition of the PCR by environmental contaminants (e.g., humic substances and metal ions) may result in false negatives. To address this problem, various methods have been devised for the purification of nucleic acids from environmental samples prior to amplification (reviewed in von Wintzingerode et al., 1997). It has also been suggested to always include an internal standard in each PCR reaction to indicate possible PCR inhibitors (Malorny et al., 2003).

The unique potential of the PCR for rapid and specific detection of species- or virulence-specific genes has been exploited in numerous assays (for example, reviewed in Straub & Chandler, 2003; Pommepuy & Le Guyader, 1998) (Table 2.2 for detection of pathogens in environmental samples). However, many protocols still utilize some form of enrichment prior to PCR amplification due to the danger of false negatives from inhibition of reaction kinetics by environmental substances. Several techniques have been utilized including filtration, centrifugation, or molecular-based separation (e.g., by magnetic beads). For example, culture-based enrichment increased the sensitivity of a PCR assay for *L. monocytogenes* in household waste samples by several orders of magnitude from 10^7 to 10 CFU per g (Burtscher & Wuertz, 2003). Horgen and colleagues detected *E. coli* at 10 cells per ml of water by concentration of the cells with magnetic beads (Foulds et al., 2002). Other authors used culture enrichment prior to PCR of putative pathogens from water or other environmental samples (Table 2.2).

Several modifications of the PCR technique hold promise for increased accuracy or high-throughput detection of pathogens. The first technique, Quantitative PCR (QPCR), allows quantification of the abundance of target gene sequences in environmental samples. QPCR is available in several formats but real-time QPCR has become the most widely used. It detects the accumulation of DNA template at the end of every cycle. This enables comparison of template accumulation kinetics between environmental samples and standards for accurate quantification (Table 2.2). For higher throughput detection of multiple pathogens in a single test tube, multiplex PCR assays have been developed (Table 2.2). These combine cocktails of specific primers for several targets and allow differentiation of individual amplicons from the mixture of products either by size or labeling of the amplification primers with different fluorors.

These examples illustrate just a few of the large number of permutations of the PCR, which have been published. Overall, PCR-based methods are among the most rapid, flexible, and cost effective of the molecular methods, and it is therefore not surprising that many laboratories have concentrated on their use.

2.5.3.3. *Molecular Typing Methods*

Nucleic-acid-based molecular typing methods allow for the differentiation of strains based on analysis of their genomes. This is important for linking specific strains to disease outbreaks but is also critical for evaluating the specificity of detection methods by providing standards for virulent and harmless strains of the same species. Molecular typing can enable identification of traits unique to virulent strains. Molecular typing methods employed for distinguishing bacterial strains include PFGE, randomly (or arbitrarily) primed PCR, analysis of DNA sequences (e.g., ribosomal genes), and MLST (reviewed in Persing et al., 2003; van Belkum, 2003). Traditionally, the “gold standard” for typing has been PFGE but newer methods such as MLST are rapidly being translated into a format suitable for routine clinical identification of pathogens.

Pulsed Field Gel Electrophoresis. PFGE differentiates genomes by cutting chromosomal DNA with “rare-cutter” restriction enzymes, which due to long recognition sequences cut infrequently. This produces few, large DNA fragments (roughly 10–800 kb), which can be separated by gel electrophoresis under a pulsed-electric field. Both variation in sequence and overall genome architecture are translated into unique patterns of DNA fragments and allow highly specific identification of strains (reviewed in Persing et al., 2003; van Belkum, 2003). PFGE is currently widely applied in food safety assessment and a number of laboratories contribute to the “Foodborne Surveillance PulseNet,” a database created by the Centers for Disease Control and Prevention and several state and national laboratories (Binder et al., 1999; Swaminathan et al., 2001). PFGE patterns of strains stored in the database can be compared to those obtained from isolates from contaminated food or clinical samples by electronically submitting images to the network. Standardized PulseNet protocols have been developed for *E. coli* (Breuer et al., 2001), *Campylobacter jejuni* (Ribot et al., 2001), *L. monocytogenes* (Graves & Swaminathan, 2001), and several more are being developed and validated (Swaminathan et al., 2001). Such approaches have high potential for better understanding of the diversity of strains responsible for disease outbreaks; however, PFGE in particular remains a challenging technique to implement reproducibly among different laboratories, and other whole genome comparative methods may ultimately replace PFGE for the routine characterization of isolates.

Multilocus Sequence Typing. One very promising alternative for characterizing bacterial isolates is MLST, which produces nucleotide sequence data that can be readily compared between laboratories and in different studies. In this technique, several defined DNA regions of each bacterial isolate are amplified by PCR and subsequently sequenced. The various sequences of about 500bp are aligned to detect nucleotide differences and sorted into allele homology groups. Since multiple genes are included in the analysis, characteristic allelic profiles can be used to identify pathogenic strains (Maiden et al., 1998; van Belkum, 2003).

2.5.3.4. *Future Nucleic Acid-Based Technologies*

DNA microarrays hold promise to improve environmental pathogen monitoring by allowing high-throughput detection of multiple pathogen populations in a single analysis. DNA

microarrays allow the differentiation of hundreds to thousands of specific sequences in a sample by simultaneous reverse dot-blot hybridizations [reviewed in (Ye et al., 2001; Call et al., 2003)]. Different, specific probes are attached to a glass slide, and fluorescently labeled target nucleic acids are hybridized to the probes in a single reaction. After stringent washes to remove non-specific hybrids, the hybridization signals are imaged using high-resolution scanners. Recently, DNA microarrays have been applied to the detection of bacteria in soil (Wu et al., 2001) and estuarine water samples (Taroncher- Oldenburg et al., 2003). However, several challenges remain before microarrays can be routinely used for pathogen detection in environmental samples or clinical specimens. Perhaps, the most critical challenge is how to optimize the stringency of the analysis conditions for simultaneous hybridization of multiple probes with different chemical properties. This limitation allows only detection of positive hybridization signals with respect to defined standards and can confound interpretation of hybridization signals from environmental samples due to non-specific cross-hybridization. Furthermore, the cost of equipment, expertise, and large-scale data analysis remains prohibitively high, relegating the use of microarray technology to a few centralized facilities. However, with recent advances in high-throughput genome analysis, microarray technology will prove to be a very valuable tool in clinical and environmental microbiology with applications for the detection and molecular typing of marine pathogens.

The ultimate form of molecular typing is whole genome sequencing. With increased analysis of diverse bacterial genomes, information on strain-to-strain variation and the transfer of virulence properties among bacterial species is becoming available. The genome sequences of over two hundred bacteria have been published, including a number of strains that are marine pathogens, or close relatives thereof (e.g., *Vibrio* spp. including *V. cholerae*, *V. parahaemolyticus*, and *V. vulnificus*, *Brucella* spp., *Clostridium* spp., *Legionella* spp., *Mycobacterium* spp. and *Shewanella* spp.) and this number is increasing rapidly with advances in high-throughput sequencing technology. Comparative genomic analysis has revealed surprising levels genomic diversity among closely-related bacterial strains (Welch, 2002) and analysis of genomes from pathogenic and non-pathogenic organisms is revealing mechanisms by which pathogenic interactions emerge while providing genetic targets to differentiate virulent from avirulent strains. Such information will be critical for the design of molecular assays to detect and monitor specific pathogens in clinical and environmental settings.

2.6. OUTLOOK

A surprisingly large number of potential human pathogens reside in the marine environment and increased risk of human exposure highlights the need to better understand their ecology and evolution. An integral part of such an effort must be the specific characterization, differentiation, and detection of pathogenic strains. Particular challenges are the potential range expansion of existing marine-indigenous pathogens (e.g., *V. cholerae*) and the emergence of new human-pathogens from marine systems. Indeed, increased reports of disease outbreaks in marine populations may evidence the emergence of new pathogens. The zoonotic potential of such outbreaks in natural or farmed marine environments needs to be recognized and approached with caution while work is done to recognize and prevent the conditions that promote marine disease. The complexity of these problems requires flexible approaches and the overview provided in this chapter attempts to represent methods, which allow both routine

monitoring of pathogens and exploration of their ecology. In the future, coordinated efforts to standardize methods and create databases for comparison will be important for a more comprehensive evaluation of the risk for human populations associated with utilization of marine environments.

REFERENCES

- Abeyta, C., Deeter, F.G., Kaysner, C.A., Stott, R.F., & Wekell, M.M. (1993). *Campylobacter-jejuni* in a Washington-state shellfish growing bed associated with illness. *J Food Protect* **56**, 323–325.
- Ahlen, C., Mandal, L.H., Johannessen, L.N., & Iversen, O.J. (2000). Survival of infectious *Pseudomonas aeruginosa* genotypes in occupational saturation diving environment and the significance of these genotypes for recurrent skin infections. *Am J Indust Med* **37**, 493–500.
- Alam, M.J., Miyoshi, S., & Shinoda, S. (2003). Studies on pathogenic *Vibrio parahaemolyticus* during a warm weather season in the Seto Inland Sea, Japan. *Environ Microbiol* **5**, 706–710.
- Almuneef, M., & Memish, Z.A. (2003). Prevalence of Brucella antibodies after acute brucellosis. *J Chemother* **15**, 148–151.
- Altekruse, S.F., Cohen, M.L., & Sberdlow, D.L. (1997). Emerging foodborne diseases. *Emerg Infect Dis* **3**, 285–293.
- Amann, R.I., Krumholz, L., & Stahl, D.A. (1990). Fluorescent-oligonucleotide probing of whole cells for determinative, phylogenetic, and environmental studies in microbiology. *J Bacteriol* **172**, 762–770.
- Anderson, I.C., Rhodes, M., & Kator, H. (1979). Sublethal stress in *Escherichia coli*: A function of salinity. *Appl Environ Microbiol* **38**, 1147–1152.
- Arzouni, J.P., Parola, P., La Scola, B., Postic, D., Brouqui, P., & Raoult, D. (2002). Human infection caused by *Leptospira fainei*. *Emerg Infect Dis* **8**, 865–868.
- Asakura, H., Watarai, M., Shirahata, T., & Makino, S. (2002). Viable but nonculturable *Salmonella* species recovery and systemic infection in morphine-treated mice. *J Infect Dis* **186**, 1526–1529.
- Aschfalk, A., Folkow, L., Rud, H., & Denzin, N. (2002). Apparent seroprevalence of *Salmonella* spp. in harp seals in the Greenland Sea as determined by enzyme-linked immunosorbent assay. *Vet Res Commun* **26**, 523–530.
- Aschfalk, A., & Muller, W. (2001). Clostridium perfringens toxin types in hooded seals in the Greenland Sea, determined by PCR and ELISA. *J Vet Med B Infect Dis Vet Public Health* **48**, 765–769.
- Ashbolt, N.J., Ball, A., Dorsch, M., Turner, C., Cox, P., Chapman, A., & Kirov, S.M. (1995). The identification and human health significance of environmental aeromonads. *Water Sci Technol* **31**, 263–269.
- Atlas, R.M. (1995). *Handbook of Media for Environmental Microbiology*. CRC Press, Boca Raton, FL.
- Austin, B., Garges, S., Conrad, B., Harding, E.E., Colwell, R.R., Simidu, U., and Taga, N. (1979). Comparative study of the aerobic, heterotrophic bacterial-flora of Chesapeake Bay and Tokyo Bay. *Appl Environ Microbiol* **37**, 704–714.
- Baker, A.S., Ruoff, K.L., & Madoff, S. (1998). Isolation of Mycoplasma species from a patient with seal finger. *Clin Infect Dis* **27**, 1168–1170.
- Barber, G.R., & Swygert, J.S. (2000). Necrotizing Fasciitis due to *Photobacterium damsela* in a man lashed by a stingray. *N Engl J Med* **342**, 824.
- Bartie, C., Venter, S.N., & Nel, L.H. (2003). Identification methods for *Legionella* from environmental samples. *Water Res* **37**, 1362–1370.
- Bates, T.C., Adams, B.L., & Oliver, J.D. (2002). The viable but nonculturable state of *Helicobacter pylori*. *Gut* **51**, A5–A6.
- Baudart, J., Coallier, J., Laurent, P., & Prevost, M. (2002). Rapid and sensitive enumeration of viable diluted cells of members of the family enterobacteriaceae in freshwater and drinking water. *Appl Environ Microbiol* **68**, 5057–5063.
- Baylor, E.R., Baylor, M.B., Blanchard, D.C., Syzdek, L.D., & Appel, C. (1977) Virus transfer from surf to wind. *Science* **198**, 575–580.
- Ben-Gigirey, B., Vieites, J.M., Kim, S.H., An, H.J., Villa, T.G., & Barros-Velazquez, J. (2002). Specific detection of *Stenotrophomonas maltophilia* strains in albacore tuna (*Thunnus alalunga*) by reverse dot-blot hybridization. *Food Control* **13**, 293–299.

- Bernardelli, A., Bastida, R., Loureiro, J., Michelis, H., Romano, M.I., Cataldi, A., & Costa, E. (1996). Tuberculosis in sea lions and fur seals from the south-western Atlantic coast. *Rev Sci Tech Off Int Epizoot* **15**, 985–1005.
- Bernardet, J.F. (1998). Cytophaga, Flavobacterium, Flexibacter and Chryseobacterium infections in cultured marine fish. *Fish Pathol* **33**, 229–238.
- Besnard, V., Federighi, M., & Cappelier, J.M. (2000). Evidence of viable but non-culturable state in *Listeria monocytogenes* by direct viable count and CTC-DAPI double staining. *Food Microbiol* **17**, 697–704.
- Bhagwat, A.A. (2003). Simultaneous detection of *Escherichia coli* O157:H7, *Listeria monocytogenes* and *Salmonella* strains by real-time PCR. *Int J Food Microbiol* **84**, 217–224.
- Biegala, I.C., Kennaway, G., Alverca, E., Lennon, J.F., Vaultot, D., & Simon, N. (2002). Identification of bacteria associated with dinoflagellates (Dinophyceae) *Alexandrium* spp. using tyramide signal amplification-fluorescent in situ hybridization and confocal microscopy. *J Phycol* **38**, 404–411.
- Binder, S., Levitt, A.M., Sacks, J.J., & Hughes, J.M. (1999). Emerging infectious diseases: Public health issues for the 21st century. *Science* **284**, 1311–1313.
- Biosca, E.G., Marco-Noales, E., Amaro, C., & Alcaide, E. (1997). An enzyme-linked immunosorbent assay for detection of *Vibrio vulnificus* biotype 2: Development and field studies. *Appl Environ Microbiol* **63**, 537–542.
- Blackstone, G.M., Nordstrom, J.L., Vickery, M.C., Bowen, M.D., Meyer, R.F., & DePaola, A. (2003). Detection of pathogenic *Vibrio parahaemolyticus* in oyster enrichments by real time PCR. *J Microbiol Methods* **53**, 149–155.
- Boehm, A.B., Grant, S.B., Kim, J.H., Mowbray, S.L., McGee, C.D., Clark, C.D., Foley, D.M., & Wellman, D.E. (2002). Decadal and shorter period variability of surf zone water quality at Huntington Beach, California. *Environmental Science & Technology* **36**, 3885–3892.
- Boerlin, P., Boerlin-Petzold, F., & Jemmi, T. (2003). Use of listeriolysin O and internalin A in a seroepidemiological study of listeriosis in Swiss dairy cows. *J Clin Microbiol* **41**, 1055–1061.
- Boulanger, C.A., & Edelstein, P.H. (1995). Precision and accuracy of recovery of *Legionella pneumophila* from seeded tap water by filtration and centrifugation. *Appl Environ Microbiol* **61**, 1805–1809.
- Boyd, E., Moyer, K., & Shi, L. (2000). Infectious CTX Phi, and the vibrio pathogenicity island prophage in *Vibrio mimicus*: Evidence for recent horizontal transfer between *V. mimicus* and *V. cholerae*. *Infect Immun* **68**, 1507–1513.
- Brayton, P.R., Bode, R.B., Colwell, R.R., MacDonell, M.T., Hall, H.L., Grimes, D.J., West, P.A., & Bryant, T.N. (1986). *Vibrio cincinnatiensis* sp. nov., a new human pathogen. *J Clin Microbiol* **23**, 104–108.
- Brayton, P.R., Tamplin, M.L., Huq, A., & Colwell, R.R. (1987). Enumeration of *Vibrio cholerae* O1 in Bangladesh waters by fluorescent-antibody direct viable count. *Appl Environ Microbiol* **53**, 2862–2865.
- Breuer, T., Benkel, D.H., Shapiro, R.L., Hall, W.N., Winnett, M.M., Linn, M.J., Neimann, J., Barrett, T.J., Dietrich, S., Downes, F.P., et al. (2001). A multistate outbreak of *Escherichia coli* O157:H7 infections linked to alfalfa sprouts grown from contaminated seeds. *Emerg Infect Dis* **7**, 977–982.
- Brew, S.D., Perrett, L.L., Stack, J.A., MacMillan, A.P., & Staunton, N.J. (1999). Human exposure to *Brucella* recovered from a sea mammal. *Vet Rec* **144**, 483.
- Brooke, C.J., & Riley, T.V. (1999). *Erysipelothrix rhusiopathiae*: Bacteriology, epidemiology and clinical manifestations of an occupational pathogen. *J Med Microbiol* **48**, 789–799.
- Buck, J.D. (1991) Recovery of *Vibrio Metschnikovii* from market seafood. *J Food Safety* **12**, 73–78.
- Burke, V., Robinson, J., Gracey, M., Peterson, D., & Partridge, K. (1984). Isolation of *Aeromonas hydrophila* from a metropolitan water supply: seasonal correlation with clinical isolates. *Appl Environ Microbiol* **48**, 361–366.
- Burke, W.A., & Tester, P.A. (2002). Skin problems related to noninfectious coastal microorganisms. *Dermatol Ther* **15**, 10–17.
- Burkhardt, W., Watkins, W.D., & Rippey, S.R. (1992). Seasonal effects on accumulation of microbial indicator organisms by *Mercenaria-Mercenaria*. *Appl Environ Microbiol* **58**, 826–831.
- Burtscher, C., & Wuertz, S. (2003). Evaluation of the use of PCR and reverse transcriptase PCR for detection of pathogenic bacteria in biosolids from anaerobic digestors and aerobic composters. *Appl Environ Microbiol* **69**, 4618–4627.
- Cabelli, V.J., Dufour, A.P., McCabe, L.J., & Levin, M.A. (1982). Swimming-associated gastroenteritis and water quality. *Am J Epidemiol* **115**, 606–616.
- Call, D.R., Borucki, M.K., & Loge, F.J. (2003). Detection of bacterial pathogens in environmental samples using DNA microarrays. *J Microbiol Methods* **53**, 235–243.
- Cappelier, J.M., Magras, C., Jouve, J.L., & Federighi, M. (1999a). Recovery of viable but non-culturable *Campylobacter jejuni* cells in two animal models. *Food Microbiol* **16**, 375–383.

- Cappelier, J.M., Minet, J., Magras, C., Colwell, R.R., & Federighi, M. (1999b). Recovery in embryonated eggs of viable but nonculturable *Campylobacter jejuni* cells and maintenance of ability to adhere to HeLa cells after resuscitation. *Appl Environ Microbiol* **65**, 5154–5157.
- Carmichael, W.W. (2001). Health effects of toxin-producing cyanobacteria: “The CyanoHABs.” *Hum Ecol Risk Assess* **7**, 1393–1407.
- Caruso, G., Zacccone, R., & Crisafi, E. (2000). Use of the indirect immunofluorescence method for detection and enumeration of *Escherichia coli* in seawater samples. *Lett Appl Microbiol* **31**, 274–278.
- Caruso, G., Crisafi, E., & Mancuso, M. (2002). Immunofluorescence detection of *Escherichia coli* in seawater: A comparison of various commercial antisera. *J Immunoassay Immunochem* **23**, 479–496.
- Caudell, M.J., and Kuhn, W.F. (1997). *Aeromonas hydrophila* soft-tissue infection: A report of two cases. *Acad Emerg Med* **4**, 157–158.
- CDC (1999). *Vibrio Surveillance System, Summary Data, 1997–1998*. Department of Health and Human Services, Atlanta, GA.
- CDC (2000). *Summary of Infections Reported to Vibrio Surveillance System, 1999*. Department of Health and Human Services, Atlanta, GA.
- Cerda-Cuellar, M., Permin, L., Larsen, J.L., & Blanch, A.R. (2001). Comparison of selective media for the detection of *Vibrio vulnificus* in environmental samples. *J Appl Microbiol* **91**, 322–327.
- Chakraborty, S., Mukhopadhyay, A.K., Bhadra, R.K., Ghosh, A.N., Mitra, R., Shimada, T., Yamasaki, S., Faruque, S.M., Takeda, Y., Colwell, R.R., et al. (2000). Virulence genes in environmental strains of *Vibrio cholerae*. *Applied and Environmental Microbiology* **66**, 4022–4028.
- Chan, S.S.W., Ng, K.C., Lyon, D.J., Cheung, W.L., Cheng, A.F.B., & Rainer, T.H. (2003). Acute bacterial gastroenteritis: A study of adult patients with positive stool cultures treated in the emergency department. *Emerg Med J* **20**, 335–338.
- Charoenc, N., & Fujioka, R.S. (1993). Assessment of *Staphylococcus bacteria* in Hawaii marine recreational waters. *Water Sci Technol* **27**, 283–289.
- Chowdhury, M.A.R., Yamanaka, H., Miyoshi, S., & Shinoda, S. (1990). Ecology of Mesophilic aeromonas spp. in aquatic environments of a Temperate region and relationship with some biotic and abiotic environmental parameters. *Zentralblatt Hygiene Umweltmedizin* **190**, 344–356.
- Chrost, R.J., & Faust, M.A. (1999). Consequences of solar radiation on bacterial secondary production and growth rates in subtropical coastal water (Atlantic coral reef of Belize, Central America). *Aquat Microb Ecol* **20**, 39–48.
- Cianciotto, N.P., & Fields, B.S. (1992). *Legionella pneumophila* mip gene potentiates intracellular infection of protozoa and human macrophages. *Proc Natl Acad Sci USA* **89**, 5188–5191.
- Cirillo, J.D., Falkow, S., Tompkins, L.S., & Bermudez, L.E. (1997). Interaction of *Mycobacterium avium* with environmental amoebae enhances virulence. *Infect Immun* **65**, 3759–3767.
- Colagross-Schouten, A.M., Mazet, J.A., Gulland, F.M., Miller, M.A., & Hietala, S. (2002). Diagnosis and seroprevalence of leptospirosis in California sea lions from coastal California. *J Wildl Dis* **38**, 7–17.
- Colburn, K.G., Kaysner, C.A., Abeyta, C., & Wekell, M.M. (1990). *Listeria* species in a California coast estuarine environment. *Appl Environ Microbiol* **56**, 2007–2011.
- Cole, J.R., Chai, B., Marsh, T.L., Farris, R.J., Wang, Q., Kulam, S.A., Chandra, S., McGarrell, D.M., Schmidt, T.M., Garrity, G.M., et al. (2003). The Ribosomal Database Project (RDP-II): previewing a new autoaligner that allows regular updates and the new prokaryotic taxonomy. *Nucleic Acids Res.* **31**, 442–443.
- Colorni, A., Diamant, A., Eldar, A., Kvitt, H., & Zlotkin, A. (2002). *Streptococcus iniae* infections in Red Sea cage-cultured and wild fishes. *Dis Aquat Organ* **49**, 165–170.
- Colwell, R.R. (1996). Global climate and infectious disease: The cholera paradigm. *Science* **274**, 2025–2031.
- Corbel, M.J. (1997). Brucellosis: an overview. *Emerg Infect Dis* **3**, 213–221.
- Corbett, S.J., Rubin, G.L., Curry, G.K., & Kleinbaum, D.G. (1993). The health effects of swimming at Sydney beaches. The Sydney Beach Users Study Advisory Group. *Am J Public Health* **83**, 1701–1706.
- Dalsgaard, A. (1998). The occurrence of human pathogenic *Vibrio* spp. and *Salmonella* in aquaculture. *Int J Food Sci Technol* **33**, 127–138.
- Dalsgaard, A., Glerup, P., Hoybye, L.L., Paarup, A.M., Meza, R., Bernal, M., Shimada, T., & Taylor, D.N. (1997). *Vibrio furnissii* isolated from humans in Peru: a possible human pathogen? *Epidemiol Infect* **119**, 143–149.
- Daum, L.T., Barnes, W.J., McAvin, J.C., Neidert, M.S., Cooper, L.A., Huff, W.B., Gaul, L., Riggins, W.S., Morris, S., Salmen, A., et al. (2002). Real-time PCR detection of salmonella in suspect foods from a gastroenteritis outbreak in Kerr county, Texas. *J Clin Microbiol* **40**, 3050–3052.

- De la Torre, C., Vega, A., Carracedo, A., and Toribio, J. (2001). Identification of *Mycobacterium marinum* in sea-urchin granulomas. *Br J Dermatol* **145**, 114–116.
- DeLong, E.F., Wickham, G.S., & Pace, N.R. (1989). Phylogenetic stains: ribosomal RNA-based probes for the identification of single cells. *Science* **243**, 1360–1363.
- DePaola, A., Kaysner, C.A., Bowers, J., & Cook, D.W. (2000). Environmental investigations of *Vibrio parahaemolyticus* in oysters after outbreaks in Washington, Texas, and New York (1997 and 1998). *Appl Environ Microbiol* **66**, 4649–4654.
- Dillon, R., Patel, T., & Ratnam, S. (1994). Occurrence of *Listeria* in hot and cold smoked seafood products. *Int J Food Microbiol* **22**, 73–77.
- Dobos, K.M., Quinn, F.D., Ashford, D.A., Horsburgh, C.R., & King, C.H. (1999). Emergence of a unique group of necrotizing mycobacterial diseases. *Emerg Infect Dis* **5**, 367–378.
- Dominguez, H., Vogel, B.F., Gram, L., Hoffmann, S., & Schaevel, S. (1996). *Shewanella alga* bacteremia in two patients with lower leg ulcers. *Clin Infect Dis* **22**, 1036–1039.
- Dorfman, M. (2003). *Testing the Waters XIII: A Guide to Water Quality at Vacation Beaches*. Natural Resources Defense Council, New York.
- Duffy, G., Kilbride, B., Sheridan, J.J., Blair, I.S., & McDowell, D.A. (2000). A membrane-immunofluorescent-viability staining technique for the detection of *Salmonella* spp. from fresh and processed meat samples. *J Appl Microbiol* **89**, 587–594.
- Dufour, A.P., Ericksen, T.H., Ballentine, R.K., Cabelli, V.J., Goldberg, M., & Fox, W.E. (1986). Bacteriological ambient water quality criteria for marine and fresh recreational waters. *Ambient Water Quality Criteria for Bacteria*. U.S. Environmental Protection Agency, Washington, DC.
- Dumontet, S., Krovacek, K., Svenson, S.B., Pasquale, V., Baloda, S.B., & Figliuolo, G. (2000). Prevalence and diversity of *Aeromonas* and *Vibrio* spp. in coastal waters of southern Italy. *Comp Immunol, Microbiol Infect Dis* **23**, 53–72.
- Dupray, E., Caprais, M.P., Derrien, A., Fach, P. (1997). *Salmonella* DNA persistence in natural seawaters using PCR analysis. *J Appl Microbiol* **82**, 507–510.
- Dutta, S., Deb, A., Chattopadhyay, U.K., & Tsukamoto, T. (2000). Isolation of Shiga toxin-producing *Escherichia coli* including O157:H7 strains from dairy cattle and beef samples marketed in Calcutta, India. *J Med Microbiol* **49**, 765–767.
- Eilers, H., Pernthaler, J., & Amann, R. (2000a). Succession of pelagic marine bacteria during enrichment: A close look at cultivation-induced shifts. *Appl Environ Microbiol* **66**, 4634–4640.
- Eilers, H., Pernthaler, J., Glockner, F.O., & Amann, R. (2000b). Culturability and in situ abundance of pelagic bacteria from the North Sea. *Appl Environ Microbiol* **66**, 3044–3051.
- Ender, P.T., & Dolan, M.J. (1997). Pneumonia associated with near-drowning. *Clin Infect Dis* **25**, 896–907.
- Endtz, H.P., Vliegthart, J.S., Vandamme, P., Weverink, H.W., vandenBraak, N.P., Verbrugh, H.A., & vanBelkum, A. (1997). Genotypic diversity of *Campylobacter lari* isolated from mussels and oysters in The Netherlands. *Int J Food Microbiol* **34**, 79–88.
- Erickson, T., Vandenhoek, T.L., Kuritza, A., & Leiken, J.B. (1992). The emergency management of moray eel bites. *Ann Emerg Med* **21**, 212–216.
- Faruque, S.M., Asadulghani, Saha, M.N., Alim, A., Albert, M.J., Islam, K.M.N., & Mekalanos, J.J. (1998). Analysis of clinical and environmental strains of nontoxigenic *Vibrio cholerae* for susceptibility to CTX Phi: Molecular basis for origination of new strains with epidemic potential. *Infect Immun* **66**, 5819–5825.
- Faruque, S.M., Khan, R., Kamruzzaman, M., Yamasaki, S., Ahmad, Q.S., Azim, T., Nair, G.B., Takeda, Y., & Sack, D.A. (2002). Isolation of *Shigella dysenteriae* type 1 and *S. flexneri* strains from surface waters in Bangladesh: comparative molecular analysis of environmental *Shigella* isolates versus clinical strains. *Appl Environ Microbiol* **68**, 3908–3913.
- Faruque, S., Rahman, M., Asadulghani, Nasirul-Islam, K.M., & Mekalanos, J.J. (1999). Lysogenic conversion of environmental *Vibrio* mimivirus strains by CTX Phi. *Infect. Immun.* **67**, 5723–5729.
- Faust, M.A., Aotaky, A.E., & Hargadon, M.T. (1975). Effect of physical parameters on the in situ survival of *Escherichia coli* MC-6 in an estuarine environment. *Appl Microbiol* **30**, 800–806.
- FDA (1994). Proposal to establish procedures for the safe processing and importing of fish and fishery products; proposed rules. *Federal Register*. Washington, DC, pp. 4142–4214.
- FDA (1997). *National Shellfish Sanitation Program Guide for the Control of Molluscan Shellfish*. Washington, DC.
- Fefer, J.J., Ratzan, K.R., Sharp, S.E., & Saiz, E. (1998). *Lactococcus garvieae* endocarditis: Report of a case and review of the literature. *Diagn Microbiol Infect Dis* **32**, 127–130.
- Feldhusen, F. (2000). The role of seafood in bacterial foodborne diseases. *Microbes Infect* **2**, 1651–1660.

- Fidalgo, S.G., Wang, Q., & Riley, T.V. (2000). Comparison of methods for detection of *Erysipelothrix* spp. and their distribution in some Australasian seafoods. *Appl Environ Microbiol* **66**, 2066–2070.
- Fields, B.S. (1996). The molecular ecology of legionellae. *Trends Microbiol* **4**, 286–290.
- Fiorentini, C., Barbieri, E., Falzano, L., Matarrese, P., Baffone, W., Pianetti, A., Katouli, M., Kuhn, I., Mollby, R., Bruscolini, F., et al. (1998). Occurrence, diversity and pathogenicity of mesophilic *Aeromonas* in estuarine waters of the Italian coast of the Adriatic Sea. *Journal of Applied Microbiology* **85**, 501–511.
- Fliermans, C.B. (1996). Ecology of *Legionella*: From data to knowledge with a little wisdom. *Microb Ecol* **32**, 203–228.
- Fliermans, C.B., Cherry, W.B., Orrison, L.H., Smith, S.J., Tison, D.L., and Pope, D.H. (1981). Ecological distribution of *Legionella pneumophila*. *Appl Environ Microbiol* **41**, 9–16.
- Foster, G., MacMillan, A.P., Godfroid, J., Howie, F.E., Ross, H.M., Cloeckaert, A., Reid, R.J., Brew, S., & Patterson, I.A. (2002). A review of *Brucella* sp. infection of sea mammals with particular emphasis on isolates from Scotland. *Veterinary Microbiology* **90**, 563–580.
- Foulds, I.V., Granacki, A., Xiao, C., Krull, U.J., Castle, A., & Horgen, P.A. (2002). Quantification of microcystin-producing cyanobacteria and *E. coli* in water by 5'-nuclease PCR. *J Appl Microbiol* **93**, 825–834.
- Fraser, S.L., Purcell, B.K., Delgado, B., Jr., Baker, A.E., & Whelen, A.C. (1997). Rapidly fatal infection due to *Photobacterium (Vibrio) damsela*. *Clin Infect Dis* **25**, 935–936.
- Fryer, J.L., & Mael, M.J. (1997). The Rickettsia: An emerging group of pathogens in fish. *Emerg Infect Dis* **3**, 137–144.
- Fujioka, R.S. (1997). Indicators of marine recreational water quality. In M. V. Walter (ed), *Manual of Environmental Microbiology*. ASM Press, Washington, DC, pp. 176–183.
- Fujioka, R.S., & Yoneyama, B.S. (2002). Sunlight inactivation of human enteric viruses and fecal bacteria. *Water Sci Technol* **46**, 291–295.
- Gallacher, S., & Smith, E.A. (1999). Bacteria and paralytic shellfish toxins. *Protist* **150**, 245–255.
- Garrett, E.S., dos Santos, C.L., & Jahncke, M.L. (1997). Public, animal, and environmental health implications of aquaculture. *Emerg Infect Dis* **3**, 453–457.
- Gerba, C.P., & McLeod, J.S. (1976). Effect of sediments on the survival of *Escherichia coli* in marine waters. *Appl Environ Microbiol* **32**, 114–120.
- Germani, Y., Begaud, E., & Desperrier, J.M. (1994a). Easy-to-perform modified Elek test to identify Shiga-like toxin-producing diarrhoeogenic *Escherichia coli*. *Res Microbiol* **145**, 333–340.
- Germani, Y., De Roquigny, H., & Begaud, E. (1994b). *Escherichia coli* heat-stable enterotoxin (STa)-biotin enzyme-linked immunosorbent assay (STa-biotin ELISA). *J Immunol Methods* **173**, 1–5.
- Glasby, C., & Hatheway, C.L. (1985). Isolation and enumeration of *Clostridium botulinum* by direct inoculation of infant fecal specimens on egg yolk agar and *Clostridium botulinum* isolation media. *J Clin Microbiol* **21**, 264–266.
- Gonzalez, C.J., Lopez-Diaz, T.M., Garcia-Lopez, M.L., Prieto, M., & Otero, A. (1999). Bacterial microflora of wild brown trout (*Salmo trutta*), wild pike (*Esox lucius*), and aquacultured rainbow trout (*Oncorhynchus mykiss*). *J Food Protect* **62**, 1270–1277.
- Gram, L., Bundvad, A., Melchiorson, J., Johansen, C., & Vogel, B.F. (1999). Occurrence of *Shewanella algae* in Danish coastal water and effects of water temperature and culture conditions on its survival. *Appl Environ Microbiol* **65**, 3896–3900.
- Grant, S.B., Sanders, B.F., Boehm, A.B., Redman, J.A., Kim, J.H., Mrse, R.D., Chu, A.K., Gouldin, M., McGee, C.D., Gardiner, N.A., et al. (2001). Generation of enterococci bacteria in a coastal saltwater marsh and its impact on surf zone water quality. *Environmental Science & Technology* **35**, 2407–2416.
- Graves, L.M., & Swaminathan, B. (2001). PulseNet standardized protocol for subtyping *Listeria monocytogenes* by macrorestriction and pulsed-field gel electrophoresis. *Int J Food Microbiol* **65**, 55–62.
- Griffin, D.W., Lipp, E.K., McLaughlin, M.R., & Rose, J.B. (2001). Marine recreation and public health microbiology: Quest for the ideal indicator. *Bioscience* **51**, 817–825.
- Grimes, D.J. (1991). Ecology of Estuarine bacteria capable of causing human-disease—a review. *Estuaries* **14**, 345–360.
- Grimes, D.J., Atwell, R.W., Brayton, P.R., Palmer, L.M., Rollins, D.M., Roszak, D.B., Singleton, F.L., Tamplin, M.L., & Colwell, R.R. (1986). The Fate of Enteric Pathogenic Bacteria in Estuarine and Marine Environments. *Microbiological Sciences* **3**, 324–329.
- Grimes, D.J., Singleton, F.L., & Colwell, R.R. (1984). Allogenic succession of marine bacterial communities in response to pharmaceutical waste. *J Appl Bacteriol* **57**, 247–261.

- Grohskopf, L.A., Roth, V.R., Feikin, D.R., Arduino, M.J., Carson, L.A., Tokars, J.I., Holt, S.C., Jensen, B.J., Hoffman, R.E., & Jarvis, W.R. (2001). *Serratia liquefaciens* Bloodstream Infections from Contamination of Epoetin Alfa at a Hemodialysis Center. *N Engl J Med* **344**, 1491–1497.
- Gulland, F.M., Koski, M., Lowenstine, L.J., Colagross, A., Morgan, L., & Spraker, T. (1996). Leptospirosis in California sea lions (*Zalophus californianus*) stranded along the central California coast, 1981–1994. *J Wildl Dis* **32**, 572–580.
- Hara-Kudo, Y., Sugiyama, K., Nishibuchi, M., Chowdhury, A., Yatsuyanagi, J., Ohtomo, Y., Saito, A., Nagano, H., Nishina, T., Nakagawa, H., et al. (2003). Prevalence of pandemic thermostable direct hemolysin-producing *Vibrio parahaemolyticus* O3:K6 in seafood and the coastal environment in Japan. *Appl Environ Microbiol* **69**, 3883–3891.
- Harb, O.S., Gao, L.Y., & Abu Kwaik, Y. (2000). From protozoa to mammalian cells: A new paradigm in the life cycle of intracellular bacterial pathogens. *Environ Microbiol* **2**, 251–265.
- Harper, C.M.G., Dangler, C.A., Xu, S.L., Feng, Y., Shen, Z.L., Sheppard, B., Stamper, A., Dewhirst, F.E., Paster, B.J., & Fox, J.G. (2000). Isolation and characterization of a *Helicobacter* sp from the gastric mucosa of dolphins, *Lagenorhynchus acutus* and *Delphinus delphis*. *Applied and Environmental Microbiology* **66**, 4751–4757.
- Hartmans, S., & DeBont, J.A.M. (1999). The Genus *Mycobacterium*-nonmedical. In M. Dworkin (ed), *The Prokaryotes*. Springer-Verlag, New York, pp. <http://link.springer-ny.com/link/service/books/10125/>
- Harvell, C.D., Kim, K., Burkholder, J.M., Colwell, R.R., Epstein, P.R., Grimes, D.J., Hofmann, E.E., Lipp, E.K., Osterhaus, A., Overstreet, R.M., et al. (1999). Review: Marine ecology—Emerging marine diseases—Climate links and anthropogenic factors. *Science* **285**, 1505–1510.
- Harvell, C.D., Mitchell, C.E., Ward, J.R., Altizer, S., Dobson, A.P., Ostfeld, R.S., & Samuel, M.D. (2002). Ecology—climate warming and disease risks for terrestrial and marine biota. *Science* **296**, 2158–2162.
- Hasan, J.A., Huq, A., Nair, G.B., Garg, S., Mukhopadhyay, A.K., Loomis, L., Bernstein, D., & Colwell, R.R. (1995). Development and testing of monoclonal antibody-based rapid immunodiagnostic test kits for direct detection of *Vibrio cholerae* O139 synonym Bengal. *J Clin Microbiol* **33**, 2935–2939.
- Heidelberg, J.F., Heidelberg, K.B., & Colwell, R.R. (2002a). Bacteria of the gamma-subclass Proteobacteria associated with zooplankton in Chesapeake Bay. *Appl Environ Microbiol* **68**, 5498–5507.
- Heidelberg, J.F., Heidelberg, K.B., & Colwell, R.R. (2002b). Seasonality of Chesapeake Bay bacterioplankton species. *Appl Environ Microbiol* **68**, 5488–5497.
- Heller, R., Holler, C., Sussmuth, R., & Gundermann, K.O. (1998). Effect of salt concentration and temperature on survival of *Legionella pneumophila*. *Lett Appl Microbiol* **26**, 64–68.
- Hernandezlopez, J., Guzmanmurillo, M.A., & Vargasalbores, F. (1995). Quantification of pathogenic marine *Vibrio* using membrane—filter technique. *J Microbiol Methods* **21**, 143–149.
- Hicks, C.L., Kinoshita, R., & Ladds, P.W. (2000). Pathology of melioidosis in captive marine mammals. *Aust Vet J* **78**, 193–195.
- Hoi, L., & Dalsgaard, A. (2000). Evaluation of a simplified semi-quantitative protocol for the estimation of *Vibrio vulnificus* in bathing water using cellobiose-colistin agar: A collaborative study with 13 municipal food controlling units in Denmark. *J Microbiol Methods* **41**, 53–57.
- Hoi, L., Larsen, J.L., Dalsgaard, I., & Dalsgaard, A. (1998). Occurrence of *Vibrio vulnificus* biotypes in Danish marine environments. *Appl Environ Microbiol* **64**, 7–13.
- Holt, H.M., Sogaard, P., & Gahrn-Hansen, B. (1997). Ear infections with *Shewanella* alga: A bacteriologic, clinical and epidemiologic study of 67 cases. *Clin Microbiol Infect* **3**, 329–334.
- House, J.K., Smith, B.P., & Kamiya, D. (2001). Serological distinction of bovine *Salmonella* carriers from vaccinated and acutely infected cows. *J Vet Diagn Invest* **13**, 483–488.
- Howard, R.J., & Bennett, N.T. (1993). Infections caused by halophilic marine *Vibrio* bacteria. *Ann Surg* **217**, 525–531.
- Howard, R.J., & Burgess, G.H. (1993). Surgical hazards posed by marine and fresh-water animals in Florida. *Am J Surg* **166**, 563–567.
- Hughes, K.A. (2003). Influence of seasonal environmental variables on the distribution of presumptive fecal coliforms around an Antarctic research station. *Appl Environ Microbiol* **69**, 4884–4891.
- Huss, H.H. (1980). Distribution of *Clostridium botulinum*. *Appl Environ Microbiol* **39**, 764–769.
- Ibekwe, A.M., Watt, P.M., Grieve, C.M., Sharma, V.K., & Lyons, S.R. (2002). Multiplex fluorogenic real-time PCR for detection and quantification of *Escherichia coli* O157:H7 in dairy wastewater wetlands. *Appl Environ Microbiol* **68**, 4853–4862.
- Itoh, H., Kuwata, G., Tateyama, S., Yamashita, K., Inoue, T., Kataoka, H., Ido, A., Ogata, K., Takasaki, M., Inoue, S., et al. (1999). *Aeromonas sobria* infection with severe soft tissue damage and segmental necrotizing gastroenteritis in a patient with alcoholic liver cirrhosis. *Pathology International* **49**, 541–546.

- Ivanova, N., Sorokin, A., Anderson, I., Galleron, N., Candelon, B., Kapatral, V., Bhattacharyya, A., Reznik, G., Mikhailova, N., Lapidus, A., et al. (2003). Genome sequence of *Bacillus cereus* and comparative analysis with *Bacillus anthracis*. *Nature* **423**, 87–91.
- Iwata, M., Tateda, K., Matsumoto, T., Furuya, N., Mizuiri, S., & Yamaguchi, K. (1999). Primary *Shewanella* alga septicemia in a patient on hemodialysis. *J Clin Microbiol* **37**, 2104–2105.
- Jackson, G.J., Merker, R.I., & Bandler, R.E. (2000). FDA's bacteriological analytical manual. <http://www.cfsan.fda.gov/~ebam/bam-toc.html>
- Jackson, J.K., Murphree, R.L., & Tamplin, M.L. (1997). Evidence that mortality from *Vibrio vulnificus* infection results from single strains among heterogeneous populations in shellfish. *J Clin Microbiol* **35**, 2098–2101.
- Jensen, S., Samuelsen, O.B., Andersen, K., Torkildsen, L., Lambert, C., Choquet, G., Paillard, C., & Bergh, O. (2003). Characterization of strains of *Vibrio splendidus* and V-tapetis isolated from corksing wrasse *Symphodus melops* suffering vibriosis. *Diseas. Aquat. Organisms* **53**, 25–31.
- Jiang, S.C., & Fu, W. (2001). Seasonal abundance and distribution of *Vibrio cholerae* in coastal waters quantified by a 16S-23S intergenic spacer probe. *Microb Ecol* **42**, 540–548.
- Johnson, R.W., & Arnett, F.C. (2001). A fatal case of *Vibrio vulnificus* presenting as septic arthritis. *Arch Intern Med* **161**, 2616–2618.
- Johnson, M.A., & Lepennec, M. (1995). Association between the mollusk bivalve *Loripes lucinalis* and a chlamydia-like organism, with comments on its pathogenic impact, life-cycle and possible mode of transmission. *Mar Biol* **123**, 523–530.
- Jones, B.L., & Wilcox, M.H. (1995). *Aeromonas* infections and their treatment. *J Antimicrob Chemother* **35**, 453–461.
- Kaneko, T., & Colwell, R.R. (1978). Annual Cycle of *Vibrio parahaemolyticus* in Chesapeake Bay. *Microb Ecol* **4**, 135–155.
- Kaper, J.B., Lockman, H., Colwell, R.R., & Joseph, S.W. (1981). *Aeromonashydrophila*—ecology and toxigenicity of isolates from an estuary. *J Appl Bacteriol* **50**, 359–377.
- Karaolis, D.K.R., Johnson, J.A., Bailey, C.C., Boedeker, E.C., Kaper, J.B., & Reeves, P.R. (1998). A *Vibrio cholerae* pathogenicity island associated with epidemic and pandemic strains. *Proc Natl Acad Sci USA* **95**, 3134–3139.
- Kent, M.L., Traxler, G.S., Kieser, D., Richard, J., Dawe, S.C., Shaw, R.W., Prosperi-Porta, G., Ketcheson, J., & Evelyn, T.P.T. (1998). Survey of salmonid pathogens in ocean-caught fishes in British Columbia, Canada. *Journal of Aquatic Animal Health* **10**, 211–219.
- Kirchhoff, H., Binder, A., Liess, B., Friedhoff, K.T., Pohlenz, J., Stede, M., & Willhaus, T. (1989). Isolation of mycoplasmas from diseased seals. *Vet Rec* **124**, 513–514.
- Koike, N., Okada, K., Yabushita, Y., Zhang, D., Yamamoto, K., Miwatani, T., & Honda, T. (1997). Rapid and differential detection of two analogous enterotoxins of *Vibrio cholerae* and enterotoxigenic *Escherichia coli* by a modified enzyme-linked immunosorbent assay. *FEMS Immunol Med Microbiol* **17**, 21–25.
- Kong, R.Y., Lee, S.K., Law, T.W., Law, S.H., & Wu, R.S. (2002). Rapid detection of six types of bacterial pathogens in marine waters by multiplex PCR. *Water Res* **36**, 2802–2812.
- Kueh, C.S.W., Kutarski, P., & Brunton, M. (1992). Contaminated marine wounds—the risk of acquiring acute bacterial-infection from marine recreational beaches. *J Appl Bacteriol* **73**, 412–420.
- Kusuda, R., & Kawai, K. (1998). Bacterial diseases of cultured marine fish in Japan. *Fish Pathol* **33**, 221–227.
- La Rosa, T., Minto, S., Marino, A., Alonzo, V., Maugeri, T.L., & Mazzola, A. (2001). Heterotrophic bacteria community and pollution indicators of mussel—farm impact in the Gulf of Gaeta (Tyrrhenian Sea). *Mar Environ Res* **52**, 301–321.
- Landers, P., Kerr, K.G., Rowbotham, T.J., Tipper, J.L., Keig, P.M., Ingham, E., & Denton, M. (2000). Survival and growth of *Burkholderia cepacia* within the free-living amoeba *Acanthamoeba polyphaga*. *Eur J Clin Microbiol Infect Dis* **19**, 121–123.
- Lee, J.H., Lee, K.H., & Choi, S.H. (2001). Enumeration of *Vibrio vulnificus* in natural samples by colony blot hybridization. *J Microbiol Biotechnol* **11**, 302–309.
- Lee, K.K., Liu, P.C., & Chuang, W.H. (2002). Pathogenesis of gastroenteritis caused by *Vibrio carchariae* in cultured marine fish. *Mar Biotechnol* **4**, 267–277.
- Lehane, L., & Rawlin, G.T. (2000). Typically acquired bacterial zoonoses from fish: A review. *Med J Aust* **173**, 256–259.
- Leong, J., Mirkazemi, M., & Kimble, F. (2000). *Shewanella putrefaciens* hand infection. *Aust NZ J Surg* **70**, 816–817.
- Levett, P.N. (2001). Leptospirosis. *Clin Microbiol Rev* **14**, 296–326.
- Lipp, E.K., Huq, A., & Colwell, R.R. (2002). Effects of global climate on infectious disease: The cholera model. *Clin Microbiol Rev* **15**, 757–770.

- Lipp, E.K., Rivera, I.N., Gil, A.I., Espeland, E.M., Choopun, N., Louis, V.R. et al. (2003). Direct detection of *Vibrio cholerae* and *ctxA* in Peruvian coastal water and plankton by PCR. *Appl Environ Microbiol* **69**, 3676–3680.
- Lipp, E.K., & Rose, J.B. (1997). The role of seafood in foodborne diseases in the United States of America. *Rev Sci Technol* **16**, 620–640.
- Lobitz, B., Beck, L., Huq, A., Wood, B., Fuchs, G., Faruque, A.S.G., & Colwell, R. (2000). Climate and infectious disease: Use of remote sensing for detection of *Vibrio cholerae* by indirect measurement. *Proc Natl Acad Sci USA* **97**, 1438–1443.
- Loge, F.J., Emerick, R.D., Thompson, D.E., Nelson, D.C., & Darby, J.L. (1999). Development of a fluorescent 16S rRNA oligonucleotide probe specific to the family Enterobacteriaceae. *Water Environ Res* **71**, 75–83.
- Lopez-Romalde, S., Magarinos, B., Ravelo, C., Toranzo, A.E., & Romalde, J.L. (2003). Existence of two O-serotypes in the fish pathogen *Pseudomonas anguilliseptica*. *Vet Microbiol* **94**, 325–333.
- Lopez-Sabater, E.I., Rodriguez-Jerez, J.J., Hernandez-Herrero, M., & Mora-Ventura, M.T. (1996). Incidence of histamine-forming bacteria and histamine content in scombroid fish species from retail markets in the Barcelona area. *Int J Food Microbiol* **28**, 411–418.
- Losonsky, G. (1991). Infections associated with swimming and diving. *Undersea Biomed Res* **18**, 181–185.
- Lotz, M.J., Tamplin, M.L., & Rodrick, G.E. (1983). Thiosulfate-citrate-bile salts-sucrose agar and its selectivity for clinical and marine vibrio organisms. *Ann Clin Lab Sci* **13**, 45–48.
- Louis, V.R., Russek-Cohen, E., Choopun, N., Rivera, I.N., Gangle, B., Jiang, S.C., Rubin, A., Patz, J.A., Huq, A., & Colwell, R.R. (2003). Predictability of *Vibrio cholerae* in Chesapeake Bay. *Appl Environ Microbiol* **69**, 2773–2785.
- Lyon, W.J. (2001). TaqMan PCR for detection of *Vibrio cholerae* O1, O139, non-O1, and non-O139 in pure cultures, raw oysters, and synthetic seawater. *Appl Environ Microbiol* **67**, 4685–4693.
- MacFarlane, R.D., McLaughlin, J.J., & Bullock, G.L. (1986). Quantitative and qualitative studies of gut flora in striped bass from estuarine and coastal marine environments. *J Wildl Dis* **22**, 344–348.
- Maiden, M.C., Bygraves, J.A., Feil, E., Morelli, G., Russell, J.E., Urwin, R., Zhang, Q., Zhou, J., Zurth, K., Caugant, D.A., et al. (1998). Multilocus sequence typing: a portable approach to the identification of clones within populations of pathogenic microorganisms. *Proc Natl Acad Sci USA* **95**, 3140–3145.
- Malorny, B., Hoorfar, J., Bunge, C., & Helmuth, R. (2003). Multicenter validation of the analytical accuracy of *Salmonella* PCR: Towards an international standard. *Appl Environ Microbiol* **69**, 290–296.
- Marolda, C.L., Hauröder, B., John, M.A., Michel, R., & Valvano, M.A. (1999). Intracellular survival and saprophytic growth of isolates from the *Burkholderia cepacia* complex in free-living amoebae. *Microbiology* **145** (7), 1509–1517.
- McDade, J.E., Shepard, C.C., Fraser, D.W., Tsai, T.R., Redus, M.A., & Dowdle, W.R. (1977). Legionnaires' disease: Isolation of a bacterium and demonstration of its role in other respiratory disease. *N Engl J Med* **297**, 1197–1203.
- Michaud, S., Levasseur, M., Doucette, G., & Cantin, G. (2002). Particle size fractionation of paralytic shellfish toxins (PSTs): Seasonal distribution and bacterial production in the St Lawrence estuary, Canada. *Toxicon* **40**, 1451–1462.
- Michel, R., & Hauröder, B. (1997). Isolation of an Acanthamoeba strain with intracellular *Burkholderia pickettii* infection. *Zentralbl Bakteriol* **285**, 541–557.
- Miliotis, M., Watkins, W., Wekell, M., DePaola, A., Cook, D., Bowers, J., Ross, M., DiNovi, M., & Burr, D. (2000). Draft risk assessment on the public health impact of *Vibrio parahaemolyticus* in raw molluscan shellfish. Rockville, MD: *Food and Drug Administration*, p. 102.
- Mitov, I., Haralambieva, I., Petrov, D., Ivanova, R., Kamarinchev, B., & Iankov, I. (2003). Cross-reactive monoclonal antibodies raised against the lipopolysaccharide antigen of *Salmonella minnesota* Re chemotype: Diagnostic relevance. *Diagn Microbiol Infect Dis* **45**, 225–231.
- Montali, R.J., Mikota, S.K., & Cheng, L.I. (2001). *Mycobacterium tuberculosis* in zoo and wildlife species. *Rev Sci Technol Off Int Epizoot* **20**, 291–303.
- Morgan, D.R., Johnson, P.C., DuPont, H.L., Satterwhite, T.K., & Wood, L.V. (1985). Lack of correlation between known virulence properties of *Aeromonas hydrophila* and enteropathogenicity for humans. *Infect Immun* **50**, 62–65.
- Moter, A., & Gobel, U.B. (2000). Fluorescence in situ hybridization (FISH) for direct visualization of microorganisms. *J Microbiol Methods* **41**, 85–112.
- Motes, M.L., DePaola, A., Cook, D.W., Veazey, J.E., Hunsucker, J.C., Garthright, W.E., Blodgett, R.J., & Chirtel, S.J. (1998). Influence of water temperature and salinity on *Vibrio vulnificus* in Northern Gulf and Atlantic Coast oysters (*Crassostrea virginica*). *Appl Environ Microbiol* **64**, 1459–1465.

- Myers, C.R., & Nealson, K.H. (1990). Respiration-linked proton translocation coupled to anaerobic reduction of Manganese(IV) and Iron(III) in *Shewanella. Putrefaciens* Mr-1. *J Bacteriol* **172**, 6232–6238.
- Naylor, R.L., Goldburg, R.J., Primavera, J.H., Kautsky, N., Beveridge, M.C., Clay, J., Folke, C., Lubchenco, J., Mooney, H., & Troell, M. (2000). Effect of aquaculture on world fish supplies. *Nature* **405**, 1017–1024.
- Nicolas, J.L., Basuyaux, O., Mazurie, J., & Thebault, A. (2002). *Vibrio carchariae*, a pathogen of the abalone *Haliotis tuberculata*. *Dis Aquat Organ* **50**, 35–43.
- Nordstrom, J.L., & DePaola, A. (2003). Improved recovery of pathogenic *Vibrio parahaemolyticus* from oysters using colony hybridization following enrichment. *J Microbiol Methods* **52**, 273–277.
- Nozue, H., Hayashi, T., Hashimoto, Y., Ezaki, T., Hamasaki, K., Ohwada, K., & Terawaki, Y. (1992). Isolation and characterization of *Shewanella*-alga from human clinical specimens and emendation of the description of *S*-alga Simidu Et-Al, 1990, 335. *Int J Syst Bacteriol* **42**, 628–634.
- Olafsen, J.A., Mikkelsen, H.V., Glaever, H.M., & Hansen, G.H. (1993). Indigenous bacteria in hemolymph and tissues of marine bivalves at low temperatures. *Appl Environ Microbiol* **59**, 1848–1854.
- Ortizroque, C.M., & Hazen, T.C. (1987). Abundance and distribution of Legionellaceae in Puerto-Rican waters. *Appl Environ Microbiol* **53**, 2231–2236.
- Oxley, A.P.A., Shipton, W., Owens, L., & McKay, D. (2002). Bacterial flora from the gut of the wild and cultured banana prawn, *Penaeus merguensis*. *J Appl Microbiol* **93**, 214–223.
- Pagani, L., Lang, A., Vedovelli, C., Moling, O., Rimenti, G., Pristera, R., & Mian, P. (2003). Soft tissue infection and bacteremia caused by *Shewanella putrefaciens*. *J Clin Microbiol* **41**, 2240–2241.
- Palumbo, J.D., Borucki, M.K., Mandrell, R.E., & Gorski, L. (2003). Serotyping of *Listeria monocytogenes* by enzyme-linked immunosorbent assay and identification of mixed-serotype cultures by colony immunoblotting. *J Clin Microbiol* **41**, 564–571.
- Parker, B.C., Ford, M.A., Gruft, H., & Falkinham, J.O., 3rd (1983). Epidemiology of infection by nontuberculous mycobacteria. IV. Preferential aerosolization of *Mycobacterium intracellulare* from natural waters. *Am Rev Respir Dis* **128**, 652–656.
- Pascual, M., Rodo, X., Ellner, S.P., Colwell, R., & Bouma, M.J. (2000). Cholera dynamics and El Nino—southern oscillation. *Science* **289**, 1766–1769.
- Pavia, A.T., Bryan, J.A., Maher, K.L., Hester, T.R., Jr., & Farmer, J.J., 3rd (1989). *Vibrio carchariae* infection after a shark bite. *Ann Intern Med* **111**, 85–86.
- Pernthaler, A., Pernthaler, J., & Amann, R. (2002a). Fluorescence in situ hybridization and catalyzed reporter deposition for the identification of marine bacteria. *Appl Environ Microbiol* **68**, 3094–3101.
- Pernthaler, A., Preston, C.M., Pernthaler, J., DeLong, E.F., & Amann, R. (2002b). Comparison of fluorescently labeled oligonucleotide and polynucleotide probes for the detection of pelagic marine bacteria and archaea. *Appl Environ Microbiol* **68**, 661–667.
- Persing, D.H., Tenover, F.C., Versalovic, J., Tang, Y.-W., Unger, E.R., Relman, D.A., & White, T.J.E. (2003). *Molecular Microbiology: Diagnostic Principles and Practice*. American Society for Microbiology Press, Washington, DC.
- PHAC (2001). Material Safety Data Sheets: Infectious substances. Office of Laboratory Security, Public Health Agency of Canada (PHAC). Population and Public Health Branch: Canadian Office of Laboratory Security.
- Polo, F., Figueras, M.J., Inza, I., Sala, J., Fleisher, J.M., & Guarro, J. (1999). Prevalence of *Salmonella* serotypes in environmental waters and their relationships with indicator organisms. *Antonie Van Leeuwenhoek* **75**, 285–292.
- Polz, M.F., & Cavanaugh, C.M. (1997). A simple method for quantification of uncultured microorganisms in the environment based on in vitro transcription of 16S rRNA. *Appl Environ Microbiol* **63**, 1028–1033.
- Polz, M.F., & Cavanaugh, C.M. (1997). Bias in template-to-product ratios in multitemplate PCR. *Appl Environ Microbiol* **64**, 3724–3730.
- Pommepuy, M., & Le Guyader, F. (1998). Molecular approaches to measuring microbial marine pollution. *Curr Opin Biotechnol* **9**, 292–299.
- Prescott, J.F. (1991). *Rhodococcus equi*: An animal and human pathogen. *Clin Microbiol Rev* **4**, 20–34.
- Rahman, I., Shahamat, M., Chowdhury, M.A., & Colwell, R.R. (1996). Potential virulence of viable but nonculturable *Shigella dysenteriae* type 1. *Appl Environ Microbiol* **62**, 115–120.
- Raidal, S.R., Ohara, M., Hobbs, R.P., & Prince, R.I. (1998). Gram-negative bacterial infections and cardiovascular parasitism in green sea turtles (*Chelonia mydas*). *Aust Vet J* **76**, 415–417.
- Ravelo, C., Magarinos, B., Lopez-Romalde, S., Toranzo, A.E., & Romalde, J.L. (2003). Molecular fingerprinting of fish-pathogenic *Lactococcus garvieae* strains by random amplified polymorphic DNA analysis. *J Clin Microbiol* **41**, 751–756.

- Rhodes, M.W., Kator, H., Kotob, S., van Berkum, P., Kaattari, I., Vogelbein, W., Floyd, M.M., Butler, W.R., Quinn, F.D., Ottinger, C., et al. (2001). A unique *Mycobacterium* species isolated from an epizootic of striped bass (*Morone saxatilis*). *Emerging Infectious Diseases* **7**, 896–899.
- Ribot, E.M., Fitzgerald, C., Kubota, K., Swaminathan, B., & Barrett, T.J. (2001). Rapid pulsed-field gel electrophoresis protocol for subtyping of *Campylobacter jejuni*. *J Clin Microbiol* **39**, 1889–1894.
- Rigsbee, W., Simpson, L.M., & Oliver, J.D. (1997). Detection of the viable but nonculturable state in *Escherichia coli* O157:H7. *J Food Safety* **16**, 255–262.
- Ritter, M.S., Mroch, H., & Burns, M.J. (1993). Soaring suppurative sea shells from the sea shore—*Pseudomonas aeruginosa* and *Klebsiella pneumoniae* septic arthritis after a marine sea shell injury. *Pediatr Emerg Care* **9**, 289–291.
- Rivera, I.N., Chun, J., Huq, A., Sack, R.B., & Colwell, R.R. (2001). Genotypes associated with virulence in environmental isolates of *Vibrio cholerae*. *Appl Environ Microbiol* **67**, 2421–2429.
- Rodgers, C.J., & Furones, M.D. (1998). Disease problems in cultured marine fish in the Mediterranean. *Fish Pathol* **33**, 157–164.
- Rompere, A., Servais, P., Baudart, J., de-Roubin, M.R., & Laurent, P. (2002). Detection and enumeration of coliforms in drinking water: current methods and emerging approaches. *J Microbiol Methods* **49**, 31–54.
- Rose, J.B., Epstein, P.R., Lipp, E.K., Sherman, B.H., Bernard, S.M., & Patz, J.A. (2001). Climate variability and change in the United States: Potential impacts on water- and foodborne diseases caused by microbiologic agents. *Environ Health Perspect* **109**, 211–221.
- Rose, N.R., Hamilton, R.G., & Detrick, B. (2002). *Manual of Clinical Laboratory Immunology*. American Society for Microbiology Press, Washington, DC.
- Rozen, Y., & Belkin, S. (2001). Survival of enteric bacteria in seawater. *FEMS Microbiol Rev* **25**, 513–529.
- Ruby, E.G., & Neelson, K.H. (1978). Seasonal-changes in species composition of luminous bacteria in nearshore seawater. *Limnol Oceanogr* **23**, 530–533.
- Ruiz, G.M., Rawlings, T.K., Dobbs, F.C., Drake, L.A., Mullady, T., Huq, A., & Colwell, R.R. (2000). Global spread of microorganisms by ships—ballast water discharged from vessels harbours a cocktail of potential pathogens. *Nature* **408**, 49–50.
- Salas, S.D., & Geesey, G.G. (1983). Surface attachment of a sediment isolate of *Enterobacter cloacae*. *Microbial Ecology* **9**, 307–315.
- Sambrook, J., & Russell, D.W. (2001). *Molecular Cloning: A Laboratory Manual*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- Saubolle, M.A., Kiehn, T.E., White, M.H., Rudinsky, M.F., & Armstrong, D. (1996). *Mycobacterium haemophilum*: Microbiology and expanding clinical and geographic spectra of disease in humans. *Clin Microbiol Rev* **9**, 435–447.
- Schlöter, M., Assmus, B., & Hartmann, A. (1995). The use of immunological methods to detect and identify bacteria in the environment. *Biotechnol Adv* **13**, 75–90.
- Sinton, L.W., Hall, C.H., Lynch, P.A., & Davies-Colley, R.J. (2002). Sunlight inactivation of fecal indicator bacteria and bacteriophages from waste stabilization pond effluent in fresh and saline waters. *Appl Environ Microbiol* **68**, 1122–1131.
- Slaven, E.M., Lopez, F.A., Hart, S.M., & Sanders, C.V. (2001). Myonecrosis caused by *Edwardsiella tarda*: A case report and case series of extraintestinal *E. tarda* infections. *Clin Infect Dis* **32**, 1430–1433.
- Smith, E.A., Mackintosh, F.H., Grant, F., & Gallacher, S. (2002). Sodium channel blocking (SCB) activity and transformation of paralytic shellfish toxins (PST) by dinoflagellate-associated bacteria. *Aquat Microb Ecol* **29**, 1–9.
- Smith, S., Taylor, G.D., & Fanning, E.A. (2003). Chronic cutaneous *Mycobacterium haemophilum* infection acquired from coral injury. *Clin Infect Dis* **37**, e100–101.
- Sohn, A.H., Probert, W.S., Glaser, C.A., Gupta, N., Bollen, A.W., Wong, J.D., Grace, E.M., & McDonald, W.C. (2003). Human neurobrucellosis with intracerebral granuloma caused by a marine mammal *Brucella* spp. *Emerging Infectious Diseases* **9**, 485–488.
- Somer, L., & Kashi, Y. (2003). A PCR method based on 16S rRNA sequence for simultaneous detection of the genus *Listeria* and the species *Listeria monocytogenes* in food products. *J Food Prot* **66**, 1658–1665.
- Songer, J.G. (1981). Methods for selective isolation of mycobacteria from the environment. *Can J Microbiol* **27**, 1–7.
- Stadtlander, C.T., & Madoff, S. (1994). Characterization of cytopathogenicity of aquarium seal mycoplasmas and seal finger mycoplasmas by light and scanning electron microscopy. *Zentralbl Bakteriol* **280**, 458–467.
- Starliper, C.E. (2001). Isolation of *Serratia liquefaciens* as a pathogen of Arctic char, *Salvelinus alpinus* (L.). *J Fish Dis* **24**, 53–56.

- Steinert, M., Birkness, K., White, E., Fields, B., & Quinn, F. (1998). *Mycobacterium avium* bacilli grow saprozoically in coculture with *Acanthamoeba polyphaga* and survive within cyst walls. *Appl Environ Microbiol* **64**, 2256–2261.
- Steinert, M., Emody, L., Amann, R., Hacker, J. (1997). Resuscitation of viable but nonculturable *Legionella pneumophila* Philadelphia JR32 by *Acanthamoeba castellanii*. *Appl Environ Microbiol* **63**, 2047–2053.
- Straub, T.M., & Chandler, D.P. (2003). Towards a unified system for detecting waterborne pathogens. *J Microbiol Methods* **53**, 185–197.
- Swaminathan, B., Barrett, T.J., Hunter, S.B., & Tauxe, R.V. (2001). PulseNet: The molecular subtyping network for foodborne bacterial disease surveillance, United States. *Emerg Infect Dis* **7**, 382–389.
- Swanson, M.S., & Hammer, B.K. (2000). *Legionella pneumophila* pathogenesis: A fateful journey from amoebae to macrophages. *Annu Rev Microbiol* **54**, 567–613.
- Tamplin, M.L. (2001). Coastal vibrios: Identifying relationships between environmental condition and human disease. *Hum Ecol Risk Assess* **7**, 1437–1445.
- Tamplin, M.L., Jackson, J.K., Buchrieser, C., Murphree, R.L., Portier, K.M., Gangar, V., Miller, L.G., & Kaspar, C.W. (1996). Pulsed-field gel electrophoresis and ribotype profiles of clinical and environmental *Vibrio vulnificus* isolates. *Applied and Environmental Microbiology* **62**, 3572–3580.
- Taroncher-Oldenburg, G., Griner, E.M., Francis, C.A., & Ward, B.B. (2003). Oligonucleotide microarray for the study of functional gene diversity in the nitrogen cycle in the environment. *Appl Environ Microbiol* **69**, 1159–1171.
- Thomas, C., & Scott, S. (1997). *All Stings Considered: First Aid and Medical Treatment of Hawaii's Marine Injuries*. University of Hawaii Press, Honolulu.
- Thompson, P.J., Cousins, D.V., Gow, B.L., Collins, D.M., Williamson, B.H., & Dagnia, H.T. (1993). Seals, Seal trainers, and Mycobacterial infection. *Am Rev Respir Dis* **147**, 164–167.
- Thompson, J.R., Randa, M.A., Marcelino, L.A., Tomita-Mitchell, A., Lim, E., and Polz, M.F. (2004). Diversity and dynamics of a north atlantic coastal *Vibrio* community. *Appl Environ Microbiol* **70**, 4103–4110.
- Thompson, J.R., Pacocha, S., Pharino, C., Klepac-Ceraj, V., Hunt, D.E., Benoit, J., Sarma-Rupavtarm, R., Distel, D.L., & Polz, M.F. (2005). Genotypic diversity within a natural coastal bacterioplankton population. *Science* **307**, 1311–1313.
- Toro, A., Gonzalez, N., Torres, J., Dvorsky, E., & Toranzos, G.A. (1995). Modified culture methods for the detection of *Vibrio* spp from Estuarine waters. *Water Sci Technol* **31**, 283–290.
- Tranvik, L.J., & Bertilsson, S. (2001). Contrasting effects of solar UV radiation on dissolved organic sources for bacterial growth. *Ecol Lett* **4**, 458–463.
- Tryland, M. (2000). Zoonoses of arctic marine mammals. *Infect Dis Rev* **2**, 55–64.
- Tryland, M., Kleivane, L., Alfredsson, A., Kjeld, M., Arnason, A., Stuen, S., & Godfroid, J. (1999). Evidence of *Brucella* infection in marine mammals in the North Atlantic Ocean. *Vet Rec* **144**, 588–592.
- Turner S.J., Lewis G.D., & Bellamy, A.R. (1997). Detection of sewage-derived *Escherichia coli* in a rural stream using multiplex PCR and automated DNA detection. *Water Sci Technol* **35**, 337–342.
- USEPA (1988). Water quality standards criteria summaries: A compilation of state/federal criteria. U.S. Environmental Protection Agency, Washington, DC.
- Uyttendaele, M., Schukkink, R., Van Gemen, B., & Debevere, J. (1995). Detection of *Campylobacter jejuni* added to foods by using a combined selective enrichment and nucleic acid sequence-based amplification (NASBA). *Appl Environ Microbiol* **61**, 1341–1347.
- Uyttendaele M, Schukkink R, Van Gemen B, & Debevere, J. (1996). Comparison of the nucleic acid amplification system NASBA(R) and agar isolation for detection of pathogenic campylobacters in naturally contaminated poultry. *J Food protect* **59**, 683–687.
- Van Belkum, A. (2003). High-throughput epidemiologic typing in clinical microbiology. *Clin Microbiol Infect* **9**, 86–100.
- Van Dolah, F.M. (2000). Marine algal toxins: Origins, health effects, and their increased occurrence. *Environ Health Perspect* **108** (Suppl 1), 133–141.
- Van Doorn, L.J., Verschuuren-Van Haperen, A., Van Belkum, A., Endtz, H.P., Vliegthart, J.S., Vandamme, P., & Quint, W.G. (1998). Rapid identification of diverse *Campylobacter lari* strains isolated from mussels and oysters using a reverse hybridization line probe assay. *J Appl Microbiol* **84**, 545–550.
- Vasquez, M., Gruttner, C., Gallacher, S., & Moore, E.R.B. (2001). Detection and characterization of toxigenic bacteria associated with *Alexandrium catenella* and *Aulacomya ater* contaminated with PSP. *J Shellfish Res* **20**, 1245–1249.
- Victor, T.C., Jordaan, A.M., van Rie, A., van der Spuy, G.D., Richardson, M., van Helden, P.D., & Warren, R. (1999). Detection of mutations in drug resistance genes of *Mycobacterium tuberculosis* by a dot-blot hybridization strategy. *Tuber Lung Dis* **79**, 343–348.

- Villari, P., Pucino, A., Santagata, N., & Torre, I. (1999). A comparison of different culture media for the membrane filter quantification of *Aeromonas* in water. *Lett Appl Microbiol* **29**, 253–257.
- Vishnubhatla, A., Oberst, R.D., Fung, D.Y., Wonglumsom, W., Hays, M.P., & Nagaraja, T.G. (2001). Evaluation of a 5'-nuclease (TaqMan) assay for the detection of virulent strains of *Yersinia enterocolitica* in raw meat and tofu samples. *J Food Prot* **64**, 355–360.
- Vogel, B.F., Holt, H.M., Gerner-Smidt, P., Bundvad, A., Sogaard, P., & Gram, L. (2000). Homogeneity of Danish environmental and clinical isolates of *Shewanella* algae. *Appl Environ Microbiol* **66**, 443–448.
- von Graevenitz, A., Bowman, J., Del Notaro, C., & Ritzler, M. (2000). Human infection with *Halomonas venusta* following fish bite. *J Clin Microbiol* **38**, 3123–3124.
- von Wintzingerode, F., Gobel, U.B., & Stackebrandt, E. (1997). Determination of microbial diversity in environmental samples: Pitfalls of PCR-based rRNA analysis. *FEMS Microbiol Rev* **21**, 213–229.
- Waage, A.S., Vardund, T., Lund, V., & Kapperud, G. (1999a). Detection of low numbers of pathogenic *Yersinia enterocolitica* in environmental water and sewage samples by nested polymerase chain reaction. *J Appl Microbiol* **87**, 814–821.
- Waage, A.S., Vardund, T., Lund, V., & Kapperud, G. (1999b). Detection of low numbers of *Salmonella* in environmental water, sewage and food samples by a nested polymerase chain reaction assay. *J Appl Microbiol* **87**, 418–428.
- Waage, A.S., Vardund, T., Lund, V., & Kapperud, G. (1999c). Detection of small numbers of *Campylobacter jejuni* and *Campylobacter coli* cells in environmental water, sewage, and food samples by a seminested PCR assay. *Appl Environ Microbiol* **65**, 1636–1643.
- Watkins, W.D. & Cabelli, V.J. (1985). Effect of fecal pollution on *Vibrio parahaemolyticus* densities in an estuarine environment. *Appl Environ Microbiol* **49**, 1307–1313.
- Weber, J.T., Hibbs, R.G., Jr., Darwish, A., Mishu, B., Corwin, A.L., Rakha, M., Hatheway, C.L., el Sharkawy, S., el-Rahim, S.A., al-Hamd, M. F., et al. (1993). A massive outbreak of type E botulism associated with traditional salted fish in Cairo. *J Infect Dis* **167**, 451–454.
- Weinstein, M., Low, D., McGeer, A., Willey, B., Rose, D., Coulter, M., Wyper, P., Borczyk, A., Lovgren, M., & Facklam, R. (1996). Invasive infection due to *Streptococcus iniae*: a new or previously unrecognized disease—Ontario, 1995–1996. *Can Commun Dis Rep* **22**, 129–31; discussion 131–132.
- Weinstein, M.R., Litt, M., Kertesz, D.A., Wyper, P., Rose, D., Coulter, M., McGeer, A., Facklam, R., Ostach, C., Willey, B.M., et al. (1997). Invasive infections due to a fish pathogen, *Streptococcus iniae*. S. iniae Study Group. *N Engl J Med* **337**, 589–594.
- Weinstock, D.M., & Brown, A.E. (2002). *Rhodococcus equi*: An emerging pathogen. *Clin Infect Dis* **34**, 1379–1385.
- Welch, R.A., Burland, V., Plunkett III, G., Redford, P., Roesch, P., Rasko, D., Buckles, E.L., Liou, S.-R., Boutin, A., Hackett, J., et al. (2002). Extensive mosaic structure revealed by the complete genome sequence of uropathogenic *Escherichia coli*. *Proc Natl Acad Sci USA* **99**, 17020–17024.
- Wendt, S.L., George, K.L., Parker, B.C., Gruft, H., & Falkinham, J.O., 3rd (1980). Epidemiology of infection by nontuberculous Mycobacteria. III. Isolation of potentially pathogenic mycobacteria from aerosols. *Am Rev Respir Dis* **122**, 259–263.
- Wenger, J.D., Hollis, D.G., Weaver, R.E., Baker, C.N., Brown, G.R., Brenner, D.J., & Broome, C.V. (1989). Infection caused by *Francisella philomiragia* (formerly *Yersinia philomiragia*). A newly recognized human pathogen. *Ann Intern Med* **110**, 888–892.
- Wilkerson, M., McAllister, S., Miller, J.M., Heiter, B.J., & Bourbeau, P.P. (1997). Comparison of five agglutination tests for identification of *Staphylococcus aureus*. *J Clin Microbiol* **35**, 148–151.
- Wright, A.C., Hill, R.T., Johnson, J.A., Roghman, M.C., Colwell, R.R., & Morris, J.G., Jr. (1996). Distribution of *Vibrio vulnificus* in the Chesapeake Bay. *Appl Environ Microbiol* **62**, 717–724.
- Wu, L., Thompson, D.K., Li, G., Hurt, R.A., Tiedje, J.M., & Zhou, J. (2001). Development and evaluation of functional gene arrays for detection of selected genes in the environment. *Appl Environ Microbiol* **67**, 5780–5790.
- Yang, C., Jiang, Y., Huang, K., Zhu, C., & Yin, Y. (2003). Application of real-time PCR for quantitative detection of *Campylobacter jejuni* in poultry, milk and environmental water. *FEMS Immunol Med Microbiol* **38**, 265–271.
- Ye, R.W., Wang, T., Bedzyk, L., Croker, K.M. (2001). Applications of DNA microarrays in microbial systems. *J Microbiol Methods* **47**, 257–272.
- Zlotkin, A., Hershko, H., & Eldar, A. (1998). Possible transmission of *Streptococcus iniae* from wild fish to cultured marine fish. *Appl Environ Microbiol* **64**, 4065–4067.
- Zo, Y.-G., Rivera, I.N.G., Russek-Cohen, E., Islam, M.S., Siddique, A.K., Yunus, M., Sack, R.B., Huq, A., & Colwell, R.R. (2002). Genomic profiles of clinical and environmental isolates of *Vibrio cholerae* O1 in cholera-endemic areas of Bangladesh. *Proc. Natl. Acad. Sci. USA* **99**, 12409–12414.

Oceans and Health:

Pathogens in the Marine Environment

Belkin, S.; Colwell, R.R. (Eds.)

2005, XIV, 464 p. 49 illus., Hardcover

ISBN: 978-0-387-23708-4