

CHAPTER 2

Phagocytosis and Immunity

Steven Greenberg

Abstract

Phagocytosis is an phylogenetically conserved mechanism utilized by many cells to ingest microbial pathogens and apoptotic or necrotic corpses. Recent studies have demonstrated that phagocytosis serves to initiate immunity mediated by both Class I and Class II MHC. Depending on the identity of the specific phagocytic receptor involved, phagocytosis can either enhance or suppress inflammation. Dysregulation of phagocytosis can lead to alterations in the immune response and may contribute to autoimmunity. Harnessing the phagocytic capacity of antigen presenting cells may ultimately lead to exploitation of phagocytosis as a therapeutic modality in intractable diseases, such as advanced cancer.

Phagocytosis is the process by which leukocytes and other cells ingest particulate ligands whose size exceeds about 1 μm . This phylogenetically ancient cellular event is critical for both innate and acquired immunity. By ingesting microbial pathogens, phagocytic leukocytes accomplish two essential immune functions. First, they initiate a microbial death pathway. They target ingested pathogens to degradative organelles, such as lysosomes and to vesicles containing components of the phagocyte oxidase complex. Second, phagocytic leukocytes, particularly dendritic cells (DCs), utilize phagocytosis to direct antigens to both MHC I and II compartments. Thus, phagocytosis serves a dual role as an effector of innate immunity and an initiator of acquired immunity.

Diversity of Phagocytic Receptors

Many receptors are capable of mediating phagocytosis (Table 1). These receptors can be broadly defined as those that recognize epitopes on the surfaces of unmodified bacteria and fungi (nonopsonic phagocytosis) or those that recognize components derived from the host (opsonic phagocytosis). Examples of the latter include receptors for the Fc portion of IgG (Fc γ Rs) and receptors that recognize various components of complement (e.g., Complement Receptor 3). It is important to distinguish between those receptors that participate solely in binding of the phagocytic target with receptors that are actually coupled to the transmembrane signaling machinery of phagocytosis. For example, many members of the Scavenger Receptor superfamily are “pattern recognition receptors,” which bind ligands, such as peptidoglycan, present on the surfaces of bacteria and fungi.¹ However, most of these receptors have not been shown to trigger distinct transmembrane signals; rather, they facilitate the recruitment of other receptors that directly engage the phagocytic machinery.

Table 1. Examples of receptors in mammalian cells that promote phagocytosis^a

Receptor	Cell Type	Target	Ligand	Refs.
FcγRI FcγRIIA FcγRIII CR1 (CD35)	Gran, Mo, MΦ, DCs, Mast, Plts PMN, Mo, MF	IgG-coated pathogens Complement-opsonized bacteria and fungi	Fc port ion of IgG C3b, C4b Mannan- binding lectin	many 84
CR3 (CD11b/CD18; αMβ3; Mac1)	PMN, Mo, MΦ, myeloid DCs	Complement-opsonized bacteria and fungi Gram-negative bacteria <i>Bordatella pertussis</i> Yeast <i>M. tuberculosis</i>	C3bi, C3d LPS Filamentous hemagglutinin β-glucan ?	85 86
CR4 (CD11c/ CD18; p150, 95) CD48 Dectin-1	MΦ, DC Mast MΦ	Enterobacteria <i>P. carinii</i> , <i>C. albicans</i>	FimH Mannosyl/fucosyl residues	87 88
Scavenger receptor AI/II Scavenger receptor BI MARCO Mer PSR CD36 CD14	MΦ Sertoli cells, Thymic Epi MΦ MΦ Many MΦ MΦ	Apoptotic lymphocytes Gram-positive cocci Apoptotic cells <i>E. coli</i> , <i>S. aureus</i> Apoptotic thymocytes Apoptotic cells Apoptotic PMN <i>P. aeruginosa</i> Apoptotic cells	?PS Leipoteichoic acid PS ? ?Gas6/PS PS PS/Thrombospondin ?LPS ?	89-91 92,93 94 95 96 97,98 99,100
β ₁ integrins α _v β ₃ α _v β ₅ E-cadherin Met	Many MΦ DC, Epi Epi Epi	<i>Yersinia</i> Apoptotic cells Apoptotic cells <i>Listeria</i> <i>Listeria</i>	Invasin ?Thrombospondin ? InIA InIB	101 97,102 103,104 34 31

^aSpecific inhibition of binding by these receptors correlates with inhibition of phagocytosis. However, with some notable exceptions (e.g., FcγRIIA and the macrophage mannose receptor), it is possible that the indicated receptor serves to enhance ligand binding, rather than to participate directly in the ingestion process; ^bDC = dendritic cells; Epi = epithelial cells; Leuk = leukocytes; Mast = mast cells; Mo = monocytes; MΦ = macrophages; PMN = polymorphonuclear leukocytes; PS = phosphatidylserine

Mechanisms of Phagocytic Signaling

The signaling mechanisms of phagocytosis are well-understood for only a handful of receptors.^{2,3} FcγRs signal by recruiting an array of tyrosine kinases in the vicinity of the ligated receptors. FcγR ligand binding receptors or their subunits contain immunoreceptor tyrosine-based activating motif (ITAMs) in their cytosolic domains. These motifs become phosphorylated by members of the Src family,⁴ which serve as high-affinity binding sites for SH2 domains of Syk, a tyrosine kinase that is expressed predominantly in hematopoietic cells. Although Syk is clearly required for phagocytosis,⁵⁻⁸ the identity of further downstream signals that are critical for phagocytosis are less certain. Phospholipid kinases (phosphatidylinositol 3-kinase; PI 3-kinase⁹⁻¹¹ and phosphatidylinositol-4-phosphate 5-kinase¹²) are clearly involved

as are various serine/threonine protein kinases (e.g., MEK1/2 and/or ERK, in the case of neutrophils¹³) and PKC.¹⁴ PLA₂ and PLD are also activated and believed to participate in the phagocytic process.^{15,16} The former may participate in vesicle trafficking during phagocytosis¹⁷ as well as contribute to the production of leukotrienes that amplify the phagocytic signal.¹⁸ Ultimately, early phagocytic signaling events culminate in net actin assembly and pseudopod extension. Actin assembly during phagocytosis is mediated by one or more Rho family GTPases.¹⁹⁻²² Other GTPases may participate in actin polymerization and/or vesicle trafficking, such as ARF6.^{23,24} In contrast, PI 3-kinase regulates pseudopod extension,⁹ in part by recruiting intracellular pools of latent phagosomal membrane,^{25,26} and in part by recruiting PH domain-containing proteins, such as myosin-X, to the phagosome.²⁷

The occurrence of ITAMs in other phagocytic receptors suggests strongly that the paradigm of phagocytic signaling utilized by FcγRs is likely to be a general one. For example, Dectin-1, a lectin that recognizes mannosyl/fucosyl residues on fungi,²⁸ and CEACAM3, a receptor that recognizes on *Neisseria*, *Moraxella*, and *Haemophilus* species contain functional ITAMs in their cytosolic domains.^{29,30}

Fate of Engulfed Pathogens—Us vs. Them

The initial host response to most bacterial and fungal pathogens is phagocytosis. The particular route of entry is a function of the nature of the pathogen being ingested and identity of the host cell receptors engaging the pathogen. For example, internalization of *Listeria* is mediated by the adhesins InlA, which binds to E-cadherin on host epithelia, and InlB, which binds to the Met tyrosine kinase and to gC1q-R on host cells.^{31,32} E-cadherin-mediated entry requires participation of catenins^{33,34} and Met-dependent signaling induces activation of PI 3-kinase. For *Yersinia*, recognition of invasin on the bacterial surface is mediated by β₁ integrins on a variety of cells; bacterial uptake requires the participation of Src-family tyrosine kinases and focal adhesion kinase.³⁵

Phagocytosis can be a highly localized event (e.g., phagocytosis mediated by “zippers”) or can be partially delocalized by virtue of a diffusible mediator. Several pathogens, such as *Salmonella* or *Shigella*, stimulate a “trigger” mechanism of invasion, inducing the assembly of actin in the host cell in the vicinity of where the bacteria interact with the host.³⁶ Using a Type III secretion system, *Salmonella* injects SopE, a protein with guanine nucleotide exchange factor for Cdc42 and Rac, into host cells.³⁷ In contrast, some pathogens, such as *Haemophilus ducreyi*³⁸ and *Yersinia*, use different strategies to evade phagocytosis. *Yersinia* secretes YopH, a tyrosine phosphatase that dephosphorylates the focal adhesion protein, Cas, as well as other potential tyrosine kinase substrates.³⁹ Another secreted product of *Yersinia*, YopE, is a RhoGAP.⁴⁰

Evolutionary pressure results in genetically stable adaptations by bacteria that serve to compromise the host. Many pathogens evade killing by inducing a delay in phagosome maturation. This can result in exclusion of active lysosomal enzymes or components of the NADPH-oxidase-containing vesicles from the phagosomes. Among the survival strategies employed by *Mycobacterium tuberculosis*, for example, is the suppression of calcium signaling, which contributes to evasion of lysosome fusion by the *Mycobacterium*-containing phagosome.⁴¹ Some of the phagosome maturation arrest activity resides in glycosylated lipids derived from the cell wall of *M. tuberculosis*.⁴²⁻⁴⁴ Other pathogens, such as *Legionella*, avoid maturation at early stages of phagosome biogenesis.⁴⁵ The *Legionella* phagocytic vacuole is highly specialized: it does not fuse with lysosomes, fails to acidify,⁴⁶ and intercepts vesicular traffic from ER exit sites to create an organelle that permits intracellular replication.^{47,48}

Phagocytosis and the MHC Class II Pathway

Once ingested, microbes that reside in phagocytic vacuoles find themselves in an increasingly hostile environment. Phagosomes undergo a maturation process, beginning with

recruitment of Rab-5-positive early endosomes, followed by fusion with Rab-7-positive late endosomes.⁴⁹ This is accompanied by further maturation and fusion with lysosomes. The capacity of lysosomes to degrade proteins is under developmental control. Mature dendritic cells demonstrate an enhanced capacity to degrade antigen, which correlates with a greater acidification of lysosomes and enhanced lysosomal vacuolar-ATPase activity.⁵⁰ Following degradation and loading onto MHC Class II, mature DCs generate tubules from lysosomal compartments, which fuse directly with the plasma membrane.⁵⁰

Phagocytosis and the MHC Class I Pathway-Intracellular Pathogens and Tumors

In the past few years, much progress has been made to explain the cellular basis for the phenomenon termed "cross-priming," first described by Bevan in 1976.⁵¹ This method of antigen presentation relies on the phagocytosis of an apoptotic cell (e.g., induced by viral infection) and the presentation of phagocytically-derived viral antigens onto MHC Class I. Previously, it had been thought that loading onto MHC Class I occurs only following loading of endogenous antigen from the cytosol to the ER. In contrast, the "phagosome-to-cytosol" pathway of antigen processing involves translocation of peptides or proteins from within the phagosome to the cytoplasm.⁵² There is precedence for proteins crossing membrane barriers; this is the principal mechanism for protein import into mitochondria, for example. How do phagosomal membranes become modified to accomplish a similar task? The observation that ER membrane has the capacity to fuse with phagosomal membrane offers one potential mechanism for this mode of antigen presentation.^{53,54} Sec61, an ER-derived protein translocation channel, was observed to become incorporated into phagosomes of dendritic cells.⁵⁴ Because this protein is capable of "reverse transport" of proteins (in the opposite direction of its recognized ability to translocate nascent chains into the ER),⁵⁵ it is possible that Sec61 provides the means by which proteins translocate across the membrane of the "phago-ER-some" into the cytosol. Once in the cytosol, the protein can be targeted to the proteasome for degradation and further MHC Class I processing by the conventional TAP-dependent pathway.

Cross-priming may be essential for immunity to viruses and other intracellular pathogens. Using a mouse model of viral immunity, Rock and colleagues demonstrated that virally infected nonhematopoietic cells are unable to stimulate primary CTL-mediated immunity directly.⁵⁶ Instead, bone-marrow-derived cells are required as antigen-presenting cells to initiate anti-viral CTL responses. From a teleological standpoint this makes sense. Viruses that typically do not infect hematopoietic cells fail to gain entrance to secondary lymphoid organs. Therefore, they are unlikely to serve as efficient antigen-presenting cells in a primary immune response. However, this interpretation has been called into question. In one recent study, poliovirus in naturally nonpermissive murine APCs acquired viral RNA *in vivo* independently of the cellular virus receptor. The polioviral RNA initiated neosynthesis of viral antigen sufficient to prime CTLs *in vivo*.⁵⁷ It thus remains an open question as to the relative importance of the endogenous pathway and cross-priming in MHC Class I-restricted immunity *in vivo*.

Cross-priming may be critical for tumor immunity. Although this pathway has been demonstrated in a variety of mononuclear phagocytes, it appears that dendritic cells are the most potent APCs in stimulating phagosome-derived MHC Class I-restricted CD8+ CTLs.⁵⁸ Many strategies for using dendritic cells loaded with tumor-derived antigens *ex vivo* have been proposed. Effective tumor immunity requires efficient loading of antigen and effective targeting of antigen-loaded DCs to reach secondary lymphoid organs. As this normally requires a maturation stimulus for the antigen-loaded DCs, various stimuli that serve as adjuvants for the anti-tumor response are under investigation.⁵⁹⁻⁶¹ These include tumor-derived heat-shock proteins,⁶² pro-inflammatory cytokines, and direct activation of costimulatory molecules on

DCs.^{63,64} In some cases, antigen-loading and maturation stimuli are triggered by the same phagocytically-competent receptor. This is the case for Fc γ receptors and complexes of tumor antigen and IgG.^{65,66}

Ingestion of Dead Cells and Regulation of Inflammation; The Power of “Negative Signaling”

Apoptotic cell death is a consequence of cellular senescence or infection of cells with various microbial pathogens. Effective removal of these cells is required for tissue remodeling, for cross-priming, and for resolution of inflammation. This occurs by phagocytosis, which accomplishes the dual role of facilitating the clearance of apoptotic bodies and completing the cell death pathway.⁶⁷ Many receptors on the phagocyte surface participate in this process; among these is the recently identified phosphatidylserine receptor (PSR), which recognizes PS exposed at the outer leaflet of the membrane of apoptotic cells and triggers the generation of TGF- β by the phagocyte. Recognition of apoptotic cells by phagocytes is enhanced by chemotactic stimuli released by apoptotic cells, such as lysophosphatidylcholine.⁶⁸

The consequences of ingestion of dead cells depends on the mechanism of cell death. Necrotic cells release a variety of pro-inflammatory substances, including heat shock proteins, which engage a subset of receptors on the phagocyte surfaces. These, in turn, bind cell surface proteins such as CD91,⁶⁹ which potentially mediates endocytosis and cross-presentation. Heat shock proteins have been proposed to serve as adjuvants by binding toll-like receptors,⁷⁰ although recent work has ascribed this function to contaminating LPS.⁷¹ In contrast, engagement of the PSR results in the production of TGF- β , which serves to dampen inflammation, especially mediated by macrophages.⁷² This may be critical in tuning the immune response; too much inflammation at an inappropriate time may result in excessive antigen presentation and, theoretically, autoimmunity (see below).

Another trigger of “negative signaling” is ligation of immunoreceptor tyrosine-based inhibitory motif (ITIM)-containing receptors. In contrast to Fc γ receptors I, IIa, and III, Fc γ RIIb contains an ITIM in its cytosolic domain. Coclustering of Fc γ RIIb with any other receptor results in an inhibition of phagocytosis and inflammation by recruitment of the SH2 domain-containing inositol 5' phosphatase (SHIP). SHIP hydrolyzes the lipid product of PI 3-kinase, phosphatidylinositol-3,4,5-trisphosphate to phosphatidylinositol-3,4-bisphosphate, thereby limiting recruitment and activation of several PH domain-containing proteins, such as members of the Tec family of tyrosine kinases. These enzymes phosphorylate and activate phospholipase C- γ ; therefore, recruitment of SHIP via ITIMs effectively inhibits signaling mediated by phospholipase C- γ , including calcium fluxes. Recruitment of SHIP via of Fc γ RIIb may be critical for the maintenance of tolerance as Fc γ RIIb^{-/-} mice develop spontaneous autoimmunity in a B cell-autonomous fashion.⁷³ Another type of ITIM bearing receptor, SIRPa, is expressed in myeloid cells and recruits the tyrosine phosphatase SHP-1 and SHP-2. One of its ligands, Surfactant Protein-A,⁷⁴ is also an opsonin for apoptotic cells.⁷⁵ Thus, apoptotic cells may generate mechanistically distinct inhibitory signals: production of TGF- β , a diffusible mediator, via the PSR, and recruitment of the cell autonomous inhibitor, SHP-1/2, by engagement of SIRPa. In other scenarios, SIRPa may serve as an “anti-phagocytic” receptor.⁷⁶ For example, the presence of its counter-receptor on red blood cells, CD47, contributes to their lack of recognition as a phagocytic target. It has been suggested that failure of cognate interactions between CD47 on red blood cells and SIRPa on phagocytic cells may contribute to autoimmune hemolytic anemia.⁷⁷

In summary, by recruitment of enzymes that acts to antagonize tyrosine kinase-mediated signaling (SHP-1/SHP-2) or PI 3-kinase-mediated signaling (SHIP), or by generation of TGF- β , the program of inflammation that accompanies phagocytosis can be curtailed or potentially reversed.

Phagocytosis and Autoimmunity

Individuals with various complement deficiencies are at greatly increased risk for developing autoimmune diseases.⁷⁸ Interestingly, these same complement components are recognized opsonins for apoptotic cells. It has been suggested that decreased clearance of apoptotic bodies leads to an abundant source of potential autoantigens and predisposes to autoimmunity.⁷⁹ This model is supported by the observation of anti-DNA antibodies in mice deficient in opsonizing complement components. In the case of C1q-deficient mice, this was accompanied by enhanced accumulation of apoptotic bodies, suggesting that C1q deficiency causes autoimmunity by impairment of the clearance of apoptotic cells.⁸⁰ A similar correlation between delayed clearance of apoptotic cells and autoimmunity was observed in mice that lacked functional c-Mer, a tyrosine kinase that engages apoptotic cells and mediates their phagocytosis.⁸¹ Thus, the development of autoimmunity is associated with a defect in the phagocytic clearance of apoptotic cells. Despite the appeal of this interpretation, it is not known whether autoimmunity and defective phagocytosis are causally related. An alternative, though not mutually exclusive explanation, is that complement is required for presentation of self-antigen, such as chromatin, by complement receptor-bearing stromal cells in the bone marrow. These are presented to potentially immature autoreactive B cells leading to tolerization through clonal deletion and/or anergy.⁸² It is also possible that lack of appropriate phagocytic clearance of apoptotic cells results in an imbalance of pro- and anti-inflammatory signals, favoring the former. While this seems counterintuitive, as the "anti-inflammatory" PSR pathway of apoptotic cell clearance would be preserved under these circumstances, other anti-inflammatory pathways of corpse clearance may exist. For example, the acute phase reactant C-reactive protein binds to apoptotic cells and serves as both an opsonin and an inducer of TGF- β in a C1q-dependent fashion.⁸³

Conclusions

The study of phagocytosis and its assigned role in immunity has come a long way since Metchnikoff. Phagocytosis is now viewed as a fundamental mechanism of antigen presentation, both through MHC Class II and Class I (via cross-priming). This has implications for immunity not only to bacteria and fungi, but also to viruses and malignant cells. The dichotomy of "pro-inflammatory" and "anti-inflammatory" phagocytosis is useful in explaining how the immune system regulates the temporal response to infection. The balance between these pathways may prove to be a key determinant of the development of autoimmunity.

Acknowledgements

I thank Benjamin Dale and Peter Henson for their critical reading of the manuscript. This work was supported by NIH HL54164.

References

1. Franc NC, White K, Ezekowitz RAB. Phagocytosis and development: Back to the future. *Curr Opin Immunol* 1999; 11(1):47-52.
2. Greenberg S, Grinstein S. Phagocytosis and innate immunity. *Curr Opin Immunol* 2002; 14(1):136-145.
3. Garcia-Garcia E, Rosales C. Signal transduction during Fc receptor-mediated phagocytosis. *J Leukoc Biol* 2002; 72(6):1092-1108.
4. Fitzer-Attas CJ, Lowry M, Crowley MT et al. Fc γ receptor-mediated phagocytosis in macrophages lacking the Src family tyrosine kinases Hck, Fgr, and Lyn. *J Exp Med* 2000; 191(4):669-682.
5. Crowley MT, Costello PS, Fitzer-Attas CJ et al. A critical role for Syk in signal transduction and phagocytosis mediated by Fc γ receptors on macrophages. *J Exp Med* 1997; 186:1027-1039.
6. Cox D, Chang P, Kurosaki T et al. Syk tyrosine kinase is required for immunoreceptor tyrosine activation motif-dependent actin assembly. *J Biol Chem* 1996; 271(28):16597-16602.

7. Matsuda M, Park JG, Wang DC et al. Abrogation of the Fc γ receptor IIA-mediated phagocytic signal by stem-loop Syk antisense oligonucleotides. *Mol Biol Cell* 1996; 7(7):1095-1106.
8. Kiefer F, Brumell J, Al-Alawi N et al. The Syk protein tyrosine kinase is essential for Fc γ receptor signaling in macrophages and neutrophils. *Mol Cell Biol* 1998; 18(7):4209-4220.
9. Cox D, Tseng C-C, Bjekic G et al. A requirement for phosphatidylinositol 3-kinase in pseudopod extension. *J Biol Chem* 1999; 274:1240-1247.
10. Ninomiya N, Hazeki K, Fukui Y et al. Involvement of phosphatidylinositol 3-kinase in Fc γ receptor signaling. *J Biol Chem* 1994; 269:22732-22737.
11. Araki N, Johnson MT, Swanson JA. A role for phosphoinositide 3-kinase in the completion of macropinocytosis and phagocytosis by macrophages. *J Cell Biol* 1996; 135:1249-1260.
12. Botelho RJ, Teruel M, Dierckman R et al. Localized biphasic changes in phosphatidylinositol 4,5-bisphosphate at sites of phagocytosis. *J Cell Biol* 2000; 151(7):1353-1368.
13. Mansfield PJ, Shayman JA, Boxer LA. Regulation of polymorphonuclear leukocyte phagocytosis by myosin light chain kinase after activation of mitogen-activated protein kinase. *Blood* 2000; 95(7):2407-2412.
14. Larsen EC, DiGennaro JA, Saito N et al. Differential requirement for classic and novel PKC isoforms in respiratory burst and phagocytosis in RAW 264.7 cells. *J Immunol* 2000; 165(5):2809-2817.
15. Lennartz MR, Brown EJ. Arachidonic acid is essential for IgG Fc receptor-mediated phagocytosis by human monocytes. *Journal of Immunology* 1991; 147(2):621-626.
16. Kusner DJ, Hall CF, Jackson S. Fc γ receptor-mediated activation of phospholipase D regulates macrophage phagocytosis of IgG-opsonized particles. *J Immunol* 1999; 162:2266-2274.
17. Lennartz MR, Yuen AFC, Masi SM et al. Phospholipase A₂ inhibition results in sequestration of plasma membrane into electron-lucent vesicles during IgG-mediated phagocytosis. *J Cell Sci* 1997; 110:2041-2052.
18. Mancuso P, Nana-Sinkam P, Peters-Golden M. Leukotriene B₄ augments neutrophil phagocytosis of *Klebsiella pneumoniae*. *Infect Immun* 2001; 69(4):2011-2016.
19. Cox D, Chang P, Zhang Q et al. Requirements for both Rac1 and Cdc42 in membrane ruffling and phagocytosis in leukocytes. *J Exp Med* 1997; 186:1487-1494.
20. Forsberg M, Druid P, Zheng L et al. Activation of Rac2 and Cdc42 on Fc and complement receptor ligation in human neutrophils. *J Leukoc Biol* 2003; 74(4):611-619, Epub 2003 Jul 2001.
21. Caron E, Hall A. Identification of two distinct mechanisms of phagocytosis controlled by different Rho GTPases. *Science* 1998; 282:1717-1721.
22. Massol P, Montcourrier P, Guillemot J-C et al. Fc receptor-mediated phagocytosis requires CDC42 and Rac1. *EMBO J* 1998; 17:6219-6229.
23. Zhang Q, Cox D, Tseng C-C et al. A requirement for ARF6 in Fc γ receptor-mediated phagocytosis in macrophages. *J Biol Chem* 1998; 273:19977-19981.
24. Niedergang F, Colucci-Guyon E, Dubois T et al. ADP ribosylation factor 6 is activated and controls membrane delivery during phagocytosis in macrophages. *J Cell Biol* 2003; 161(6):1143-1150, Epub 2003 Jun 1116.
25. Cox D, Lee DJ, Dale BM et al. A Rab11-containing rapidly recycling compartment in macrophages that promotes phagocytosis. *Proc Natl Acad Sci USA* 2000; 97:680-685.
26. Hackam DJ, Rotstein OD, Sjolin C et al. v-SNARE-dependent secretion is required for phagocytosis. *Proc Natl Acad Sci USA* 1998; 95(20):11691-11696.
27. Cox D, Berg JS, Cammer M et al. Myosin X is a downstream effector of PI(3)K during phagocytosis. *Nature Cell Biology* 2002; 4(7):469-477.
28. Steele C, Marrero L, Swain S et al. Alveolar macrophage-mediated killing of *Pneumocystis carinii* f. sp. *muris* involves molecular recognition by the Dectin-1 β -glucan receptor. *J Exp Med* 2003; 198(11):1677-1688.
29. Schmitter T, Agerer F, Peterson L et al. Granulocyte CEACAM3 Is a phagocytic receptor of the innate immune system that mediates recognition and elimination of human-specific pathogens. *J Exp Med* 2004; 199(1):35-46.
30. McCaw SE, Schneider J, Liao EH et al. Immunoreceptor tyrosine-based activation motif phosphorylation during engulfment of *Neisseria gonorrhoeae* by the neutrophil-restricted CEACAM3 (CD66d) receptor. *Mol Microbiol* 2003; 49(3):623-637.
31. Shen Y, Naujokas M, Park M et al. InIB-dependent internalization of *Listeria* is mediated by the Met receptor tyrosine kinase. *Cell* 2000; 103(3):501-510.

32. Braun L, Ghebrehiwet B, Cossart P. gC1q-R/p32, a C1q-binding protein, is a receptor for the InlB invasion protein of *Listeria monocytogenes*. *EMBO J* 2000; 19(7):1458-1466.
33. Lecuit M, Hurme R, Pizarro-Cerda J et al. A role for α - and β -catenins in bacterial uptake. *Proc Natl Acad Sci USA* 2000; 97(18):10008-10013.
34. Mengaud J, Ohayon H, Gounon P et al. E-cadherin is the receptor for internalin, a surface protein required for entry of *L. monocytogenes* into epithelial cells. *Cell* 1996; 84(6):923-932.
35. Alrutz MA, Isberg RR. Involvement of focal adhesion kinase in invasin-mediated uptake. *Proc Natl Acad Sci USA* 1998; 95(23):13658-13663.
36. Tran Van Nhieu G, Caron E, Hall A et al. IpaC induces actin polymerization and filopodia formation during *Shigella* entry into epithelial cells. *EMBO J* 1999; 18(12):3249-3262.
37. Hardt W-D, Chen L-M, Schuebel KE et al. *S. typhimurium* encodes an activator of Rho GTPases that induces membrane ruffling and nuclear responses in host cells. *Cell* 1998; 93:815-826.
38. Vakevainen M, Greenberg S, Hansen EJ. Inhibition of phagocytosis by *Haemophilus ducreyi* requires expression of the LspA1 and LspA2 proteins. *Infect Immun* 2003; 71(10):5994-6003.
39. Black DS, Bliska JB. Identification of p130^{Cas} as a substrate of *Yersinia* YopH (Yop51), a bacterial protein tyrosine phosphatase that translocates into mammalian cells and targets focal adhesions. *EMBO J* 1997; 16(10):2730-2744.
40. Black DS, Bliska JB. The RhoGAP activity of the *Yersinia pseudotuberculosis* cytotoxin YopE is required for antiphagocytic function and virulence. *Mol Microbiol* 2000; 37(3):515-527.
41. Malik ZA, Denning GM, Kusner DJ. Inhibition of Ca^{2+} signaling by *Mycobacterium tuberculosis* is associated with reduced phagosome-lysosome fusion and increased survival within human macrophages. *J Exp Med* 2000; 191(2):287-302.
42. Fratti RA, Chua J, Vergne I et al. *Mycobacterium tuberculosis* glycosylated phosphatidylinositol causes phagosome maturation arrest. *Proc Natl Acad Sci USA* 2003; 100(9):5437-5442, Epub 2003 Apr 5417.
43. Russell DG. Phagosomes, fatty acids and tuberculosis. *Nat Cell Biol* 2003; 5(9):776-778.
44. Anes E, Kuhnlel MP, Bos E et al. Selected lipids activate phagosome actin assembly and maturation resulting in killing of pathogenic mycobacteria. *Nat Cell Biol* 2003; 5(9):793-802, Epub 2003 Aug 2024.
45. Clemens DL, Lee BY, Horwitz MA. Deviant expression of Rab5 on phagosomes containing the intracellular pathogens *Mycobacterium tuberculosis* and *Legionella pneumophila* is associated with altered phagosomal fate. *Infect Immunity* 2000; 68(5):2671-2684.
46. Horwitz MA, Maxfield FR. *Legionella pneumophila* inhibits acidification of its phagosome in human monocytes. *J Cell Biol* 1984; 99:1936-1943.
47. Horwitz M. Phagocytosis of the Legionnaires' disease bacterium (*Legionella pneumophila*) occurs by a novel mechanism: Engulfment within a pseudopod coil. *Cell* 1984; 36:27-33.
48. Kagan JC, Roy CR. *Legionella* phagosomes intercept vesicular traffic from endoplasmic reticulum exit sites. *Nat Cell Biol* 2002; 4(12):945-954.
49. Harrison RE, Bucci C, Vieira OV et al. Phagosomes fuse with late endosomes and/or lysosomes by extension of membrane protrusions along microtubules: Role of Rab7 and RILP. *Mol Cell Biol* 2003; 23(18):6494-6506.
50. Trombetta ES, Ebersold M, Garrett W et al. Activation of lysosomal function during dendritic cell maturation. *Science* 2003; 299(5611):1400-1403.
51. Bevan MJ. Cross-priming for a secondary cytotoxic response to minor H antigens with H-2 congenic cells which do not cross-react in the cytotoxic assay. *J Exp Med* 1976; 143(5):1283-1288.
52. Thery C, Amigorena S. The cell biology of antigen presentation in dendritic cells. *Curr Opin Immunol* 2001; 13(1):45-51.
53. Guermontprez P, Saveanu L, Kleijmeer M et al. ER-phagosome fusion defines an MHC Class I cross-presentation compartment in dendritic cells. *Nature* 2003; 425(6956):397-402.
54. Houde M, Bertholet S, Gagnon E et al. Phagosomes are competent organelles for antigen cross-presentation. *Nature* 2003; 425(6956):402-406.
55. Wiertz EJ, Tortorella D, Bogoy M et al. Sec61-mediated transfer of a membrane protein from the endoplasmic reticulum to the proteasome for destruction. *Nature* 1996; 384(6608):432-438.
56. Sigal LJ, Crotty S, Andino R et al. Cytotoxic T-cell immunity to virus-infected nonhaematopoietic cells requires presentation of exogenous antigen. *Nature* 1999; 398(6722):77-80.

57. Freigang S, Egger D, Bienz K et al. Endogenous neosynthesis vs. Cross-presentation of viral antigens for cytotoxic T cell priming. *Proc Natl Acad Sci USA* 2003; 100(23):13477-13482, Epub 2003 Oct 13431.
58. Albert ML, Sauter B, Bhardwaj N. Dendritic cells acquire antigen from apoptotic cells and induce class I- restricted CTLs. *Nature* 1998; 392(6671):86-89.
59. Ochsenbein AF. Principles of tumor immunosurveillance and implications for immunotherapy. *Cancer Gene Ther* 2002; 9(12):1043-1055.
60. Arina A, Tirapu I, Alfaro C et al. Clinical implications of antigen transfer mechanisms from malignant to dendritic cells. Exploiting cross-priming. *Exp Hematol* 2002; 30(12):1355-1364.
61. Ward S, Casey D, Labarthe MC et al. Immunotherapeutic potential of whole tumour cells. *Cancer Immunol Immunother* 2002; 51(7):351-357, Epub 2002 Jun 2014.
62. Suto R, Srivastava PK. A mechanism for the specific immunogenicity of heat shock protein-chaperoned peptides. *Science* 1995; 269(5230):1585-1588.
63. Mackey MF, Gunn JR, Maliszewsky C et al. Dendritic cells require maturation via CD40 to generate protective antitumor immunity. *J Immunol* 1998; 161(5):2094-2098.
64. Labeur MS, Roters B, Pers B et al. Generation of tumor immunity by bone marrow-derived dendritic cells correlates with dendritic cell maturation stage. *J Immunol* 1999; 162(1):168-175.
65. Regnault A, Lankar D, Lacabanne V et al. Fcγ receptor-mediated induction of dendritic cell maturation and major histocompatibility complex class I-restricted antigen presentation after immune complex internalization. *J Exp Med* 1999; 189(2):371-380.
66. Rafiq K, Bergtold A, Clynes R. Immune complex-mediated antigen presentation induces tumor immunity. *J Clin Invest* 2002; 110(1):71-79.
67. Hoeppner DJ, Hengartner MO, Schnabel R. Engulfment genes cooperate with ced-3 to promote cell death in *Caenorhabditis elegans*. *Nature* 2001; 412(6843):202-206.
68. Lauber K, Bohn E, Krober SM et al. Apoptotic cells induce migration of phagocytes via caspase-3-mediated release of a lipid attraction signal. *Cell* 2003; 113(6):717-730.
69. Binder RJ, Han DK, Srivastava PK. CD91: A receptor for heat shock protein gp96. *Nat Immunol* 2000; 1(2):151-155.
70. Vabulas RM, Wagner H, Schild H. Heat shock proteins as ligands of toll-like receptors. *Curr Top Microbiol Immunol* 2002; 270:169-184.
71. Gao B, Tsan MF. Endotoxin contamination in recombinant human heat shock protein 70 (Hsp70) preparation is responsible for the induction of tumor necrosis factor alpha release by murine macrophages. *J Biol Chem* 2003; 278(1):174-179.
72. Fadok VA, Bratton DL, Konowal A et al. Macrophages that have ingested apoptotic cells in vitro inhibit proinflammatory cytokine production through autocrine/paracrine mechanisms involving TGF-β, PGE2, and PAF. *J Clin Invest* 1998; 101(4):890-898.
73. Bolland S, Ravetch JV. Spontaneous autoimmune disease in FcγRIIB-deficient mice results from strain-specific epistasis. *Immunity* 2000; 13(2):277-285.
74. Gardai SJ, Xiao YQ, Dickinson M et al. By binding SIRPα or calreticulin/CD91, lung collectins act as dual function surveillance molecules to suppress or enhance inflammation. *Cell* 2003; 115(1):13-23.
75. Vandivier RW, Ogden CA, Fadok VA et al. Role of surfactant proteins A, D, and C1q in the clearance of apoptotic cells in vivo and in vitro: Calreticulin and CD91 as a common collectin receptor complex. *J Immunol* 2002; 169(7):3978-3986.
76. Oldenborg PA, Gresham HD, Lindberg FP. CD47-signal regulatory protein alpha (SIRP alpha) regulates Fc gamma and complement receptor-mediated phagocytosis. *J Exp Med* 2001; 193(7):855-861.
77. Oldenborg PA, Zheleznyak A, Fang YF et al. Role of CD47 as a marker of self on red blood cells. *Science* 2000; 288(5473):2051-2054.
78. Hart SP, Smith JR, Dransfield I. Phagocytosis of opsonized apoptotic cells: Roles for 'old-fashioned' receptors for antibody and complement. *Clin Exp Immunol* 2004; 135(2):181-185.
79. Taylor PR, Carugati A, Fadok VA et al. A hierarchical role for classical pathway complement proteins in the clearance of apoptotic cells in vivo. *J Exp Med* 2000; 192(3):359-366.
80. Bickerstaff MC, Botto M, Hutchinson WL et al. Serum amyloid P component controls chromatin degradation and prevents antinuclear autoimmunity. *Nat Med* 1999; 5(6):694-697.

81. Cohen PL, Caricchio R, Abraham V et al. Delayed apoptotic cell clearance and lupus-like autoimmunity in mice lacking the c-met membrane tyrosine kinase. *J Exp Med* 2002; 196(1):135-140.
82. Paul E, Carroll MC. SAP-less chromatin triggers systemic lupus erythematosus. *Nat Med* 1999; 5(6):607-608.
83. Gershov D, Kim S, Brot N et al. C-reactive protein binds to apoptotic cells, protects the cells from assembly of the terminal complement components, and sustains an antiinflammatory innate immune response: Implications for systemic autoimmunity. *J Exp Med* 2000; 192(9):1353-1363.
84. Ghiran I, Barbashov SF, Klickstein LB et al. Complement receptor 1/CD35 is a receptor for mannan-binding lectin. *J Exp Med* 2000; 192(12):1797-1808.
85. Ross GD. Regulation of the adhesion versus cytotoxic functions of the Mac- 1/CR3/ α M β 2-integrin glycoprotein. *Crit Rev Immunol* 2000; 20(3):197-222.
86. Zaffran Y, Zhang L, Ellner JJ. Role of CR4 in Mycobacterium tuberculosis-human macrophages binding and signal transduction in the absence of serum. *Infect Immun* 1998; 66(9):4541-4544.
87. Malaviya R, Gao Z, Thankavel K et al. The mast cell tumor necrosis factor alpha response to FimH-expressing Escherichia coli is mediated by the glycosylphosphatidylinositol-anchored molecule CD48. *Proc Natl Acad Sci USA* 1999; 96(14):8110-8115.
88. Linehan SA, Martinez-Pomares L, Gordon S. Macrophage lectins in host defence. *Microbes Infect* 2000; 2(3):279-288.
89. Suzuki H, Kurihara Y, Takeya M et al. A role for macrophage scavenger receptors in atherosclerosis and susceptibility to infection. *Nature* 1997; 386(6622):292-296.
90. Platt N, Suzuki H, Kurihara Y et al. Role for the class A macrophage scavenger receptor in the phagocytosis of apoptotic thymocytes in vitro. *Proc Natl Acad Sci USA* 1996; 93(22):12456-12460.
91. Thomas CA, Li Y, Kodama T et al. Protection from lethal gram-positive infection by macrophage scavenger receptor-dependent phagocytosis. *J Exp Med* 2000; 191(1):147-156.
92. Shiratsuchi A, Kawasaki Y, Ikemoto M et al. Role of class B scavenger receptor type I in phagocytosis of apoptotic rat spermatogenic cells by Sertoli cells. *J Biol Chem* 1999; 274(9):5901-5908.
93. Imachi H, Murao K, Hiramane C et al. Human scavenger receptor B1 is involved in recognition of apoptotic thymocytes by thymic nurse cells. *Lab Invest* 2000; 80(2):263-270.
94. Palecanda A, Paulauskis J, Al-Mutairi E et al. Role of the scavenger receptor MARCO in alveolar macrophage binding of unopsonized environmental particles. *J Exp Med* 1999; 189(9):1497-1506.
95. Scott RS, McMahon EJ, Pop SM et al. Phagocytosis and clearance of apoptotic cells is mediated by MER. *Nature* 2001; 411(6834):207-211.
96. Fadok VA, Bratton DL, Rose DM et al. A receptor for phosphatidylserine-specific clearance of apoptotic cells. *Nature* 2000; 405(6782):85-90.
97. Savill J, Hogg N, Ren Y et al. Thrombospondin cooperates with CD36 and the vitronectin receptor in macrophage recognition of neutrophils undergoing apoptosis. *Journal of Clinical Investigation* 1992; 90(4):1513-1522.
98. Fadok VA, Warner ML, Bratton DL et al. CD36 is required for phagocytosis of apoptotic cells by human macrophages that use either a phosphatidylserine receptor or the vitronectin receptor $\alpha_v\beta_3$. *J Immunol* 1998; 161(11):6250-6257.
99. Heale J-P, Pollard AJ, Crookall K et al. Two distinct receptors mediate nonopsonic phagocytosis of different strains of Pseudomonas aeruginosa. *J Infect Dis* 2001; 183(8):1214-1220.
100. Devitt A, Moffatt OD, Raykundalia C et al. Human CD14 mediates recognition and phagocytosis of apoptotic cells. *Nature* 1998; 392(6675):505-509.
101. Isberg RR, Leong JM. Multiple β_1 chain integrins are receptors for invasins, a protein that promotes bacterial penetration into mammalian cells. *Cell* 1990; 60(5):861-871.
102. Savill J, Dransfield I, Hogg N et al. Vitronectin receptor-mediated phagocytosis of cells undergoing apoptosis. *Nature* 1990; 343:170-173.
103. Albert ML, Pearce SF, Francisco LM et al. Immature dendritic cells phagocytose apoptotic cells via $\alpha_v\beta_5$ and CD36, and cross-present antigens to cytotoxic T lymphocytes. *J Exp Med* 1998; 188(7):1359-1368.
104. Finnemann SC, Rodriguez-Boulan E. Macrophage and retinal pigment epithelium phagocytosis: Apoptotic cells and photoreceptors compete for $\alpha_v\beta_3$ and $\alpha_v\beta_5$ integrins, and protein kinase C regulates $\alpha_v\beta_5$ binding and cytoskeletal linkage. *J Exp Med* 1999; 190(6):861-874.



<http://www.springer.com/978-0-387-25419-7>

Molecular Mechanisms of Phagocytosis

Rosales, C. (Ed.)

2005, XI, 154 p., Hardcover

ISBN: 978-0-387-25419-7