

■ Table of Contents

1	Mature T-cell and NK-cell Neoplasms	
■ 1.1	Mycosis Fungoides	2
■	Familial mycosis fungoides; involving a father and daughter T. HORIBE, H. YAGI, M. TAKIGAWA and K. TOMITA	2
■	A case of mycosis fungoides with polyarthritis showing clonal identity in skin and synovial tissue Y. SANDBERG, A.W. LANGERAK and F. HEULE	4
■	Vesicular mycosis fungoides M.H. VERMEER, M.W. BEKKENK, P.M. JANSSEN, C.J.L.M. MEIJER and R. WILLEMZE	6
■	Unusual case of mycosis fungoides clinically mimicking a dermatophytosis subsequently evolving in aggressive CD30+ cutaneous lymphoma L. CITARELLA and S. CHIMENTI	8
■	Sclero-atrophic mycosis fungoides with a cytotoxic atypical phenotype M.G. BERNENGO, P. QUAGLINO and M. NOVELLI	10
■	CD56+ early mycosis fungoides L. CERRONI and H. KERL	12
■ 1.2	Sézary Syndrome	14
■	Sézary syndrome with t(2;5) translocation, as developed after prolonged cyclosporine therapy for actinic reticuloid A. RANKI, L. KARENKO and L. JESKANEN	14
■ 1.3	CD30+ T-cell Lymphoproliferative Disorders of the Skin	16
■	CD8+ lymphomatoid papulosis with simultaneous Type A, Type B, and Type C lesions J. J. ABBOTT, R.H. WEENIG and L.E. GIBSON	16

■	Lymphomatoid papulosis with a NK-cell phenotype	18
	M. W. BEKKENK, C. J. L. M. MEIJER and R. WILLEMZE	
■	Anaplastic large cell lymphoma associated with acquired ichthyosis	20
	H. MATSUURA and S. MORIZANE	
■	Anaplastic large T-cell lymphoma	22
	A. SEKULIC, R. H. WEENIG, I. AHMED and M. R. PITTELKOW	
■	Primary cutaneous histiocyte and neutrophil rich CD30+/CD56+ anaplastic large cell lymphoma of the scalp with prominent angio- and neuroinvasion	24
	L. BOUDOVA, D. V. KAZAKOV, P. MUKENŠNABL, V. KUNTSCHER and M. MICHAL	
■	Late relapse of primary cutaneous CD30+ anaplastic large cell lymphoma confirmed by T-cell receptor (TCR) PCR analysis	26
	Y. SANDBERG, A. W. LANGERAK and F. HEULE	
■	1.4 Other Non-cytotoxic and Cytotoxic Peripheral T-cell Lymphomas	28
■	Peripheral CD4+ T-cell lymphoma with cytotoxic phenotype	28
	A. SEKULIC, M. R. PITTELKOW and R. H. WEENIG	
■	CD8+ disseminated small and medium pleomorphic T-cell lymphoma with blood involvement and secondary hemophagocytic syndrome	30
	M. BAGOT and J. WECHSLER	
■	CD8+ cutaneous T-cell lymphoma with an indolent course	32
	B. LAETSCH, L. SCHÄRER, G. BURG and R. DUMMER	
■	Primary cutaneous pleomorphic small/medium-sized T-cell lymphoma revealed by a plantar callus	34
	G. QUEREUX and B. DRÉNO	
■	Atypical poorly differentiated cutaneous T-cell lymphoma with an angiocentric growth pattern	36
	B. LAETSCH and R. DUMMER	
■	Angiocentric non-Hodgkin's lymphoma of the leg	38
	G. S. WOOD and M. A. HINSHAW	
■	γ/δ cutaneous T-cell lymphoma	40
	R. H. WEENIG and L. E. GIBSON	
■	Epidermotropic and subcutaneous γ/δ T-cell lymphoma	42
	L. CERRONI and H. KERL	

■	Cutaneous T-cell lymphoma presenting as reticular erythematous mucinosis	44
	J.M. TWERSKY and D.F. MUTASIM	
■	1.5 Extranodal NK/T-cell Lymphoma	48
■	Nasal NK-cell lymphoma preceded by a puffy eyelid and swollen cheek due to intramuscular infiltration of Epstein-Barr virus-infected cells	48
	K. IWATSUKI and M. OHTSUKA	
■	Epstein-Barr virus-associated lymphoma mimicking hydro vacciniforme: Final diagnosis of a case reported in 1986	50
	T. OONO	
■	Epstein-Barr virus-positive blastoid NK-cell lymphoma	52
	T. SALE, G.S. WOOD and B.J. LONGLEY	
2	Mature B-cell Neoplasms	
■	2.1 Cutaneous Marginal Zone B-cell Lymphoma (MALT Type)	56
■	Primary cutaneous marginal zone B-cell lymphoma with secondary anetoderma in a patient with Sjögren's syndrome	56
	M.H. VERMEER, J. HOEFNAGEL, P.M. JANSSEN, C.J.L.M. MEIJER and R. WILLEMZE	
■	2.2 Primary Cutaneous Follicle Center Lymphoma Variants: Follicular, Follicular and Diffuse, Diffuse	58
■	Follicular lymphoma with follicular dendritic cell overgrowth	58
	D.V. KAZAKOV, L. BOUDOVA and M. MICHAL	
■	2.3 Cutaneous Diffuse Large B-cell Lymphoma Variants: DLBCL of the Leg, DLBCL Others	60
■	Diffuse large B-cell lymphoma with cutaneous and ocular involvement	60
	W. KEMPE, PH. GOLLING, B. CHRISTEN, E. MESSMER, S. MICHAELIS, G. BURG and R. DUMMER	
■	2.4 Other B-cell Lymphomas	62
■	EBV-positive cutaneous B-cell lymphoproliferative disease after imatinib mesylate (Glivec)	62
	M.W. BEKKENK, M.H. VERMEER, C.J.L.M. MEIJER, P.M. JANSSEN, J.M. MIDDELDORP, S.J.C. STEVENS and R. WILLEMZE	

■ 2.5 Chronic Lymphocytic Leukemia	64
■ B-cell chronic lymphocytic leukemia revealed by an erythroderma .. G. QUEREUX and B. DRÉNO	64
■ B-cell chronic lymphocytic leukemia (B-CLL) at the site of <i>Borrelia burgdorferi</i> infection	66
L. CERRONI and H. KERL	
■ Scarring leukemia cutis	68
I.E. BELOUSOVA and D.V. KAZAKOV	
■ 2.6 Mantle Cell Lymphoma	70
■ Lymphoma-associated insect bite-like reaction arising in a patient with mantle cell lymphoma	70
K. ASAGOE	
3 Immature Hematopoietic Malignancies	
■ 3.1 Blastic NK-cell Lymphoma	74
■ Blastic NK-cell lymphoma	74
H. HASHIZUME	
■ 3.2 Related Disorders	76
■ Subcutaneous splenosis of the abdominal wall	76
L. BOUDOVA, D.V. KAZAKOV, P. MUKENŠNABL, O. HES and M. MICHAL	
■ Post-transplant lymphoproliferative disorder	78
N.J. SAMOLITIS and R.M. HARRIS	

Cutaneous Lymphomas

Unusual Cases 2

Burg, G.; Kempf, W.; Michaelis, S.; Feit, J. (Eds.)

2006, XV, 96 p. 236 illus. in color., Hardcover

ISBN: 978-3-7985-1609-0