

# Expression and Nutritional Regulation of Lipogenic Genes in the Ruminant Lactating Mammary Gland

L. Bernard, C. Leroux and Y. Chilliard

**Abstract** The effect of nutrition on milk fat yield and composition has largely been investigated in cows and goats, with some differences for fatty acid (FA) composition responses and marked species differences in milk fat yield response. Recently, the characterization of lipogenic genes in ruminant species allowed *in vivo* studies focused on the effect of nutrition on mammary expression of these genes, in cows (mainly fed milk fat-depressing diets) and goats (fed lipid-supplemented diets). These few studies demonstrated some similarities in the regulation of gene expression between the two species, although the responses were not always in agreement with milk FA secretion responses. A central role for *trans*-10 C18:1 and *trans*-10, *cis*-12 CLA as regulators of milk fat synthesis has been proposed. However, *trans*-10 C18:1 does not directly control milk fat synthesis in cows, despite the fact that it largely responds to dietary factors, with its concentration being negatively correlated with milk fat yield response in cows and, to a lesser extent, in goats. Milk *trans*-10, *cis*-12 CLA is often correlated with milk fat depression in cows but not in goats and, when postruminally infused, acts as an inhibitor of the expression of key lipogenic genes in cows. Recent evidence has also proven the inhibitory effect of the *trans*-9, *cis*-11 CLA isomer. The molecular mechanisms by which nutrients regulate lipogenic gene expression have yet to be well identified, but a central role for SREBP-1 has been outlined as mediator of FA effects, whereas the roles of PPARs and STAT5 need to be determined. It is expected that the development of *in vitro* functional systems for lipid synthesis and secretion will allow future progress toward (1) the identification of the inhibitors and activators of fat synthesis, (2) the knowledge of cellular mechanisms, and (3) the understanding of differences between ruminant species.

**Keywords:** nutrition · gene expression · lipogenesis · mammary gland · lactating ruminant

---

L. Bernard

Adipose Tissue and Milk Lipid Laboratory, Herbivore Research Unit, INRA-Theix,  
63 122 St Genès-Champanelle, France  
e-mail: laurence.bernard@clermont.inra.fr

## Introduction

In ruminants, the major constituents of milk (lipids, proteins, carbohydrates, and salts) and their concentrations are linked to intrinsic or extrinsic (nutritional and environmental) factors (Coleman et al., 2000). Among these, the genetic factor, through the animal species, and the nutritional factor are the two major factors determining milk composition. Indeed, nutrition has a considerable effect on the composition of the lipids (Jensen, 2002), conversely to the protein fraction, which generally is only marginally affected by this factor (Coulon et al., 2001). Moreover, milk fat is an important component of the nutritional quality of dairy products, with the saturated fatty acids (FA) (mainly C12, C14, and C16) commonly considered to have a negative effect on human health when consumed in excess (Williams, 2000), whereas other FA, such as oleic and linolenic acids, have positive effects by a direct vascular antiatherogenic action (Massaro et al., 1999). Besides this, *cis*-9, *trans*-11 C18:2, the major conjugated linoleic acid (CLA) isomer found in ruminant products including milk, was shown both in animals studies and *in vitro* experiments to exert a number of advantageous physiological effects (Pariza et al., 2001). In addition, milk fat content and composition is one of the most important components of the technological and sensorial qualities of dairy products. Thus, modification of milk fat content and FA composition by dietary manipulation has been investigated in cows and goats, with particular attention on the effects of fat supplementation of the diet.


In bovines, the consequences of lipid supplementation on the milk yield and fat and protein contents have been well described, with an increase in milk production (for most lipid supplements) and a slight but systematic reduction in the protein and casein contents. In dairy cows, due to important interactions between dietary forages and concentrates and their components (fibers, starch, lipids), the supplements given do not all have the same efficiency regarding fat content modulation. Thus, concentrate-rich diets, concentrate-rich diets supplemented with vegetable oils, and diets supplemented with fish oil lead to a decrease in fat content, while encapsulated lipids lead to a large increase in fat content (Chilliard & Ferlay, 2004; Palmquist et al., 1993). Conversely, in goats almost all types of lipid supplements induce a marked increase in the milk fat content without systematic modification of milk production or protein content (Chilliard et al., 2003a). These modifications in the yield of fat are observed together with an important modification of milk FA composition, which is well documented both in cows (Chilliard et al., 2000, 2001; Palmquist et al., 1993) and in goats (Chilliard et al., 2003a, 2006a).

The mechanisms underlying these intra- and inter-species-specific responses are not yet well understood. Nevertheless, recently, thanks to the characterization of the lipogenic genes involved in milk synthesis and secretion and the development of molecular biology tools, few studies have been undertaken to relate the effects of diet on the milk FA profile to mammary gland lipid metabolism. These studies mainly considered lipogenic genes, in particular genes for the enzymes involved in the uptake, *de novo* synthesis, desaturation, and esterification of FA

in order to relate the effects of the diet on the abundance of their transcripts and/or enzymatic activities. This chapter reviews the present knowledge on the main lipogenic genes, in particular the known effects of nutritional factors, especially those of fat supplementation, on ruminant mammary lipogenic gene expression, together with milk fat content and FA composition. In addition, we present the putative molecular mechanisms underlying these regulations.

## Milk Fatty Acid Origin

Milk fat is composed of ca. 98% triglycerides, of which ca. 95% is FA and less than 1% is phospholipids, with small amounts of cholesterol, 1,2-diacylglycerol, monoacylglycerol, and free FA. Milk FA have a dual origin: (1) They are either *de novo* synthesized in the mammary gland (Fig.1) from acetate and



**Fig. 1** Milk fat synthesis in the ruminant mammary epithelial cell. Abbreviations used: ACC = acetyl-CoA carboxylase; AGPAT = acyl glycerol phosphate acyl transferase; CD36 = cluster of differentiation 36; CLD = cytoplasmic lipid droplet; CoA = coenzyme A; CM = chylomicron; DGAT = diacyl glycerol acyl transferase; ER = endoplasmic reticulum; FA = fatty acid; FABP = fatty acid binding protein; FAS = fatty acid synthase; Glut 1 = glucose transporter1; GPAT = glycerol-3 phosphate acyl transferase; LPL = lipoprotein lipase; MFG = milk fat globule; SCD = stearoyl-CoA desaturase; TG = triglyceride; VLDL = very low density-lipoprotein


3-hydroxybutyrate, produced by ruminal fermentation of carbohydrates and by rumen epithelium from absorbed butyrate, respectively, thus resulting in short- and medium-chain FA (C4:0 to C16:0) that represent 40–50% of the FA secreted in milk, or (2) they are imported from the plasma, where they are either released by the enzyme lipoprotein lipase (LPL) (Barber et al., 1997) from triglycerides circulating in chylomicra or very low-density lipoprotein (VLDL), or derived from the plasma nonesterified fatty acids (NEFA) that circulate bound to albumin, for long-chain FA ( $\geq$ C18) as well as ca. one-half of the C16:0, depending on the diet composition. These long-chain FA originate mainly from dietary lipid absorption from the digestive tract (with the dietary FA undergoing total or partial hydrogenation in the rumen) and from body reserves mobilization (especially at the beginning of lactation). Commonly, mobilization of body fat accounts for less than 10% of milk FA, with this proportion increasing in ruminants in negative energy balance in direct proportion to the extent of the energy deficit (Bauman & Griinari, 2001). Furthermore, FA may be desaturated, but not elongated, in the secretory mammary epithelial cells (MEC) (Chilliard et al., 2000).

## Gene Characterization and Mechanisms of Mammary Lipogenesis

The acetyl-CoA carboxylase (ACC) and fatty acid synthase (FAS) enzymes (encoded by the *ACACA* and *FASN* genes, respectively) are involved in the metabolic pathway for *de novo* FA synthesis in the mammary tissue, whereas the LPL enzyme is involved in the uptake of plasma FA. These FA could be desaturated by the stearoyl-CoA desaturase (SCD), resulting in synthesis of *cis*-9 unsaturated FA, and then esterified to glycerol sequentially via glycerol-3 phosphate acyl transferase (GPAT), acyl glycerol phosphate acyl transferase (AGPAT), and diacyl glycerol acyl transferase (DGAT). Then the triglycerides are secreted as milk fat globules (Fig. 1). The genes specifying these enzymes, implicated in the key processes of lipogenesis within the mammary gland, are candidate genes whose regulation has been studied first.

In ruminants, thanks to the recent knowledge of the cDNA sequences of several lipogenic genes (see Fig. 2)—*LPL* (Bonnet et al., 2000a; Senda et al., 1987), *ACACA* (Barber & Travers, 1998; Mao et al., 2001), *FASN* (Leroux et al., submitted; Roy et al., 2005), SCD (Bernard et al., 2001; Keating et al., 2005; Ward et al., 1998), *DGAT1* (Winter et al., 2002), *AGPAT* (Mistry & Medrano, 2002), and recently *GPAT* (Roy et al., 2006b)—molecular tools for studying their expression have been developed to allow the quantification of their mRNA by northern blot or real-time RT-PCR and/or the activity of the corresponding enzymes.

In addition, the recent development of high-throughput techniques such as microarrays allows us to complete this candidate gene approach. These new methods simultaneously provide data for the expression of thousands of genes



\* tissue preferential expression

**Fig. 2** Characterization (size and structure) of the transcripts of the main lipogenic genes in ruminant species, lipoprotein lipase (*LPL*; Senda et al., 1987; Bonnet et al., 2000), stearoyl-CoA desaturase (*SCD*; Bernard et al., 2001; Keating et al., 2005), fatty acid synthase (*FASN*; Roy et al., 2005; Leroux et al., in preparation), acetyl-CoA carboxylase (*ACACA*; Barber et al., 1998, 2005), glycerol-3 phosphate acyl transferase (*GPAT*; Roy et al., 2006b), acyl glycerol phosphate acyl transferase (*AGPAT*; Mistry & Medrano, 2002), and diacyl glycerol acyl transferase (*DGAT*; Grisart et al., 2002; Winter et al., 2002)

and allow a larger understanding of different mammary functions including milk lipid synthesis and secretion.

### *Uptake of Fatty Acids*

The hydrolysis of lipoprotein triacylglycerol is catalyzed by LPL, which selectively releases FA esterified at the *sn*-1 (-3) position. In bovines, mammary tissue expresses three *LPL* transcripts, which are 1.7, 3.4, and 3.6 Kb in size (Bonnet et al., 2000a; Senda et al., 1987) due to the alternative use of the polyadenylation site. In this species, a predominance of the 3.4-Kb transcript in the mammary gland (Senda et al., 1987) as in ovine adipose tissue and of the 3.6-Kb transcript in muscles (Bonnet et al., 2000b) has been reported. In ovine species, the total cDNA sequence has been reported (Bonnet et al., 2000a). A complex regulation, by dietary and hormonal factors, modulates LPL activity via transcriptional, posttranscriptional, and posttranslational mechanisms. Immediately prior to parturition, mammary LPL activity increases markedly,

remains high throughout lactation, and is simultaneously downregulated in adipose tissue in cows (Shirley et al. 1973) and goats (Chilliard et al., 2003a).

From immunochemical and biochemical studies, it was shown that LPL is located both on (or near) the surface and within the cell of the major cell types of the different tissues as well as on the luminal surface of vascular endothelial cells. Mammary tissue contains various cell types in addition to parenchymal secretory MEC, including adipocytes, in varying proportions to MEC depending on the developmental and physiological status of the mammary gland. Discrepancies concerning LPL localization were reported from histological studies made on the rodent mammary gland. Thus, some studies (Camps et al., 1990) demonstrated the presence of LPL mRNA and protein in MEC using *in situ* hybridization and immunofluorescence techniques, respectively, and concluded that the origin of mammary LPL is the secretory MEC. Conversely, others (Jensen et al., 1991) demonstrated that LPL protein and mRNA are located in mammary depleted adipocytes or adipocyte precursors located in interstitial cells, suggesting that mammary LPL could originate partly from mammary adipocytes to be subsequently secreted and transported by cellular uptake and transcytosis, both to its final site of action on the capillary endothelial cell and through the secretory MEC into milk (Jensen et al., 1991; Neville & Picciano, 1997).

Arteriovenous difference measurements in lactating goats have shown that the utilization of triglycerides and NEFA for milk lipid synthesis is related to their plasma concentrations (Annison et al., 1968). Otherwise, experiments demonstrated that the availability of the substrate determines its utilization by the mammary gland with either a large NEFA utilization when plasma triglycerides are low and NEFA are high, as in fasting animals (West et al., 1972), or no net utilization of NEFA at plasma concentrations below 0.2 mM (Nielsen & Jakobsen, 1994). Similarly, triglyceride utilization increases when its plasma concentration increases. Thus, duodenal lipid infusion (Gagliostro et al., 1991) increased plasma triglyceride concentration and apparent mammary uptake, and there was simultaneously a net production of NEFA by the mammary gland, due to their release in the vascular bed during LPL action (Fig. 1).

The mechanism by which FA crosses the capillary endothelium and interstitial space to reach the MEC has not yet been identified. After arriving at the MEC, FA could cross the plasma membrane by diffusion or via a saturable transport system. In mammals, the acyl-CoA binding proteins (ACBP) (Knudsen et al., 2000) that bind long-chain acyl-CoA have an important role in regulating the FA transport and concentration in the cytosol. Nevertheless, in ruminants, Mikkelsen and Knudsen (1987) found lower concentrations of these ACBP in the mammary gland and muscles compared to the liver cytosol. Elsewhere it has been suggested in rodents and ruminants that another FA binding protein, CD36 (cluster of differentiation 36), expressed in the lactating MEC, and found in the milk fat globule membrane, heart, platelets, and adipocytes, may function as a transporter of long-chain FA (Abumrad et al., 2000). The detection of CD36 in mammary tissue could be linked to the

presence of adipocytes. Nevertheless, the presence of *CD36* mRNA in rodent MEC lines has been shown, with a slight enhancement of its gene expression after the addition of lactogenic hormones (Aoki et al., 1997).

Furthermore, fatty acid binding proteins (FABP; a family of intracellular lipid binding proteins found across numerous species) are involved in the uptake and intracellular trafficking of FA in many tissues (Lehner & Kuksis, 1996). In the bovine mammary gland, co-expression of FABP and *CD36* has been shown, which increases during lactation and decreases during involution (Spitsberg et al., 1995), demonstrating that their expression is related to physiological variations of lipid transport and metabolism within the cell. In the same way, simultaneous elevations of *CD36* mRNA expression, of cytosolic TAG, and of lipid droplets were observed in primary bovine MEC (Yonezawa et al., 2004b). Barber et al. (1997) proposed a role for *CD36* in the transport of FA across the secretory MEC membrane, working in conjunction with intracellular FABP. In the bovine lactating mammary gland, the presence of two forms of FABP has been demonstrated (Specht et al., 1996), identified as A-FABP and H-FABP, thus named according to the tissue of their first detection, adipose tissue and heart, respectively. Studying the proteins' cellular location, Specht et al. (1996) showed that A-FABP and H-FABP were present in myoepithelial cells and MEC, respectively. The expression of *FABP* types is generally interpreted in terms of specialized functions in FA metabolism, with *H-FABP* predominantly found in cells where FA are used as an energy source and probably involved in their  $\beta$ -oxidation. The significance of its abundance in mammary tissue in which active triglyceride synthesis and low FA oxidation occur during lactation remains to be understood.

In addition, ATP-binding cassette (ABC) transporters (a family of membrane proteins) are involved in the transport, against concentration gradients, of a wide variety of compounds, including ions, peptides, sugars, and lipids, at the cost of ATP energy (Klein et al., 1999). The *ABCG5* and *ABCG8* members of this family play an important role in cholesterol homeostasis and have been described specifically in intestine and liver cells in rodents (Mutch et al., 2004; Yu et al., 2002). The recent identification and expression of *ABCG5* and *ABCG8* transporters in the bovine mammary gland open a wide range of future investigations on their potential role in lipid trafficking and excretion during lactation and control of sterol concentrations in milk (Viturro et al., 2006).

## ***De Novo Fatty Acid Synthesis***

### **Acetyl-CoA Carboxylase Gene (*ACAC*)**

The *ACACA* and *ACACB* genes are distinct genes that respectively encode the isoenzymic ACC proteins ACC $\alpha$  and ACC $\beta$ . The *ACACA* gene is expressed in all cell types but is found at its highest levels in the lipogenic tissues

(Lopez-Casillas et al., 1991), where its protein product ACC $\alpha$  provides cytoplasmic malonyl-CoA for FA synthesis. The *ACACB* gene is the major form expressed in heart and skeletal muscles (Abu-Elheiga et al., 1997), where its protein product ACC $\beta$  is implicated in the regulation of the  $\beta$ -oxidation of FA in the mitochondria (Abu-Elheiga et al., 2000). Expression of the ACC $\alpha$  isoenzyme is regulated in a complex fashion in the short term, through allosteric mechanisms with cellular metabolites possessing a positive (citrate) or negative effect (malonyl-CoA and long-chain acyl-CoA), and reversible phosphorylation on a number of specific serine residues, as well as chronically, through the regulation of transcription of the gene (Kim, 1997).

In ruminants, the *ACACA* cDNA sequence was first reported by Barber and Travers (1995) and corresponds to the synthesis of a protein with 2,346 amino acids. More recently, the *ACACA* gene was characterized in sheep (Barber & Travers, 1998) and cattle (Mao et al., 2001). Initially, the existence of three promoters, PI, PII, and PIII, was demonstrated, and their use together with alternative splicing of the primary transcripts from promoters I and II results in the generation of a heterogeneous population of transcripts differing in the sequence of their 5'UTR. These promoters are used in tissue-differential fashion. PIII use is limited to lung, liver, kidney, brain, and predominantly the lactating mammary gland in bovine (Mao et al., 2002) and ovine (Barber et al., 2003) species. PII expression is ubiquitous, with an elevated expression in the lactating mammary gland. PI is preferentially used in adipose tissue and liver under lipogenic conditions and in lactating bovine mammary gland. PI generate ~30% of *ACACA* mRNA (Mao et al., 2001), while in the ovine mammary gland, its contribution is low (2%; Molenaar et al., 2003). The reason for this difference of promoter usage between the bovine and the ovine is unknown. In addition, Barber et al. (2005) demonstrated the existence of a fourth promoter in human, rodent, and ruminant species and mainly expressed in brain, which again underlines the complexity of the structure and regulation of the *ACACA* gene.

### **Fatty Acid Synthase Gene (*FASN*)**

The *FASN* gene encodes the protein FAS, which, under a complex homodimeric form, is responsible for the synthesis of short- and medium-chain FA (C4-C16) in the mammary gland during lactation (Wakil, 1989). In ruminants, the FAS enzyme contains six catalytic activity domains on a single protein of 2,513 amino acids. Contrary to what is observed in rodent mammary gland and duck uropygial gland, ruminant FAS synthesizes medium-chain FA without the implication of a thioesterase II (Barber et al., 1997). In addition to being able to load acetyl-CoA, malonyl-CoA, and butyryl-CoA, ruminant FAS contains a loading acyltransferase whose substrate specificity extends to up to C12, with the result that it is able to load and also release these medium-chain FA (Knudsen & Grunnet, 1982). This way of medium-chain FA synthesis is specific to the

lactating ruminant mammary gland, whereas the product of FAS in other ruminant tissues is predominantly C16:0, as in non-ruminant tissues (Christie, 1979). *FASN* mRNA ranges in size from 8.4 to 9.3 Kb, depending on the species: In several human tissues (Jayakumar et al., 1995), bovine mammary gland (Beswick & Kennelly, 1998), and ovine (Bonnet et al., 1998) and porcine (Ding et al., 2000) adipose tissues, only one transcript has been detected by northern blot. Conversely, two mRNA, generated by the use of two alternative polyadenylation signals, have been detected in rat adipose tissue (Guichard et al., 1992) and mammary gland (Schweizer et al., 1989). The gene, termed *FASN*, has recently been cloned in bovine (Roy et al., 2005), and an alternate transcript was discovered without part of exon 9 (minus 358 bp). In caprine, the cDNA was recently characterized with only one transcript described (Leroux et al., submitted). In addition, Roy et al. (2006c) identified several single-nucleotide polymorphisms (SNPs) in the bovine *FASN* gene, and the analysis of two of them, located respectively in exon 1 and 34, suggested an association of these polymorphisms with variations in milk fat content.

### ***Stearoyl-CoA Desaturase***

The *SCD* gene encodes a protein of 359 amino acid residues located in the endoplasmic reticulum that catalyzes the  $\Delta$ -9 desaturation, introducing a *cis* double bond, of a spectrum of fatty acyl-CoA substrates, mainly from C14 to C19. In rodents, *SCD* relies on different genes whose expression and regulation by polyunsaturated FA (PUFA) are tissue-specific (Ntambi, 1999). Conversely, in ruminants there is only one *SCD* gene (Bernard et al., 2001), generating a 5-Kb transcript that was characterized in sheep (Ward et al., 1998), cows (Chung et al., 2000), and goats (Bernard et al., 2001). In goats, the 3'-UTR sequence derives from a single exon and is unusually long (3.8 Kb), as observed for humans (Zhang et al., 1999), rats (Mihara, 1990), and mice (Ntambi et al., 1988). In addition, the caprine 3'-UTR is characterized by the presence of several AU-rich elements, which could be mRNA destabilization sequences, and presents a genetic polymorphism with the presence or absence of a triplet nucleotide (TGT) in position 3178-3180 (Bernard et al., 2001, and GenBank accession number AF325499). Immediately after parturition, *SCD* mRNA in ovine (Ward et al., 1998) and activity in bovine (Kinsella, 1970) increased in the mammary gland. In lactating goats, the *SCD* gene is highly expressed in the mammary gland and subcutaneous adipose tissue, compared to perirenal adipose tissue (Bernard et al., 2005b). In the lactating mammary gland, palmitoleoyl-CoA and oleoyl-CoA are synthesized from palmitoyl-CoA and stearoyl-CoA by the action of the SCD enzyme (Enoch et al., 1976). In addition, in bovine mammary gland, SCD is responsible for the synthesis of the major part of *cis*-9, *trans*-11- (Corl et al., 2001; Griinari et al., 2000; Loores et al., 2005d; Shingfield et al., 2003) and of *trans*-7, *cis*-9- (Corl et al., 2002) CLA isomers.

The promoter region of the bovine *SCD* gene has recently been characterized (Keating et al., 2006), and a region of critical importance, designated stearyl-CoA desaturase transcriptional enhancer element (STE) and containing three binding complexes, was identified in the MAC-T cell. In addition, this STE region was shown to play a key role in the inhibitory effect on *SCD* gene transcription of *trans*-10, *cis*-12 CLA and, to a lesser extent, of *cis*-9, *trans*-11 CLA (Keating et al., 2006).

### ***Esterification of FA to Glycerol***

In mammals, FA are not distributed randomly on the *sn*-1, *sn*-2, and *sn*-3 positions of the glycerol backbone of the milk triglycerides; this nonrandom distribution determines functional and nutritional attributes (German et al., 1997). In bovine, a description of the stereospecific position of the major FA in TAG has been reviewed by Jensen (2002). A high proportion (56–62%) of FA esterified at positions *sn*-1 and *sn*-2 of the glycerol backbone are medium- and long-chain saturated FA (C10:0 to C18:0), with C16:0 equally distributed among *sn*-1 and *sn*-2, C8:0, C10:0, C12:0, and C14:0 more located at *sn*-2, and C18:0 more located at *sn*-1. In addition, about 24% of FA esterified at position *sn*-1 is C18:1. Finally, a high proportion of FA esterified at position *sn*-3 is short-chain FA (C4:0, C6:0, C8:0; 44% on a molar basis) and oleic acid (27%). When consumed by humans, milk TAG are hydrolyzed by pancreatic lipase specifically in the *sn*-1 and *sn*-3 positions, allowing the FA present at position *sn*-2 to be preferentially absorbed because they remain in the monoacyl glycerol form (Small, 1991). The first step in triglyceride biosynthesis is the esterification of glycerol-3-phosphate in the *sn*-1 position, which is catalyzed by the glycerol-3-phosphate acyl transferase (GPAT). Two isoforms of GPAT have been identified in mammals, which can be distinguished by subcellular localization (mitochondrial vs. endoplasmic reticulum) and sensitivity to sulfhydryl group modifying agent N-ethylmaleimide (NEM). The mitochondrial isoform is resistant to NEM, and the endoplasmic reticulum isoform is sensitive to NEM. In rodents, both isoforms have a role in the TAG synthesis in the liver and adipose tissue (Coleman & Lee, 2004). Regarding the mitochondrial *GPAT* gene, the genomic structure and cDNA sequence were recently determined in ruminants, with the presence of two transcripts differing in their 5'-UTR (Roy et al., 2006b).

The second step of triglyceride synthesis is committed by AGPAT (or lysophosphatidic acid acyltransferase, LPAAT). AGPAT has a greater affinity for saturated fatty acyl-CoA (Mistry & Medrano, 2002) in the order C16 > C14 > C12 > C10 > C8 (Marshall & Knudsen, 1977), which is in accordance with the observed high proportion of medium- and long-chain saturated FA at the *sn*-2 position in milk, with palmitate as the major FA (representing 43% of the total palmitate found in triacylglycerol). Consequently, a possible regulation of

substrate specificity of the enzymes of FA esterification should be of major importance for both the mammary cell and human nutrition. In addition, substrate availability in the bovine and ovine mammary gland is also a factor for the *sn*-2 position FA composition, allowing its manipulation to some extent by nutritional factors, in interaction with the aforementioned substrate affinity. Bovine and ovine *AGPAT* genes were characterized, cloned, and located on bovine chromosome 23 (Mistry & Medrano, 2002). Bovine and ovine AGPAT are proteins made up of 287 amino acids that differ by only one amino acid residue.

The third enzyme, DGAT, is located on the endoplasmic reticulum membrane. DGAT is the only protein that is specific to triacylglycerol synthesis and therefore may play an important regulatory role (Mayorek et al., 1989). However, little is known about the regulation of *DGAT* expression, whereas its gene has been particularly well studied in ruminants due to its genetic variability. The complete bovine *DGAT1* gene (Grisart et al., 2002; Winter et al., 2002) and the near-complete coding region of the caprine *DGAT1* gene (Angiolillo et al., 2006) have been sequenced. A quantitative trait loci (QTL) QTL for milk fat content has been detected in the centromeric region of cattle chromosome 14, and *DGAT1* was proposed as a positional and functional candidate for this trait (Winter et al., 2002). These studies found a nonconservative substitution of lysine by alanine (Grisart et al., 2004) in *DGAT1* caused by AA to GC dinucleotide substitution at position 10434 of the gene sequence, in exon 8 (GenBank accession number AY065621). This polymorphism was related to milk composition and yield variations. The K allele was recently shown *in vitro* to be associated with increased activity of the enzyme in agreement with its positive link with bovine milk fat percentage (Grisart et al., 2004). In addition, in the German Holstein population, Kühn et al. (2004) described five alleles at a variable number tandem repeat (VNTR) polymorphism in the *DGAT1* promoter, which showed an effect on fat content additional to the *DGAT1* K232A mutation. Due to the presence of a potential transcription factor binding site in the 18nt element of the VNTR, the variation in the number of tandem repeats of the 18nt element might be causal for the variability in the transcription level of the *DGAT1* gene. In sheep, *DGAT1* is an obvious candidate gene for milk fat content, as a QTL was detected in chromosome 9, which is homologous to the bovine chromosome 14 region (Barillet et al., 2005).

## **Regulation of Mammary Lipogenic Gene Expression by Dietary Factors**

The response of gene expression to nutrient changes involves the control of events that could occur at transcriptional (e.g., through transcription factors), posttranscriptional (e.g., such as mRNA stability), translational (e.g., its initiation, etc.), and posttranslational (e.g., via turnover or activation of enzymatic protein) levels. However, it is often unclear whether the regulatory factors are

the dietary components themselves or their metabolites or are hormonal changes produced in response to the nutritional changes. Moreover, in most of the studies, as in those reported in this chapter, it is difficult to determine the level of regulation involved since, for a given gene, measurements of the relevant mRNA, the enzyme protein content, and the activity are not often studied simultaneously. Only few data are indeed available in ruminants on the nutritional regulation of mammary lipogenic gene expression either *in vivo* or *in vitro*.

The *in vivo* trials have been carried out in midlactation cows and goats. Data in cows come from five studies that were undertaken mainly (four of the five studies) with milk fat-depressing (MFD) diets:

- **Study 1:** a high level of concentrate (containing 76% cracked corn, 19% heat-treated soybean meal, 1% sunflower oil) with a forage- (alfalfa hay) to-concentrate ratio of 16/84, compared to a control diet with a high-forage (composed of 83% corn silage and 17% alfalfa hay) content and a forage-to-concentrate ratio of 53/47 (Peterson et al., 2003; 3 cows),
- **Study 2:** a high level of concentrate (containing 52% ground corn and 15% soybean meal) with a forage- (corn silage) to-concentrate ratio of 25/70, supplemented with 5% of soybean oil, compared to a high level of forage (containing 76% corn silage and 24% alfalfa hay) with a forage-to-concentrate (containing mainly of 57% ground corn and 34% soybean meal) ratio of 60/40 (Piperova et al., 2000; 10 cows),
- **Study 3:** a diet based on grass silage (19%), corn silage (19%), and rolled barley (44%) supplemented, or not, with either 1.7% of glutaraldehyde-protected fish oil or 2.7% unprotected fish oil (Ahnadi et al., 2002; 16 cows),
- **Study 4:** a diet based on grass (27%), corn silage (30%), and rolled barley (22%) supplemented with 3.3% unprotected canola seeds plus 1.5% canola meal, or 4.8% formaldehyde-protected canola seeds, compared to the same diet with 4.8% of canola meal, which, for the two treatment diets, represents a supply of 0.7% and 0.6% of extra lipids compared to the control diet, respectively (Delbecchi et al., 2001; 6 cows),
- **Study 5:** a forage-based diet (containing 25% corn silage, 17% alfalfa silage, and 3% alfalfa hay) with a forage-to-concentrate (22% of ground corn and 28% of grain mix) ratio of 45/55 and supplemented with 3% of soybean oil and 1.5% of fish oil, compared to a forage-based diet (containing 35% corn silage, 23% alfalfa silage, and 5% alfalfa hay) with a forage-to-concentrate (15% of ground corn and 20% of grain mix) ratio of 65/35 (Harvatiné & Bauman, 2006; 16 cows).

In goats, data are from two nutritional studies undertaken with dietary lipid supplementations varying in their nature, form of presentation, and dose:

- **Study 6:** a 54% hay diet supplemented, or not, with 3.6% of lipids from oleic sunflower oil or formaldehyde-treated linseeds, with the concentrate fraction containing 40% rolled barley, 17% dehydrated sugar beet pulp, 10% pelleted dehydrated Lucerne, and 17% soybean meal (Bernard et al., 2005c; 14 goats),

- **Study 7:** two diets with 47% of either hay (H;13 goats) or corn silage (CS;14 goats) supplemented, or not, with 5.8% of either linseed or linoleic sunflower oil, with the concentrate fraction containing 24(H)–24(CS)% rolled barley, 39(H)–34(CS) dehydrated sugar beet pulp, 12(H)–0(CS) pelleted dehydrated Lucerne, and 25(H)–42(CS)% soybean meal, respectively (Bernard et al., unpublished 27 goats).

In these two goat studies, conversely to what is generally observed in dairy cows (Bauman & Griinari, 2003), dietary lipid supplementation always increased the milk fat content and did not increase (or only slightly increased) milk yield, in accordance with previously published goat studies (Chilliard et al., 2003a, 2006a). The seven cow and goat studies presented above have been used as a database to allow us to evaluate how dietary factors may change metabolic pathways and lipogenic gene expression, in interaction with animal species peculiarities.

### ***Regulation of Lipoprotein Lipase***

Studies in cows on MFD diets reported either no change in the abundance of mammary *LPL* mRNA (study 3 with 1.7% “protected” fish oil ; Ahnadi et al., 2002) or a tendency to decrease (study 1; Peterson et al., 2003) or a strong decrease (study 5; Harvatine & Bauman, 2006, and study 3 with 2.7% unprotected fish oil; Ahnadi et al., 2002). In addition, still with an MFD diet, no change in the abundance of *FABP* has been observed (Peterson et al., 2003). However, in the two former studies (1 and 3), milk long-chain FA (> C16) yield (g/day) was significantly decreased, by 43% after “protected” fish oil supplementation (study 3) and by 28% with a high-concentrate diet (study 1). Elsewhere, in suckling beef cows fed hay supplemented with 14% high-linoleate safflower seeds, compared to 12% corn and 6% safflower seed meal, a trend toward a greater *LPL* mRNA was observed ( $P = 0.09$ ; Murrieta et al., 2006) together with an increase in long-chain FA percentage in milk fat. Furthermore, in goats fed a hay-based diet supplemented with oleic sunflower oil (study 6; Bernard et al., 2005c), no significant effect was observed on mammary *LPL* activity, whereas *LPL* mRNA content was increased, together with a large increase (83%) in long-chain FA (C18) secretion. In goats fed with hay or corn silage supplemented with either linseed or sunflower oil (study 7), a large increase in the secretion of milk long-chain FA (> 100%) was observed without any effect on mammary *LPL* activity and mRNA except when corn silage was supplemented with sunflower oil, in which case *LPL* activity increased. In addition, in late-lactating goats fed an alfalfa hay-based diet, supplementation with 3.8% lipids from soybeans had no effect on mammary *LPL* mRNA abundance, while the secretion of long-chain FA (C18) in milk was increased by 58% (Bernard et al., 2005b). Thus, most of the results available in ruminants are not in agreement with those reported in rodents, where a very high dietary

lipid intake (20%) enhanced both mammary gland LPL activity and lipid uptake (Del Prado et al., 1999).

The aforementioned results led to the hypothesis that either *LPL* mRNA or LPL activity measured *in vitro* in optimal conditions generally does not limit the uptake of long-chain FA by the mammary gland in ruminants. Other factors such as the availability of plasma triglyceride FA (Gagliostro et al., 1991) and the location of LPL (capillary endothelial cells, MEC, or depleted adipocytes) could play an important role. These results are also in accordance with those from previous studies in lactating goats, demonstrating that mammary utilization of plasma triglycerides and NEFA is related to their arterial concentration (see earlier section). Moreover, the existence of positive correlations between milk stearic acid percentage and milk fat content was observed (Chilliard et al., 2003b), related both to the response to dietary lipids and to individual variations within each dietary treatment (Bernard et al., 2006). This suggests that a significant availability and uptake of this FA is an important factor for milk fat secretion in goats, as in cows (Loor et al., 2005a, b).

### ***Regulation of Genes Involved in de Novo Lipid Synthesis (ACCA and FASN)***


In cows, an MFD diet induced joint reductions in *ACACA* mRNA abundance and ACC activity and in FAS activity in mammary tissue, together with a dramatic 60% decrease in C4–C16 FA secretion (g/day) (study 2; Piperova et al., 2000). Furthermore, Ahnadi et al. (2002; study 3) observed a decrease in mammary *ACACA* and *FASN* mRNA levels along with a 38% decrease in C4–C16 FA secretion with fish oil-supplemented diets. Moreover, in cows with a high-concentrate diet (study 1; Peterson et al., 2003), the observed reductions of milk fat secretion (27% decrease) and of C4–C16 (30% decrease) match a reduction of mammary mRNA abundance of several genes involved in milk lipid synthesis pathways, in particular *ACACA* and *FASN*, whereas no effect on milk  $\kappa$ -casein mRNA was observed. In addition, in cows fed an MFD diet, a downregulation of *FASN* and *ACACA* as well as other lipogenic genes such as *SCD*, *LPL*, and *FABP3*, was observed using a bovine oligonucleotide microarray (Loor et al., 2005c). Similarly, in cows fed a low-forage diet supplemented with soybean and fish oils, Harvatine and Bauman (2006; study 5) observed a decrease in the yield of milk fat (38%) and short- and medium chain-FA (50%), together with a significant reduction in the expression of *FASN* and other genes involved in the regulation of lipid metabolism (*SREBP1*, *INSIG-1*, *S14*). However, in suckling beef cows, Murrieta et al. (2006) observed no effect from a diet supplemented with high-linoleate safflower seeds on mammary *ACACA* and *FASN* mRNA levels, despite a 33% decrease in the weight percentage of C10–C14 FA.

Elsewhere, food deprivation for 48 hours in dairy goats was shown to change mammary transcriptome profil, using a bovine 8,379-gene microarray (Ollier et al., 2007): in particular, six genes involved in lipid metabolism and transport such as *FASN*, *ACSBG1* (acyl-CoA synthetase “bubblegum”), and *AZGP1* (zinc-alpha-2-glycoprotein precursor) were downregulated simultaneously with a decrease in milk lactose, protein, and fat secretion. The regulation of these last two genes needs further studies to evaluate their impact on mammary lipogenesis.

In goats (studies 6 and 7), the observed slight decrease of 15–22% in milk C4–C16 FA secretion (g/day) after supplementation of hay or corn silage diets with vegetable lipids was not accompanied by any significant variation of *ACACA* and *FASN* mRNA levels or activities. However, a positive relationship ( $r = +0.62$ ) was observed (Fig. 3a) between *ACACA* mRNA variation of and C4–C16 secretion response to lipid supplementation, calculated from 81 individual values obtained by biopsies from goats fed hay or corn silage diets supplemented with 5.8% of either linseed or sunflower oil (study 7). Altogether, results from cow and goat studies demonstrated a positive relationship between both *ACACA* ( $r = +0.66$ ; Fig. 4a) and *FASN* ( $r = +0.73$ ; Fig. 4b) mRNA abundance and C4–C16 secretion responses to dietary treatment. These results suggest that variations of *ACACA* and *FASN* mRNA abundance play a role in the response of milk short- and medium-chain FA to dietary manipulation, especially the addition of lipids to the diet. Furthermore, the responses to lipid supplementation of *ACACA* and *FASN* mRNA were correlated ( $r = +0.56$ ; Fig. 3b) in study 7, suggesting the existence of a similar regulatory mechanism for these genes.


The differences observed between cows (Ahnadi et al., 2002; Piperova et al., 2000) and goats (Bernard et al., 2005c) in the magnitude of responses of *ACACA* (Fig. 4a) and *FASN* (Fig. 4b) mRNA and/or activity to dietary PUFA supplementation may be partly explained by factors linked to the diet, such as the level of starchy concentrate in the diet, the nature (fish oil vs. vegetable oil) and presentation (seeds vs. free oil) of the lipid supplements, as well as species differences. The fact that lipid supplementation in goats always induces an increase in milk fat content and secretion whereas in cows it generally decreases them (see earlier section) supports species-differences implications.

Altogether, from *in vivo* studies where *ACACA* and *FASN* mRNA levels and activities in the mammary gland were measured (studies 6 and 7 and study 2 for ACC in cows), positive relationships between mRNA and activities variations in response to dietary treatment were observed for *ACACA* and *FASN* (Bernard et al., 2006). This suggests a regulation at a transcriptional level, at least, for these two genes in the ruminant mammary gland, which is in accordance with data obtained in rat adipose tissue (Girard et al., 1997) and liver (Girard et al., 1994) in response to nutritional factors. For *ACACA*, this level of regulation occurs in addition to the well-documented posttranslational



**Fig. 3** Relationships between (a) milk short- and medium-chain fatty acid secretion and mammary acetyl-CoA carboxylase (*ACACA*) mRNA level responses to different dietary treatments (adapted from Bernard et al., 2006), and (b) mammary fatty acid synthase (*FASN*) and acetyl-CoA carboxylase (*ACACA*) mRNA-level responses to different dietary treatments. Results from 54 individual responses (lipid-supplemented = control) from 81 mammary biopsies in two  $3 \times 3$  Latin squares, in 27 goats (study 7; see text). Lipid supplements were linseed oil or sunflower oil

regulation, in particular through covalent modification via phosphorylation/dephosphorylation under hormonal control as well as by allosteric activation or inhibition by cellular metabolites (see earlier section).



**Fig. 4** Relationships between milk short- and medium-chain fatty acid secretion and mammary acetyl-CoA carboxylase (*ACACA*) (a) and fatty acid synthase (*FASN*) and (b) mRNA-level responses to different dietary treatments in goats and cows. Results are expressed as treatment mean response (vs. control group). Dietary treatments are described in the text and consisted of either lipid supplementation in goats (●; studies 6 and 7;  $n = 41$ ; Bernard et al., 2005c, unpublished) and cows (△; study 4; Delbecchi et al., 2001) or milk fat-depressed (MFD) diets (◇ studies 1–3 and study 5; Peterson et al., 2003; Piperova et al., 2000; Ahnadi et al., 2002; Harvatine & Bauman, 2006), and postprandial *trans*-10, *cis*-12 CLA infusion (○ 13.6 g/day and 10 g/day, respectively, in Baumgard et al., 2002, and Harvatine & Bauman, 2006) in cows

### Regulation of Stearoyl-CoA Desaturase

In rodents, nutritional regulation of SCD activity mainly occurs in the liver and has been studied extensively (Ntambi, 1999). Conversely, in the ruminant lactating mammary gland, only a few studies have investigated the nutritional regulation of *SCD* mRNA abundance and/or protein activity.

From the five nutritional studies performed in cows, four reported mammary *SCD* expression (studies 1, 3–5), and only feeding 1.7% of protected

fish oil supplement (rich in long-chain *n*-3 FA) (Ahnadi et al., 2002) significantly decreased ( $P < 0.05$ ) *SCD* mRNA abundance.


In goats fed hay-based diets supplemented with lipids, mammary *SCD* mRNA abundance decreased with formaldehyde-treated linseed and enzyme activity decreased with oleic sunflower oil, linseed oil, and sunflower oil, whereas lipid supplementation on *SCD* mRNA or activity had no effect on a corn silage diet (studies 6 and 7). Elsewhere, in late-lactating goats fed an alfalfa hay-based diet, mammary *SCD* mRNA was not affected by the dietary addition of 3.8% lipids from soybeans (Bernard et al., 2005b). Altogether, results from goats suggest an interaction between the basal diet and the dietary lipids used and predict a negative transcriptional or posttranscriptional regulation by dietary PUFA and/or by their ruminal biohydrogenation products may occur.

*In vivo*  $\Delta$ -9 desaturase activity has often been estimated by the milk ratios for the pairs of FA that represent a product/substrate relationship for *SCD* (*cis*-9 C14:1/C14:0, *cis*-9 C16:1/C16:0, *cis*-9 C18:1/C18:0, *cis*-9, *trans*-11 CLA/*trans*-11 C18:1; Bauman et al., 2001) due to the fact that the *in vitro* activity assay needs fresh materials, is laborious (Legrand et al., 1997), and is done in optimal conditions (pH, substrate, cofactors) that differ from *in vivo* conditions.

From the goat studies, we saw that the four FA pair ratios that represent a proxy for *SCD* activity were more or less related to the *SCD* activity itself, across 10 dietary groups (Bernard et al., 2005c; Chilliard et al., 2006b; Fig. 5). However, in terms of response to dietary lipids (six comparisons), the milk ratio of myristoleic acid to myristic acid (*cis*-9 C14:1/C14:0) gave the best estimation for the response of mammary *SCD* activity. This is due to the fact that almost all the myristoleic acid present in the milk is likely to be synthesized in the mammary gland by *SCD*. Indeed, myristic acid originates almost exclusively from *de novo* synthesis within the mammary gland (C14:0 is poorly represented in feedstuffs used for ruminants, including lipid supplements). Conversely, variable proportions of palmitic, palmitoleic, stearic, oleic, vaccenic, and ruminic acids come from absorption from the digestive tract and/or mobilization of body fat reserves. Then the ratios that involve these latter FA are less indicative of *SCD* activity. In addition, differential uptake, turnover, and use of the different FA of the four pair ratios by the mammary tissue itself may occur. Moreover, the four FA pair ratios could be influenced by other factors than *SCD* activity, such as a differential accuracy in the quantification of the *cis*-9 isomers in milk as well as, as stated above, by differences between *in vivo* (effective) and *in vitro* (potential) *SCD* activity.

## **Regulation of Acyltransferases**

Only one study in cows reported data on genes encoding acyltransferases with an observed reduction of mammary *GPAT* and *AGPAT* mRNA abundance



**Fig. 5** Relationships between milk FA desaturation ratios and stearoyl-CoA desaturase (SCD) activity in 43 goats fed hay-based diet supplemented, or not, with lipids. The lipid supplements were either 3.6% of lipids from oleic sunflower oil (OSO) or formaldehyde-treated linseed (FLS) (study 6; Bernard et al., 2005c), or 5.8% lipids from linseed oil (LO) or sunflower oil (SO) (study 7; Bernard et al., unpublished), or 6.5% whole rapeseed (RS) or 4.5% of sunflower oil (CSO) (Chilliard et al., 2006b). (a) Results are means of the six lipid-supplemented groups (black symbols) and the four control groups (white symbols). (b) Relationship between the responses (% of the control group) of SCD activity and milk *cis*-9 C14:1/C14:0 to the six lipid supplements

together with a reduction of milk fat yield (by 27%) with a high-concentrate diet (Peterson et al., 2003). In goats, food deprivation (FD) 48 hours before slaughter was shown to increase mammary *AGPAT* and *DGAT1* mRNA content, whereas milk fat secretion decreased (Ollier et al., 2006). This apparent discrepancy could be due to a known posttranscriptional regulation of these two genes. Indeed, phosphorylation/dephosphorylation mechanisms have been suggested for *AGPAT* activity by Mistry and Medrano (2002), whereas a translational control has been reported for *DGAT1* expression (Yu et al., 2002).

## Molecular Mechanisms Involved in the Regulation of Mammary Lipogenesis by Nutrition

### A Key Regulatory Role for Specific Fatty Acids

#### In Vivo Results in Nutritional Studies

From earlier *in vivo* studies with postruminal infusions of plant oils (e.g., Chilliard et al., 1991), a role for long-chain saturated FA and/or PUFA in decreasing


mammary *de novo* FA synthesis has been suggested. Nevertheless, only recently was a central role proposed for specific *trans*-FA as potent inhibitors of mammary lipid synthesis, from studies with specific dietary conditions inducing a dramatic MFD (Bauman & Griinari, 2003). In the same way, an impairment of mammary lipid synthesis has been observed in rats fed a diet containing a mixture of *trans*-isomers (Assumpcao et al., 2002).

Indeed, the so-called low-milk fat syndrome in cows seems to be mainly due to specific PUFA biohydrogenation products formed in the rumen. Diets that induce MFD belong to three groups:

1. Diets rich in readily digestible carbohydrates and poor in fibrous components, without the addition of lipid supplements (e.g., high-grain/low-forage diets; Peterson et al., 2003), but containing a minimal amount of PUFA in dietary feedstuffs (Griinari et al., 1998),
2. Low-fiber diets associated with supplemental PUFA of plant origin (Piperova et al., 2000),
3. Diets associated with dietary supplements of marine oils (fish oils, fish meals, oils from marine mammals and/or algae) that induce MFD regardless of the level of starch or fiber in the diet (Ahnadi et al., 2002; Chilliard et al., 2001).

In the past, a number of theories have been proposed to explain diet-induced MFD, with the starting point for all these theories being an alteration in ruminal fermentations (reviews by Bauman & Griinari, 2001, 2003). One of them is that rumen production of acetate and butyrate was too low to support milk fat synthesis. Another one is that ruminal production of propionate increased, enhancing the hepatic rate of gluconeogenesis and the levels of circulating glucose and insulin and adipose tissue lipogenesis, thus inducing a shortage of nutrients available for the mammary gland. Another theory that now prevails was first proposed by Davis and Brown (1970) and, as mentioned above, suggests that mammary fat synthesis is inhibited by specific *trans*-FA, which results from alterations in rumen PUFA biohydrogenation.

In cows, *in vivo* trials support the *trans*-FA theory because a wide range of MFD diets are accompanied by an increase in the *trans*-C18:1 percentage of milk fat (review by Bauman & Griinari, 2003). An important development of this theory was the discovery that MFD was associated with a specific increase in *trans*-10 C18:1 rather than total *trans*-C18:1 isomer (Griinari et al., 1998). This finding was confirmed in several studies in cows from which a curvilinear response curve between *trans*-10 C18:1 and fat yield responses (Fig. 6) was evidenced (Loor et al., 2005a). Elsewhere, a curvilinear relationship between the decrease in milk fat percentage and small increases in milk *trans*-10, *cis*-12 CLA (Bauman & Griinari, 2003) was also observed in some, but not all, studies. For example, the reduction of milk fat (27% decrease) and of C4–C16 (30% decrease) secretion observed in cows with a high-concentrate diet (study 1; Peterson et al., 2003) was accompanied by a small but significant increase in milk fat *trans*-10, *cis*-12 CLA secretion (+0.5 g/day).



**Fig. 6** Relationships between milk *trans*-10 C18:1 and fat yield responses in goats and cows. (a) Goat studies: data from 24 lipid-supplemented groups compared to 12 control groups (353 goats). The forages were either hay (■), fresh grass (◇), or corn silage (△). The lipids supplements were either sunflower oil, oleic sunflower oil, linseed oil, extruded linseeds, or rapeseeds (4–6% lipids in diet DM) and were given for 3 to 5 weeks (adapted from Chilliard et al., 2006a, b). (b) Cow studies: data from 31 lipid- or concentrate-supplemented groups, compared to control groups (13 studies) (adapted from Loor et al., 2005a)

The *trans*-10 C18:1 isomer synthesis in the rumen is probably due to the reduction of *trans*-10, *cis*-12 CLA resulting from an alternative pathway for biohydrogenation of linoleic acid, which increases when rumen pH decreases. Indeed, this “*trans*-10 pathway” seems to increase with diets rich in concentrate and/or corn silage and to occur after a one- to two-week period of latency after the start of dietary PUFA supplementation, following an earlier but transient increase in the “*trans*-11 pathway” (Chilliard & Ferlay, 2004; Roy et al., 2006a; Shingfield et al., 2006).

We must emphasize that in goats as in cows, a negative and curvilinear relationship between the responses of milk *trans*-10 C18:1 percentage and milk fat yield is observed (Chilliard et al., 2006a; Fig. 6), despite the fact that the fat yield response was always positive in goats, but always negative or null in cows. Then, in goats, the highest increases in *trans*-10 C18:1 were observed with either corn silage or fresh grass diets (Fig. 6) and matched with

the lowest increases in milk fat yield. In addition, the maximum observed value of milk *trans*-10 C18:1 in goats was much lower than in cows, and no significant increases in goat milk *trans*-10, *cis*-12 CLA occurred in the 24 diet comparisons in Fig. 6. These observed similarities and differences among ruminants suggest a species specificity of FA ruminal and/or mammary metabolism. The positive effect of lipid supplementation on goat milk fat yield could be due in part to the mammary sensitivity to increased availability of stearic acid arising from dietary PUFA biohydrogenation in the rumen (Bernard et al., 2006).

Regarding the impact of PUFA on lipogenic gene expression and, in particular, on SCD, still little is known in ruminants. Conversely, in rodents the downregulation of *SCD* gene expression by (*n*-6) and (*n*-3) PUFA has been largely described in liver and AT (Ntambi, 1999), whereas little is known in the lactating mammary gland (Lin et al., 2004; Singh et al., 2004). Indeed, Singh et al. (2004) observed negative effects from olive oil or safflower oil fed to lactating mice on both *SCD* mRNA and activity in the mammary gland and *SCD* mRNA in the liver. Similarly, in the mouse mammary gland, Lin et al. (2004) reported a decrease in *SCD*, *ACACA*, and *FASN* mRNA abundance and SCD activity by the dietary addition of either *trans*-11 C18:1, *cis*-9, *trans*-11 CLA, or *trans*-10, *cis*-12 CLA, whereas these treatments had no effect on these transcripts in liver. These results on lactating mice indicate a tissue-specific regulation of lipogenic gene expression by *trans*-FA and outline the possibility to manipulate mammary *SCD* gene expression by nutrition.

In goats, the observed negative effect of the addition of oleic sunflower oil, sunflower oil, and linseed oil to hay diet (trials 6 and 7) on SCD activity (Fig. 5b) and that of formaldehyde-treated linseed on *SCD* mRNA might be partly attributed to dietary *cis*-9 C18:1, C18:2 (*n*-6), and/or C18:3 (*n*-3) escaping from the rumen and/or to *trans*-isomers formed during ruminal metabolism of these three FA (Chilliard et al., 2003c; Ferlay et al., 2003; Rouel et al., 2004). In addition, besides PUFA biohydrogenation processes, oleic acid could be isomerized in several *trans*-C18:1 isomers, including *trans*-10, as shown in microbial cultures from bovine rumen (Mosley et al., 2002). This finding is in agreement with the observed increase of several *trans*-C18:1 isomers in milk from goats fed oleic sunflower oil (Bernard et al., 2005c; Ferlay et al., 2003; Chilliard et al., 2006a). Nevertheless, data in Fig. 5b suggest that  $\alpha$ -linolenic acid from formaldehyde-treated linseeds should be more efficient than dietary oleic acid-rich oil to decrease SCD activity.

### Duodenal or Intravenous Infusion of Specific Fatty Acids

Duodenal infusion trials of pure-CLA isomers demonstrated that *trans*-10, *cis*-12 CLA inhibits milk fat synthesis in dairy cows, whereas the *cis*-9, *trans*-11 CLA isomer has no effect (Baumgard et al., 2000; Loor & Herbein, 2003). The severe reduction (48%) in milk fat yield due to the infusion of a high

dose (13.6 g/day) of *trans*-10, *cis*-12 CLA (Baumgard et al., 2000) was accompanied by a dramatic reduction (> 35%) of mRNA abundance of enzymes involved in mammary uptake and intracellular trafficking of FA (*LPL* and *FABP*), *de novo* FA synthesis (*ACACA* and *FASN*), desaturation (*SCD*), and esterification (*GPAT* and *AGPAT*). Similarly, intravenous administration of *trans*-10, *cis*-12 CLA, either from 2 to up to 6 g/day (Viswanadha et al., 2003) or 10 g/day (Harvatine & Bauman, 2006), depressed milk fat yield, with, for the latter study, a joint decrease in the expression of genes involved in mammary uptake (*LPL*), *de novo* FA synthesis (*FASN*), and the regulation of lipid metabolism (*SREBP1*, *S14*, *INSIG-1*).

However, in the absence of duodenal infusion, the levels of *trans*-10, *cis*-12 CLA in the rumen, duodenal fluid, or milk always remained very low compared to the levels used in infusion studies (see above) or reached by the *trans*-10 C18:1 isomer (ratio between *trans*-10, *cis*-12 CLA, and *trans*-10 C18:1 of ~0.01; Bauman & Griinari, 2003; Loor et al., 2004a, b, 2005a, d; Piperova et al., 2000). In addition, in cows fed with marine oil for which an MFD was observed, little or only traces of milk *trans*-10, *cis*-12 CLA were detected, whereas substantial increases in *trans*-10 C18:1 were observed (Loor et al., 2005b; Offer et al., 2001). Furthermore, species-specific responses to *trans*-10, *cis*-12 CLA duodenal infusion have been noted, with no effect on milk fat secretion in goats (Andrade & Schmidely, 2005), contrary to what is observed in cows (see above), whereas in lactating sheep, lipid-encapsulated CLA supplement containing *trans*-10, *cis*-12 CLA significantly reduced milk fat synthesis (Lock et al., 2006).

Due to the lack of pure material, it was not possible until recently to investigate a direct effect of *trans*-10 C18:1 on milk fat synthesis, whereas the potent inhibitory effect of *trans*-10, *cis*-12 CLA was clearly established by postruminal infusion trials in dairy cows (Bauman & Griinari, 2003). However, a recent study (Lock et al., 2007) using chemically synthesized *trans*-10 C18:1 infused postruminally over 4 days at 42.6 g/day/cow showed that despite the fact that this isomer was absorbed, taken up by the mammary gland, and transferred to milk fat, it had no effect on milk fat synthesis. As suggested by the authors (Lock et al., 2007), it is likely that the formation of *trans*-10 C18:1 and of *trans*-10, *cis*-12-CLA due to alterations in the rumen environment is accompanied by the formation of other biohydrogenation intermediates that could as well inhibit (or co-inhibit with *trans*-10, *cis*-12-CLA) milk fat synthesis. Thus, from nutritional studies, several other rumen-derived FA were proposed recently as potential inhibitors of cow milk fat synthesis due to high negative correlations between their milk fat concentrations and milk fat content and secretion. These proposed FA were several *cis*- or *trans*-C18:1 isomers, and *cis*-9, *trans*-13 C18:2, *cis*-9, *trans*-12 C18:2, *trans*-11, *cis*-13 CLA, *trans*-11, *cis*-15 C18:2 (Loor et al., 2005a), and *trans*-9, *cis*-11 CLA (Roy et al., 2006a; Shingfield et al., 2005, 2006). To confirm the potential role of these FA as well as of other minor CLA isomers found in milk, when available, few of them were postruminally infused. Whereas *trans*-8, *cis*-10 CLA, *cis*-11, *trans*-13 CLA, and *trans*-10, *trans*-12 CLA did not

inhibit milk fat yield (Perfield et al., 2004, 2006; Sæbo et al., 2005), it was demonstrated that *cis*-10, *trans*-12 CLA (Sæbo et al., 2005) and *trans*-9, *cis*-11 CLA (Perfield et al., 2005) did reduce milk fat synthesis. However, the efficiency of the *trans*-9, *cis*-11 CLA was much lower than that of *trans*-10, *cis*-12 CLA (Perfield et al., 2005). This last result is in agreement with feeding trials on cows with or without MFD, with a greater slope of the equation between milk fat yield and *trans*-10, *cis*-12 CLA than *trans*-9, *cis*-11 CLA (Roy et al., 2006a).

Again, some of these isomers (*trans*-7, *cis*-9 CLA, *trans*-11, *cis*-13 CLA, *cis*-9, *trans*-13 C18:2, *trans*-11, *cis*-15 C18:2, *cis*-9, *trans*-13 C18:2) increased in goats supplemented with dietary PUFA, whereas no MFD was observed (Chilliard & Ferlay, 2004; Chilliard et al., 2003c; Ferlay et al., 2003).

### **In Vitro Studies on the Effect of Fatty Acids on Lipogenesis in Mammary Epithelial Cells**

Studies several years ago on dispersed bovine MEC (Hansen et al., 1986; Hansen & Knudsen, 1987) demonstrated that the addition of C18:0, *cis*-9 C18:1, or C18:2 (*n*-6) inhibited *de novo* synthesis of FA with 16 carbons or less, except C4. More recently, looking at the effects of specific FA on the bovine mammary cell line (MAC-T cells), Jayan and Herbein (2000) showed that, compared to stearic acid, *trans*-11 C18:1 and, to a lesser extent, *cis*-9 C18:1 reduced ACC and FAS activities.

Furthermore, 35 years ago, Bickerstaffe and Annison (1970) observed negative effects of oleic, linoleic, and linolenic acids on goat mammary SCD activity measured *in vitro*, which have been partly confirmed by our *in vivo* studies on goats (see earlier section). The addition of *trans*-11 C18:1 increased SCD mRNA abundance in bovine MEC (Matitashvili & Bauman, 2000) and SCD activity in the bovine MAC-T cell line (Jayan & Herbein, 2000). Recently, Keating et al. (2006) demonstrated that treatment of bovine MAC-T cells with *trans*-10, *cis*-12 CLA (and, to a lesser extent, *cis*-9, *trans*-11 CLA) caused a significant reduction in SCD transcriptional activity, with this effect mediated through the stearoyl-CoA desaturase transcriptional enhancer element region (STE; see earlier section). The same study also showed that bovine SCD promoter was upregulated by insulin and downregulated by oleic acid, whereas linoleic, linolenic, stearic, and vaccenic acids had no effect. However, the *in vitro* effects of other specific *trans*-C18:1 and C18:2 isomers on mammary SCD gene expression are still unknown.

Elsewhere, in bovine mammary epithelial cell (BME-UV) cultures, the addition of *trans*-10, *cis*-12 CLA inhibited the stimulatory effect of prolactin on the cytosolic NADP<sup>+</sup>-dependent isocitrate dehydrogenase (*IDH1*) gene expression, involved in the generation of NADPH required for *de novo* fatty acid synthesis, whereas the *cis*-9, *trans*-11 CLA isomer had no effect (Liu et al., 2006).

Further research is necessary in ruminants to identify the more important inhibitors of fat synthesis either *in vivo* (i.e., postruminal infusion of exogenous

FA) or *in vitro*, which is hampered by the lack of pure *trans*-C18:1 and C18:2 isomers and by the difficulty in obtaining an *in vitro* functional model for lipid synthesis and secretion (Barber et al., 1997).

### ***Signaling Pathways Mediating Nutritional Regulation of Gene Expression***

Whereas the signaling mechanisms involved in the regulation of lipogenic gene expression in rodent liver and adipose tissue have been comprehensively described (Clarke, 2001), little is known about these mechanisms in ruminants, particularly in the mammary gland. However, it was suggested that several genes involved in milk fat synthesis in the bovine mammary gland may share a common regulatory mechanism because of their coordinated downregulation observed in response to a postruminal infusion of *trans*-10, *cis*-12 CLA (Baumgard et al., 2002) and to an MFD diet (Peterson et al., 2003). Clarke (2001) reviewed rodent data and proposed that PUFA control the main metabolic pathways of lipid metabolism by governing the DNA-binding activity and nuclear abundance of selected transcription factors regulating the expression of key genes. The major transcription factors involved are sterol regulatory binding protein-1 (SREBP-1) and peroxisome proliferator-activated receptors (PPARs), with the FA or cholesterol acting by binding to the nuclear receptors PPAR (and LXR, HNF-4 $\alpha$ ), whereas the FA induce changes in the nuclear abundance of SREBP. As only few data are available in ruminants on these transcription factors, the following sections describe the state of knowledge on transcription factors mediating nutritional regulation of gene expression in rodents as well.

#### **SREBP-1**

SREBPs are basic-helix-loop-helix-leucine zipper (bHLH-LZ) transcription factors that belong to a family of transcription factors that regulate lipid homeostasis by controlling the expression of several enzymes required for endogenous cholesterol, FA, triacylglycerol, and phospholipid synthesis. Three SREBP forms have been characterized in rodents—SREBP-1a, -1c, and -2—differing in their roles in lipid synthesis. The SREBP-1a form is mainly expressed in cultured cells and tissues with a high cell proliferation capacity. The SREBP-1c form is expressed in many organs (mainly adipose tissue, brain, muscle, etc.) (Shimomura et al., 1997). These two forms derive from a single gene through the use of alternate promoters that give rise to different first exons (Brown & Goldstein, 1997). SREBP-2 derives from a different gene and is involved in the transcription of cholesterologenic enzymes.

SREBPs are synthesized as ~1,150 amino acid inactive precursors bound to the membrane of the endoplasmic reticulum (ER) through a tight association with SREBP cleavage activating protein (SCAP) (Miller et al., 2001). Upon sterol deprivation, the SREBP-SCAP complex moves to the Golgi apparatus, where two functionally active distinct proteases, site-1 and site-2 proteases (S1P and S2P), sequentially cleave the precursor protein SREBP to release the NH<sub>2</sub>-terminal active domain (Sakai et al., 1996). It has been shown that this sterol-dependent trafficking requires an intact sterol-sensing domain located in the SCAP protein (Brown & Goldstein, 1997), demonstrating the dual role of SCAP as escorter and sensor. Recently, the insulin-induced gene (*INSIG-1*) protein that binds SCAP and thus facilitates retention of the SCAP/SREBP complex in the ER was identified (Yang et al., 2002). Upon appropriate conditions (low sterol concentrations or possibly insulin action; Eberlé et al., 2004), the interaction between INSIG and SCAP decreases and allows the SCAP to escort SREBPs to the Golgi apparatus for the cleavage activation process.

Finally, the mature SREBP, i.e., the NH<sub>2</sub>-terminal portion (domain containing the bHLH-LZ), is translocated to the nucleus, where it binds (as a homodimeric form) its target genes on sterol binding elements or on palindromic sequences called E-boxes within their promoter regions (Wang et al., 1993). These target genes are implicated in cholesterol, FA, and lipid synthesis, including LPL, ACC, FAS, and SCD (Shimomura et al., 1998).

In rodents, FA downregulate the nuclear abundance of SREBP-1 by two described mechanisms: either an inhibition of the proteolytic activation process of SREBP-1 protein (under cholesterol dependence) or an inhibition of the *SREBP-1* gene transcription. Recently, in the ovine lactating mammary gland, Barber et al. (2003) identified SREBP-1 as a major regulator of *de novo* lipid synthesis through the activation of ACC $\alpha$  PIII, achieved together with NF-Y, USF-1, and USF-2 transcription factors. In addition, SREBP-1 binding motifs were also identified in the proximal promoter of ACC $\alpha$  PII, which is upregulated during lactation, indicating that SREBP-1 could play an important role in the joint regulation of PII and PIII in mammary tissue (Barber et al., 2003). Elsewhere it was shown that the addition of *trans*-10, *cis*-12 CLA to the bovine MEC line (MAC-T) had no effect on *SREBP-1* mRNA or SREBP-1 precursor protein content but reduced the abundance of the activated nuclear fragment of the protein (Peterson et al., 2004). This was accompanied by a reduction in transcriptional activation of the lipogenic genes *ACACA*, *FASN*, and *SCD*. These findings suggest that the inhibitory effect of this CLA isomer on lipid synthesis could be due to an inhibition of the proteolytic activation of SREBP1. Similarly, the use of microarray tools to characterize mammary gene profiling in cows fed an MFD diet (composed of 70% forage, 25% concentrate, and 5% soybean oil) showed a downregulation of several genes associated with fatty acid metabolism (see previous section) and of eight transcription factors without modification of *SREBP1* gene expression (Loor et al., 2005c). In addition, Harvatine and Bauman (2006) demonstrated the existence of a joint decrease in the mammary expression of *SREBP1*, *Spot 14* (*S14*), *INSIG-1*, *FASN*, and *LPL*.

that could explain MFD in lactating cows either fed a low-forage/high-oil diet or infused with *trans*-10, *cis*-12 CLA. Again, the decreased expression of SREBP-1 and proteins associated with SREBP-1 activation, together with SREBP-1 responsive lipogenic enzymes, provides strong support for the central role of SREBP-1 in the regulation of milk fat synthesis. Furthermore, this study outlined a possible involvement of S14 in the regulation of FA synthesis in the bovine mammary gland, as shown in rodent liver and adipose tissue (Cunningham et al., 1998).

## PPARs

PPARs belong to a superfamily of hormone receptors with, as for all of the members of this family, a DNA-binding domain, a gene-activating domain, and a ligand-binding domain. They regulate the transcription of genes involved in different lipid metabolism pathways including the transport of plasma triglycerides, the cellular FA uptake, and the peroxisomal and mitochondrial  $\beta$ -oxidation (Schoonjans et al., 1996a). The activating ligands of PPARs are peroxisome proliferators, including chemical molecules as fibrates, thiazolidinedione as well as molecules as FA, including PUFA, and their metabolites. PPARs heterodimerize with the *cis*-9 retinoic acid receptor (RXR) to bind to specific response elements located in the promoter region of the target genes.

Three PPAR subtypes have been identified: PPAR $\alpha$ , expressed mainly in liver as well as in heart, kidney, intestinal mucosa, and brown adipose tissue, involved in FA transport and  $\beta$ - and  $\omega$ -oxidation; PPAR $\beta$  abundantly and ubiquitously expressed but mainly found in heart, lung, and kidney; PPAR $\gamma$ , most abundant in adipose tissue, stimulating adipocyte differentiation and lipogenesis of the mature adipocyte (Schoonjans et al., 1996b). PPAR $\gamma$  is also expressed in a number of epithelial tissues (breast, prostate, and colon), in which it seems to favor less malignant phenotype cells in human cancer (Sarraf et al., 1999). The *PPAR* $\gamma$  gene generates two transcripts, designated *PPAR* $\gamma$ 1 and  $\gamma$ 2, resulting from differential mRNA splicing and promoter usage (Yeldandi et al., 2000), and leading to two protein isoforms, with PPAR $\gamma$ 2 having 30 additional amino acid residues at the N terminal extremity. Among the activating compounds of the *PPAR* genes, (*n*-3) and (*n*-6) PUFA and mainly their metabolites (eicosanoids and oxidized FA) are the major natural activators of PPAR $\alpha$  (Clarke, 2001), while 15-deoxy- $\Delta^{12,14}$ -prostaglandin J2 is the activator of the PPAR $\gamma$  subtype (Rosen & Spiegelman, 2001). In addition, *in vitro* studies on mature adipocytes revealed that *trans*-10, *cis*-12 CLA downregulates *PPAR* $\gamma$  gene expression, that could be a mechanism by which this CLA isomer prevents lipid accumulation in adipocytes (Granlund et al., 2003).

The few data available in bovine show similarities with those from rodent species. Thus, in bovine subcutaneous adipose tissue, the observed joint upregulation of PPAR $\gamma$  and FAS and *ACACA* and *LPL* gene expression by

propionate infusion (Lee & Hossner, 2002) suggests an implication of PPAR $\gamma$  in the nutritional or insulin activation of lipogenesis. Elsewhere, bovine *PPAR $\gamma$ 1* and *PPAR $\gamma$ 2* cDNAs have been characterized (Sundvold et al., 1997) with expression of the two isoforms in adipose tissue, whereas only PPAR $\gamma$ 2 was expressed in the mammary gland. Recently, in primary cultured bovine MEC, the expression of PPAR $\gamma$ 2 in response to the addition of acetate and octanoate was increased while ACC activity decreased (Yonezawa et al., 2004a), conversely to previous observations in adipose tissue (Lee & Hossner, 2002). Elsewhere, an upregulation of mammary *PPAR $\gamma$*  gene expression was shown in dairy cows between -14 and +14 days relative to parturition, using a bovine cDNA microarray (Loor et al., 2004c).

### Other Transcription Factors

The molecular mechanisms that control milk protein and lipogenic gene expression are not fully understood and probably involve undiscovered proteins within the mammary gland. Thus, in the nuclear extract of bovine mammary gland, the transcription factors Sp1 and NF-1 were identified (Wheeler et al., 1997), which are already known in rodents to act in conjunction with other proteins such as SREBP1, as well as six other proteins whose abundance was positively related with lactation or pregnancy status. Four of these proteins were identified as lactoferrin, annexin II, vimentin, and heavy-chain immunoglobulin. The presence of lactoferrin in the nuclear extracts is consistent with a study demonstrating that lactoferrin binds to DNA in a sequence-specific manner and activates transcription (He & Furmanski, 1995). Nevertheless, the function of lactoferrin as a transcription factor has not yet been confirmed.

Elsewhere, over the past few years, response elements to lactogenic hormones have been mapped within the promoters of milk protein genes; in some cases, the proteins that mediate the lactational signals are known. Signal transducer(s) and activator(s) of transcription (STAT) form a family of cytoplasmic proteins that are activated in response to a large number of cytokines, growth factors, and hormones (Hennighausen, 1997). The STAT proteins are activated via a cascade of phosphorylation events in which Janus protein tyrosine kinases (Jak2) are first phosphorylated. Then the activated Jak2 phosphorylate STAT proteins. In turn, STAT detach from the receptor complex, form homo- or heterodimers, and translocate from the cytoplasm to the nucleus, where they interact with specific promoter regions and regulate gene expression (Hennighausen, 1997). Until now, seven bovine *STAT* genes have been identified, *STAT1-6*, *5a*, and *5b*, the latter two of which have already been sequenced (Seyfert et al., 2000). STAT5 was originally identified as a "mammary gland factor" mediating the prolactin signal to establish galactopoiesis (Rosen et al., 1999). Recently, Mao et al. (2002) demonstrated that STAT5 binding (at position -797) contributes to the lactational stimulation of

promoter III (PIII) of the *ACACA* gene in the mammary gland. Hence, prolactin, acting through STAT5, contributes to the activation of ACC expression in milk-producing cells. Similarly, in BME-UV bovine mammary epithelial cell, Liu et al. (2006) reported that prolactin enhances *IDH1* mRNA and protein expression, but the molecular mechanisms of this regulation were not investigated. Elsewhere, in bovine mammary gland explant culture, Yang et al. (2000a) showed a rapid stimulation of STAT5 DNA binding activity by prolactin, growth hormone, and IGF1. In addition, the same authors demonstrated *in vivo* that STAT5 protein level and DNA binding activity are modulated by several physiological signals, including GH infusion and milking frequency (Yang et al., 2000b). Altogether these findings suggest that STAT5 might be important in determining the milk composition by coordinating FA and protein synthesis during lactation and that STAT5 transcription factor may represent part of the common route by which different extracellular signals (linked to hormonal status as well as to milking frequency) could converge and be transduced intracellularly to coordinate cell functions in the mammary gland. Recently, a study reported an association between *STAT1* variants and milk fat and protein yield and percentages in Holstein dairy cattle, implicating the *STAT1* gene in the regulation of milk protein and fat synthesis (Cobanoglu et al., 2006).

Despite the recent increased knowledge in ruminants on the characterization of transcription factors in the mammary gland, many questions still remain unanswered, in particular the role of STAT and PPARs in the regulation of lipid metabolism.

## Conclusions

Over the last several years, the biochemical pathways of lipid synthesis in the mammary gland have been elucidated, and many of the enzymatic proteins and their cDNAs have been characterized. This has allowed the development of studies on the nutritional regulation of a few “candidate”, genes involved in mammary FA uptake (*LPL*), *de novo* synthesis (*ACACA* and *FASN*), and desaturation (*SCD*). These studies showed that the responses of mammary “candidate” gene expression to nutritional factors do not always match milk FA secretion responses. In goats and cows, data suggest that the availability of substrates rather than the LPL activity is the limiting factor in the uptake of long-chain FA, except with extreme MFD diets fed to cows, in which both mammary LPL mRNA and activity decreased. In cows and goats, data converged to demonstrate that *ACACA* and *FASN* gene expressions are key factors of short- and medium-chain FA synthesis, even though they are not always repressed by the addition of PUFA to the diet, in goats at least. In this species, *ACACA* and *FASN* gene expressions are regulated by dietary factors at a transcriptional level at least, and *SCD* is regulated at a transcriptional and/or

posttranscriptional level, depending on the lipid supplements. Conversely, in cows, the level of *SCD* mRNA varied little with the nutritional factors studied so far, except for a decrease when “protected” fish oil was fed. A fine balance between the exogenous unsaturated FA and the *SCD* desaturation products must be maintained within the mammary gland in order to preserve the fluidity of cellular membranes and milk fat (Chilliard et al., 2000; Parodi, 1982). In addition to its role on milk nutritional quality via the synthesis in *cis*-9, *trans*-11 CLA and *cis*-9-monounsaturated fatty acids, the impact of *SCD* on membrane fluidity underlines the importance of this enzyme.

The regulatory systems governing the nutritional response of mammary gene expression, in particular the intracellular signaling systems involved in these regulations, need to be further investigated in the future. The basis of the effects of nutrients and particularly the identification of specific *trans*-FA controlling lipogenic gene expression are obvious targets. Milk *trans*-10, *cis*-12 CLA is sometimes correlated with milk fat depression in cows (but not in goats) and, when infused postruminally at high doses, acts as a potent inhibitor of the expression of all the lipogenic genes. Conversely, *trans*-10 C18:1 does not directly control milk fat synthesis in dairy cows, although it largely responds to dietary factors, with its concentration being negatively related to milk fat response in cows and, to a lesser extent, in goats. Nevertheless, marked differences are observed between the milk fat yield responses of these two species, with few differences in milk FA profile responses, in particular lower increases in *trans*-10 C18:1 in goats. Elsewhere, more information on the promoter of the lipogenic genes should be acquired, which would help to clarify the roles and mechanisms of the action of PUFA and/or *trans*-FA, in order to better understand the molecular mechanisms involved in dietary- and/or species-related responses. Few data on transcription factors are available, and a central role for SREBP-1 has been outlined as mediator of FA effects, and STAT5 for hormonal and physiological effects at least, whereas the roles of PPARs need to be determined. It is expected that the development of *in vitro* functional systems for lipid synthesis and secretion would allow future progress in the identification of the inhibitors and activators of fat synthesis and in understanding differences between ruminant species.

This chapter reviewed studies focusing on the nutritional regulation of the expression of a few candidate genes controlling lipid synthesis. Nevertheless, the expression of specific milk fat globule membrane proteins (Mather, 2000) such as butyrophilin, xanthine oxidoreductase, and CD36, which intervene in milk lipid secretion, is also likely to have consequences on milk fat yield and composition (Ogg et al., 2004). The recent development of tools for studying the mammary transcriptome (macro- and microarrays; e.g., Bernard et al., 2005a; Leroux et al., 2003; Looor et al., 2004c, 2005c; Suchyta et al., 2003) and proteome (Daniels et al., 2006) will allow us to study the effect of nutritional changes on the expression (mRNA and protein) of a large number of genes putatively involved in mammary gland function, including lipid synthesis and secretion, in relationship to milk FA profile (Ollier et al., 2007). Such tools will allow us to

identify new “candidate” genes or proteins whose expression is regulated by nutrition, and to understand their regulation pathways. In addition, further investigations on ruminal digestion and body metabolism of nutrients (absorption, partitioning between tissues) and mammary metabolic flows will also contribute to highlight the mechanisms underlying the *in vivo* responses to dietary factors and the differences among ruminant species.

**Acknowledgment** The authors thank S. Ollier and M. Bonnet for helpful discussions during the preparation of the manuscript and P. Béraud for secretarial assistance.

## References

- Abu-Elheiga, L., Almarza-Ortega, D. B., Baldini, A., & Wakil, S. J. (1997). Human acetyl-CoA carboxylase 2. Molecular cloning, characterization, chromosomal mapping, and evidence for two isoforms. *Journal of Biological Chemistry*, 272, 10669–10677.
- Abu-Elheiga, L., Brinkley, W. R., Zhong, L., Chirala, S. S., Woldegiorgis, G., & Wakil, S. J. (2000). The subcellular localization of acetyl-CoA carboxylase 2. *Proceedings of the National Academy of Sciences USA*, 97, 1444–1449.
- Abumrad, N. A., el-Maghrabi, M. R., Amri, E. Z., Lopez, E., & Grimaldi, P. A. (1993). Cloning of a rat adipocyte membrane protein implicated in binding or transport of long-chain fatty acids that is induced during preadipocyte differentiation. Homology with human CD36. *Journal of Biological Chemistry*, 268, 17665–17668.
- Ahnadi, C. E., Beswick, N., Delbecchi, L., Kennelly, J. J., & Lacasse, P. (2002). Addition of fish oil to diets for dairy cows. II. Effects on milk fat and gene expression of mammary lipogenic enzymes. *Journal of Dairy Research*, 69, 521–531.
- Andrade, P. V. D., & Schmidely, P. (2005). Effect of duodenal infusion of *trans*10, *cis*12-CLA on milk performance and milk fatty acid profile in dairy goats fed high or low concentrate diet in combination with rolled canola seed. *Reproduction Nutrition Development*, 46, 31–48.
- Angiolillo, A., Amills, M., Urrutia, B., Domenech, A., Sastre, Y., Badaoui, B., & Jordana, J. (2006). Identification of a single nucleotide polymorphism at intron 16 of the caprine acyl-coenzyme A: diacylglycerol acyltransferase 1 (DGAT1) gene. *Journal of Dairy Research*, in press.
- Annisson, E. F., Linzell, J. L., & West, C. E. (1968). Mammary and whole animal metabolism of glucose and fatty acids in fasting lactating goats. *Journal of Physiology*, 197, 445–459.
- Aoki, N., Ishii, T., Ohira, S., Yamaguchi, Y., Negi, M., Adachi, T., Nakamura, R., & Matsuda, T. (1997). Stage specific expression of milk fat globule membrane glycoproteins in mouse mammary gland: Comparison of MFG-E8, butyrophilin, and CD36 with a major milk protein, beta-casein. *Biochimica et Biophysica Acta*, 1334, 182–190.
- Assumpcao, R. P., Santos, F. D., Setta, C. L., Barreto, G. F., Matta, I. E., Estadella, D., Azeredo, V. B., & Tavares do Carmo, M. G. (2002). *Trans* fatty acids in maternal diet may impair lipid biosynthesis in mammary gland of lactating rats. *Annals of Nutrition and Metabolism*, 46, 169–175.
- Barber, M. C., & Travers, M. T. (1995). Cloning and characterisation of multiple acetyl-CoA carboxylase transcripts in ovine adipose tissue. *Gene*, 154, 271–275.
- Barber, M. C., & Travers, M. T. (1998). Elucidation of a promoter activity that directs the expression of acetyl-CoA carboxylase alpha with an alternative N-terminus in a tissue-restricted fashion. *Biochemical Journal*, 333 (Pt 1), 17–25.
- Barber, M. C., Clegg, R. A., Travers, M. T., & Vernon, R. G. (1997). Lipid metabolism in the lactating mammary gland. *Biochimica et Biophysica Acta*, 1347, 101–126.

- Barber, M. C., Vallance, A. J., Kennedy, H. T., & Travers, M. T. (2003). Induction of transcripts derived from promoter III of the acetyl-CoA carboxylase- $\alpha$  gene in mammary gland is associated with recruitment of SREBP-1 to a region of the proximal promoter defined by a DNase I hypersensitive site. *Biochemical Journal*, 375, 489–501.
- Barber, M. C., Price, N. T., & Travers, M. T. (2005). Structure and regulation of acetyl-CoA carboxylase genes of metazoa. *Biochimica et Biophysica Acta*, 1733, 1–28.
- Barillet, F., Arranz, J. J., & Carta, A. (2005). Mapping quantitative trait loci for milk production and genetic polymorphisms of milk proteins in dairy sheep. *Genetics Selection Evolution*, 37 (Suppl 1), S109–123.
- Bauman, D. E., & Griinari, J. M. (2001). Regulation and nutritional manipulation of milk fat: Low-fat milk syndrome. *Livestock Production Science*, 70, 15–29.
- Bauman, D. E., & Griinari, J. M. (2003). Nutritional regulation of milk fat synthesis. *Annual Review of Nutrition*, 23, 203–227.
- Bauman D. E., Corl, B. A., Baumgard, L. H., & Griinari, J. M. (2001). Conjugated linoleic acid (CLA) and the dairy cow. In J. Wisman and P. C. Garnsworthy (Eds.), *Recent Advances in Animal Nutrition* (pp. 221–250). Nottingham: Nottingham University Press.
- Baumgard, L. H., Corl, B. A., Dwyer, D. A., Sæbo, A., & Bauman, D. E. (2000). Identification of the conjugated linoleic acid isomer that inhibits milk fat synthesis. *American Journal of Physiology—Regulatory, Integrative and Comparative Physiology*, 278, R179–184.
- Baumgard, L. H., Corl, B. A., Dwyer, D. A., & Bauman, D. E. (2002). Effects of conjugated linoleic acids (CLA) on tissue response to homeostatic signals and plasma variables associated with lipid metabolism in lactating dairy cows. *Journal of Animal Science*, 80, 1285–1293.
- Bernard, L., Leroux, C., Hayes, H., Gautier, M., Chilliard, Y., & Martin, P. (2001). Characterization of the caprine stearoyl-CoA desaturase gene and its mRNA showing an unusually long 3'-UTR sequence arising from a single exon. *Gene*, 281, 53–61.
- Bernard, C., Degrelle, S., Ollier, S., Campion, E., Cassar-Malek, I., Charpigny, G., Dhorne-Pollet, S., Hue, I., Hocquette, J. F., Le Provost, F., Leroux, C., Piump, F., Rolland, G., Uzbekova, S., Zalachas, E., & Martin, P. (2005a). A cDNA macro-array resource for gene expression profiling in ruminant tissues involved in reproduction and production (milk and beef) traits. *Journal of Physiology and Pharmacology*, 56 (Suppl 3), 215–224.
- Bernard, L., Leroux, C., Bonnet, M., Rouel, J., Martin P., & Chilliard Y. (2005b). Expression and nutritional regulation of lipogenic genes in mammary gland and adipose tissues of lactating goats. *Journal of Dairy Research*, 72, 250–255.
- Bernard, L., Rouel, J., Leroux, C., Ferlay, A., Faulconnier, Y., Legrand P., & Chilliard, Y. (2005c). Mammary lipid metabolism and fatty acid secretion in Alpine goats fed vegetable lipids. *Journal of Dairy Science*, 88, 1478–1489.
- Bernard, L., Leroux, C., & Chilliard, Y. (2006). Characterisation and nutritional regulation of the main lipogenic genes in the ruminant lactating mammary gland. In K. Sejrsen, T. Hvelplund, M. O. Nielsen (Eds.), *Ruminant Physiology: Digestion, metabolism and impact of nutrition on gene expression, immunology and stress* (pp. 295–362). Wageningen, The Netherlands: Wageningen Academic Publishers.
- Beswick, N. S., & Kennelly, J. J. (1998). The influence of bovine growth hormone and growth hormone releasing factor on acetyl-CoA carboxylase and fatty acid synthase in primiparous Holstein cows. *Comparative Biochemistry and Physiology. Part C, Pharmacology, Toxicology and Endocrinology*, 120, 241–249.
- Bickerstaffe, R., & Anison, E. F. (1970). Lipid metabolism in the perfused chicken liver. The uptake and metabolism of oleic acid, elaidic acid, *cis*-vaccenic acid, *trans*-vaccenic acid and stearic acid. *Biochemical Journal*, 118, 433–442.
- Bonnet, M., Faulconnier, Y., Fléchet, J., Hocquette, J. F., Leroux, C., Langin, D., Martin, P., & Chilliard, Y. (1998). Messenger RNAs encoding lipoprotein lipase, fatty acid synthase and hormone-sensitive lipase in the adipose tissue of underfed-refed ewes and cows. *Reproduction, Nutrition, Development*, 38, 297–307.

- Bonnet, M., Leroux, C., Chilliard, Y., & Martin, P. (2000a). Rapid communication: Nucleotide sequence of the ovine lipoprotein lipase cDNA. *Journal of Animal Science*, 78, 2994–2995.
- Bonnet, M., Leroux, C., Faulconnier, Y., Hocquette, J. F., Bocquier, F., Martin, P., & Chilliard, Y. (2000b). Lipoprotein lipase activity and mRNA are up-regulated by refeeding in adipose tissue and cardiac muscle of sheep. *Journal of Nutrition*, 130, 749–756.
- Brown, M. S., & Goldstein, J. L. (1997). The SREBP pathway: Regulation of cholesterol metabolism by proteolysis of a membrane-bound transcription factor. *Cell*, 89, 331–340.
- Camps, L., Reina, M., Llobera, M., Vilaro, S., & Olivecrona, T. (1990). Lipoprotein lipase: Cellular origin and functional distribution. *American Journal of Physiology*, 258, C673–681.
- Chilliard, Y., & Ferlay, A. (2004). Dietary lipids and forages interactions on cow and goat milk fatty acid composition and sensory properties. *Reproduction, Nutrition, Development*, 44, 467–492.
- Chilliard, Y., Gagliostro, G., Fléchet, J., Lefaivre, J., & Sebastian, I. (1991). Duodenal rapeseed oil infusion in early and midlactation cows. 5. Milk fatty acids and adipose tissue lipogenic activities. *Journal of Dairy Science*, 74, 1844–1854.
- Chilliard, Y., Ferlay, A., Mansbridge, R. M., & Doreau, M. (2000). Ruminant milk fat plasticity: Nutritional control of saturated, polyunsaturated, *trans* and conjugated fatty acids. *Annales de Zootechnie*, 49, 181–205.
- Chilliard, Y., Ferlay, A., & Doreau, M. (2001). Effect of different types of forages, animal fat or marine oils in cow's diet on milk fat secretion and composition, especially conjugated linoleic acid (CLA) and polyunsaturated fatty acids. *Livestock Production Science*, 70, 31–48.
- Chilliard, Y., Ferlay, A., Rouel, J., & Lamberet, G. (2003a). A review of nutritional and physiological factors affecting goat milk lipid synthesis and lipolysis. *Journal of Dairy Science*, 86, 1751–1770.
- Chilliard, Y., Rouel, J., Capitan, P., Chabosseau, J. M., Raynal-Ljutovac, K., & Ferlay, A. (2003b). Correlations between milk fat content and fatty acid composition in goats receiving different combinations of forages and lipid supplements. In Y. van der Honing (Ed.), *Book of Abstracts of the 54th Annual Meeting of European Association for Animal Production* (p. 343), Rome, Italy, August 31–September 3, 2003. Wageningen, The Netherlands: Wageningen Academic Publishers.
- Chilliard, Y., Rouel, J., Chabosseau, J. M., Capitan, P., Gaborit, P., & Ferlay, A. (2003c). Interactions between raygrass preservation and linseed oil supplementation on goat milk yield and composition, including *trans* and conjugated fatty acids. In Y. van der Honing (Ed.), *Book of Abstracts of the 54th Annual Meeting of European Association for Animal Production* (p. 343), Rome, Italy, August 31–September 3, 2003. Wageningen, The Netherlands: Wageningen Academic Publishers.
- Chilliard, Y., Rouel, J., Ferlay, A., Bernard, L., Gaborit, P., Raynal-Ljutovac, K., Lauret, A., & Leroux, C. (2006a). Optimising goat's milk and cheese fatty acid composition. In C. Williams and J. Buttriss (Eds.), *Improving the Fat Content of Foods* (pp. 281–312). Cambridge: Woodhead Publishing Ltd.
- Chilliard, Y., Ollier, S., Rouel, J., Bernard, L., & Leroux, C. (2006b). Milk fatty acid profile in goats receiving high forage or high concentrate diets supplemented, or not, with either whole rapeseeds or sunflower oil. In Y. van der Honing (Ed.), *Book of Abstracts of the 57th Annual Meeting of European Association for Animal Production* (p. 296), Antalya, Turkey, September 17–20, 2006. Wageningen, The Netherlands: Wageningen Academic Publishers.
- Christie, W. W. (1979). The effects of diet and other factors on the lipid composition of ruminant tissues and milk. *Progress in Lipid Research*, 17, 245–277.
- Chung, M., Ha, S., Jeong, S., Bok, J., Cho, K., Baik, M., & Choi, Y. (2000). Cloning and characterization of bovine stearoyl CoA desaturase cDNA from adipose tissues. *Bioscience, Biotechnology, and Biochemistry*, 64, 1526–1530.

- Clarke, S. D. (2001). Nonalcoholic steatosis and steatohepatitis. I. Molecular mechanism for polyunsaturated fatty acid regulation of gene transcription. *American Journal of Physiology—Gastrointestinal and Liver Physiology*, 281, G865–869.
- Cobanoglu, O., Zaitoun, I., Chang, Y. M., Shook, G. E., & Khatib, H. (2006). Effects of the signal transducer and activator of transcription 1 (STAT1) gene on milk production traits in Holstein dairy cattle. *Journal of Dairy Science*, 89, 4433–4437.
- Coleman, R. A., & Lee, D. P. (2004). Enzymes of triacylglycerol synthesis and their regulation. *Progress in Lipid Research*, 43, 134–176.
- Coleman, R. A., Lewin, T. M., & Muoio, D. M. (2000). Physiological and nutritional regulation of enzymes of triacylglycerol synthesis. *Annual Review of Nutrition*, 20, 77–103.
- Corl, B. A., Baumgard, L. H., Dwyer, D. A., Griinari, J. M., Phillips, B. S., & Bauman, D. E. (2001). The role of Delta(9)-desaturase in the production of *cis*-9, *trans*-11 CLA. *Journal of Nutritional Biochemistry*, 12, 622–630.
- Corl, B. A., Baumgard, L. H., Griinari, J. M., Delmonte, P., Morehouse, K. M., Yurawecz, M. P., & Bauman, D. E. (2002). *Trans*-7, *cis*-9 CLA is synthesized endogenously by delta9-desaturase in dairy cows. *Lipids*, 37, 681–688.
- Coulon, J. B., Dupont, D., Pochet, S., Pradel, P., & Duployer, H. (2001). Effect of genetic potential and level of feeding on milk protein composition. *Journal of Dairy Research*, 68, 569–577.
- Cunningham, B. A., Moncur, J. T., Huntington, J. T., & Kinlaw, W. B. (1998). ?Spot 14? protein: A metabolic integrator in normal and neoplastic cells. *Thyroid*, 8, 815–825.
- Daniels, K. M., Webb, K. E., Jr., McGilliard, M. L., Meyer, M. J., Van Amburgh, M. E., & Akers, R. M. (2006). Effects of body weight and nutrition on mammary protein expression profiles in Holstein heifers. *Journal of Dairy Science*, 89, 4276–4288.
- Davis, C. L., & Brown, R. E. (1970). Low-fat milk syndrome. In A.T. Phillipson (Ed.) *Physiology of Digestion and Metabolism in Ruminants* (pp. 545–565). Newcastle Upon Tyne: Oriel.
- Delbecchi, L., Ahnadi, C. E., Kennelly J. J., & Lacasse, P. (2001). Milk fatty acid composition and mammary lipid metabolism in Holstein cows fed protected or unprotected canola seeds. *Journal of Dairy Science*, 84, 1375–1381.
- Del Prado, M., Villalpando, S., Gordillo, J., & Hernandez-Montes, H. (1999). A high dietary lipid intake during pregnancy and lactation enhances mammary gland lipid uptake and lipoprotein lipase activity in rats. *Journal of Nutrition*, 129, 1574–1578.
- Ding, S. T., Schinckel, A. P., Weber, T. E., & Mersmann, H. J. (2000). Expression of porcine transcription factors and genes related to fatty acid metabolism in different tissues and genetic populations. *Journal of Animal Science*, 78, 2127–2134.
- Eberlé, D., Hegarty, B., Bossard, P., Ferré, P., & Foufelle, F. (2004). SREBP transcription factors: Master regulators of lipid homeostasis. *Biochimie*, 86, 839–848.
- Enoch, H. G., Catala, A., & Strittmatter, P. (1976). Mechanism of rat liver microsomal stearyl-CoA desaturase. Studies of the substrate specificity, enzyme-substrate interactions, and the function of lipid. *Journal of Biological Chemistry*, 251, 5095–5103.
- Ferlay, A., Rouel, J., Chabosseau, J. M., Capitan, P., Raynal-Ljutovac, K., & Chilliard, Y. (2003). Interactions between raygrass preservation and high-oleic sunflower oil supplementation on goat milk composition, including *trans* and conjugated fatty acids. In Y. van der Honing (Ed.), *Book of Abstracts of the 55th Annual Meeting of European Association for Animal Production* (p. 350), Bled, Slovenia, September 5–9, 2004. Wageningen, The Netherlands: Wageningen Academic Publishers.
- Gagliostro, G., Chilliard, Y., & Davicco, M. J. (1991). Duodenal rapeseed oil infusion in early and midlactation cows. 3. Plasma hormones and mammary apparent uptake of metabolites. *Journal of Dairy Science*, 74, 1893–1903.
- German, J. B., Morand, C., Dillard, C. J., & Xu, R. (1997). Milk fat composition: Targets for alteration of function and nutrition. In R. A. S. Welch, D. J. W. Burns, S. R. Davis, A. I. Popay, and C. G. Prosser (Eds.), *Milk Composition, Production and Biotechnology* (pp. 39–72). New York: CAB International.

- Girard, J., Perdereau, D., Foufelle, F., Prip-Buus, C., & Ferré, P. (1994). Regulation of lipogenic enzyme gene expression by nutrients and hormones. *FASEB Journal*, 8, 36–42.
- Girard, J., Ferré, P., & Foufelle, F. (1997). Mechanisms by which carbohydrates regulate expression of genes for glycolytic and lipogenic enzymes. *Annual Review of Nutrition*, 17, 325–352.
- Granlund, L., Juvet, L. K., Pedersen, J. I., & Nebb, H. I. (2003). *Trans*<sup>10</sup>, *cis*<sup>12</sup>-conjugated linoleic acid prevents triacylglycerol accumulation in adipocytes by acting as a PPAR $\gamma$  modulator. *Journal of Lipid Research*, 44, 1441–1452.
- Griinari, J. M., Dwyer, D. A., McGuire, M. A., Bauman, D. E., Palmquist, D. L., & Nurmela, K. V. (1998). *Trans*-octadecenoic acids and milk fat depression in lactating dairy cows. *Journal of Dairy Science*, 81, 1251–1261.
- Griinari, J. M., Corl, B. A., Lacy, S. H., Chouinard, P. Y., Nurmela, K. V., & Bauman, D. E. (2000). Conjugated linoleic acid is synthesized endogenously in lactating dairy cows by  $\Delta^9$ -desaturase. *Journal of Nutrition*, 130, 2285–2291.
- Grisart, B., Coppieters, W., Farnir, F., Karim, L., Ford, C., Berzi, P., Cambisano, N., Mni, M., Reid, S., Simon, P., Spelman, R., Georges, M., & Snell, R. (2002). Positional candidate cloning of a QTL in dairy cattle: Identification of a missense mutation in the bovine DGAT1 gene with major effect on milk yield and composition. *Genome Research*, 12, 222–231.
- Grisart, B., Farnir, F., Karim, L., Cambisano, N., Kim, J. J., Kvasz, A., Mni, M., Simon, P., Frere, J. M., Coppieters, W., & Georges, M. (2004). Genetic and functional confirmation of the causality of the DGAT1 K232A quantitative trait nucleotide in affecting milk yield and composition. *Proceedings of the National Academy of Sciences USA*, 101, 2398–2403.
- Guichard, C., Dugail, I., Le Liepvre, X., & Lavau, M. (1992). Genetic regulation of fatty acid synthetase expression in adipose tissue: Overtranscription of the gene in genetically obese rats. *Journal of Lipid Research*, 33, 679–687.
- Hansen, H. O., & Knudsen, J. (1987). Effect of exogenous long-chain fatty acids on lipid biosynthesis in dispersed ruminant mammary gland epithelial cells: Esterification of long-chain exogenous fatty acids. *Journal of Dairy Science*, 70, 1344–1349.
- Hansen, H. O., Tornehave, D., & Knudsen, J. (1986). Synthesis of milk specific fatty acids and proteins by dispersed goat mammary-gland epithelial cells. *Biochemical Journal*, 238, 167–172.
- Harvatine, K. J., & Bauman, D. E. (2006). SREBP1 and thyroid hormone responsive spot 14 (S14) are involved in the regulation of bovine mammary lipid synthesis during diet-induced milk fat depression and treatment with CLA. *Journal of Nutrition*, 136, 2468–2474.
- He, J., & Furmanski, P. (1995). Sequence specificity and transcriptional activation in the binding of lactoferrin to DNA. *Nature*, 373, 721–724.
- Hennighausen, L. (1997). Molecular mechanisms of hormone controlled gene expression in the breast. *Molecular Biology Reports*, 24, 169–174.
- Jayakumar, A., Tai, M. H., Huang, W. Y., al-Feel, W., Hsu, M., Abu-Elheiga, L., Chirala, S. S., & Wakil, S. J. (1995). Human fatty acid synthase: Properties and molecular cloning. *Proceedings of the National Academy of Sciences USA*, 92, 8695–8699.
- Jayan, G. C., & Herbein, J. H. (2000). “Healthier” dairy fat using *trans*-vaccenic acid. *Nutrition and food Science*, 30, 304–309.
- Jensen, D. R., Bessesen, D. H., Etienne, J., Eckel, R. H., & Neville, M. C. (1991). Distribution and source of lipoprotein lipase in mouse mammary gland. *Journal of Lipid Research*, 32, 733–742.
- Jensen, R. G. (2002). The composition of bovine milk lipids: January 1995 to December 2000. *Journal of Dairy Science*, 85, 295–350.
- Keating, A. F., Stanton, C., Murphy, J. J., Smith, T. J., Ross, R. P., & Cairns, M. T. (2005). Isolation and characterization of the bovine stearoyl-CoA desaturase promoter and analysis of polymorphisms in the promoter region in dairy cows. *Mammalian Genome*, 16, 184–193.
- Keating, A. F., Kennelly, J. J., & Zhao, F. Q. (2006). Characterization and regulation of the bovine stearoyl-CoA desaturase gene promoter. *Biochemical and Biophysical Research Communications*, 344, 233–240.

- Kim, K. H. (1997). Regulation of mammalian acetyl-coenzyme A carboxylase. *Annual Review of Nutrition* 17, 77–99.
- Kinsella, J. E. (1970). Stearic acid metabolism by mammary cells. *Journal of Dairy Science*, 53, 1757–1765.
- Klein, I., Sarkadi, B., & Varadi, A. (1999). An inventory of the human ABC proteins. *Biochimica et Biophysica Acta*, 1461, 237–262.
- Knudsen, J., & Grunnet, I. (1982). Transacylation as a chain-termination mechanism in fatty acid synthesis by mammalian fatty acid synthetase. Synthesis of medium-chain-length (C8–C12) acyl-CoA esters by goat mammary-gland fatty acid synthetase. *Biochemical Journal*, 202, 139–143.
- Knudsen, J., Neergaard, T. B., Gaigg, B., Jensen, M. V., & Hansen, J. K. (2000). Role of acyl-CoA binding protein in acyl-CoA metabolism and acyl-CoA-mediated cell signaling. *Journal of Nutrition*, 130, 294S–298S.
- Kühn, C., Thaller, G., Winter, A., Bininda-Emonds, O. R., Kaupe, B., Erhardt, G., Bennewitz, J., Schwerin, M., & Fries, R. (2004). Evidence for multiple alleles at the DGAT1 locus better explains a quantitative trait locus with major effect on milk fat content in cattle. *Genetics*, 167, 1873–1881.
- Lee, S. H., & Hossner, K. L. (2002). Coordinate regulation of ovine adipose tissue gene expression by propionate. *Journal of Animal Science*, 80, 2840–2849.
- Légrand, P., Catheline, D., Fichot, M. C., & Lemarchal, P. (1997). Inhibiting delta9-desaturase activity impairs triacylglycerol secretion in cultured chicken hepatocytes. *Journal of Nutrition*, 127, 249–256.
- Lehner, R., & Kuksis, A. (1996). Biosynthesis of triacylglycerols. *Progress in Lipid Research*, 35, 169–201.
- Leroux, C., Le Provost, F., Petit, E., Bernard, L., Chilliard, Y., & Martin, P. (2003). Real-time RT-PCR and cDNA macroarray to study the impact of the genetic polymorphism at the alphas1-casein locus on the expression of genes in the goat mammary gland during lactation. *Reproduction, Nutrition, Development*, 43, 459–469.
- Leroux, C., Laubier, J., Giraud-Delville, C., Le Provost, F., Chadi, S., Bernard, L., Bonnet, M., Chilliard, Y., & Martin, P. (2007). Sharp increase of goat fatty acid synthase gene expression in mammary gland during the last part of pregnancy. *Journal of Dairy Research*, submitted.
- Lin, X., Looor, J. J., & Herbein, J. H. (2004). *Trans*10,*cis*12-18:2 is a more potent inhibitor of *de novo* fatty acid synthesis and desaturation than *cis*9,*trans*11-18:2 in the mammary gland of lactating mice. *Journal of Nutrition*, 134, 1362–1368.
- Liu, W., Degner, S. C., & Romagnolo, D. F. (2006). *Trans*-10, *cis*-12 conjugated linoleic acid inhibits prolactin-induced cytosolic NADP<sup>+</sup>-dependent isocitrate dehydrogenase expression in bovine mammary epithelial cells. *Journal of Nutrition*, 136, 2743–2747.
- Lock, A. L., Teles, B. M., Perfield, J. W. II, Bauman, D. E., & Sinclair, L. A. (2006). A conjugated linoleic acid supplement containing *trans*-10, *cis*-12 reduces milk fat synthesis in lactating sheep. *Journal of Dairy Science*, 89, 1525–1532.
- Lock, A. L., Tyburczy, C., Dwyer, D. A., Harvatine, K. J., Destailats, F., Mouloungui, Z., Candy, L., & Bauman, D. E. (2007). *Trans*-10 octadecenoic acid does not reduce milk fat synthesis in dairy cows. *Journal of Nutrition*, 137, 71–76.
- Looor, J. J., & Herbein, J. H. (2003). Reduced fatty acid synthesis and desaturation due to exogenous *trans*10,*cis*12-CLA in cows fed oleic or linoleic oil. *Journal of Dairy Science*, 86, 1354–1369.
- Looor, J. J., Ueda, K., Ferlay, A., Chilliard, Y., & Doreau, M. (2004a). Biohydrogenation, duodenal flow, and intestinal digestibility of *trans* fatty acids and conjugated linoleic acids in response to dietary forage:concentrate ratio and linseed oil in dairy cows. *Journal of Dairy Science*, 87, 2472–2485.
- Looor, J. J., Ueda, K., Ferlay, A., Chilliard, Y., & Doreau, M. (2004b). Short communication: Diurnal profiles of conjugated linoleic acids and *trans* fatty acids in ruminal fluid from

- cows fed a high concentrate diet supplemented with fish oil, linseed oil, or sunflower oil. *Journal of Dairy Science*, 87, 2468–2471.
- Loor, J. J., Dann, H. M., Everts, R. E., Rodriguez-Zas, S. L., Lewin, H. A., & Drackley, J. K. (2004c). Mammary and hepatic gene expression analysis in periparturient dairy cows using a bovine cDNA microarray. *Journal of Animal Science*, 82 (Suppl 1), 196.
- Loor, J. J., Ferlay, A., Ollier, A., Doreau, M., & Chilliard, Y. (2005a). Relationship among *trans* and conjugated fatty acids and bovine milk fat yield due to dietary concentrate and linseed oil. *Journal of Dairy Science*, 88, 726–740.
- Loor, J. J., Doreau, M., Chardigny, J. M., Ollier, A., Sébédio, J. L., & Chilliard, Y. (2005b). Effects of ruminal or duodenal supply of fish oil on milk fat secretion and profiles of *trans*-fatty acids and conjugated linoleic acid isomers in dairy cows fed maize silage. *Animal Feed Science and Technology*, 119, 227–246.
- Loor, J. J., Piperova, L. S., Everts, R. E., Rodriguez-Zas, S. L., Drackley, J. K., Erdman, R. A., & Lewin, H. A. (2005c). Mammary gene expression profiling in cows fed a milk fat-depressing diet using a bovine 13,000 oligonucleotide microarray. *Journal of Animal Science*, 83 (Suppl 1), 120.
- Loor, J. J., Ferlay, A., Ollier, A., Ueda, K., Doreau, M., & Chilliard, Y. (2005d). High-concentrate diets and polyunsaturated oils alter *trans* and conjugated isomers in bovine rumen, blood, and milk. *Journal of Dairy Science*, 88, 3986–3999.
- Lopez-Casillas, F., Ponce-Castaneda, M. V., & Kim, K. H. (1991). *In vivo* regulation of the activity of the two promoters of the rat acetyl coenzyme-A carboxylase gene. *Endocrinology*, 129, 1049–1058.
- Mao, J., Marcos, S., Davis, S. K., Burzlaff, J., & Seyfert, H. M. (2001). Genomic distribution of three promoters of the bovine gene encoding acetyl-CoA carboxylase alpha and evidence that the nutritionally regulated promoter I contains a repressive element different from that in rat. *Biochemical Journal*, 358, 127–135.
- Mao, J., Molenaar, A. J., Wheeler, T. T., & Seyfert, H. M. (2002). STAT5 binding contributes to lactational stimulation of promoter III expressing the bovine acetyl-CoA carboxylase alpha-encoding gene in the mammary gland. *Journal of Molecular Endocrinology*, 29, 73–88.
- Marshall, M. O., & Knudsen, J. (1977). The specificity of 1-acyl-sn-glycerol 3-phosphate acyltransferase in microsomal fractions from lactating cow mammary gland towards short, medium and long chain acyl-CoA esters. *Biochimica et Biophysica Acta*, 489, 236–241.
- Massaro, M., Carluccio, M. A., & De Caterina, R. (1999). Direct vascular antiatherogenic effects of oleic acid: A clue to the cardioprotective effects of the Mediterranean diet. *Cardiologia*, 44, 507–513.
- Mather, I. H. (2000). A review and proposed nomenclature for major proteins of the milk-fat globule membrane. *Journal of Dairy Science*, 83, 203–247.
- Matitashvili, E., & Bauman, D. E. (2000). Effect of different isomers of C18:1 and C18:2 fatty acids on lipogenesis in bovine mammary epithelial cells. *Journal of Animal Science*, 78, 165.
- Mayorek, N., Grinstein, I., & Bar-Tana, J. (1989). Triacylglycerol synthesis in cultured rat hepatocytes. The rate-limiting role of diacylglycerol acyltransferase. *European Journal of Biochemistry*, 182, 395–400.
- Mihara, K. (1990). Structure and regulation of rat liver microsomal stearoyl-CoA desaturase gene. *Journal of Biochemistry*, 108, 1022–1029.
- Mikkelsen, J., & Knudsen, J. (1987). Acyl-CoA-binding protein from cow. Binding characteristics and cellular and tissue distribution. *Biochemical Journal*, 248, 709–714.
- Miller, R. T., Scappino, L. A., Long, S. M., & Corton, J. C. (2001). Role of thyroid hormones in hepatic effects of peroxisome proliferators. *Toxicologic Pathology*, 29, 149–155.
- Mistry, D. H., & Medrano, J. F. (2002). Cloning and localization of the bovine and ovine lysophosphatidic acid acyltransferase (LPAAT) genes that codes for an enzyme involved in triglyceride biosynthesis. *Journal of Dairy Science*, 85, 28–35.

- Molenaar, A., Mao, J., Oden, K., & Seyfert, H. M. (2003). All three promoters of the acetyl-coenzyme A-carboxylase alpha-encoding gene are expressed in mammary epithelial cells of ruminants. *Journal of Histochemistry and Cytochemistry*, 51, 1073–1081.
- Mosley, E. E., Powell, G. L., Riley, M. B., & Jenkins, T. C. (2002). Microbial biohydrogenation of oleic acid to *trans* isomers *in vitro*. *Journal of Lipid Research*, 43, 290–296.
- Murrieta, C. M., Hess, B. W., Scholljegerdes, E. J., Engle, T. E., Hossner, K. L., Moss, G. E., & Rule, D. C. (2006). Evaluation of milk somatic cells as a source of mRNA for study of lipogenesis in the mammary gland of lactating beef cows supplemented with dietary high-linoleate safflower seeds. *Journal of Animal Science*, 84, 2399–2405.
- Mutch, D. M., Anderle, P., Fiaux, M., Mansourian, R., Vidal, K., Wahli, W., Williamson, G., & Roberts, M. A. (2004). Regional variations in ABC transporter expression along the mouse intestinal tract. *Physiological Genomics*, 17, 11–20.
- Neville, M. C., & Picciano, M. F. (1997). Regulation of milk lipid secretion and composition. *Annual Review of Nutrition*, 17, 159–183.
- Nielsen, M. O., & Jakobsen, K. (1994). Changes in mammary uptake of free fatty acids, triglyceride, cholesterol and phospholipid in relation to milk synthesis during lactation in goats. *Comparative Biochemistry and Physiology. Part A, Physiology*, 109, 857–867.
- Ntambi, J. M. (1999). Regulation of stearoyl-CoA desaturase by polyunsaturated fatty acids and cholesterol. *Journal of Lipid Research*, 40, 1549–1558.
- Ntambi, J. M., Buhrow, S. A., Kaestner, K. H., Christy, R. J., Sibley, E., Kelly, T. J., Jr., & Lane, M. D. (1988). Differentiation-induced gene expression in 3T3-L1 preadipocytes. Characterization of a differentially expressed gene encoding stearoyl-CoA desaturase. *Journal of Biological Chemistry*, 263, 17291–17300.
- Offer, N. W., Speake, B. K., Dixon, J., & Marsden, M. (2001). Effect of fish-oil supplementation on levels of (N-3) poly-unsaturated fatty acids in the lipoprotein fractions of bovine plasma. *Animal Science*, 73, 523–531.
- Ogg, S. L., Weldon, A. K., Dobbie, L., Smith, A. J., & Mather, I. H. (2004). Expression of butyrophilin (Btn1a1) in lactating mammary gland is essential for the regulated secretion of milk-lipid droplets. *Proceedings of the National Academy of Sciences USA*, 101, 10084–10089.
- Ollier, S., Robert-Granié, C., Bes, S., Goutte, M., Faulconnier, Y., Chilliard, Y., & Leroux, C. (2006). Impact of nutrition on mammary transcriptome and its interaction with the CSN1S1 genotype in lactating goats. In Y. van der Honing (Ed.) *Book of Abstracts, 57th Annual Meeting of European Association for Animal Production* (p. 49), Antalya, Turkey, September 17–20, 2006. Wageningen, The Netherlands: Wageningen Academic Publishers.
- Ollier, S., Robert-Granié, C., Bernard, L., Chilliard, Y., & Leroux, C. (2007). Mammary transcriptome analysis of food-deprived lactating goats highlights genes involved in milk secretion and programmed cell death. *Journal of Nutrition*, 137, 560–567.
- Palmquist, D. L., Beaulieu, A. D., & Barbano, D. M. (1993). Feed and animal factors influencing milk fat composition. *Journal of Dairy Science*, 76, 1753–1771.
- Pariza, M. W., Park, Y., & Cook, M. E. (2001). The biologically active isomers of conjugated linoleic acid. *Progress in Lipid Research*, 40, 283–298.
- Parodi, P. W. (1982). Positional distribution of fatty acids in triglycerides from milk of several species of mammals. *Lipids*, 17, 437–442.
- Perfield, J. W. II, Sæbo, A., & Bauman, D. E. (2004). Use of conjugated linoleic acid (CLA) enrichments to examine the effects of *trans*-8, *cis*-10 CLA, and *cis*-11, *trans*-13 CLA on milk-fat synthesis. *Journal of Dairy Science*, 87, 1196–1202.
- Perfield, J. W. II, Lock, A. L., Sæbo, A., Grinari, J. M., & Bauman, D. E. (2005). *Trans*-9,*cis*-11 conjugated linoleic acid (CLA) reduces milk fat synthesis in lactating dairy cows. *Journal of Dairy Science*, 88 (Suppl 1), 211.

- Perfield, J. W. II, Delmonte, P., Lock, A. L., Yurawecz, M. P., & Bauman, D. E. (2006). *Trans*-10, *trans*-12 conjugated linoleic acid does not affect milk fat yield but reduces delta9-desaturase index in dairy cows. *Journal of Dairy Science*, 89, 2559–2566.
- Peterson, D. G., Matitashvili, E. A., & Bauman, D. E. (2003). Diet-induced milk fat depression in dairy cows results in increased *trans*-10, *cis*-12 CLA in milk fat and coordinate suppression of mRNA abundance for mammary enzymes involved in milk fat synthesis. *Journal of Nutrition*, 133, 3098–3102.
- Peterson, D. G., Matitashvili, E. A., & Bauman, D. E. (2004). The inhibitory effect of *trans*-10, *cis*-12 CLA on lipid synthesis in bovine mammary epithelial cells involves reduced proteolytic activation of the transcription factor SREBP-1. *Journal of Nutrition*, 134, 2523–2527.
- Piperova, L. S., Teter, B. B., Bruckental, I., Sampugna, J., Mills, S. E., Yurawecz, M. P., Fritsche, J., Ku, K., & Erdman, R. A. (2000). Mammary lipogenic enzyme activity, *trans* fatty acids and conjugated linoleic acids are altered in lactating dairy cows fed a milk fat-depressing diet. *Journal of Nutrition*, 130, 2568–2574.
- Rosen, E. D., & Spiegelman, B. M. (2001). PPAR $\gamma$ : A nuclear regulator of metabolism, differentiation, and cell growth. *Journal of Biological Chemistry*, 276, 37731–37734.
- Rosen, E. D., Sarraf, P., Troy, A. E., Bradwin, G., Moore, K., Milstone, D. S., Spiegelman, B. M., & Mortensen, R. M. (1999). PPAR $\gamma$  is required for the differentiation of adipose tissue *in vivo* and *in vitro*. *Molecular Cell*, 4, 611–617.
- Rouel, J., Ferlay, A., Bruneteau, E., Capitan, P., Raynal-Ljutovac, K., & Chilliard, Y. (2004). Interactions between starchy concentrate and linseed oil supplementation on goat milk yield and composition, including *trans* and conjugated fatty acids (FA). In Y. van der Honing (Ed.), *Book of Abstracts of the 55th Annual Meeting of European Association for Animal Production* (p. 124), Bled, Slovenia, September 5–9, 2004. Wageningen, The Netherlands: Wageningen Academic Publishers.
- Roy, R., Taourit, S., Zaragoza, P., Eggen, A., & Rodellar, C. (2005). Genomic structure and alternative transcript of bovine fatty acid synthase gene (FASN): Comparative analysis of the FASN gene between monogastric and ruminant species. *Cytogenetic and Genome Research*, 111, 65–73.
- Roy, A., Ferlay, A., Shingfield, K. J., & Chilliard, Y. (2006a). Examination of the persistency of milk fatty acid composition responses to plant oils in cows fed different basal diets, with particular emphasis on *trans*-C<sub>18:1</sub> fatty acids and isomers of conjugated linoleic acid. *Animal Science*, 82, 479–492.
- Roy, R., Ordoas, L., Taourit, S., Zaragoza, P., Eggen, A., & Rodellar, C. (2006b). Genomic structure and an alternative transcript of bovine mitochondrial glycerol-3-phosphate acyltransferase gene (GPAM). *Cytogenetics and Genome Research*, 112, 82–89.
- Roy, R., Ordoas, L., Zaragoza, P., Romero, A., Moreno, C., Altarriba, J., & Rodellar, C. (2006c). Association of polymorphisms in the bovine FASN gene with milk-fat content. *Animal Genetics*, 37, 215–218.
- Sæbo, A., Sæbo, P. C., Griinari, J. M., & Shingfield, K. J. (2005). Effect of abomasal infusions of geometric isomers of 10,12 conjugated linoleic acid on milk fat synthesis in dairy cows. *Lipids*, 40, 823–832.
- Sakai, J., Duncan, E. A., Rawson, R. B., Hua, X., Brown, M. S., & Goldstein, J. L. (1996). Sterol-regulated release of SREBP-2 from cell membranes requires two sequential cleavages, one within a transmembrane segment. *Cell*, 85, 1037–1046.
- Sarraf, P., Mueller, E., Smith, W. M., Wright, H. M., Kum, J. B., Aaltonen, L. A., de la Chapelle, A., Spiegelman, B. M., & Eng, C. (1999). Loss-of-function mutations in PPAR gamma associated with human colon cancer. *Molecular Cell*, 3, 799–804.
- Schoonjans, K., Peinado-Onsurbe, J., Lefebvre, A. M., Heyman, R. A., Briggs, M., Deeb, S., Staels, B., & Auwerx, J. (1996a). PPAR $\alpha$  and PPAR $\gamma$  activators direct a distinct tissue-specific transcriptional response via a PPRE in the lipoprotein lipase gene. *EMBO Journal*, 15, 5336–5348.

- Schoonjans, K., Staels, B., & Auwerx, J. (1996b). Role of the peroxisome proliferator-activated receptor (PPAR) in mediating the effects of fibrates and fatty acids on gene expression. *Journal of Lipid Research*, 37, 907–925.
- Schweizer, M., Takabayashi, K., Laux, T., Beck, K. F., & Schreglmann, R. (1989). Rat mammary gland fatty acid synthase: Localization of the constituent domains and two functional polyadenylation/termination signals in the cDNA. *Nucleic Acids Research*, 17, 567–586.
- Senda, M., Oka, K., Brown, W. V., Qasba, P. K., & Furuichi, Y. (1987). Molecular cloning and sequence of a cDNA coding for bovine lipoprotein lipase. *Proceedings of the National Academy of Sciences USA*, 84, 4369–4373.
- Seyfert, H. M., Pitra, C., Meyer, L., Brunner, R. M., Wheeler, T. T., Molenaar, A., McCracken, J. Y., Herrmann, J., Thiesen, H. J., & Schwerin, M. (2000). Molecular characterization of STAT5A- and STAT5B-encoding genes reveals extended intragenic sequence homogeneity in cattle and mouse and different degrees of divergent evolution of various domains. *Journal of Molecular Evolution*, 50, 550–561.
- Shimomura, I., Shimano, H., Horton, J. D., Goldstein, J. L., & Brown, M. S. (1997). Differential expression of exons 1a and 1c in mRNAs for sterol regulatory element binding protein-1 in human and mouse organs and cultured cells. *Journal of Clinical Investigation*, 99, 838–845.
- Shimomura, I., Hammer, R. E., Richardson, J. A., Ikemoto, S., Bashmakov, Y., Goldstein, J. L., & Brown, M. S. (1998). Insulin resistance and diabetes mellitus in transgenic mice expressing nuclear SREBP-1c in adipose tissue: Model for congenital generalized lipodystrophy. *Genes and Development*, 12, 3182–3194.
- Shingfield, K. J., Ahvenjarvi, S., Toivonen, V., Ärölä, A., Nurmela, K. V., Huhtanen, P., & Griinari, J. M. (2003). Effect of dietary fish oil on biohydrogenation of fatty acids and milk fatty acid content in cows. *Animal Science*, 77, 165–179.
- Shingfield, K. J., Reynolds, C. K., Lupoli, B., Toivonen, V., Yurawecz, M. P., Delmonte, P., Griinari, J. M., Grandison, A. S., & Beever, D. E. (2005). Effect of forage type and proportion of concentrate in diet on milk fatty acid composition in cows fed sunflower oil and fish oil. *Animal Science*, 80, 225–238.
- Shingfield, K. J., Reynolds, C. K., Hervas, G., Griinari, J. M., Grandison, A. S., & Beever, D. E. (2006). Examination of the persistency of milk fatty acid composition responses to fish oil and sunflower oil in the diet of dairy cows. *Journal of Dairy Science*, 89, 714–732.
- Shirley, J. E., Emery, R. S., Convey, E. M., & Oxender, W. D. (1973). Enzymic changes in bovine adipose and mammary tissue, serum and mammary tissue hormonal changes with initiation of lactation. *Journal of Dairy Science*, 56, 569–574.
- Singh, K., Hartley, D. G., McFadden, T. B., & Mackenzie, D. D. (2004). Dietary fat regulates mammary stearoyl coA desaturase expression and activity in lactating mice. *Journal of Dairy Research*, 71, 1–6.
- Small, D. M. (1991). The effects of glyceride structure on absorption and metabolism. *Annual Review of Nutrition*, 11, 413–434.
- Specht, B., Bartetzko, N., Hohoff, C., Kuhl, H., Franke, R., Borchers, T., & Spener, F. (1996). Mammary derived growth inhibitor is not a distinct protein but a mix of heart-type and adipocyte-type fatty acid-binding protein. *Journal of Biological Chemistry*, 271, 19943–19949.
- Spitsberg, V. L., Matitashvili, E., & Gorewit, R. C. (1995). Association and coexpression of fatty-acid-binding protein and glycoprotein CD36 in the bovine mammary gland. *European Journal of Biochemistry*, 230, 872–878.
- Suchyta, S. P., Sipkovsky, S., Halgren, R. G., Kruska, R., Elftman, M., Weber-Nielsen, M., Vandelaar, M. J., Xiao, L., Tempelman, R. J., & Coussens, P. M. (2003). Bovine mammary gene expression profiling using a cDNA microarray enhanced for mammary-specific transcripts. *Physiological Genomics*, 16, 8–18.

- Sundvold, H., Brzozowska, A., & Lien, S. (1997). Characterisation of bovine peroxisome proliferator-activated receptors gamma 1 and gamma 2: Genetic mapping and differential expression of the two isoforms. *Biochemical and Biophysical Research Communications*, 239, 857–861.
- Viswanadha, S., Giesy, J. G., Hanson, T. W., & McGuire, M. A. (2003). Dose response of milk fat to intravenous administration of the *trans*-10, *cis*-12 isomer of conjugated linoleic acid. *Journal of Dairy Science*, 86, 3229–3236.
- Vituro, E., Farke, C., Meyer, H. H., & Albrecht, C. (2006). Identification, sequence analysis and mRNA tissue distribution of the bovine sterol transporters ABCG5 and ABCG8. *Journal of Dairy Science*, 89, 553–561.
- Wakil, S. J. (1989). Fatty acid synthase, a proficient multifunctional enzyme. *Biochemistry*, 28, 4523–4530.
- Wang, X., Briggs, M. R., Hua, X., Yokoyama, C., Goldstein, J. L., & Brown, M. S. (1993). Nuclear protein that binds sterol regulatory element of low density lipoprotein receptor promoter. II. Purification and characterization. *Journal of Biological Chemistry*, 268, 14497–14504.
- Ward, R. J., Travers, M. T., Richards, S. E., Vernon, R. G., Salter, A. M., Buttery, P. J., & Barber, M. C. (1998). Stearoyl-CoA desaturase mRNA is transcribed from a single gene in the ovine genome. *Biochimica et Biophysica Acta*, 1391, 145–156.
- West, C. E., Bickerstaffe, R., Annison, E. F., & Linzell, J. L. (1972). Studies on the mode of uptake of blood triglycerides by the mammary gland of the lactating goat. The uptake and incorporation into milk fat and mammary lymph of labelled glycerol, fatty acids and triglycerides. *Biochemical Journal*, 126, 477–490.
- Wheeler, T. T., Broadhurst, M. K., Rajan, G. H., & Wilkins, R. J. (1997). Differences in the abundance of nuclear proteins in the bovine mammary gland throughout the lactation and gestation cycles. *Journal of Dairy Science*, 80, 2011–2019.
- Williams, C. M. (2000). Dietary fatty acids and human health. *Annales de Zootechnie*, 49, 165–180.
- Winter, A., Kramer, W., Werner, F. A., Kollers, S., Kata, S., Durstewitz, G., Buitkamp, J., Womack, J. E., Thaller, G., & Fries, R. (2002). Association of a lysine-232/alanine polymorphism in a bovine gene encoding acyl-CoA:diacylglycerol acyltransferase (DGAT1) with variation at a quantitative trait locus for milk fat content. *Proceedings of the National Academy of Sciences USA*, 99, 9300–9305.
- Yang, J., Kennelly, J. J., & Baracos, V. E. (2000a). The activity of transcription factor Stat5 responds to prolactin, growth hormone, and IGF-I in rat and bovine mammary explant culture. *Journal of Animal Science*, 78, 3114–3125.
- Yang, J., Kennelly, J. J., & Baracos, V. E. (2000b). Physiological levels of Stat5 DNA binding activity and protein in bovine mammary gland. *Journal of Animal Science*, 78, 3126–3134.
- Yang, T., Espenshade, P. J., Wright, M. E., Yabe, D., Gong, Y., Aebersold, R., Goldstein, J. L., & Brown, M. S. (2002). Crucial step in cholesterol homeostasis: Sterols promote binding of SCAP to INSIG-1, a membrane protein that facilitates retention of SREBPs in ER. *Cell*, 110, 489–500.
- Yeldandi, A. V., Rao, M. S., & Reddy, J. K. (2000). Hydrogen peroxide generation in peroxisome proliferator-induced oncogenesis. *Mutation Research*, 448, 159–177.
- Yonezawa, T., Yonekura, S., Kobayashi, Y., Hagino, A., Katoh, K., & Obara, Y. (2004a). Effects of long-chain fatty acids on cytosolic triacylglycerol accumulation and lipid droplet formation in primary cultured bovine mammary epithelial cells. *Journal of Dairy Science*, 87, 2527–2534.
- Yonezawa, T., Yonekura, S., Sanosaka, M., Hagino, A., Katoh, K., & Obara, Y. (2004b). Octanoate stimulates cytosolic triacylglycerol accumulation and CD36 mRNA expression but inhibits acetyl coenzyme A carboxylase activity in primary cultured bovine mammary epithelial cells. *Journal of Dairy Research*, 71, 398–404.

- Yu, L., Li-Hawkins, J., Hammer, R. E., Berge, K. E., Horton, J. D., Cohen, J. C., & Hobbs, H. H. (2002). Overexpression of ABCG5 and ABCG8 promotes biliary cholesterol secretion and reduces fractional absorption of dietary cholesterol. *Journal of Clinical Investigation*, 110, 671–680.
- Zhang, L., Ge, L., Parimoo, S., Stenn, K., & Prouty, S. M. (1999). Human stearyl-CoA desaturase: Alternative transcripts generated from a single gene by usage of tandem polyadenylation sites. *Biochemical Journal*, 340 (Pt 1), 255–264.



<http://www.springer.com/978-0-387-74086-7>

Bioactive Components of Milk

Bosze, Z. (Ed.)

2008, XVI, 496 p., Hardcover

ISBN: 978-0-387-74086-7