
4 Blood Supply, Meninges and Cerebrospinal Fluid Circulation

Introduction	95
Arteries of the Brain	95
Meninges, Cisterns and Cerebrospinal Fluid Circulation	110
Circumventricular Organs	126
Veins of the Brain	126
Vessels and Meninges of the Spinal Cord	128

Introduction

The vascularization and the circulation of the cerebrospinal fluid (liquor cerebrospinalis, CSF) of the brain and the spinal cord are of great clinical importance. The main vascular syndromes are summarized in Table 4.1. In this chapter, the anatomy of blood vessels, meninges and circumventricular organs will be discussed. The central nervous system, which is of ectodermal origin (Chap. 2), is surrounded by mesodermal structures. A system of three connective tissue layers, the meninges, and a fluid compartment containing CSF are located between the bony skull and vertebral column and the nervous tissue of the brain and the spinal cord. Blood vessels, themselves of mesodermal origin, are surrounded by derivatives of the meninges over their full extent, until the interface between the capillary wall and the glial basal membrane makes exchange of substances possible. CSF is produced by the choroid plexus of the ventricles. It circulates from the interstitial spaces of the nervous tissue and the choroid plexus, through the ventricles and their apertures in the roof of the fourth ventricle, to the CSF compartment of the subarachnoid space and its exit


through the arachnoid villi to the venous system. The nervous tissue of the central nervous system and the CSF spaces remain segregated from the rest of the body by barrier layers in the meninges (the barrier layer of the arachnoid), the choroid plexus (the blood-CSF barrier) and the capillaries (the blood-brain barrier). The circulation of the CSF plays an important role in maintaining the environment of the nervous tissue; moreover, the subarachnoid space forms a bed that absorbs external shocks.

Arteries of the Brain

The arteries of the brain originate from two of the greater vessels in the neck: the paired internal carotid and vertebral arteries (Fig. 4.1). The *internal carotid artery* enters the skull through the petrosal bone in the carotid canal. It loops through the sinus cavernosus (carotis syphon), where it emits the *ophthalmic artery*. Immediately beyond the origin of the posterior communicating artery it splits into the middle and anterior cerebral arteries. The vertebral arteries enter the skull through the foramen magnum. After their passage through the dura, the arteries become located within connective tissue derived from the pia mater and the arachnoid.

The middle cerebral artery supplies the convexity of the hemisphere (Figs. 4.3, 4.4, 4.5) and the anterior cerebral artery vascularizes approximately the anterior and upper half of the medial aspect of the hemisphere, up to the precuneus (Fig. 4.2). The vertebral arteries unite into the basilar artery at the ventral aspect of the medulla oblongata. Its terminal

- 1 Cerebromeningeal anastomosis*
- 2 Calvaria (outer and inner surfaces)
- 3 Cerebrum (outer surface)
- 4 Callosomarginal artery
- 5 Pericallosal artery
- 6 Corpus callosum
- 7 Anterior cerebral artery
- 8 Supratrochlear artery
- 9 Dorsal nasal artery***
- 10 Frontal foramen




- | | | |
|--|---|---------------------------------------|
| 11 Anterior meningeal artery | 25 Infraorbital canal | 41 Mastoid foramen |
| 12 Lacrimal artery | 26 Infraorbital foramen*** | 42 Posterior meningeal artery |
| 13 Anterior ethmoidal foramen | 27 Angular artery | 43 Junction of the vertebral arteries |
| 14 Anterior ethmoidal artery*** | 28 Facial artery | 44 Jugular foramen |
| 15 Posterior ethmoidal foramen | 29 Maxillary artery | 45 Superficial temporal artery |
| 16 Posterior ethmoidal artery | 30 Middle meningeal artery | 46 Ascending pharyngeal artery |
| 17 Ophthalmic artery | 31 Foramen spinosum | 47 External carotid artery |
| 18 Superior orbital fissure | 32 Internal carotid artery, petrous part | 48 Internal carotid artery |
| 19 Middle meningeal artery, anastomotic branch**** | 33 Middle meningeal artery, parietal branch | 49 Common carotid artery |
| 20 Middle meningeal artery, frontal branch | 34 Middle cerebral artery, insular part | 50 Vertebral artery |
| 21 Optic canal | 35 Posterior communicating artery | |
| 22 Superior conchal artery (anastomosis)*** | 36 Basilar artery | |
| 23 Sphenopalatine artery | 37 Posterior cerebral artery | |
| 24 Infraorbital artery | 38 Parietal foramen** | |
| | 39 Occipital artery | |
| | 40 Occipital artery, mastoid branch** | |

Anastomoses


- | | | |
|---------|-----------------------|------|
| 1 | Cerebromeningeal | * |
| 38+40 | Extracranio-meningeal | ** |
| 9+22+26 | Extracranial-orbital | *** |
| 14+19 | Orbitomeningeal | **** |

Fig. 4.1. Collateral circulation in the arterial system of the head; semidiagrammatic lateral view (2/3x). *Black:* external carotid artery with extracranial branches; *black hatched:* system of the vertebral artery (main trunk); *solid red:* meningeal arteries; *red hatched:* internal carotid artery with orbital and lateral cortical branches; *open red:* medial cortical branches of internal carotid artery




- | | |
|--|---|
| 1 Central sulcus | 20 Posterior cerebral artery, precommunicating part |
| 2 Marginal branch of the cingulate sulcus | 21 Posteromedial central arteries |
| 3 Precuneus | 22 Posterior medial choroidal branch |
| 4 Artery of the precuneus | 23 Posterior cerebral artery, postcommunicating branch |
| 5 Pericallosal artery, posterior branch
(anastomosis with 28) | 24 Posterior thalamic branches |
| 6 Paracentral artery | 25 Medial occipital artery |
| 7 Cingulate sulcus | 26 Cingulatethalamic artery |
| 8 Posteromedial frontal artery | 27 Superior thalamic branch |
| 9 Intermediomedial frontal artery | 28 Dorsal branch of the corpus callosum
(anastomosis with 5) |
| 10 Anteromedial frontal artery | 29 Parietal branch |
| 11 Callosomarginal artery | 30 Parieto-occipital sulcus |
| 12 Pericallosal artery | 31 Parieto-occipital branch |
| 13 Median artery of the corpus callosum | 32 Calcarine branch (in calcarine sulcus) |
| 14 Anterior cerebral artery, postcommunicating part | 33 Posterior temporal branches |
| 15 Anterior communicating artery | 34 Medial intermediate temporal branch |
| 16 Medial frontobasal artery | 35 Lateral occipital artery |
| 17 Temporopolar artery | 36 Collateral sulcus |
| 18 Internal carotid artery | 37 Anterior temporal branches |
| 19 Posterior communicating artery | |

Fig. 4.2. The arteries of the medial hemisphere; the anterior and posterior cerebral arteries (1/1×). Some central branches of the posterior cerebral artery are also shown. End branches of the anterior cerebral artery that reach the lateral side of the superior frontal gyrus are illustrated in Fig. 4.3. Figures 4.2–4.6 are all derived from the same specimen



- | | |
|---|---|
| 1 Central sulcus | 15 Transverse occipital sulcus |
| 2 Posteromedial frontal branch | 16 Temporo-occipital artery |
| 3 Intermediomedial frontal branch | 17 Superior temporal sulcus |
| 4 Anteromedial frontal branch | 18 Posterior temporal artery |
| 5 Medial frontobasal artery | 19 Middle temporal artery |
| 6 Lateral frontobasal artery | 20 Cistern of the lateral cerebral fossa |
| 7 Prefrontal artery | 21 Anterior temporal artery |
| 8 Inferior frontal sulcus | 22 Pontine cistern |
| 9 Artery of the precentral sulcus | 23 Abducens nerve |
| 10 Artery of the central sulcus | 24 Pontocerebellar cistern |
| 11 Artery of the postcentral sulcus
(anterior parietal artery) | 25 Medullary cistern |
| 12 Posterior parietal artery | 26 Vertebral artery |
| 13 Artery of the angular gyrus | 27 Cerebellomedullary cistern (cisterna magna) |
| 14 Intraparietal sulcus | 28 Posterior inferior cerebellar artery, lateral branch |

Fig. 4.3. The arteries of the lateral cerebral cortex: the middle cerebral artery (1/1×). In this figure the lateral and medullary cisterns are left intact. On the lateral surface of the cerebellum one inferior and two superior cerebellar branches are illustrated (see Fig. 4.11). On the superior frontal gyrus some end branches of the anterior cerebral artery can be seen




- 1 Central sulcus
- 2 Artery of the central sulcus (branches)
- 3 Postcentral gyrus
- 4 Precentral gyrus
- 5 Artery of the precentral sulcus
- 6 Inferior frontal sulcus
- 7 Inferior frontal gyrus, triangular part
- 8 Prefrontal artery (candelabrum artery)
- 9 Lateral frontobasal artery (branched)
- 10 Anterior trunk of the middle cerebral artery (ascending frontal artery)
- 11 Anterior temporal artery (branches)
- 12 Temporopolar artery
- 13 Middle trunk of the middle cerebral artery
- 14 Posterior trunk of the middle cerebral artery
- 15 Middle temporal artery
- 16 Posterior temporal artery
- 17 Superior temporal sulcus

- 18 Temporo-occipital artery
- 19 Lateral sulcus, posterior branch
- 20 Artery of the postcentral sulcus (anterior parietal artery)
- 21 Posterior parietal artery
- 22 Artery of the angular gyrus
- 23 Angular gyrus
- 24 Intraparietal sulcus
- 25 Parieto-occipital sulcus
- 26 Lunate sulcus
- 27 Anterior occipital sulcus

Alternative subdivision

- 11+15 Anterior temporal artery
- 16 Middle temporal artery
- 18 Posterior temporal artery
- 20 Parietal artery
- 21+22 Artery of the angular gyrus

Fig. 4.4. The branches of the middle cerebral artery seen at their full extent: lateral view (1/1×). In this specimen, as in most cases, a trifurcation can be seen of the artery. The branches of the anterior (frontal) trunk are shown in *black* and *red*; the branches of the middle (parietal) trunk are in *black* only; the branches of the posterior (temporal) trunk are in *red*. The candelabrum-like branching, especially of the anterior trunk, is a common phenomenon



- | | |
|--|---|
| 1 Cistern of the lamina terminalis | 18 Medial frontobasal artery |
| 2 Optic nerve | (branch of the anterior cerebral artery) |
| 3 Cistern of the vallevula cerebri | 19 Lateral frontobasal artery |
| 4 Cistern of the chiasm | (branch of the middle cerebral artery) |
| 5 Internal carotid artery | 20 Inferior frontal gyrus, orbital part |
| (cerebral part) | 21 Temporopolar artery |
| 6 Oculomotor nerve | 22 Anterior temporal artery |
| 7 Hypophysis | 23 Inferior temporal sulcus |
| 8 Interpeduncular cistern | 24 Collateral sulcus with lateral occipital artery |
| 9 Abducens nerve | 25 Anterior temporal branches |
| 10 Trochlear nerve | 26 Occipitotemporal sulcus |
| 11 Pontine cistern | 27 Vertebral artery |
| 12 Trigeminal cistern | 28 Posterior inferior cerebellar artery, lateral branches |
| 13 Cistern of the internal acoustic meatus with facial and | 29 Posterior inferior cerebellar artery, medial branches |
| vestibulocochlear nerves | 30 Horizontal fissure of the cerebellum |
| 14 Glossopharyngeal, vagal and accessory nerves | 31 Medial middle temporal branch |
| 15 Pontocerebellar cistern | 32 Posterior temporal branches |
| 16 Medullary cistern | 33 Occipitotemporal sulcus |
| 17 Cerebellomedullary cistern | 34 Collateral sulcus |
| | 35 Lateral occipital artery |

Fig. 4.5. The arteries of the brain viewed from the basal side (1/1×). In this figure the basal, cerebellar and medullary cisterns are left intact




Fig. 4.6. The cerebral arteries viewed from the basal side (1/1x). Part of the left temporal lobe has been removed to show the sphenoid part of the middle cerebral artery and the arterial supply of the choroid plexus of the lateral ventricle. The lateral occipital artery has been interrupted to gain a clear view of the diencephalic, mesencephalic and retrosplenic branches of the posterior cerebral artery

branches are the left and right posterior cerebral arteries, which supply the posterior, medial and basal aspects of the cerebral hemisphere. The vertebro-basilar arteries also supply the brain stem and the cerebellum. It gives rise to the inferior, middle and superior cerebellar arteries (Fig. 4.11). Frontal and lateral projections of the arterial system are shown in Figs. 4.9 and 4.10.

A system of communicating arteries, known as the *circle of Willis* [18, 36], interconnects the anterior and middle cerebral arteries of both sides with the vertebro-basilar system (Figs. 4.6 and 4.14). It is located at the base of the brain and surrounds the infundibulum and the optic chiasm. It includes the *anterior communicating artery*, which interconnects the anterior cerebral arteries, immediately in front of the optic chiasm, and the two *posterior communicating arteries*, which form an anastomosis between the most distal part of the internal carotid and the posterior cerebral artery near their origin from the basilar artery.

The initial segments of the middle and anterior cerebral arteries give rise to *central arteries* (Figs. 4.6, 4.8, 4.12), which enter the brain in the anterior perforated substance (Fig. 3.4). Together with branches from the posterior communicating artery, they supply the basal ganglia, the internal capsule and the thalamus. The *middle cerebral artery* enters the sulcus lateralis. Just before this point it emits the anterior choroidal artery, which also supplies a branch to the globus pallidus (Fig. 4.8). At the surface of the insula the middle cerebral artery branches into anterior, middle and posterior trunks. The branches of the middle cerebral artery loop over the opercula and ramify over the surface of the cerebral hemisphere to supply the cerebral cortex and the adjacent white matter (Figs. 4.3, 4.4). The *anterior cerebral artery* enters the longitudinal fissure to branch on the medial aspect of the hemisphere. The anterior communicating artery, which connects the two anterior cerebral arteries, is located immediately rostral to the optic chiasm (Fig. 4.2).

The *vertebral arteries* enter the skull through the foramen magnum. They give rise to the *anterior spinal artery*, which descends in the anterior median fissure of the cord, and emit the *posterior*

or inferior cerebellar arteries. The vertebral arteries unite into the *basilar artery* at the ventral aspect of the brain stem. The basilar artery gives origin to the *anterior inferior* and *superior cerebellar arteries* and splits into the posterior cerebral arteries. The oculomotor nerve emerges between the superior cerebellar and the posterior cerebral arteries and thus marks the bifurcation of the basilar artery (Fig. 4.11). The posterior cerebellar artery makes a characteristic, caudally directed curve before it reaches the cerebellum. Both the posterior inferior and superior cerebellar arteries contribute branches to the dorsolateral brain stem.

The *posterior cerebral artery* supplies the medial aspect of the temporal and occipital lobes. The border region of the vascularization territories of the posterior and middle cerebral arteries include the temporal and occipital poles. The latter contains the posterior portion of the primary visual (striate) cortex with the representation of the fovea. Occlusion of the posterior cerebral artery thus leads to loss of peripheral vision, with maintained central vision ("tunnel vision") (Table 4.1). The borders of the arterial territories of the cerebral hemisphere do not correspond to the borders of the four major lobes distinguished in the gross anatomy of the cerebral hemispheres (Fig. 4.7). Asymmetries of the brain's arterial system are frequently observed, most often in the calibre of the vertebral, the posterior cerebral and the posterior communicating arteries.

The vertebral, basilar and posterior cerebral arteries also give rise to smaller branches, which enter the brain stem in the median sulcus and more laterally (Fig. 4.11). Branches from the basilar and posterior cerebral arteries (Fig. 4.18) enter the mesencephalon in the posterior perforated substance, located in the floor of the interpeduncular fossa (Fig. 3.12). The vascularization territories of these arteries have been thoroughly studied by Duvernoy [12]. These territories are illustrated in a number of transverse sections in which both arterial supply and venous drainage are visualized (Figs. 4.18–4.20). These figures also document the important contributions of the cerebellar arteries to the vascularization of the brain stem.





Fig. 4.7. Cortical territories of the three cerebral arteries; semidiagrammatic lateral and medial views of the left cerebral hemisphere (2/3×). The territories correspond to the vascularization pattern illustrated in Figs. 4.2–4.4. *Stippled areas:* sites of possible cerebrocerebral arterial anastomoses, mostly according to Gillilan [13]



- | | |
|--|--|
| 1 Caudate nucleus | 17 Internal carotid artery, cavernous part |
| 2 Putamen | 18 Internal carotid artery, petrous part |
| 3 Globus pallidus, external segment | 19 Internal carotid artery, cervical part |
| 4 Globus pallidus, internal segment | 20 Medial nucleus of the thalamus |
| 5 Thalamus | 21 Midline nuclei of the thalamus |
| 6 Anterior perforated substance | 22 Anterior nucleus of the thalamus |
| 7 Anterolateral central arteries, lateral branches | 23 Globus pallidus, internal segment |
| 8 Anterolateral central arteries, medial branches | 24 Tail of the caudate nucleus |
| 9 Long central artery (Heubner [16]) | 25 Anterior choroidal artery |
| 10 Anteromedial central arteries | 26 Subthalamus with posteromedial central arteries |
| 11 Anterior cerebral artery | 27 Hypothalamus with hypothalamic branch |
| 12 Posterior perforated substance | 28 Amygdaloid nucleus |
| 13 Middle cerebral artery, sphenoid part | 29 Posterior cerebral artery |
| 14 Superior hypophyseal artery | 30 Posterior communicating artery |
| 15 Inferior hypophyseal artery | 31 Basilar artery |
| 16 Internal carotid artery, cerebral part | 32 Vertebral artery |

Fig. 4.8. The central arteries from the carotid and vertebral system in a frontal view (1/1×). Substrate based on a reconstruction. The frontal section is perpendicular to the horizontal plane of Frankfurt, passing through the centre of the insula. The central arteries have been derived from different sources




Fig. 4.9. Orthogonal frontal projection of the cerebral and cerebellar arteries in situ, together with some bony landmarks and the lateral ventricles (2/3×). The projection was made parallel to the horizontal plane of Frankfurt by using a graphical reconstruction from the frontal slices of one specimen, and by cross-reference with Fig. 4.10. In this figure and the next, ample use has been made of indications by Thijssen [29]. Most vessels are illustrated only in one half of the skull; the vertebral artery is shown bilaterally. *OH*, Upper horizontal plane (Krönlein): tangential to supraorbital margin; *FH*, Horizontal plane of Frankfurt (Reid): tangential to infraorbital margin; *double arrow*, sulcus lateralis; *single arrow*: foramen magnum




Fig. 4.10. Orthogonal lateral projection of the cerebral and cerebellar arteries, together with external and bony landmarks, in a schematized composition of data from different specimens and publications ($2/3\times$). Some neural structures are also illustrated in their outlines: the left hemisphere, cerebellum, left insula, corpus callosum and ventricular system. Within the outlines of the orbita the bulbus oculi and the optic nerve are indicated. On the outer side of the figure a number of reference lines are added. In the centre, two lines tangential to the anterior (AC) and posterior (PC) commissures can be seen: the one passing above the AC and beneath the PC is part of the bicommissural line of Talairach [27] (BC); the other tangent is part of the upper horizontal line of Krönllein (OH); CM, canthus-meatus line; FH, horizontal line or plane of Frankfurt (Reid); GI, glabella-inion line; VCA, vertical tangential to anterior commissure; VCP, vertical tangential to posterior commissure




Fig. 4.11. The arteries of cerebellum, brain stem, thalamus and the corpus striatum in a lateral view (3/2×). Some arteries are slightly simplified in order to show their course and relations more clearly. The three arrow points indicate the choroïdal branches of the three choroïdal arteries. The same simplifications as in Figs. 4.2–4.6, with some slight simplifications





Fig. 4.12. Sinuses and veins of the diencephalon, brain stem and cerebellum in a lateral view (3/2×). Composite drawing from two specimens with additions from other sources. The cortical origins of the basal vein have been added, i.e. the insular veins, the deep middle cerebral vein and the anterior cerebellar veins. The tentorium has been made fully transparent and the cavernous sinus has been deprived of its lateral dural wall. The inner lateral wall of the trigeminal space has also been removed. The orbit has been opened by a sagittal cut through its centre



- | | |
|---|---|
| 1 Diploic veins | 26 Sigmoid sinus |
| 2 Superior sagittal sinus | 27 Superior petrosal sinus |
| 3 Superior cerebral veins | 28 Inferior petrosal sinus |
| 4 Parietal emissary vein | 29 Basilar plexus |
| 5 Superficial temporal veins (parietal branch) | 30 Middle meningeal veins |
| 6 Superior anastomotic vein (Trolard [30]) | 31 Cavernous sinus |
| 7 Inferior sagittal sinus | 32 Pterygoid plexus |
| 8 Superior thalamostriate vein | 33 Superior ophthalmic vein |
| 9 Superior choroidal vein | 34 Angular vein |
| 10 Internal cerebral vein | 35 Inferior ophthalmic vein |
| 11 Superficial middle cerebral vein | 36 Infraorbital foramen |
| 12 Deep middle cerebral vein | 37 Infraorbital vein |
| 13 Inferior choroidal vein | 38 Deep facial vein |
| 14 Basal vein | 39 Facial vein |
| 15 Lateral mesencephalic vein and petrosal vein | 40 Palatine vein |
| 16 Inferior anastomotic vein (Labbé [4]) | 41 Maxillary veins |
| 17 Great cerebral vein | 42 Superficial temporal veins (see no. 5) |
| 18 Straight sinus (sinus rectus) | 43 Internal jugular vein |
| 19 Inferior cerebral veins | 44 Retromandibular vein |
| 20 Confluens of the sinuses | 45 External jugular vein |
| 21 Occipital emissary vein | 46 Deep cervical vein |
| 22 Transverse sinus | 47 Internal vertebral venous plexus |
| 23 Occipital sinus | 48 Occipital vein |
| 24 Mastoid emissary vein | |
| 25 Condylar emissary vein | |

Fig. 4.13. Collateral circulation in the venous system of the head; semidiagrammatic lateral view (2/3×). Unpaired sinuses in the median plane are drawn *without outlines*; the extracranial veins draining into the internal and external jugular veins are in *black*; between the intravertebral venous plexuses a fragment of the spinal medulla can be seen. The *arrows* indicate the continuity of the superficial temporal veins

The existence of a collateral circulation is of great significance for the vascularization of the CNS. There are different types and different sites of anastomoses; moreover, the diameter of these anastomoses may differ considerably. Anastomoses between arteries can be found in relation to three arterial systems, i.e. between the two main arterial systems of the carotid and vertebral arteries and between the arterial systems of the brain and the external carotid artery.

Apart from the main arterial anastomosis between the systems of the internal carotid and the vertebral-basilar arterial system in the arterial circle of Willis, cerebro-cerebral anastomoses are present between the branches of the middle cerebral artery (Figs. 4.2, 4.14). Anastomoses between the cerebellar arteries are documented in Figs. 4.9 and 4.11.

Anastomoses with the external carotid artery occur both with meningeal and extracranial branches of this artery. Four types of anastomoses with branches of the external carotid artery are indicated with asterisks in Fig. 4.1. Orbital anastomoses with branches of the ophthalmic artery are enumerated as two special categories.

Meninges, Cisterns and Cerebrospinal Fluid Circulation

The brain is completely enclosed by three connective tissue layers: the meninges. These are, starting from the brain's surface, the *pia mater*, the *arachnoid* and the *dura mater*. The dura is also known as the pachymeninx, due to its strength and thickness, which is imparted by multiple layers of collagen tissue. The thin and loose tissue of the pia mater and the arachnoid is collectively known as the leptomeninx.

The cranial dura is merged with the periosteum of the inner table of the skull. As a consequence, the dura is firmly attached to the skull, especially at the sites of the sutures. Dural septa are located between the main divisions of the brain. In the midline, the *falx cerebri* is located between the cerebral hemispheres and the tentorium cerebelli extends between the occipital and temporal lobes of the hemisphere and the cerebellum. Venous sinuses occupy the inner and outer margins of the falx (superior and inferior sagittal sinus), the junction of the falx and the tentorium (straight sinus, or sinus rectus), and the attachment of the tentorium to the skull transverse sinus and superior petrosal sinus (Figs. 4.14–4.16). The *pia mater* closely covers the surface of the brain and intrudes into its sulci and depressions. The *arachnoid* covers the brain at a variable distance, thus creating a subarachnoidal space between the pia and the arachnoid. This space is bridged by many trabeculae. It contains the CSF. Widening of the subarachnoid space are known as the cisterns.

For the understanding of the production, circulation and drainage of CSF, the fine structure of the interface of the CSF compartments, the nervous tissue and the mesenchymal tissue of the meninges is important. The central nervous system is isolated from the rest of the body by a series of cellular barriers, which limit the flux of hydrophilic molecules between these cells. These barriers generally consist of extensive tight junctions between the cells, where the outer leaflets of the plasma membranes of two opposing cells are fused. These barriers are found in the epithelium of the choroid plexus (blood-CSF barrier), the outer (barrier) layer of the arachnoid and in the endothelium of capillaries located within arachnoid and the pia mater and nervous tissue (blood-brain barrier).

The Human Central Nervous System

A Synopsis and Atlas

Nieuwenhuys, R.; Voogd, J.; Huijzen, C. van

2008, XIV, 967 p., Hardcover

ISBN: 978-3-540-34684-5