

Contents

Part I Introduction

| | |
|---|-----------|
| 1 Introduction | 1 |
| U. Schiewer | |
| 1.1 The Baltic Sea | 1 |
| 1.1.1 Geological Background | 3 |
| 1.1.2 Climate, Hydrology and Chemistry | 4 |
| 1.1.3 Catchment Area, Eutrophication and Pollution | 6 |
| 1.1.4 Ecological Structure and Function | 10 |
| 1.2 Pelagial | 10 |
| 1.3 Benthic | 12 |
| 1.4 Conversion of Substances | 16 |
| 1.5 Spatial and Temporal Variations in the Baltic Sea | 18 |
| 1.6 Protection | 20 |
| References | 20 |
| 2 The Baltic Coastal Zones | 23 |
| U. Schiewer | |
| 2.1 Coastal Structures and Types | 23 |
| 2.2 Coastal Dynamics | 25 |
| 2.3 Ecology of Coastal Waters | 29 |
| 2.4 Anthropogenic Loads and Protection Measures | 30 |
| References | 33 |

Part II Ecological Case Studies

Part II.A Southern Baltic Coast

| | |
|---|-----------|
| 3 Darß-Zingst Boddens, Northern Rügener Boddens and Schlei | 35 |
| U. Schiewer | |
| 3.1 Introduction | 35 |
| 3.2 Darß-Zingst Boddens | 35 |

| | | |
|----------|---|------------|
| 3.2.1 | Environmental Characteristics | 35 |
| 3.2.2 | Planktonic Communities | 44 |
| 3.2.3 | Benthic Communities | 52 |
| 3.2.4 | Benthic–Pelagic Coupling and Eutrophication | 59 |
| 3.3 | Northern Rügener Boddens | 63 |
| 3.3.1 | Environmental Characteristics | 63 |
| 3.3.2 | Planktonic Communities | 65 |
| 3.3.3 | Benthic Communities | 68 |
| 3.3.4 | Benthic–Pelagic Coupling and Eutrophication | 70 |
| 3.4 | Schlei | 71 |
| 3.4.1 | Environmental Characteristics | 71 |
| 3.4.2 | Planktonic Communities | 73 |
| 3.4.3 | Benthic Communities | 76 |
| 3.4.4 | Benthic–Pelagic Coupling and Eutrophication | 78 |
| 3.5 | Conclusions and Protection Measures | 80 |
| | References | 81 |
| 4 | Greifswalder Bodden, Wismar-Bucht and Salzhaff | 87 |
| | U. Schiewer | |
| 4.1 | Introduction | 87 |
| 4.2 | Greifswalder Bodden | 87 |
| 4.2.1 | Environmental Characteristics | 87 |
| 4.2.2 | Planktonic Communities | 96 |
| 4.2.3 | Benthic Communities | 100 |
| 4.2.4 | Benthic–Pelagic Coupling and Eutrophication | 102 |
| 4.3 | Wismar-Bucht and Salzhaff | 104 |
| 4.3.1 | Environmental Characteristics | 106 |
| 4.3.2 | Planktonic Communities | 108 |
| 4.3.3 | Benthic Communities | 109 |
| 4.3.4 | Benthic–Pelagic Coupling and Eutrophication | 111 |
| 4.3.5 | Protection Measures | 111 |
| 4.4 | Conclusions and Summary | 111 |
| | References | 112 |
| 5 | The Szczecin (Oder-) Lagoon | 115 |
| | T. Radziejewska and G. Schernewski | |
| 5.1 | Introduction | 115 |
| 5.2 | Environmental Characteristics | 116 |
| 5.3 | Pelagic Communities | 120 |
| 5.4 | Benthic Communities | 121 |
| 5.5 | Eutrophication and Benthic–Pelagic Coupling | 123 |
| 5.6 | Summary and Conclusions | 125 |
| | References | 126 |

| | |
|--|------------|
| 6 Near-Shore Zones: Koserow and Tromper Wiek. | 131 |
| U. Schiewer | |
| 6.1 Introduction | 131 |
| 6.2 Environmental Characteristics | 131 |
| 6.3 Results | 132 |
| 6.4 Discussion and Conclusions | 133 |
| References. | 138 |

Part II.B Eastern Baltic Coast

| | |
|--|------------|
| 7 Gulf of Gdańsk and Puck Bay | 139 |
| L. Kruk-Dowgiałło and A. Szaniawska | |
| 7.1 Introduction | 139 |
| 7.2 Gulf of Gdańsk. | 140 |
| 7.2.1 Environmental characteristics | 140 |
| 7.2.2 Planktonic Communities | 147 |
| 7.2.3 Benthic Communities | 150 |
| 7.3 Puck Bay | 154 |
| 7.3.1 Environmental Characteristics. | 154 |
| 7.3.2 Planktonic Communities | 156 |
| 7.3.3 Benthic Communities | 157 |
| 7.3.4 Fish Communities | 159 |
| 7.4 Eutrophication and Benthic–Pelagic Coupling | 160 |
| 7.5 Conclusions | 162 |
| References. | 162 |
| 8 The Vistula Lagoon. | 167 |
| B. Chubarenko and P. Margoński | |
| 8.1 Environmental Characteristics | 167 |
| 8.1.1 Lagoon Morphometry | 167 |
| 8.1.2 Lagoon Hydrography. | 168 |
| 8.1.3 Transboundary Aspects | 169 |
| 8.1.4 Lagoon Hydrology. | 169 |
| 8.1.5 Lagoon Water Dynamics | 172 |
| 8.1.6 Hydrological–Hydrodynamical Typology of the Vistula Lagoon System | 173 |
| 8.1.7 Sediment Regime. | 174 |
| 8.2 Eutrophication | 174 |
| 8.2.1 Water Quality. | 174 |
| 8.2.2 Limitation of Primary Production | 178 |
| 8.2.3 Trophic State Index | 178 |
| 8.3 Planktonic Communities | 179 |

| | | |
|-----------|---|------------|
| 8.3.1 | Primary Production | 179 |
| 8.3.2 | Bacterioplankton | 179 |
| 8.3.3 | Phytoplankton | 180 |
| 8.3.4 | Zooplankton | 182 |
| 8.4 | Benthic Communities | 184 |
| 8.4.1 | Macrophytes | 184 |
| 8.4.2 | Macrozoobenthos | 185 |
| 8.5 | Fish and Fisheries | 187 |
| 8.6 | Conclusions and Summary | 190 |
| | References | 191 |
| 9 | The Curonian Lagoon | 197 |
| | Z.R. Gasiūnaitė, D. Daunys, S. Olenin, and A. Razinkovas | |
| 9.1 | Environmental Characteristics | 197 |
| 9.2 | Historical Overview | 201 |
| 9.3 | Phytoplankton | 202 |
| 9.4 | Zooplankton | 204 |
| 9.5 | Fish | 206 |
| 9.6 | Benthic Communities | 207 |
| 9.7 | Biological Processes and Interactions | 209 |
| 9.8 | Protection | 210 |
| 9.9 | Summary | 211 |
| | References | 211 |
| 10 | Gulf of Riga and Pärnu Bay | 217 |
| | J. Kotta, V. Lauringson, G. Martin, M. Simm, I. Kotta, K. Herkül, and H. Ojaveer | |
| 10.1 | Environmental Characteristics | 217 |
| 10.2 | Plankton Communities | 222 |
| 10.3 | Benthic Communities | 228 |
| 10.4 | Benthic–Pelagic Coupling and Eutrophication | 233 |
| 10.5 | Conclusions and Summary | 239 |
| | References | 239 |
| 11 | Haapsalu and Matsalu Bays | 245 |
| | J. Kotta, A. Jaanus, and I. Kotta | |
| 11.1 | Environmental Characteristics | 245 |
| 11.2 | Plankton Communities | 249 |
| 11.3 | Benthic Communities | 252 |
| 11.4 | Benthic–Pelagic Coupling and Eutrophication | 253 |
| 11.5 | Conclusions and Summary | 257 |
| | References | 257 |

Part II.C Northeastern Baltic Coast

| | |
|--|------------|
| 12 The Neva Estuary Ecosystem. | 259 |
| I.V. Telesh, S.M. Golubkov, and A.F. Alimov | |
| 12.1 Environmental Characteristics. | 259 |
| 12.2 Planktonic Communities | 264 |
| 12.2.1 Phytoplankton | 264 |
| 12.2.2 Bacterioplankton | 267 |
| 12.2.3 Zooplankton. | 268 |
| 12.3 Fish Communities | 271 |
| 12.4 Benthic Communities | 272 |
| 12.5 Benthic–Pelagic Coupling, “Biological Pollution” and Eutrophication. | 276 |
| 12.6 Functional Response of Aquatic Communities to Anthropogenic Stress. | 278 |
| 12.7 Conclusions | 280 |
| References. | 281 |
| 13 The Gulf of Finland | 285 |
| H. Pitkänen, J. Lehtoranta, and H. Peltonen | |
| 13.1 Introduction | 285 |
| 13.2 An Infinite Estuary Sensitive to Physical Forcing. | 286 |
| 13.3 Geomorphology and Hydrodynamics | 287 |
| 13.4 External Nutrient Loading. | 290 |
| 13.5 Water Column Nutrients | 291 |
| 13.5.1 Strong Vertical and Seasonal Variations | 291 |
| 13.5.2 Temporal Trends: Increasing for P, Decreasing for N | 294 |
| 13.5.3 Trends in Eutrophication | 296 |
| 13.6 Internal Nutrient Fluxes. | 298 |
| 13.6.1 Sediment–Water Exchange and Denitrification. | 298 |
| 13.6.2 The Role of Benthic Macrofauna. | 300 |
| 13.7 Nutrient Dynamics are Reflected in the Phytoplankton | 301 |
| 13.8 Fish Assemblages and Environmental Conditions | 302 |
| 13.9 Conclusions – Is Recovery Possible? | 304 |
| References. | 304 |

Part II.D Western Baltic Coast

| | |
|---|------------|
| 14 The Stockholm Archipelago | 309 |
| C. Hill and K. Wallström | |
| 14.1 Environmental Characteristics | 309 |
| 14.2 The Pelagic Zone | 314 |
| 14.2.1 Nutrient Cycles. | 314 |
| 14.2.2 Plankton | 315 |
| 14.2.3 Fish. | 316 |
| 14.2.4 Seabirds | 318 |
| 14.2.5 Marine Mammals | 318 |
| 14.3 Benthic Communities. | 318 |
| 14.3.1 Bottom Substrate | 318 |
| 14.3.2 Vegetation of Hard Bottoms. | 319 |
| 14.3.3 Macrozoobenthos of Hard Bottoms | 320 |
| 14.3.4 Vegetation of Soft Bottoms | 320 |
| 14.3.5 Macrozoobenthos of Soft Bottoms. | 322 |
| 14.4 Effects of Eutrophication and Other Anthropogenic Disturbances | 322 |
| 14.5 Environmental Pollutants. | 325 |
| 14.6 Physical Exploitation and Erosion of Shorelines. | 327 |
| 14.7 The Spread of Alien Species | 327 |
| 14.8 Recruitment Failure in Coastal Fish. | 328 |
| 14.9 Abnormal Mortality in Seabirds. | 328 |
| 14.10 Nature Conservation. | 328 |
| 14.11 Summary and Conclusions. | 329 |
| 14.11.1 Final Remarks | 331 |
| References. | 331 |
| 15 The Askö Area and Himmerfjärden | 335 |
| H. Kautsky | |
| 15.1 Askö Area | 335 |
| 15.1.1 Environmental Characteristics. | 335 |
| 15.1.2 Planktonic Communities | 339 |
| 15.1.3 Benthic Communities | 342 |
| 15.1.4 Long-Term Change | 348 |
| 15.1.5 No Recent Increase in Eutrophication. | 348 |
| 15.1.6 Production and Energy Flow | 352 |
| 15.2 Himmerfjärden Area | 355 |
| 15.2.1 Environmental Characteristics. | 355 |
| 15.2.2 Eutrophication | 356 |
| References. | 357 |

| | | |
|-----------|---|------------|
| 16 | Odense Fjord and Kerteminde Fjord/Kertinge Nor | 361 |
| | H.U. Riisgård, M.H. Jensen, and N. Rask | |
| 16.1 | Introduction | 361 |
| 16.2 | Case Study: Odense Fjord | 361 |
| 16.2.1 | Basic Characteristics | 361 |
| 16.2.2 | Nutrient Loads | 367 |
| 16.2.3 | Nutrients in Water and Sediment | 367 |
| 16.2.4 | Phytoplankton | 371 |
| 16.2.5 | Macrophytes | 372 |
| 16.2.6 | Benthic Filter-Feeders and Grazing Impact | 373 |
| 16.2.7 | Fish | 377 |
| 16.2.8 | Impacts of Hazardous Substances | 377 |
| 16.3 | Case Study: Kerteminde Fjord/Kertinge Nor | 378 |
| 16.3.1 | Basic Characteristics | 378 |
| 16.3.2 | Nutrients and Phytoplankton | 379 |
| 16.3.3 | Biological Structure and Nutrient Dynamics in Kertinge Nor | 380 |
| 16.3.4 | Implications of Density-Driven Currents for Predation Impact by Jellyfish | 385 |
| 16.3.5 | Density-Driven Currents and Grazing Impact of Filter-Feeding Benthos | 386 |
| 16.3.6 | Biogeochemistry, Microphytobenthos and Macrozoobenthos | 388 |
| 16.3.7 | Fish | 390 |
| 16.4 | Odense Fjord and Kerteminde Fjord/Kertinge Nor: Protection Measures and Environmental Objectives | 390 |
| 16.4.1 | International Protection Measures | 390 |
| 16.4.2 | National Protection Measures | 391 |
| | References | 392 |

Part III Synthesis

| | | |
|-----------|--|------------|
| 17 | Synthesis | 395 |
| | U. Schiewer | |
| 17.1 | Common Features and Differences | 395 |
| 17.1.1 | Type 1: Open Coast/Near Shore | 397 |
| 17.1.2 | Type 2: Big Bays | 398 |
| 17.1.3 | Type 3: Good Water Exchange with the Baltic | 399 |
| 17.1.4 | Type 4: Impeded Exchange with the Baltic | 399 |
| 17.1.5 | Type 5: Coastal Waters Dominated by River Inflow | 399 |
| 17.1.6 | Characteristics of Baltic Coastal Waters | 400 |

| | | |
|--------|--|------------|
| 17.2 | Eutrophication | 405 |
| 17.2.1 | Zooplankton | 406 |
| 17.2.2 | Protozooplankton | 406 |
| 17.2.3 | Fish Populations | 407 |
| 17.2.4 | Submersed Macrophytes | 407 |
| 17.2.5 | Sediments | 408 |
| 17.2.6 | Microbial Food Webs and Aggregates/ Fluffy-Sediment-Layer | 408 |
| 17.2.7 | Benthos | 410 |
| 17.3 | Other Threats | 410 |
| 17.4 | Buffer and Filter Capacity | 411 |
| 17.5 | Ecosystem Stability | 412 |
| 17.6 | Knowledge Gaps | 413 |
| 17.7 | Protection Measures | 413 |
| 17.8 | Conclusions and Outlook | 415 |
| | References | 416 |
| | Index | 419 |



<http://www.springer.com/978-3-540-73523-6>

Ecology of Baltic Coastal Waters

Schiewer, U. (Ed.)

2008, XIX, 430 p., Hardcover

ISBN: 978-3-540-73523-6