

Extensions

2.1 Extensions of Unipotent Groups and Isogenies

Let G_1 and G_2 be connected algebraic k -groups. A k -homomorphism $\eta : G_1 \rightarrow G_2$ is a *k-isogeny* if it is an epimorphism with finite kernel. By [77], Corollary p. 412, any element of the kernel of a separable k -isogeny is defined over the separable closure k_s of k . The groups SL_2 and PSL_2 give an example for the fact that the existence of an isogeny is not a symmetric relation. But if G_1 and G_2 are connected commutative unipotent algebraic group and $\eta : G_1 \rightarrow G_2$ is an isogeny, then there always exists an isogeny $\theta : G_2 \rightarrow G_1$ (see [89], Proposition 10, p. 176). The equivalence relation generated by isogenies can be defined as follows: G_1 and G_2 are *isogenous* if there exists a group G_3 and two isogenies $\eta_i : G_3 \rightarrow G_i$ ($i = 1, 2$), see [77], section 3, p. 417.

A main result in the context of commutative unipotent algebraic groups, defined over a perfect field k , states that any such connected k -group is k -isogenous to a direct product of Witt groups \mathfrak{W}_n (cf. [18], 6.11 p. 595). The class of Witt groups \mathfrak{W}_n is thoroughly described in [18], Chapitre V, or in [89], VII, 8. The class of unipotent algebraic groups isogenous to a Witt group is therefore the class of commutative unipotent chains in positive characteristic.

In this section we treat the theory of extensions of algebraic groups, as developed in a series of papers by Weil [98], Rosenlicht [77, 79, 82, 83], Serre [89]. The purpose is to describe an algebraic group G if a normal connected algebraic subgroup A and the factor group G/A are given.

A bit more general than the direct product are the *direct product with amalgamated central subgroup* $G = G_1 \vee G_2$ and the *direct product with amalgamated factor group* $G = G_1 \wedge G_2$, defined as follows:

2.1.1 Definition. Let G_1, G_2 be connected algebraic groups, let $Z_i \in G_i$ be isogenous connected central algebraic subgroups and let $\mu_i : H \rightarrow Z_i$ be two isogenies. The factor group $(G_1 \times G_2)/\Delta$, where

$$\Delta = \{(g_1, g_2^{-1}) \in G_1 \times G_2 : g_i = \mu_i(x) \text{ for } i = 1, 2 \text{ with } x \in H\}$$

is called the direct product with amalgamated central subgroup and is denoted by $(G_1 \vee G_2)_H$ or simply by $G_1 \vee G_2$, if H is understood.

2.1.2 Definition. Let G_1, G_2 be connected algebraic groups, let $N_i \in G_i$ be connected normal algebraic subgroups such that G_1/N_1 is isogenous to G_2/N_2 and let $\mu_i : H \rightarrow G_i/N_i$ be two isogenies. The subgroup

$$(G_1 \wedge G_2)_H = \{(g_1, g_2) \in G_1 \times G_2 : g_i N_i = \mu_i(x) \text{ for } i = 1, 2 \text{ with } x \in H\}$$

of $G_1 \times G_2$ is called the direct product with amalgamated factor group and is denoted by $(G_1 \wedge G_2)_H$ or simply by $G_1 \wedge G_2$, if H is understood.

The following characterisation of the above products follows by standard arguments.

2.1.3 Proposition. The connected algebraic group G contains two connected algebraic subgroups G_1, G_2 such that $G = G_1 G_2$, $[G_1, G_2] = 1$ and $(G_1 \cap G_2)^\circ = H$ if and only if G is isogenous to the direct product with amalgamated central subgroup $(\overline{G}_1 \vee \overline{G}_2)_{\overline{H}}$, where \overline{G}_i is isogenous to G_i and \overline{H} is isogenous to H .

The connected algebraic group G contains two connected normal algebraic subgroups N_1, N_2 such that the homomorphism $\pi : G \rightarrow G/N_1 \times G/N_2$, $\pi(x) = (xN_1, xN_2)$ is an isogeny if and only if G is isogenous to the direct product with amalgamated factor group $(\overline{G}_1 \wedge \overline{G}_2)_H$, where \overline{G}_i is isogenous to G/N_i and H is isogenous to $G/(N_1 N_2)$. \square

Now we summarize some results concerning the theory of extensions of algebraic groups. These are slightly different from the ones in the general case of abstract groups, especially in the matter of the field of definition and in the non-affine case. One of the purposes is to show that the description of an algebraic group by means of coordinate functions cries for questions of separability of the field of definition. At the end of this section we will therefore abandon the attempt of describing algebraic chains in the context of algebraic k -groups for a general field k and we will assume (mainly) that k is perfect. The principal reference are the papers [82] and [77] of Rosenlicht.

Let A and B be two connected algebraic k -groups, with A commutative and affine, and let $\phi : B \times B \rightarrow A$ be a k -rational regular factor system, i.e. an everywhere defined k -rational map satisfying the equation $\delta^2 \phi = 0$, where one defines

$$\delta^2 \phi : (x, y, z) \mapsto \phi(x, y) + \phi(xy, z) - \phi(y, z) - \phi(x, yz). \quad (2.1)$$

The following multiplication

$$(b_1, a_1)(b_2, a_2) = (b_1 b_2, a_1 + a_2 + \phi(b_1, b_2)) \quad (2.2)$$

makes $B \times A$ an algebraic k -group G_ϕ , where $1 \times A$ is a central algebraic k -subgroup of G_ϕ , the factor group $G_\phi/(1 \times A)$ is k -isomorphic to B and it is possible to establish an exact sequence

$$1 \longrightarrow A \xrightarrow{\iota} G_\phi \xrightarrow{\pi} B \longrightarrow 1 \quad (2.3)$$

of separable k -homomorphisms (cf. [77], Theorem 4, p. 413), where $\iota(a) = (1, a - \phi(1, 1))$ and $\pi(b, a) = b$. Since ι and π are separable, it is possible to identify A with the k -subgroup $1 \times A$ of G_ϕ and B with the factor group $G_\phi/(1 \times A)$. We say that G_ϕ is an *explicit central extension* of A by B , emphasizing that A is, up to a separable k -isomorphism, a *central* k -subgroup of G_ϕ . The set $C_k^2(B, A)$ of all k -rational regular factor systems from $B \times B$ to A is a commutative group with respect to the addition of maps. For any k -rational regular map $\psi : B \longrightarrow A$, the map

$$\delta^1 \psi : (x, y) \mapsto -\psi(y) + \psi(xy) - \psi(x)$$

is a k -rational regular factor system, usually called trivial. The trivial k -rational regular factor systems form a subgroup $B_k^2(B, A)$ of $C_k^2(B, A)$ and the factor group $C_k^2(B, A)/B_k^2(B, A)$ is usually denoted by $H_k^2(B, A)$.

Two explicit central extensions G_{ϕ_1}, G_{ϕ_2} of A by B , given by ϕ_1, ϕ_2 , respectively, are *k-equivalent* if there exists a rational k -isomorphism $\gamma : G_{\phi_1} \longrightarrow G_{\phi_2}$ such that the diagram

$$\begin{array}{ccccccc} 1 & \longrightarrow & A & \longrightarrow & G_{\phi_1} & \longrightarrow & B \longrightarrow 1 \\ & & id_A \downarrow & & \gamma \downarrow & & id_B \downarrow \\ 1 & \longrightarrow & A & \longrightarrow & G_{\phi_2} & \longrightarrow & B \longrightarrow 1 \end{array}$$

is commutative (see [50], 6.10, p. 363ff). This happens if and only if $(\phi_1 - \phi_2) \in B_k^2(B, A)$, hence two extensions G_{ϕ_1} and G_{ϕ_2} are equivalent if and only if ϕ_1 and ϕ_2 differ by a trivial factor system. In particular, the extension defined by a trivial factor system $\phi \in B_k^2(B, A)$ is equivalent to the direct product $G_\phi = A \times B$, which corresponds to the factor system $\varphi(x, y) = 0$ for all $x, y \in B$. In this case we say that the extension *splits*.

Thus we have a bijection between the classes of equivalent explicit central extensions and the classes of $H_k^2(B, A)$. (This bijection is indeed an isomorphism of groups, if one defines a group multiplication on the set of extensions G_ϕ , as described by the well-known methods of Baer [5]).

Now we turn to the problem of describing a connected algebraic k -group G by means of factor systems, once we know a connected central affine k -subgroup A and the corresponding factor group $B = G/A$.

Let G be a connected algebraic k -group and let A be a connected central *affine* algebraic k -subgroup of G . Then the embedding of A in G and the canonical projection π of G onto $B = G/A$ give an exact sequence

$$1 \longrightarrow A \longrightarrow G \xrightarrow{\pi} B \longrightarrow 1$$

of separable rational \mathbf{k} -homomorphisms (see [77], Theorem 4, p. 413). The question whether the group G is birationally \mathbf{k} -isomorphic to an explicit central extension G_ϕ for a suitable \mathbf{k} -rational regular factor system ϕ is answered by the existence of a \mathbf{k} -rational regular cross section. A rational *cross section* is a rational map from B into G such that $\pi\sigma = id$. Note that a rational cross section is not necessarily a regular one, indeed it could be defined only on an open dense subset of B .

Let σ be a \mathbf{k} -rational regular cross section. Since $\pi\sigma = id$, one has that $\delta^1\sigma$ is a \mathbf{k} -rational regular function from $B \times B$ to A . As A is a central subgroup of G , writing the multiplication in G additively we have

$$\begin{aligned}
& [\sigma(y) - \sigma(xy) + \sigma(x)] + [\sigma(z) - \sigma(xyz) + \sigma(xy)] + \\
& -[\sigma(z) - \sigma(yz) + \sigma(y)] - [\sigma(yz) - \sigma(xyz) + \sigma(x)] = \\
& [\sigma(y) - \sigma(xy) + \sigma(x)] + [\sigma(z) - \sigma(xyz) + \sigma(xy)] + \\
& [-\sigma(y) + \sigma(yz) - \sigma(z)] + [-\sigma(x) + \sigma(xyz) - \sigma(yz)] = \\
& [\sigma(y) - \sigma(xy) + \sigma(x)] + [-\sigma(x) + \sigma(xyz) - \sigma(yz)] + \\
& [\sigma(z) - \sigma(xyz) + \sigma(xy)] + [-\sigma(y) + \sigma(yz) - \sigma(z)] = \\
& \sigma(y) - \sigma(xy) + \sigma(xyz) - \sigma(yz) + \\
& [\sigma(z) - \sigma(xyz) + \sigma(xy)] + [-\sigma(y) + \sigma(yz) - \sigma(z)] = \\
& \sigma(y) + [\sigma(z) - \sigma(xyz) + \sigma(xy)] - \sigma(xy) + \sigma(xyz) - \sigma(yz) + [-\sigma(y) + \sigma(yz) - \sigma(z)] = \\
& \sigma(y) + \sigma(z) - \sigma(yz) + [-\sigma(y) + \sigma(yz) - \sigma(z)] = \\
& \sigma(y) + [-\sigma(y) + \sigma(yz) - \sigma(z)] + \sigma(z) - \sigma(yz) = 0.
\end{aligned}$$

Therefore $\phi = -\delta^1\sigma$ is a \mathbf{k} -rational regular factor system which makes G birationally \mathbf{k} -isomorphic to the explicit central extension G_ϕ defined by (2.3). In fact, define a \mathbf{k} -rational regular map $\rho : G_\phi \longrightarrow G$ by $\rho(b, a) = \sigma(b)a$. This turns out to be a birational isomorphism, the inverse of which is $\rho^{-1}(g) = (\pi(g), g(\sigma\pi(g))^{-1})$.

For any other \mathbf{k} -rational regular cross section τ we have, by definition, $\pi\sigma = \pi\tau = id$. This forces $(\sigma - \tau)(B) \subseteq \ker \pi$, hence $\sigma - \tau : B \longrightarrow A$ and $\delta^1(\sigma - \tau)$ is a trivial factor system. Thus the factor systems $\delta^1\sigma$ and $\delta^1\tau$ defined by two cross sections give equivalent extensions, and, up to exchanging σ with $\sigma' : x \mapsto \sigma(x)\sigma(1)^{-1}$, we can always assume that $\sigma(1) = 1$. Moreover, a group G having a \mathbf{k} -rational regular cross section σ characterises a unique class of

equivalent factor systems of $H_k^2(B, A)$, and G is birationally k -isomorphic to the direct product $A \times B$ if and only if there exists a k -rational regular section $\sigma : B \rightarrow G$ which is a homomorphism (injective since $\pi\sigma = id$).

2.1.4 Remark. As shown, the possibility that G is birationally k -isomorphic to a suitable explicit central extension G_ϕ depends on the existence of a k -rational regular cross section. In [77], Theorem 10, p. 426 (see also [83]), it is proved that, if A is k -split, then a k -rational cross section σ exists. In general, however, σ is not a regular map. In this case $\phi = -\delta^1\sigma$ is a k -rational, but not regular, factor system, defining, according to Weil's construction in [98], a *pre-group* $G(\phi)$ which is not a group, the law of composition being not defined everywhere. However, a birational map exists which transforms the law of composition of $G(\phi)$ into the one of G . (cf. [98], Théorème p. 375 or [101], Théorème 15, p. 136, [89], Lemme 8, p. 89). The group G is therefore not explicitly described, since the factor system giving $G(\phi)$ is not defined everywhere. A natural candidate to describe this situation is a *toroidal group* in the sense of Rosenlicht [80], i.e. a connected algebraic group containing no unipotent element. Tori, abelian varieties and algebraic groups with a torus as the maximal connected affine subgroup are all toroidal. By [80], Theorem 2, p. 986, any regular map $\phi : V \times W \rightarrow A$, where V and W are varieties and A is a toroidal group, has the shape $\phi(v, w) = \phi_1(v) + \phi_2(w)$ for suitable regular mappings $\phi_1 : V \rightarrow A$ and $\phi_2 : W \rightarrow A$. If ϕ is a k -rational regular factor system from $B \times B$ to A , where B is an algebraic group, then

$$0 = \phi(v, w) + \phi(vw, z) - \phi(w, z) - \phi(v, wz) =$$

$$\phi_1(v) + \phi_2(w) + \phi_1(vw) + \phi_2(z) - \phi_1(w) - \phi_2(z) - \phi_1(v) - \phi_2(wz)$$

for all $v, w, z \in B$, and from this it follows that ϕ_1 and ϕ_2 are constant maps. This shows that there exist no regular non-trivial factor systems into a toroidal group.

Let for instance E be a smooth elliptic curve defined over a finite field and let J_m be the generalized Jacobian of E defined, according to [76], Theorem 7, p. 518 in § 3, by the modulus $m = (M) + (N)$, where M, N are two distinct non-zero points of E . This means that J_m is a connected commutative algebraic group having no non-trivial affine image and containing a one-dimensional torus T as maximal affine subgroup such that J_m/T is isomorphic to E . By Proposition 1.3.3 the group J_m cannot be defined over a finite field.

According to [17], Theorem 5, one can define on the set $T \times E$ a pre-group operation by

$$(k_1, P_1) + (k_2, P_2) = (k_1 \cdot k_2 \cdot \phi(P_1, P_2), P_1 + P_2)$$

putting

$$\phi(P_1, P_2) = \frac{\ell_{P_1, P_2}(M)}{\ell_{P_1 + P_2, O}(M)} \cdot \frac{\ell_{P_1 + P_2, O}(N)}{\ell_{P_1, P_2}(N)}$$

where $\ell_{P,Q}(X) = 0$ is the equation of the line through P and Q (tangent at E if $P = Q$) and O is the zero of E . A birational map exists which transforms the law of composition of the pre-group $T \times E$ into the one of J_m , but we observe that ϕ is defined only if $P_1, P_2, \pm(P_1 + P_2) \notin \{M, N\}$. Therefore this is an example of an extension of the one-dimensional torus T by the elliptic curve E defined by a rational factor system, which cannot be defined by a regular factor system. \square

2.1.5 Remark. In [82], Theorem 1, p. 99 or Corollary 1, p. 100, Rosenlicht shows that sufficient conditions for the existence of k -rational regular cross sections are that A is k -split and unipotent and that B is affine. In contrast to this, let G be a connected nilpotent linear algebraic group defined over a separably algebraically closed non-perfect field k such that its unipotent part G_u is not defined over k . The maximal torus T of G is defined over k and G is a T -principal fiber space over G/T . Then there is no regular cross section $\sigma : G/T \rightarrow G$ that is defined over k (see [82], p. 100).

Now we give a concrete example for the situation taking for $T = \mathbf{G}_m$ a one-dimensional torus defined over a non-perfect field k of characteristic $p > 0$. Let E be a purely inseparable extension of k of degree $[E : k] = n = p^t$. We consider the connected commutative k -group ΠT of [91], Section 12.4, that we recalled in Remark 1.3.9. The group ΠT has dimension n and contains T up to a birational k -isomorphism. There exists a surjective E -homomorphism $\rho : \Pi T \rightarrow T$, the kernel $\ker \rho$ of which is the unipotent radical of ΠT , is connected, has dimension $n - 1$ and does not contain non-trivial algebraic k -subgroups of ΠT . In particular, $\ker \pi$ is not defined over k . As a connected commutative algebraic group, over E the group ΠT is isomorphic to the direct sum of its unipotent radical ΠT_u with its maximal torus T (see [8] Theorem 10.6, p. 137). Assume by contradiction that the exact sequence

$$1 \longrightarrow T \longrightarrow \Pi T \xrightarrow{\pi} \Pi T/T \longrightarrow 1$$

has a k -rational regular cross section. Then there exists also a k -regular cross section σ with $0 = \sigma\pi(0)$, and the map $\psi : g \mapsto g(\sigma\pi(g))^{-1}$ is a morphism $\Pi T \rightarrow T$ sending 0 into 0. Hence ψ is a rational homomorphism, which is separable since ψ is the identity on T (cf. [45], Theorem, p. 44) and defined over k . This implies that its kernel ΠT_u is defined over k , which is a contradiction. \square

Before leaving the questions of rationality and turning to the connections between factor systems and isogenies, we want to remark once more that the existence of a k -rational regular factor system is guaranteed if A is a unipotent group defined over a perfect field k and B is affine.

For any factor system $\phi \in \mathcal{C}^2(B, A)$ there is precisely one factor system ϕ_0 , equivalent to ϕ , satisfying $\phi_0(1, 1) = 0$. In the group G_{ϕ_0} , equivalent to G_ϕ , we have the useful identity $(b, a) = (b, 0)(1, a)$. Given a factor system $\phi \in \mathcal{C}^2(B, A)$, one can construct others in the following way. If $f : A \rightarrow A$ and

$g : B \longrightarrow B$ are rational epimorphisms we define $f\phi$ by $f\phi(x, y) = f(\phi(x, y))$ and we define ϕg by $\phi g(x, y) = \phi(g(x), g(y))$. The maps $f\phi$ and ϕg are factor systems and we have:

$$f(\phi + \psi) = f\phi + f\psi, \quad (\phi + \psi)g = \phi g + \psi g, \quad (f\phi)g = f(\phi g).$$

Moreover, we get an induced epimorphism \hat{f} from G_ϕ onto $G_{f\phi}$ and an induced epimorphism \hat{g} from $G_{\phi g}$ onto G_ϕ by:

$$\hat{f}(b, a) = (b, f(a)) \quad \text{and} \quad \hat{g}(b, a) = (g(b), a).$$

We note that f (respectively g) is an isogeny if and only if \hat{f} (respectively \hat{g}) is one.

One has $fB^2(B, A) \subseteq B^2(B, A)$ and $B^2(B, A)g \subseteq B^2(B, A)$. Thus we obtain actions of the rational endomorphisms of A , respectively B , on $H^2(B, A)$. We denote by $[\phi]$ the coset $\phi + B^2(B, A)$ and by $G_{[\phi]}$ the set of extensions equivalent to G_ϕ . For rational endomorphisms $f : A \longrightarrow A$, $g : B \longrightarrow B$ and $[\phi] \in H^2(A, B)$, one has the actions given by $f \cdot [\phi] = [f\phi]$, $[\phi] \cdot g = [\phi g]$.

As the group A is commutative, the set $\text{End}(A)$ of rational endomorphisms of A is a ring and the action of $\text{End}(A)$ on $H^2(B, A)$ just defined makes $H^2(B, A)$ an $\text{End}(A)$ -module. It must be observed, however, that in general $H^2(B, A)$ is not an $\text{End}(B)$ -module, even if B is commutative, because in general the element $\phi(g_1 + g_2) - \phi g_1 - \phi g_2$ does not belong to $B^2(B, A)$, as the following Remark shows.

2.1.6 Remark. To see concretely that the right action of $\text{End}(B)$ on $H^2(B, A)$ does not define a module structure, let $A = B = \mathbf{G}_a$ be the connected unipotent one-dimensional additive group, over a perfect field of characteristic p . It is shown in [18], II, § 3, 4.6, that $H^2(\mathbf{G}_a, \mathbf{G}_a)$ is a free left $\text{End}(\mathbf{G}_a)$ -module, having the following family of polynomials as a basis (modulo $B^2(\mathbf{G}_a, \mathbf{G}_a)$):

$$\begin{aligned} \Phi_1(x, y) &= \sum_i \frac{(p-1)!}{i!(p-i)!} x^i y^{p-i}, \\ \eta_j(x, y) &= xy^{p^j} \quad (j = 1, 2, \dots). \end{aligned}$$

Put $g_1(t) = t$ and $g_2(t) = t^p$ and consider the factor system $\theta = \eta_1(g_1 + g_2) - (\eta_1 g_1 + \eta_1 g_2)$. Then we have

$$\theta(x, y) = (x + x^p)(y + y^p)^p - xy^p - x^p y^{p^2} = x^p y^p + xy^{p^2}.$$

Since $\theta(x, y) \neq \theta(y, x)$, we infer that $\theta \notin B^2(B, A)$, i.e. $H^2(\mathbf{G}_a, \mathbf{G}_a)$ is not a right $\text{End}(\mathbf{G}_a)$ -module. Moreover, if $p > 2$ then we have $x^p y^p = \frac{1}{2}[(x + y)^{2p} - x^{2p} - y^{2p}] \in B^2(B, A)$, thus θ is equivalent to η_2 . On the other hand, for $p = 2$ we have $x^2 y^2 = \Phi_1(x, y)^2$, thus θ is equivalent to $\Phi_1^2 + \eta_2$. \square

2.1.7 Remark. The above computation shows that in characteristic 2 it can happen that the factor set $\eta_k g$ does not belong to the *left* $\text{End}(A)$ -submodule

\mathbf{M} generated by the set $\{\eta_j : j = 1, 2, \dots\}$. In odd characteristic this is not possible, because any group G_{η_j} has exponent p , whereas for $\phi \notin \mathbf{M}$ the group G_ϕ has exponent p^2 . More details on the right action of $\text{End}(B)$ on $H^2(\mathbf{G}_a, \mathbf{G}_a)$ can be found in Proposition 2.1.9. \square

2.1.8 Remark. For the basis element Φ_1 and an arbitrary p -polynomial $g(t) = \sum_i a_i t^{p^i}$ it is easy to check that

$$[\Phi_1 g] = [\tilde{g} \Phi_1],$$

where $\tilde{g}(t) = \sum_i a_i^p t^{p^i}$. Therefore the submodule of $H^2(\mathbf{G}_a, \mathbf{G}_a)$ consisting of symmetric factor systems is a two-sided module over the ring $\text{End}(\mathbf{G}_a)$, and this is basic for the fact that, given an isogeny $\gamma_1 : G_1 \rightarrow G_2$ of two-dimensional commutative unipotent algebraic groups, one finds an isogeny $\gamma_2 : G_2 \rightarrow G_1$ (see Proposition 2.1.12 (i) or [89], § VII, n. 10). More generally this is possible by the same reason for n -dimensional commutative unipotent algebraic groups. But it is by no means possible for non-commutative factor systems, as Example 2.1.14 shows.

If however we restrict our attention to the subspace generated a monomial $g(t) = at^{p^k}$ we obtain

$$\eta_j g(x, y) = ax^{p^k} \cdot (ay^{p^k})^{p^j} = a^{1+p^j} (xy^{p^j})^{p^k} = a^{1+p^j} (\eta_j(x, y))^{p^k}.$$

Putting $\tilde{g}_j(t) = a^{1+p^j} t^{p^k}$ we get $\eta_j g = \tilde{g}_j \eta_j$. \square

In the following proposition we give a general formula for the factor systems $\eta_j \in H^2(\mathbf{G}_a, \mathbf{G}_a)$, a special case of which has been used in the above Remark 2.1.6. We recall that the ring $\text{End}_k(\mathbf{G}_a)$ of k -endomorphisms of the additive group \mathbf{G}_a is isomorphic to the non-commutative ring $k[\mathbf{F}]$ of p -polynomials, where \mathbf{F} is the Frobenius homomorphism and

$$\sum_i \alpha_i \mathbf{F}^i : x \mapsto \sum_i \alpha_i x^{p^i}.$$

The following proposition shows that in odd characteristic any factor system can be derived by Φ_1 and η_1 only.

2.1.9 Proposition. *If the characteristic of the ground field is greater than 2, then in the free left $\text{End}(\mathbf{G}_a)$ -module $H^2(\mathbf{G}_a, \mathbf{G}_a)$ we have*

$$\begin{aligned} [\eta_{2k}] &= \sum_{i=0}^{k-1} \mathbf{F}^i [\eta_1 (1 + \mathbf{F}^{2(k-i)-1})] - \left(\sum_{i=0}^{2k-1} \mathbf{F}^i \right) [\eta_1] \\ [\eta_{2k+1}] &= \sum_{i=0}^k \mathbf{F}^i [\eta_1 (1 + \mathbf{F}^{2(k-i)})] + \mathbf{F}^k [\eta_1] - \sum_{i=1}^{k-1} \mathbf{F}^i [\eta_1] - \sum_{i=1}^k \mathbf{F}^{k+i} [\eta_1]. \end{aligned}$$

Proof. Put $\alpha(x) = \frac{1}{2}x^{2p}$, hence $\delta^1\alpha(x, y) = \frac{1}{2}((x+y)^{2p} - x^{2p} - y^{2p}) = x^p y^p$. The assertion follows from the fact that

$$\eta_2 = \eta_1(1 + \mathbf{F}) - (1 + \mathbf{F})\eta_1 + \delta^1\alpha$$

whereas, for any $k > 1$, we have

$$\eta_k(1 + \mathbf{F}) = (1 + \mathbf{F})\eta_k + \eta_{k+1} + \mathbf{F}\eta_{k-1}.$$

□

2.1.10 Corollary. *If the characteristic of the ground field is greater than 2, for any $\varphi \in H^2(\mathbf{G}_a, \mathbf{G}_a)$ there exist $f_0, f_1, \dots, f_n, g_1, \dots, g_n \in k[\mathbf{F}]$ such that*

$$\varphi = f_0\Phi_1 + \sum_{k=1}^n f_k\eta_1g_k.$$

□

The following Remark 2.1.11, which makes Propositions 2.1.12 and 2.1.13 particularly meaningful, plays a certain rôle in Section 4.2.

2.1.11 Remark. We illustrate here the fact that the functor $H^2(B, A)$ is contra-variant in B and co-variant in A in the special case of isogenies. The arguments and the notations are essentially those of [89], VII, 1. p. 164-165.

1) For any explicit central extension G_{ϕ_1}

$$1 \longrightarrow A_1 \longrightarrow G_{\phi_1} \longrightarrow B \longrightarrow 1$$

of A_1 by B , defined by the factor system $\phi_1 : B \times B \longrightarrow A_1$, and any isogeny $\alpha : A_1 \longrightarrow A_2$ there exists a unique (up to equivalence) explicit central extension G_{ϕ_2}

$$1 \longrightarrow A_2 \longrightarrow G_{\phi_2} \longrightarrow B \longrightarrow 1$$

and an isogeny $\alpha_* : G_{\phi_1} \longrightarrow G_{\phi_2}$, such that the following diagram commutes

$$\begin{array}{ccccccc} 1 & \longrightarrow & A_1 & \longrightarrow & G_{\phi_1} & \longrightarrow & B \longrightarrow 1 \\ & & \alpha \downarrow & & \alpha_* \downarrow & & id_B \downarrow \\ 1 & \longrightarrow & A_2 & \longrightarrow & G_{\phi_2} & \longrightarrow & B \longrightarrow 1. \end{array}$$

Explicitly we have $[\phi_2] = \alpha[\phi_1]$ and α_* is defined by $\alpha_*(x, y) = (x, \alpha(y))$. Moreover, the group G_{ϕ_2} is the factor group of $G_{\phi_1} \times A_2$ modulo the algebraic subgroup $\Delta = \{(-a, \alpha(a)) : a \in A_1\}$.

2) For any explicit central extension G_{ϕ_1}

$$1 \longrightarrow A \longrightarrow G_{\phi_1} \xrightarrow{\pi} B_1 \longrightarrow 1$$

of A by B_1 , defined by the factor system $\phi_1 : B_1 \times B_1 \longrightarrow A$, and any isogeny $\beta : B_2 \longrightarrow B_1$ there exists a unique explicit central extension G_{ϕ_2}

$$1 \longrightarrow A \longrightarrow G_{\phi_2} \xrightarrow{\pi} B_2 \longrightarrow 1$$

and an isogeny $\beta^* : G_{\phi_2} \longrightarrow G_{\phi_1}$, such that the following diagram commutes

$$\begin{array}{ccccccc} 1 & \longrightarrow & A & \longrightarrow & G_{\phi_2} & \xrightarrow{\pi} & B_2 \longrightarrow 1 \\ & & id_A \downarrow & & \beta^* \downarrow & & \beta \downarrow \\ 1 & \longrightarrow & A & \longrightarrow & G_{\phi_1} & \xrightarrow{\pi} & B_1 \longrightarrow 1. \end{array}$$

Explicitly we have $[\phi_2] = [\phi_1]\beta$ and the isogeny β^* is defined by $\beta^*(x, y) = (\beta(x), y)$. Moreover, the group G_{ϕ_2} is the algebraic subgroup of the direct product $B_2 \times G_{\phi_1}$ defined by

$$G_{\phi_2} = \{(b, g) \in B_2 \times G_{\phi_1} : \beta(b) = \pi(g)\}.$$

□

In the case where G_{ϕ_1} and G_{ϕ_2} are two central extensions

$$1 \longrightarrow A_1 \longrightarrow G_{\phi_1} \longrightarrow B_1 \longrightarrow 1$$

$$1 \longrightarrow A_2 \longrightarrow G_{\phi_2} \longrightarrow B_2 \longrightarrow 1$$

we have:

2.1.12 Proposition. *Let G_{ϕ_1} (respectively G_{ϕ_2}) be a central extension of the algebraic affine group A_1 (respectively A_2) by the (not necessarily commutative) algebraic group B_1 (respectively B_2).*

- (i) *There is an isogeny $i : G_{\phi_1} \longrightarrow G_{\phi_2}$ such that $i(A_1) = A_2$ if and only if there exist isogenies $f : A_1 \longrightarrow A_2$ and $g : B_1 \longrightarrow B_2$ with $f[\phi_1] = [\phi_2]g$.*
(ii) *If there are isogenies $f : A_1 \longrightarrow A_2$ and $g : B_2 \longrightarrow B_1$ such that*

$$[\phi_2] = f[\phi_1]g \tag{2.4}$$

then the groups G_{ϕ_1} and G_{ϕ_2} are isogenous.

Proof. (i) Let $i : G_{\phi_1} \longrightarrow G_{\phi_2}$ be an isogeny such that $i(A_1) = A_2$. Then i is given by

$$i(x_0, x_1) = (g(x_0), f(x_1) + h(x_0)),$$

where $f : A_1 \longrightarrow A_2$, $g : B_1 \longrightarrow B_2$ are isogenies and $h : B_1 \longrightarrow A_2$ is a rational regular map satisfying the equation $f\phi_1 = \phi_2g + \delta^1h$.

Conversely, given isogenies g, f as above and a rational regular map $h : B_1 \longrightarrow A_2$ satisfying $f\phi_1 = \phi_2g + \delta^1h$, the mapping $i : G_{\phi_1} \longrightarrow G_{\phi_2}$

$$(x_0, x_1) \mapsto (g(x_0), f(x_1) + h(x_0))$$

is an isogeny, since

$$\ker(i) = \{(b, a) : b \in \ker(g), f(a) + h(b) = 0\}$$

is finite, and $i(A_1)$ is clearly equal to A_2 .

(ii) By (i) we find that G_{ϕ_1} is isogenous to $G_{f_1\phi_1}$ which in turn is isogenous to $G_{f_1\phi_1g_1} = G_{\phi_2}$. \square

The next proposition shows the crucial rôle played by the Ore condition in the context of extensions of algebraic groups and isogenies.

2.1.13 Proposition. *Let A (respectively A_1, A_2) be either the Witt group \mathfrak{W}_m or the vector group $(\mathbf{G}_a)^m$. Let G_ψ (respectively G_{ϕ_1}, G_{ϕ_2}) be a central extension of A (respectively A_1, A_2) by the (not necessarily commutative) algebraic group B (respectively B_1, B_2). If $\eta_i : G_\psi \rightarrow G_{\phi_i}$ are isogenies with $\eta_i(A) = A_i$, then there exist $h_i : A_i \rightarrow A$ and $g_i : B \rightarrow B_i$ such that $h_2[\phi_2]g_2 = h_1[\phi_1]g_1$.*

Proof. By (i) there exist isogenies $f_i : A \rightarrow A_i$ and $g_i : B \rightarrow B_i$, ($i = 1, 2$), such that $f_1[\psi] = [\phi_1]g_1$ and $f_2[\psi] = [\phi_2]g_2$. It is shown in [18], V, § 3, 6.9, p. 593, that in the semigroup of isogenies of a Witt group the Ore condition holds. For a vector group this follows from [54], § 10, p. 313. Hence can find two isogenies $h_i : A_i \rightarrow A$ such that $h_1f_1 = h_2f_2$ and we obtain $h_2[\phi_2]g_2 = h_1[\phi_1]g_1$. \square

We have already mentioned that the groups SL_2 and PSL_2 show that the existence of an isogeny is not a symmetric relation. However, if there exists an isogeny from a connected commutative unipotent group G_1 onto G_2 then an isogeny from G_2 onto G_1 exists as well (see [89], Proposition 10, p. 176). Already for unipotent connected non-commutative algebraic groups this is not any more the case as the following example shows.

2.1.14 Example. In Remark 2.1.6 we denoted by $\eta_1 : \mathbf{G}_a \times \mathbf{G}_a \rightarrow \mathbf{G}_a$ the factor system defined by $\eta_1(x, y) = xy^p$. As soon as the p -polynomial g is not monomial, the factor system η_1g is no longer contained in the left $\text{End}(\mathbf{G}_a)$ -submodule generated by η_1 . Therefore a necessary condition to have the equality $f[\eta_1] = [\eta_1]g$, for some p -polynomial f , is that g is monomial, that is $g = a\mathbf{F}^k$ (see Remark 2.1.8). But in this case we have $\eta_1g = \tilde{g}\eta_1$, where $\tilde{g} = a^{1+p}\mathbf{F}^k$. This shows that, if f is a p -polynomial which is not a monomial, it cannot happen that $f[\eta_1] = [\eta_1]g$, because the left $\text{End}(\mathbf{G}_a)$ -submodule generated by η_1 is free. By Proposition 2.1.12 (i), there cannot exist an isogeny from $G_{f[\eta_1]}$ to $G_{[\eta_1]}$ whereas by the same Proposition we have an isogeny from $G_{[\eta_1]}$ to $G_{f[\eta_1]}$. \square

2.1.15 Remark. Non-central extensions of a group A by a group B are in general described by an action of B as a non-trivial group of automorphisms of A

$$a \mapsto a^b \quad (a \in A, b \in B),$$

and a mapping $F : B \times B \longrightarrow A$, satisfying

$$F(b_1 b_2, b_3) \cdot F(b_1, b_2)^{b_3} = F(b_1, b_2 b_3) \cdot F(b_2, b_3). \quad (2.5)$$

For the trivial action of B on a commutative group A , the equation (2.5) just reduces to the functional equation of a factor system describing a central extension, as in Section 2.1. With a slight abuse, we call a mapping F satisfying (2.5) a factor system. If A and B are algebraic groups, it is necessary to assume that the factor system F and all the automorphisms $a \mapsto a^b$ for all $b \in B$ are rational maps, in order to have the extension of A by B as an algebraic group.

Let B_α be the central extension

$$1 \longrightarrow B_1 \longrightarrow B_\alpha \longrightarrow B_2 \longrightarrow 1$$

defined on $B_2 \times B_1$ by the product

$$(b_0, b_1)(b'_0, b'_1) = (b_0 \cdot b'_0, b_1 + b'_1 + \alpha(b_0, b'_0))$$

and let G_ϕ be the central extension

$$1 \longrightarrow A \longrightarrow G_\phi \longrightarrow B_\alpha \longrightarrow 1$$

defined on $B \times A$ by the product

$$((b_0, b_1), a) \cdot ((b'_0, b'_1), a') = ((b_0 \cdot b'_0, b_1 + b'_1 + \alpha(b_0, b'_0)), a + a' + \phi(b_0, b_1, b'_0, b'_1)). \quad (2.6)$$

Let $H = \{(b_0, b_1, a) \in G_\phi : b_0 = 1\}$. Under the assumption that $[G, H] \leq A$ we want to find the factor system γ corresponding to the section $\tau : B_2 \longrightarrow G_\phi$, $\tau(b_0) = (b_0, 0, 0)$ of the non-central extension

$$1 \longrightarrow H \longrightarrow G_\phi \longrightarrow B_2 \longrightarrow 1$$

and we want to compare this factor system with ϕ . With the same argument mentioned in Section 2.1 for central extension, one can easily see that such a factor system $\gamma = (\gamma_1, \gamma_2) : B_2 \times B_2 \longrightarrow H$ is $\gamma = -\delta^1 \tau$. (It is remarkable that the effects of changing the section for a non-central extension are not those of adding a trivial factor system, because in this case $\delta^1(\tau - \tau') \neq \delta^1 \tau - \delta^1 \tau'$. For a concrete example see the proof of Theorem 6.4.7). A direct computation shows now that

$$\gamma(b_0, b'_0) = (b_0 \cdot b'_0, 0, 0)^{-1} \cdot (b_0, 0, 0) \cdot (b'_0, 0, 0) = (1, \alpha(b_0, b'_0), \beta(b_0, b'_0))$$

where α is the map appearing in (2.6) and

$$\begin{aligned} \beta(b_0, b'_0) &= -\phi(b_0 b'_0, 0; (b_0 b'_0)^{-1}, -\alpha(b_0 b'_0, (b_0 b'_0)^{-1})) + \\ &\phi(b_0, 0; b'_0, 0) + \phi((b_0 b'_0)^{-1}, -\alpha(b_0 b'_0, (b_0 b'_0)^{-1}); b_0 b'_0, \alpha(b_0, b'_0)). \end{aligned}$$

Therefore the group G_ϕ is isomorphic to the group defined on $B_2 \times H$ by the multiplication

$$\begin{aligned} ((b_0, b_1), a) \cdot ((b'_0, b'_1), a') &= (b_0, 0, 0) \cdot (1, b_1, a) \cdot (b'_0, 0, 0) \cdot (1, b'_1, a') = \\ &= (b_0, 0, 0) \cdot (b'_0, 0, 0) \cdot (1, b_1, a)^{(b_0, 0, 0)} \cdot (1, b'_1, a') = \\ &= (b_0 b'_0, 0, 0) \cdot (1, \alpha(b_0, b'_0), \beta(b_0, b'_0)) \cdot (1, b_1, a + \sigma_{b_0}(b_1)) \cdot (1, b'_1, a') = \\ &= (b_0 b'_0, b_1 + b'_1 + \alpha(b_0, b'_0), a + a' + \rho(b_0, b_1, b'_0, b'_1)) \end{aligned}$$

where

$$\rho(b_0, b_1, b'_0, b'_1) = \sigma_{b_0}(b_1) + \beta(b_0, b'_0) + \phi(1, b_1, 1, b'_1) + \phi(1, \alpha(b_0, b'_0), 1, b_1 + b'_1)$$

whereas for any $b_0 \in B_2$ and for any $(1, b_1, a) \in H$ the map $\sigma_{b_0} : B_1 \rightarrow A$ is a homomorphism such that $(1, b_1, a)^{b_0} = (1, b_1, a + \sigma_{b_0}(b_1))$.

Comparing the representation given by γ with the one given by ϕ we find the remarkable fact that $\gamma_1 = \alpha$ whereas ρ is in general different from ϕ . \square

The universal covering \mathbb{C}^n of an arbitrary connected commutative complex Lie group G is a decisive tool for the description of homomorphisms and extensions of connected commutative complex Lie groups. It plays a similar rôle as the Witt group \mathfrak{W}_n for connected commutative unipotent groups. For non-commutative unipotent groups unfortunately no similar tool is available.

2.2 Extensions of Commutative Lie Groups

Since any commutative connected complex Lie group is (holomorphically) isomorphic to the direct product of a linear torus $(\mathbb{C}^*)^m$, a vector group \mathbb{C}^l and a toroidal group X , the theory of commutative extensions of such Lie groups reduces to the case of extensions which are toroidal groups. These groups play a similar rôle as the connected algebraic group $G = D(G)$ with no non-trivial affine epimorphic image.

Homomorphisms and extensions of complex tori X are completely described in [7], Ch. 1, Section 5, by means of *period matrices*, the columns of which are the vectors of the lattice Λ of a suitable representation of $X = \mathbb{C}^n/\Lambda$. This method works also for connected commutative complex Lie groups $G = \mathbb{C}^n/\Lambda$ such that the complex rank of Λ is n , which we will treat now.

Let $X = \mathbb{C}^n/\Lambda$ be a connected commutative complex Lie group. If the complex rank of Λ is $m < n$, then Λ is contained in a complex subspace V of dimension m of \mathbb{C}^n . Up to a change of basis and a canonical identification of V with \mathbb{C}^m , we can see then that X is isomorphic to $\mathbb{C}^{n-m} \oplus \mathbb{C}^m/\Lambda$. From now on we assume therefore that the complex rank of Λ is n , and we say that such groups have *maximal complex rank*. Let the real rank of Λ be $n+q$, where $0 \leq q \leq n$.

Up to a change of basis we can assume that $\Lambda = \mathbb{Z}^n \oplus \Gamma$. The corresponding column matrix is

$$P = (I_n, G) = \begin{pmatrix} I_q & 0 & \hat{T} \\ 0 & I_{n-q} & \tilde{T} \end{pmatrix} \in M_{n,n+q}(\mathbb{C})$$

where the columns of $G = \begin{pmatrix} \hat{T} \\ \tilde{T} \end{pmatrix}$ are \mathbb{R} -independent generators of Γ .

In accordance to [7], p. 2, we call P the *period matrix* of X . The imaginary part of G has real rank q , because the columns of P are \mathbb{R} -independent. Up to a permutation of the vectors of the basis we can assume that the imaginary part of \hat{T} is invertible.

For $q = 0$ we have $\Lambda = \mathbb{Z}^n$, hence the group $X = \mathbb{C}^n / \mathbb{Z}^n \cong (\mathbb{C}^*)^n$ is a linear torus, whereas for $q = n$ the group X is a complex torus by definition ([7], p. 1). According to [1], 1.1.11, p. 9, if $P = (I_n, G)$ is the matrix of a \mathbb{R} -basis of the lattice Λ , the group \mathbb{C}^n / Λ is toroidal if and only the following *irrationality condition* holds:

$$\text{for any non-zero } \mathbf{v} \in \mathbb{Z}^n \text{ the vector } \mathbf{v}G \text{ is never contained in } \mathbb{Z}^q. \quad (2.7)$$

Homomorphisms of connected commutative complex Lie groups of maximal complex rank can be described in terms of period matrices. In fact, a homomorphism $f : X_1 = \mathbb{C}^{n_1} / \Lambda_1 \longrightarrow X_2 = \mathbb{C}^{n_2} / \Lambda_2$ lifts to a unique homomorphism $\hat{f} : \mathbb{C}^{n_1} \longrightarrow \mathbb{C}^{n_2}$ of \mathbb{C} -vector spaces such that $\hat{f}(\Lambda_1) \leq \Lambda_2$. This lifting defines therefore two homomorphisms

$$\rho_a : \text{Hom}(X_1, X_2) \longrightarrow \text{Hom}(\mathbb{C}^{n_1}, \mathbb{C}^{n_2}) \cong M_{n_2, n_1}(\mathbb{C})$$

$$\rho_r : \text{Hom}(X_1, X_2) \longrightarrow \text{Hom}(\Lambda_1, \Lambda_2) \cong M_{n_2+q_2, n_1+q_1}(\mathbb{Z})$$

such that

$$\rho_a(f)P_1 = P_2\rho_r(f), \quad (2.8)$$

where P_i is a period matrix of X_i ($i = 1, 2$) and where we have identified $\rho_a(f)$ and $\rho_r(f)$ with the matrices corresponding to the chosen basis of \mathbb{C}^{n_i} . The homomorphisms ρ_a and ρ_r are called the *analytic* and the *rational representation* of $\text{Hom}(X_1, X_2)$ and the equations in (2.8) are called *Hurwitz relations* (cf. [1], p. 8).

2.2.1 Proposition. *A homomorphism $f : X_1 \longrightarrow X_2$ is an isogeny if and only if $\rho_a(f)$ and $\rho_r(f)$ are square matrices with non-zero determinant. In this case there exists an isogeny $g : X_2 \longrightarrow X_1$ with $fg = \text{id}_{X_2}$ and $gf = l \text{id}_{X_1}$ where $l = |\rho_r(f)|$. In particular, the isogeny f is an isomorphism if and only if $|\rho_r(f)| = \pm 1$.*

Proof. If f is an isogeny, then $\hat{f} = \rho_a(f)$ is bijective for dimensional reasons, hence $|\rho_a(f)| \neq 0$. If we put $\Gamma = \hat{f}^{-1}(\Lambda_2)$, then $\Lambda_1 \leq \Gamma$ and Γ / Λ_1 is the kernel

of the isogeny f . If the real rank of Λ_2 were greater than the real rank of Λ_1 , then Γ/Λ_1 would be infinite. As $\rho_a(f)P_1 = P_2\rho_r(f)$ we find $\Lambda_1 = \Gamma\rho_r(f)$. Hence we have: 1) the real rank of Λ_1 is not greater than the real rank of Λ_2 , since Λ_2 has the same real rank as Γ , 2) $\rho_r(f)$ is a square matrix with non-zero determinant.

Conversely, if $\rho_a(f)$ and $\rho_r(f)$ are square matrices with non-zero determinant, then f is surjective, its kernel is discrete and Λ_1 and $\Gamma = \hat{f}^{-1}(\Lambda_2)$ have the same real rank. As the factor group Γ/Λ_1 is the kernel of f , it has to be finite, proving that f is an isogeny.

Finally, let $l = |\rho_r(f)|$ and let $R = l\rho_r(f)^{-1}$, hence R has integral entries. Since $l\rho_a(f)^{-1}P_2 = P_1R$ we can define a homomorphism $g : X_2 \rightarrow X_1$ such that $\rho_a(g) = l\rho_a(f)^{-1}$ and $\rho_r(g) = R$. Since $\rho_a(g)$ and $\rho_r(g)$ are square matrices with non-zero determinant, the homomorphism g is an isogeny and it is easy to see that $fg = \text{id}_{X_2}$ and $gf = \text{id}_{X_1}$.

In particular, if $l = \pm 1$ the isogeny f is an isomorphism. Conversely, if f is an isomorphism, then the rational representation $\rho_r(f) : \Lambda_1 \rightarrow \Lambda_2$ is an isomorphism of lattices having $\rho_r(f^{-1})$ as the inverse, hence $|\rho_r(f)| = \pm 1$. \square

Now we want to study closed subgroups and factor groups of toroidal groups as well as holomorphic commutative extensions of toroidal groups by toroidal groups.

2.2.2 Proposition. *Let $X \cong \mathbb{C}^n/\Lambda$ be a connected commutative complex Lie group of maximal rank n . For any k -dimensional connected closed commutative complex subgroup $X_1 = \mathbb{C}^k/\Lambda_1$ of maximal rank k of X there exists a period matrix P such that*

$$P = \begin{pmatrix} P_1 & \Sigma \\ 0 & P_2 \end{pmatrix}$$

where P_1 is a period matrix of X_1 and P_2 is a period matrix of the factor group X/X_1 .

Proof. Let P_1 be a period matrix of the closed subgroup $X_1 \cong \mathbb{C}^k/\Lambda_1$ of X . As X_1 is a closed subgroup of X we can construct an exact sequence

$$0 \longrightarrow \mathbb{C}^k/\Lambda_1 \xrightarrow{\iota} \mathbb{C}^n/\Lambda \xrightarrow{\pi} \mathbb{C}^{n-k}/\Lambda_2 \longrightarrow 0,$$

where Λ_2 is a lattice corresponding to the connected complex commutative Lie group $X/X_1 \cong \mathbb{C}^{n-k}/\Lambda_2$. Consider the linear maps $\hat{\iota} = \rho_a(\iota)$ and $\hat{\pi} = \rho_a(\pi)$. As $\ker \iota = \hat{\iota}^{-1}(\Lambda)/\Lambda_1$ and ι is injective we have $\hat{\iota}^{-1}(\Lambda) = \Lambda_1$ from which it follows that also $\hat{\iota}$ is injective. Furthermore by the relation $\ker \pi = \hat{\pi}^{-1}(\Lambda_2)/\Lambda = \iota(\mathbb{C}^k/\Lambda_1) = (\hat{\iota}(\mathbb{C}^k) + \Lambda)/\Lambda$ we have $\hat{\pi}^{-1}(\Lambda_2) = \hat{\iota}(\mathbb{C}^k) + \Lambda$, which yields $\ker \hat{\pi} = \hat{\iota}(\mathbb{C}^k)$. Consequently $\hat{\iota}$ and $\hat{\pi}$ define an exact sequence

$$0 \longrightarrow \mathbb{C}^k \xrightarrow{\hat{\iota}} \mathbb{C}^n \xrightarrow{\hat{\pi}} \mathbb{C}^{n-k} \longrightarrow 0.$$

By the relation $\hat{\pi}^{-1}(\Lambda_2) = \hat{\imath}(\mathbb{C}^k) + \Lambda$ we have that the homomorphism $\hat{\pi}|_\Lambda : \Lambda \longrightarrow \Lambda_2$ is surjective and $\ker \hat{\pi}|_\Lambda = \hat{\imath}(\Lambda_1)$. This defines an exact sequence

$$0 \longrightarrow \Lambda_1 \xrightarrow{\hat{\imath}} \Lambda \xrightarrow{\hat{\pi}} \Lambda_2 \longrightarrow 0.$$

Since Λ_2 is a free commutative group we get $\Lambda = \imath(\Lambda_1) \oplus \Gamma$ where $\Gamma \cong \Lambda_2$. Up to a change of basis of the spaces \mathbb{C}^k , \mathbb{C}^n and \mathbb{C}^{n-k} we can assume that $\hat{\imath} = \rho_a(1) = \begin{pmatrix} I_k \\ 0 \end{pmatrix}$, $\hat{\pi} = \rho_a(\pi) = \begin{pmatrix} 0 & I_{n-k} \end{pmatrix}$, and we can choose a period matrix P of X such that $P = \begin{pmatrix} P_1 & \Sigma \\ 0 & A \end{pmatrix}$, where the columns of the matrix $\begin{pmatrix} \Sigma \\ A \end{pmatrix}$ are \mathbb{R} -independent \mathbb{Z} -generators of Γ . Furthermore, fixing a period matrix P_2 of X/X_1 , up to a change of generators of Γ we can assume that $\rho_r(\pi) = \begin{pmatrix} 0 & I_{n+q-k-q_1} \end{pmatrix}$, where $n+q$ (respectively n_1+q_1) is the real rank of X (respectively of X_1). Now, by the Hurwitz relations we have

$$\begin{pmatrix} 0 & I_{n-k} \end{pmatrix} \begin{pmatrix} P_1 & \Sigma \\ 0 & A \end{pmatrix} = P_2 \begin{pmatrix} 0 & I_{n+q-k-q_1} \end{pmatrix}$$

from which it follows that $A = P_2$. □

Now we look for closed linear subtori of a connected commutative complex Lie group $X \cong \mathbb{C}^n/\Lambda$ of maximal rank n with period matrix $P = (I_n \ G)$. Denote by $H = H(l_1, \dots, l_{n-m})$ the m -dimensional subspace of \mathbb{C}^n defined by

$$H = \{(z_1, \dots, z_n) \in \mathbb{C}^n : z_{l_k} = 0 \text{ for } l_k \in \{1, \dots, n\} \text{ and } k = 1, \dots, n-m\}$$

and let $C_H(P)$ be the matrix obtained from P in the following way: we cancel in P any row with exception of those labeled by l_1, \dots, l_{n-m} as well as any of the first n columns with exception of those labeled by l_1, \dots, l_{n-m} . Clearly $C_H(P) = (I_{n-m} \ G')$, with $G' \in M_{n-m,q}(\mathbb{C})$.

2.2.3 Proposition. *Let $X \cong \mathbb{C}^n/\Lambda$ be a connected commutative complex Lie group of maximal rank n and let $P = (I_n \ G)$ be a period matrix of X . If the columns of $C_H(P)$ are \mathbb{R} -independent, then $X_1 = (H + \Lambda)/\Lambda$ is a closed linear subtorus of X .*

Proof. Let $X_2 \cong \mathbb{C}^{n-m}/\Lambda_2$ be the connected commutative complex Lie group of maximal rank $n-m$ having $C_H(P)$ as a period matrix and let $\hat{f} : \mathbb{C}^n \longrightarrow \mathbb{C}^{n-m}$ be the homomorphism defined by $\hat{f}(z_1, \dots, z_n) = (z_{l_1}, \dots, z_{l_{n-m}})$. Since $\hat{f}(\Lambda) \leq \Lambda_2$, a homomorphism $f : X \longrightarrow X_2$ is induced such that X_1 is the kernel. This proves that X_1 is a closed subgroup. In order to prove that X_1 is a linear torus we show that $H \cap \Lambda$ has real rank m . This follows from the fact that the columns of $C_H(P)$ are \mathbb{R} -independent, hence no non-trivial linear combination of the columns l_1, \dots, l_{n-m} of the matrix P with integral (or even real) coefficients enters in H . □

2.2.4 Remark. The above proposition shows that a toroidal group X with period matrix

$$P = (I_n \ G) = \begin{pmatrix} I_q & 0 & \hat{T} \\ 0 & I_{n-q} & \tilde{T} \end{pmatrix} \in M_{n,n+q}(\mathbb{C})$$

(such that the imaginary part of \hat{T} is invertible) contains a closed linear subtorus L of dimension $n - q$ corresponding to the submatrix $P_1 = I_{n-q}$, because the submatrix $C_H(P) = (I_q \ \hat{T})$ is the period matrix of a complex torus. Hence L is a maximal closed linear subtorus of X . For instance, in the three-dimensional toroidal group X having

$$P = \begin{pmatrix} 1 & 0 & 0 & i & i \\ 0 & 1 & 0 & i\sqrt{2} & 0 \\ 0 & 0 & 1 & 0 & i\sqrt{2} \end{pmatrix}$$

as a period matrix, the three subgroups $H(2, 3)$, $H(1, 3)$ and $H(1, 2)$ are one-dimensional maximal closed linear subtori. Thus X is a \mathbb{C}^* -fiber bundle over the complex tori defined by the period matrices

$$C_{H(2,3)} = \begin{pmatrix} 1 & 0 & i\sqrt{2} & 0 \\ 0 & 1 & 0 & i\sqrt{2} \end{pmatrix}, \quad C_{H(1,3)} = \begin{pmatrix} 1 & 0 & i & i \\ 0 & 1 & 0 & i\sqrt{2} \end{pmatrix},$$

$$C_{H(1,2)} = \begin{pmatrix} 1 & 0 & i & i \\ 0 & 1 & i\sqrt{2} & 0 \end{pmatrix}$$

□

Let $X_1 = \mathbb{C}^{n_1}/\Lambda_1$, $X_2 = \mathbb{C}^{n_2}/\Lambda_2$ be connected commutative complex Lie groups of maximal ranks n_1, n_2 and let P_1, P_2 be the corresponding period matrices. Let

$$0 \longrightarrow X_1 \longrightarrow X \longrightarrow X_2 \longrightarrow 0$$

be an exact sequence of connected commutative complex Lie groups. By Proposition 2.2.2 we find a basis such that the corresponding period matrix is

$$P = \begin{pmatrix} P_1 & \Sigma \\ 0 & P_2 \end{pmatrix} \in M_{n,n+q_1+q_2}(\mathbb{C}).$$

Conversely to each matrix of this form there corresponds a toroidal group X containing a closed subgroup X_1 having P_1 as a period matrix and such that X/X_1 is isomorphic to a toroidal group X_2 having P_2 as a period matrix. In fact, X_1 is the kernel of the homomorphism $f : X \longrightarrow X_2$ which lifts to $\hat{f}(z_1, \dots, z_n) = (z_{n_1+1}, \dots, z_n)$.

As a consequence of Hurwitz relations we find that $P = \begin{pmatrix} P_1 & \Sigma \\ 0 & P_2 \end{pmatrix}$ and $Q = \begin{pmatrix} P_1 & \Sigma \\ 0 & P_2 \end{pmatrix}$ define equivalent extensions if and only if a matrix $A \in M_{n_1, n_2}(\mathbb{C})$ and a matrix $M \in M_{n_1+q_1, n_2+q_2}(\mathbb{Z})$ exists such that

$$\begin{pmatrix} I_{n_1} & A \\ 0 & I_{n_2} \end{pmatrix} \begin{pmatrix} P_1 & \Sigma \\ 0 & P_2 \end{pmatrix} = \begin{pmatrix} P_1 & \Sigma' \\ 0 & P_2 \end{pmatrix} \begin{pmatrix} I_{n_1+q_1} & M \\ 0 & I_{n_2+q_2} \end{pmatrix}.$$

The period matrix $P = \begin{pmatrix} P_1 & \Sigma \\ 0 & P_2 \end{pmatrix}$ defines therefore a split extension of X_1 by X_2 if and only if $\Sigma = P_1 M - A P_2$ with $A \in M_{n_1, n_2}(\mathbb{C})$ and $M \in M_{n_1+q_1, n_2+q_2}(\mathbb{Z})$. Moreover, if the period matrix $P = \begin{pmatrix} P_1 & \Sigma \\ 0 & P_2 \end{pmatrix}$ is such that $\Sigma = P_1 M - A P_2$ with $M \in M_{n_1+q_1, n_2+q_2}(\mathbb{Q})$, then P defines an extension of X_1 by X_2 which is isogenous to a split one. An isogeny $f : X_1 \rightarrow X_2$ is given by $\rho_a(f) = \begin{pmatrix} l I_{n_1} & 0 \\ 0 & I_{n_2} \end{pmatrix}$ and $\rho_r(f) = \begin{pmatrix} l I_{n_1+q_1} & 0 \\ 0 & I_{n_2+q_2} \end{pmatrix}$, where $l \in \mathbb{Z}$ is such that lM has integral entries.

Hence we have the following

2.2.5 Proposition. *Let X_1, X_2 be connected commutative complex Lie groups of maximal rank n_1, n_2 and let P_1, P_2 be the corresponding period matrices. The period matrix*

$$P = \begin{pmatrix} P_1 & \Sigma \\ 0 & P_2 \end{pmatrix} \in M_{n, n+q_1+q_2}(\mathbb{C}) \quad (n = n_1 + n_2)$$

defines an extension of X_1 by X_2 which is isogenous to a split one, via an isogeny f such that $\rho_a(f) = \begin{pmatrix} l I_{n_1} & 0 \\ 0 & I_{n_2} \end{pmatrix}$ and $\rho_r(f) = \begin{pmatrix} l I_{n_1+q_1} & 0 \\ 0 & I_{n_2+q_2} \end{pmatrix}$, if and only if $\Sigma = P_1 M - A P_2$ with $A \in M_{n_1, n_2}(\mathbb{C})$ and $M \in M_{n_1+q_1, n_2+q_2}(\mathbb{Q})$, where $l \in \mathbb{Z}$ is such that lM has integral entries. \square

Extensions of complex tori X_1 and X_2 which are not isogenous to a split analytic extension $X_1 \oplus X_2$ are called *Shafarevich extensions* in [7], Ch. 1, § 6, p. 23. Hence it seems for us to be natural to call also non-split analytic extensions of a toroidal group by a toroidal group Shafarevich extensions.

If X_1 and X_2 are abelian varieties, Shafarevich extensions of X_1 by X_2 are not abelian varieties and hence provide a wide class of non-projective complex tori, since an abelian variety is a complex torus admitting a holomorphic embedding into some projective space (cf. [7], p. xiii).

Algebraic Groups and Lie Groups with Few Factors

Di Bartolo, A.; Falcone, G.; Plaumann, P.; Strambach, K.

2008, XVI, 212 p., Softcover

ISBN: 978-3-540-78583-5