
Amygdala and Fear

2

The Amygdala and the Neural Pathways of Fear

Jacek Dębiec and Joseph LeDoux

CONTENTS

INTRODUCTION
THE INTRA-AMYGDALA MICROCIRCUITRY OF FEAR
SYNAPTIC PLASTICITY AND ITS MOLECULES
FEAR MEMORIES: THE QUESTION OF PERSISTENCE
EXTINCTION OF FEAR: THE POWER OF CORRECTIVE
EXPERIENCE
ACTIVELY COPING WITH FEAR
FROM ANIMAL TO HUMAN AMYGDALA
CONCLUSIONS
REFERENCES

Abstract

Fear and anxiety are evolutionarily developed responses to perceived or anticipated threat. They involve behavioral, autonomic, and endocrine alterations aimed at increasing an organism's chances of survival. Excessive or uncontrolled fear and anxiety may lead to anxiety disorders. Animal and human studies indicate the critical role of the amygdala in adaptive and maladaptive fear. Recent advances elucidating the organization of the neural circuitry and molecular mechanisms of fear provide new insights in normal as well as pathological fear. In this chapter, we review the microcircuitry of the amygdala with a special emphasis on its relevance to fear processing and fear learning. We also discuss recent developments in understanding the basic molecular mechanism of fear. Finally, we address some of the implications of amygdala research for developing novel therapeutic approaches to maladaptive fear and anxiety.

From: *Post-Traumatic Stress Disorder: Basic Science and Clinical Practice*

Edited by: P. J. Shiromani et al., DOI: 10.1007/978-1-60327-329-9_2

© Humana Press, a part of Springer Science+Business Media, LLC 2009

Key Words: Amygdala, anxiety, extinction, fear, fear conditioning, learning, memory, memory consolidation, memory reconsolidation, synaptic plasticity.

INTRODUCTION

Research on neural mechanisms of fear and anxiety has advanced significantly in recent decades. Animal models have been particularly useful in characterizing the microanatomy as well as the cellular and molecular mechanisms of fear and anxiety. More recently, animal studies have been complemented by a growing body of human research, especially involving functional imaging. Both avenues of investigation point at the key role of the amygdala in processing fear.

Fear is a natural, evolutionarily developed response to environmental threats. It includes autonomic and endocrine changes supporting defensive behaviors, such as fighting, fleeing, or immobility (freezing). Physiological adjustments allow increased blood flow and energy supply to skeletal muscles and the brain. These alterations support actions aimed at increasing the organism's chances of survival. Information about natural threats have been evolutionarily hardwired into animal brains, which appear to selectively respond to relevant environmental factors, such as sights, sounds, or odors of common predators; specific social behaviors of conspecifics; and painful or intense stimuli (e.g., sound of thunder, etc.). These natural factors elicit innate fear. However, an individual has to learn through experience about a variety of other possible threats. While innate pre-programmed fear reactions are inherited, acquired fear responses result from a capacity of an organism to learn and remember cues associated with danger experienced throughout life. Whereas fear is considered to be a response to an actual danger and is typically triggered by specific stimuli, anxiety is a state of preparation for a predicted threat, which can be real or imaginary.

Maladaptive fear and anxiety occur in anxiety disorders. Although anxiety disorders may involve innate mechanisms that unfold during life, such as a tendency for extreme shyness (1), fear learning contributes significantly to many anxiety pathologies (2–6). Thus, defining neural networks as well as cellular and molecular pathways underlying fear learning is crucial for better understanding of pathogenesis of anxiety disorders and for development of new treatment approaches.

One of the most commonly and successfully used experimental models of fear learning is Pavlovian fear conditioning (7). In this procedure, a neutral event (a conditioned stimulus, CS), such as tone, is paired with an aversive event (an unconditioned stimulus, US), such as a mild electric shock (Fig. 1) (6,8,9). Once the CS and the US are paired, the CS acquires an ability to elicit behavioral, autonomic, and endocrine fear responses. These responses are expressed automatically on subsequent exposures to the CS. Fear conditioning has been observed in a variety of species, ranging from insects and worms to birds and mammals.

Animal studies using fear conditioning demonstrate a unique and powerful character of fear learning. First, fear conditioning occurs very quickly. Usually, a single pairing of the CS with the US is sufficient to establish a memory. Second, once learned, conditioned fear responses persist, often remaining throughout the




Fig. 1. Auditory fear conditioning. In auditory fear conditioning, a neutral stimulus (conditioned stimulus, CS), such as tone, is paired with an aversive event (unconditioned stimulus, US), often a mild electric shock to the foot pads. As a consequence, subsequent presentations of the CS trigger fear responses, such as immobility or freezing

life of an organism. Third, defensive responses to stimuli previously associated with aversive events may weaken or extinguish through experiences that show that the CS no longer predicts harm. However, the original conditioning can frequently be recovered either spontaneously or as a result of a new stressful experience months or years after it has been extinguished. Fourth, fear motivates other kinds of behaviors, such as approach and avoidance. Avoidance can be adaptive, but in anxiety disorders avoidance often takes on a maladaptive role, with the patient successfully avoiding fear and anxiety but at the expense of failing to perform routine life roles.

THE INTRA-AMYGDALA MICROCIRCUITRY OF FEAR

The amygdala was named by the nineteenth century German anatomist Karl Burdach for the almond-like (in Greek almond: *amygdale*) shape of one of its subregions (10). Although the amygdala is a complex structure involved in a variety of functions, overwhelming evidence shows the critical role of the amygdala in fear (6,11–18), as well as in fear and anxiety pathologies (19–22). The role of the amygdala in fear is ubiquitous in vertebrate species (6).

The amygdala is located bilaterally deep inside the temporal lobe. It consists of several distinct groups of cells organized in nuclei (23) (Fig. 2). The regions




Fig. 2. Amygdala inputs, internal connections, and outputs. The amygdala consists of a group of nuclei. The nuclei most relevant to fear conditioning are the lateral (LA), basal (B), and central (CE) nuclei of the amygdala, as well as the subgroup of neurons known as intercalated cells (ITCs). The LA is the main sensory input area. It receives inputs from the thalamus and the cortex; thus, it is responsible for linking the CS with the US. The basal nucleus receives projections from the hippocampus and entorhinal and polymodal associative cortices, areas that may convey information about the environmental context in which the fearful event is occurring. The CE is the common output area controlling expression of behavioral, autonomic, and endocrine fear responses. The LA is connected with the CE in direct and indirect (via the basal nucleus and the ITCs) ways. The ITCs have inhibitory control over the CE. Major identified neuromodulators involved in fear regulation include NE norepinephrine, DA dopamine, *ACh* acetylcholine, *5HT* 5-hydroxytryptophan (serotonin)

most relevant to fear conditioning are the lateral (LA), basal (B), and central (CE) nuclei, as well as a distinct subgroup of neurons known as the massa intercalata or intercalated cells (ITCs) (6,18,23–26)

The LA is considered to be a sensory gateway of the amygdala. It receives inputs from all sensory modalities, including visual, auditory, tactile, olfactory, and gustatory, as well as from fibers transmitting pain (11). Although other amygdala nuclei also receive some sensory afferents, the LA is the main region where the sensory pathways converge. Studies using fear conditioning demonstrate that the LA is responsible for linking information about the neutral cue (CS) with that of the noxious stimulus (US). One of the most thoroughly investigated variants of fear conditioning is auditory fear conditioning, whereas a CS consists of a single tone. Fear conditioning studies reveal that auditory information reaches the LA through two distinct sensory inputs: thalamic and cortical (6,11). The thalamic pathway conveys a rapid but imprecise auditory signal from extralemnic areas, whereas the cortical pathway delivers a refined representation




Fig. 3. Two roads to fear. Sensory information reaches the LA (lateral nucleus) through two independent sensory inputs: The “low road” or thalamic pathway provides the amygdala with a rapid but imprecise signal, while the “high road” delivers a more complex and detailed representation derived from cortical association areas responsible in humans for conscious processing. The two-road model of signal transmission illustrates how fear responses can be initiated (by the direct low road) before we are aware of the eliciting stimulus

from associative cortices, which in humans are responsible for conscious processing. This last pathway, however, includes additional synaptic connections, which results in longer transmission. The “two-roads” model of signal transmission is used to illustrate how fear responses can be triggered (by thalamic road) before we are aware of the initiating stimulus (Fig. 3) (6).

While the lateral nucleus is believed to be the main sensory gateway, the central nucleus is considered to be the major output region (27–29). The CE projects to brain stem regions and through these projections controls expression of fear responses, including some behavioral responses, such as freezing, as well as autonomic and endocrine reactions. In addition, the CE is responsible for activating amine modulatory systems, such as adrenergic, serotonergic, dopaminergic, and cholinergic systems (11,18,23) (Fig. 2).

The major input and output amygdala regions, the lateral and the central nuclei, respectively, are connected through direct and indirect routes (23,30). The indirect routes are believed to be major communication channels between the both nuclei and involve connections from LA to the basal nucleus and the ITCs, both of which project to CE. The basal nucleus also receives inputs from the hippocampus and entorhinal and polymodal associative cortices and delivers

information about the environmental context in which the threat is occurring (23,24). In addition to fibers descending to the CE, the basal nucleus also projects to striatal areas. These outputs are believed to control instrumental behaviors (23,31,32). The ITC network extends from the rostral amygdala to the anterior commissure (33) and consists of clusters of γ -aminobutyric acid (GABA) or GABA-ergic neurons. These neurons control the flow of activity from the LA and basal nucleus to the CE by way of forward inhibition (34).

SYNAPTIC PLASTICITY AND ITS MOLECULES

Fear conditioning results in alterations in the strength of synaptic signaling in the amygdala (Fig. 4). Although synaptic plasticity occurs in other amygdala regions receiving some sensory inputs, such as the CE (35) and basal nucleus




Fig. 4. Signal transduction pathways involved in fear conditioning in the lateral amygdala. Consolidation of fear conditioning is initiated by the activation of receptors at the postsynaptic neuron (1). This in turn triggers cascades of intracellular events that regulate the activity of mitogen-activated protein kinase (MAPK) (2). MAPK controls transcription factors, such as CREB (cyclic adenosine monophosphate response element-binding) proteins, which regulate transcription of certain genes (3). Gene activation leads to new RNA (ribonucleic acid) and new protein synthesis and thus structural changes in the postsynaptic neuron (4). Plausible mechanisms of retrograde signaling (5) from the postsynaptic neuron induce changes in the presynaptic neuron (6). cAMP cyclic adenosine monophosphate, NO nitric oxide, PKA protein kinase A, TrkB neurotrophic tyrosine kinase receptor 2, mGluR5 metabotropic glutamate 5 receptor, NMDAR N-methyl D-aspartate receptor, AMPAR alpha-amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid receptor, PKC protein kinase C, CaMKII calcium/calmodulin-dependent protein kinase II, NOS nitric oxide system. L-VGCC L-type voltage-gated calcium channel

(36), the main site of synaptic changes underlying learning and memory is the LA (13,24,25,37–41). The LA is a site where the CS and the US pathways converge. Most of the inputs projecting onto the amygdala are excitatory and release glutamate binding to NMDA (*N*-methyl *D*-aspartate) and AMPA (α -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid) receptors localized on principal neurons. These neurons in turn transmit the information to other amygdala regions.

The cellular hypothesis of fear conditioning posits that relatively weak CS inputs become strengthened by the cooccurrence of the US, which is capable of eliciting robust responses in the LA. The activation of postsynaptic NMDA receptors on principal neurons by the CS signaling pathways in the context of the strong depolarization by the US inputs triggers calcium influx (42) (Fig. 4). This initiates cascades of intracellular processes involving second messengers and protein kinases, which lead to activation of transcription factors, gene expression, new protein synthesis, and synaptic alterations (24,37,39–41). Newly synthesized proteins strengthen synaptic connections. For example, it has been demonstrated that fear conditioning stimulates trafficking of AMPA receptors to synapses in LA (43). The insertion of new glutamatergic receptors increases the postsynaptic response elicited by presynaptic inputs.

Synaptic plasticity and learning in the amygdala are regulated by a variety of modulatory systems. Identified neuromodulators include the catecholamines noradrenaline (44–46) and dopamine (47,48) as well as glucocorticoids (49,50), serotonin (51), nitric oxide (52), BDNF (brain-derived neurotrophic factor) (53), endocannabinoids (54), neuropeptide Y (55,56), and others. These neuromodulators may facilitate changes on principal neurons or inhibitory GABA-ergic interneurons or both.

FEAR MEMORIES: THE QUESTION OF PERSISTENCE

Synaptic changes triggered by experience lead to the consolidation of the new learning. However, before consolidation is completed, interference with any of the stages of underlying intracellular events, from activation of the receptors to new protein synthesis, impairs the memory. According to the standard view of memory consolidation (57), once synaptic alterations are stabilized, memories become immune to interference.

The standard memory consolidation model has been successfully used in clinical studies. For example, it is well established that noradrenergic signaling enhances learning, especially emotional learning (57), and two pilot studies have demonstrated that administration of the β -adrenergic receptor antagonist propranolol shortly after trauma decreases the risk of post-traumatic stress disorder (PTSD) (58,59).

The traditional view of consolidation, that once memories are consolidated they persist in an unaltered state (57), was first challenged by studies in the 1960s showing that reactivation of a consolidated memory through retrieval renders this memory susceptible to amnesic treatments (60–62). Recent studies

Reconsolidation Processes Enable Modification of the Memory




Fig. 5. Reconsolidation processes enable modification of the memory. Reactivation of the consolidated memory renders it labile and susceptible to interferences. This enables the modification of the existing memory trace by the circumstances accompanying the memory retrieval

ultimately challenged the consolidation theory, providing compelling evidence that memory reactivation triggers another round of synaptic plasticity, a phenomenon known as *memory reconsolidation* (Fig. 5) (62–71). It has been demonstrated that reconsolidation of auditory fear conditioning in the LA involves NMDA receptor activation (66,72), protein kinases (73–75), expression of immediate early gene *Zif268* (76), and new protein synthesis (63,77,78) and is modulated by noradrenergic signaling (45).

One of the implications of reconsolidation research is that even well-consolidated memories may be altered. In particular, a reconsolidation model may be helpful in developing new treatments of fear pathologies, such as PTSD or specific phobias (62,65,69,70,79–81). Blocking reconsolidation may be helpful in attenuating learned fear responses and thus reducing the debilitating impact of traumatic experiences.

EXTINCTION OF FEAR: THE POWER OF CORRECTIVE EXPERIENCE

Learned fear responses may be attenuated by repeated exposure to the fear-arousing stimulus in a safe or neutral context (82,83). This phenomenon, referred to as fear extinction, forms a theoretical basis of exposure therapies (84–86). Extinction is a form of learning in which an organism learns that cues previously associated with a fearful event no longer predict danger. Extinction learning involves an interacting circuitry that includes the amygdala, medial prefrontal cortex, and the hippocampus (87–96). Studies in humans have found involvement of the amygdala and prefrontal cortex in extinction (97).

As a form of learning, extinction shares with fear conditioning similar molecular mechanisms. Specifically, glutamatergic signaling is required to initiate synaptic plasticity processes. Animal studies have demonstrated that enhancing glutamatergic stimulation by using the NMDA receptor agonist d-cycloserine facilitates extinction learning via the amygdala (98). This was further applied in human studies showing that exposure therapy in conjunction with d-cycloserine facilitates fear extinction (99–101).

ACTIVELY COPING WITH FEAR

Successfully extinguished fears often spontaneously recover or may be reinstated by a new traumatic event. Extinction learning is based on a passive exposure to fear-related stimuli. In contrast, a new study in rodents showed that active coping with fear may produce enduring reduction of fear. This new learning paradigm is referred to as *escape from fear* (102,103). In escape-from-fear learning, an organism learns to perform active behaviors that eliminate a fearful stimulus and thus reduce fear. The circuitry underlying escape-from-fear learning involves a circuit switch in the amygdala that, instead of transmitting signaling from the LA to the CE, directs information to the basal nucleus, which through its projections to the striatum and cortex controls actions (14). A recent study in humans using an active coping task found evidence for the involvement of not only the same emotion regulation circuit as extinction—the amygdala and medial prefrontal cortex—but also the striatum (Schiller D, Cain CK, Kuhlman K, LeDoux JE, Phelps EA, unpublished data). One implication of these findings is that therapies actively engaging patients may produce more enduring effects (104–106).

FROM ANIMAL TO HUMAN AMYGDALA

Recent studies using brain-imaging techniques have supported earlier animal research depicting the amygdala as a key structure involved in fear and anxiety (Fig. 6). The amygdala has been implied in adaptive fear (97,107,108) as well as in pathological fear and anxiety (21,109–112). In particular, the amygdala has been implied in PTSD (109–111) and phobias (112,113).

In addition, pharmacological approaches developed using animal models offer promising results in treating human fear and anxiety pathologies (58,59,81, 99,101).

CONCLUSIONS

Recent studies significantly advanced our knowledge about the organization of neural circuits and cellular and molecular mechanisms underlying fear and fear learning. This has been accomplished in major part thanks to the use of animal models, which allow insights into the microcircuitry and molecular mechanism of fear. Animal research forms a foundation for further human studies and provides clues about possible future therapeutic approaches.




Fig. 6. Functional magnetic resonance imaging (fMRI) of fear learning in the human brain. *Top:* Structural magnetic resonance image (MRI) of the human brain (the box delineates an area containing the amygdala). **a** Conditioned fear: fMRI showing amygdala activation by the conditioned stimulus (CS) following pairing with the unconditioned stimulus (US). **b** Instructed fear: fMRI showing amygdala activation by a CS that was not directly paired with the US but instead the subjects were instructed about the US. **c** Observational fear learning: fMRI showing amygdala activation by a CS after the subjects observed someone else undergoing fear conditioning in which this CS was paired with a US. (Images provided by Elizabeth Phelps)

REFERENCES

1. Kagan, J., Reznick, J. S., and Snidman, N. (1988) Biological bases of childhood shyness. *Science* 240, 167–71.
2. Dollard, J., and Miller, N. E. (1950) *Personality and Psychotherapy; An Analysis in Terms of Learning, Thinking, and Culture*. New York: McGraw-Hill.
3. Eysenck, M. W. (1997) *Anxiety and Cognition: A Unified Theory*. Hove, U.K.: Erlbaum.
4. Bandura, A. (1986) *Social Foundations of Thought and Action: A Social Cognitive Theory*. Englewood Cliffs, NJ: Prentice Hall.
5. Rosen, J. B., and Schulkin, J. (1998) From normal fear to pathological anxiety. *Psychol Rev* 105, 325–50.
6. LeDoux, J. E. (1996) *The Emotional Brain*. New York: Simon and Schuster.
7. Pavlov, I. P. (1927) *Conditioned Reflexes; An Investigation of the Physiological Activity of the Cerebral Cortex*. Oxford, U.K.: Oxford University Press.
8. Bouton, M. E., and Bolles, R. C. (1980) Conditioned fear assessed by freezing and by the suppression of three different baselines. *Anim Learn Behav* 8, 429–34.
9. Fanselow, M. S. (1980) Conditional and unconditional components of postshock freezing. *Pavlov J Biol Sci* 15, 177–82.
10. Swanson, L. W., and Petrovich, G. D. (1998) What is the amygdala? *Trends Neurosci* 21, 323–31.
11. LeDoux, J. E. (2000) Emotion circuits in the brain. *Annu Rev Neurosci* 23, 155–84.
12. Davidson, R. J., Jackson, D. C., and Kalin, N. H. (2000) Emotion, plasticity, context, and regulation: perspectives from affective neuroscience. *Psychol Bull* 126, 890–909.
13. Maren, S. (2001) Neurobiology of Pavlovian fear conditioning. *Annu Rev Neurosci* 24, 897–931.
14. Cardinal, R.N., Parkinson, J.A., Hall, J., and Everitt, B.J. (2002) Emotion and motivation: The role of the amygdala, ventral striatum, and prefrontal cortex. *Neurosci Biobehav Rev* 26, 321–52.
15. McGaugh, J.L. (2004) The amygdala modulates the consolidation of memories of emotionally arousing experiences. *Annu Rev Neurosci* 27, 1–28.
16. Paré, D. (2002) Mechanisms of Pavlovian fear conditioning: Has the engram been located? *Trends Neurosci* 25, 436–37; discussion 437–38.
17. Phelps, E.A., and LeDoux, J.E. (2005) Contributions of the amygdala to emotion processing: from animal models to human behavior. *Neuron* 48, 175–87.
18. Lang, P.J., and Davis, M. (2006) Emotion, motivation, and the brain: reflex foundations in animal and human research. *Prog Brain Res* 156, 3–29.
19. Rauch, S.L., Shin, L.M., and Wright, C.I. (2003) Neuroimaging studies of amygdala function in anxiety disorders. *Ann N Y Acad Sci* 985, 389–410.
20. Charney, D.S. (2004) Psychobiological mechanisms of resilience and vulnerability: implications for successful adaptation to extreme stress. *Am J Psychiatry* 161, 195–216.
21. Etkin, A., and Wager, T.D. (2007) Functional neuroimaging of anxiety: a meta-analysis of emotional processing in PTSD, social anxiety disorder, and specific phobia. *Am J Psychiatry* 164, 1476–88.
22. Liberzon, I., and Sripada, C.S. (2008) The functional neuroanatomy of PTSD: a critical review. *Prog Brain Res* 167, 151–69.
23. LeDoux, J.E. (2007) The amygdala. *Curr Biol* 17, R868–74.

24. Maren, S. (2005) Synaptic mechanisms of associative memory in the amygdala. *Neuron* 47, 783–86.
25. Fanselow, M.S., and Poulos, A.M. (2005) The neuroscience of mammalian associative learning. *Annu Rev Psychol* 56, 207–34.
26. Paré, D., Quirk, G.J., and LeDoux, J.E. (2004) New vistas on amygdala networks in conditioned fear. *J Neurophysiol* 92, 1–9.
27. Kapp, B.S., Whalen, P.J., Supple, W.F., and Pascoe, J.P. (1992) Amygdaloid contributions to conditioned arousal and sensory information processing. In: Aggleton, J.P., ed., *The Amygdala: Neurobiological Aspects of Emotion, Memory, and Mental Dysfunction*. New York: Wiley-Liss: 229–54.
28. Davis, M. (1992) The role of the amygdala in fear and anxiety. *Annu Rev Neurosci* 15, 353–75.
29. LeDoux, J.E. (1992) Emotion and the amygdala. In: Aggleton, J.P., ed., *The Amygdala: Neurobiological Aspects of Emotion, Memory, and Mental Dysfunction*. New York: Wiley-Liss: 339–51.
30. Pitkanen, A., Savander, V., and LeDoux, J.E. (1997) Organization of intra-amygdaloid circuitries in the rat: an emerging framework for understanding functions of the amygdala. *Trends Neurosci* 20, 517–23.
31. Everitt, B.J., Parkinson, J.A., Olmstead, M.C., Arroyo, M., Robledo, P., and Robbins, T.W. (1999) Associative processes in addiction and reward. The role of amygdala-ventral striatal subsystems. *Ann N Y Acad Sci* 877, 412–38.
32. Balleine, B.W., and Killcross, S. (2006) Parallel incentive processing: an integrated view of amygdala function. *Trends Neurosci* 29, 272–79.
33. Millhouse, O.E. (1986) The intercalated cells of the amygdala. *J Comp Neurol* 247, 246–71.
34. Paré, D., Royer, S., Smith, Y., and Lang, E.J. (2003) Contextual inhibitory gating of impulse traffic in the intra-amygdaloid network. *Ann N Y Acad Sci* 985, 78–91.
35. Samson, R.D., Duvarci, S., and Paré, D. (2005) Synaptic plasticity in the central nucleus of the amygdala. *Rev Neurosci* 16, 287–302.
36. Chapman, P.F., Ramsay, M.F., Krezel, W., and Knevet, S.G. (2003) Synaptic plasticity in the amygdala: comparisons with hippocampus. *Ann N Y Acad Sci* 985, 114–24.
37. Schafe, G.E., Nader, K., Blair, H.T., and LeDoux, J.E. (2001) Memory consolidation of Pavlovian fear conditioning: a cellular and molecular perspective. *Trends Neurosci* 24, 540–46.
38. Sah, P., Faber, E.S., Lopez De Armentia, M., and Power, J. (2003) The amygdaloid complex: anatomy and physiology *Physiol Rev* 83, 803–34.
39. Rodrigues, S.M., Schafe, G.E., and LeDoux, J.E. (2004) Molecular mechanisms underlying emotional learning and memory in the lateral amygdala. *Neuron* 44, 75–91.
40. Dityatev, A.E., and Bolshakov, V.Y. (2005) Amygdala, long-term potentiation, and fear conditioning. *Neuroscientist* 11, 75–88.
41. Sigurdsson, T., Doyère, V., Cain, C.K., and LeDoux, J.E. (2007) Long-term potentiation in the amygdala: a cellular mechanism of fear learning and memory. *Neuropharmacology* 52, 215–27.
42. Blair, H.T., Schafe, G.E., Bauer, E.P., Rodrigues, S.M., and LeDoux, J.E. (2001) Synaptic plasticity in the lateral amygdala: a cellular hypothesis of fear conditioning. *Learn Mem* 8, 229–42.
43. Rumpel, S., LeDoux, J., Zador, A., and Malinow, R. (2005) Postsynaptic receptor trafficking underlying a form of associative learning. *Science* 308, 83–88.
44. Huang, C.C., Wang, S.J., and Gean, P.W. (1998) Selective enhancement of p-type calcium currents by isoproterenol in the rat amygdala. *J Neurosci* 18, 2276–82.

45. Debiec, J., and LeDoux, J.E. (2004) Disruption of reconsolidation but not consolidation of auditory fear conditioning by noradrenergic blockade in the amygdala. *Neuroscience* 129, 267–72.
46. Tully, K., Li, Y., Tsvetkov, E., and Bolshakov, V.Y. (2007) Norepinephrine enables the induction of associative long-term potentiation at thalamo-amygdala synapses. *Proc Natl Acad Sci U S A* 104, 14146–50.
47. Guarraci, F.A., Frohardt, R.J., Falls, W.A., and Kapp, B.S. (2000) The effects of intra-amygdaloid infusions of a d2 dopamine receptor antagonist on Pavlovian fear conditioning. *Behav Neurosci* 114, 647–51.
48. Kröner, S., Rosenkranz, J.A., Grace, A.A., and Barrionuevo, G. (2005) Dopamine modulates excitability of basolateral amygdala neurons in vitro. *J Neurophysiol* 93, 1598–1610.
49. Johnson, L.R., Farb, C., Morrison, J.H., McEwen, B.S., and LeDoux, J.E. (2005) Localization of glucocorticoid receptors at postsynaptic membranes in the lateral amygdala. *Neuroscience* 136, 289–99.
50. Duvarci, S., and Paré, D. (2007) Glucocorticoids enhance the excitability of principal basolateral amygdala neurons. *J Neurosci* 27, 4482–91.
51. Huang, Y.Y., and Kandel, E.R. (2007) 5-hydroxytryptamine induces a protein kinase a/mitogen-activated protein kinase-mediated and macromolecular synthesis-dependent late phase of long-term potentiation in the amygdala. *J Neurosci* 27, 3111–19.
52. Schafe, G.E., Bauer, E.P., Rosis, S., Farb, C.R., Rodrigues, S.M., and LeDoux, J.E. (2005) Memory consolidation of Pavlovian fear conditioning requires nitric oxide signaling in the lateral amygdala. *Eur J Neurosci* 22, 201–11.
53. Rattiner, L.M., Davis, M., and Ressler, K.J. (2005) Brain-derived neurotrophic factor in amygdala-dependent learning. *Neuroscientist* 11, 323–33.
54. Azad, S.C., Monory, K., Marsicano, G., et al. (2004) Circuitry for associative plasticity in the amygdala involves endocannabinoid signaling. *J Neurosci* 24, 9953–61.
55. Krysiak, R., Obuchowicz, E., and Herman, Z.S. (2000) Conditioned fear-induced changes in neuropeptide Y-like immunoreactivity in rats: the effect of diazepam and buspirone. *Neuropeptides* 34, 148–57.
56. Cui, H., Sakamoto, H., Higashi, S., and Kawata, M. (2008) Effects of single-prolonged stress on neurons and their afferent inputs in the amygdala. *Neuroscience* 152, 703–12.
57. McGaugh, J.L. (2000) Memory—a century of consolidation. *Science* 287, 248–51.
58. Pitman, R.K., Sanders, K.M., Zusman, R.M., et al. (2002) Pilot study of secondary prevention of posttraumatic stress disorder with propranolol. *Biol Psychiatry* 51, 189–92.
59. Vaiva, G., Ducrocq, F., Jezequel, K., et al. (2003) Immediate treatment with propranolol decreases posttraumatic stress disorder two months after trauma. *Biol Psychiatry* 54, 947–49.
60. Misanin, J.R., Miller, R.R., and Lewis, D.J. (1968) Retrograde amnesia produced by electroconvulsive shock after reactivation of a consolidated memory trace. *Science* 160, 554–55.
61. Schneider, A.M., and Sherman, W. (1968) Amnesia: a function of the temporal relation of footshock to electroconvulsive shock. *Science* 159, 219–21.
62. Sara, S.J. (2000) Retrieval and reconsolidation: toward a neurobiology of remembering. *Learn Mem* 7, 73–84.
63. Nader, K., Schafe, G.E., and LeDoux, J.E. (2000) Fear memories require protein synthesis in the amygdala for reconsolidation after retrieval. *Nature* 406, 722–26.
64. Debiec, J., LeDoux, J.E., and Nader, K. (2002) Cellular and systems reconsolidation in the hippocampus. *Neuron* 36, 527–38.
65. Nader, K. (2003) Memory traces unbound. *Trends Neurosci* 26, 65–72.
66. Lee, J.L., Milton, A.L., and Everitt, B.J. (2006) Reconsolidation and extinction of conditioned fear: inhibition and potentiation. *J Neurosci* 26, 10051–56.

67. Suzuki, A., Josselyn, S.A., Frankland, P.W., Masushige, S., Silva, A.J., and Kida, S. (2004) Memory reconsolidation and extinction have distinct temporal and biochemical signatures. *J Neurosci* 24, 4787–95.
68. Alberini, C.M., Milekic, M.H., and Tronel, S. (2006) Mechanisms of memory stabilization and de-stabilization. *Cell Mol Life Sci* 63, 999–1008.
69. Dudai, Y. (2006) Reconsolidation: the advantage of being refocused. *Curr Opin Neurobiol* 16, 174–78.
70. Tronson, N.C., and Taylor, J.R. (2007) Molecular mechanisms of memory reconsolidation. *Nat Rev Neurosci* 8, 262–75.
71. Alberini, C.M. (2008) The role of protein synthesis during the labile phases of memory: revisiting the skepticism *Neurobiol Learn Mem* 89, 234–46.
72. Ben Mamou, C., Gamache, K., and Nader, K. (2006) NMDA receptors are critical for unleashing consolidated auditory fear memories. *Nat Neurosci* 9, 1237–39.
73. Duvarci, S., Nader, K., and LeDoux, J.E. (2005) Activation of extracellular signal-regulated kinase- mitogen-activated protein kinase cascade in the amygdala is required for memory reconsolidation of auditory fear conditioning. *Eur J Neurosci* 21, 283–89.
74. Doyère, V., Dębiec, J., Monfils, M.H., Schafe, G.E., and LeDoux, J.E. (2007) Synapse-specific reconsolidation of distinct fear memories in the lateral amygdala. *Nat Neurosci* 10, 414–16.
75. Tronson, N.C., Wiseman, S.L., Olausson, P., and Taylor, J.R. (2006) Bidirectional behavioral plasticity of memory reconsolidation depends on amygdalar protein kinase a. *Nat Neurosci* 9, 167–69.
76. Lee, J.L., Di Ciano, P., Thomas, K.L., and Everitt, B.J. (2005) Disrupting reconsolidation of drug memories reduces cocaine-seeking behavior. *Neuron* 47, 795–801.
77. Duvarci, S., & Nader, K. (2004) Characterization of fear memory reconsolidation. *J Neurosci* 24, 9269–75.
78. Dębiec, J., Doyère, V., Nader, K., and LeDoux, J.E. (2006) Directly reactivated, but not indirectly reactivated, memories undergo reconsolidation in the amygdala. *Proc Natl Acad Sci U S A* 103, 3428–33.
79. Dębiec, J., and Altemus, M. (2006) Toward a new treatment for traumatic memories. *Cerebrum*, Sep, 2–11.
80. Dębiec, J., and LeDoux, J.E. (2006) Noradrenergic signaling in the amygdala contributes to the reconsolidation of fear memory: treatment implications for PTSD. *Ann N Y Acad Sci* 1071, 521–24.
81. Brunet, A., Orr, S.P., Tremblay, J., Robertson, K., Nader, K., and Pitman, R.K. (2008) Effect of post-retrieval propranolol on psychophysiologic responding during subsequent script-driven traumatic imagery in post-traumatic stress disorder. *J Psychiatr Res* 42, 503–6.
82. Bouton, M.E. (1988) Context and ambiguity in the extinction of emotional learning: implications for exposure therapy. *Behav Res Ther* 26, 137–49.
83. Myers, K.M., and Davis, M. (2002) Behavioral and neural analysis of extinction. *Neuron* 36, 567–84.
84. Rudd, M.D., and Joiner, T. (1998) The role of symptom induction in the treatment of panic and anxiety. Identifiable domains, conditional properties, and treatment targets. *Behav Modif* 22, 96–107.
85. Shear, M.K., and Beidel, D.C. (1998) Psychotherapy in the overall management strategy for social anxiety disorder. *J Clin Psychiatry* 59(suppl 17), 39–46.
86. Foa, E.B. (2000) Psychosocial treatment of posttraumatic stress disorder. *J Clin Psychiatry* 61(suppl 5), 43–48; discussion 49–51.

87. Morgan, M.A., Romanski, L.M., and LeDoux, J.E. (1993) Extinction of emotional learning: contribution of medial prefrontal cortex. *Neurosci Lett* 163, 109–13.
88. Morgan, M.A., and LeDoux, J.E. (1995) Differential contribution of dorsal and ventral medial prefrontal cortex to the acquisition and extinction of conditioned fear in rats. *Behav Neurosci* 109, 681–88.
89. Milad, M.R., and Quirk, G.J. (2002) Neurons in medial prefrontal cortex signal memory for fear extinction. *Nature* 420, 70–74.
90. Sotres-Bayon, F., Bush, D.E., and LeDoux, J.E. (2004) Emotional perseveration: an update on prefrontal-amygdala interactions in fear extinction. *Learn Mem* 11, 525–35.
91. Sotres-Bayon, F., Cain, C.K., and LeDoux, J.E. (2006) Brain mechanisms of fear extinction: historical perspectives on the contribution of prefrontal cortex. *Biol Psychiatry* 60, 329–36.
92. Sotres-Bayon, F., Bush, D.E., and LeDoux, J.E. (2007) Acquisition of fear extinction requires activation of NR2B-containing NMDA receptors in the lateral amygdala. *Neuropsychopharmacology* 32, 1929–40.
93. Sotres-Bayon, F., Diaz-Mataix, L., Bush, D.E., and Ledoux, J.E. (2008) Dissociable roles for the ventromedial prefrontal cortex and amygdala in fear extinction: NR2B contribution. *Cereb Cortex* Jun 17 [Epub ahead of print].
94. Quirk, G.J., and Beer, J.S. (2006) Prefrontal involvement in the regulation of emotion: convergence of rat and human studies. *Curr Opin Neurobiol* 16, 723–27.
95. Corcoran, K.A., and Quirk, G.J. (2007) Recalling safety: cooperative functions of the ventromedial prefrontal cortex and the hippocampus in extinction. *CNS Spectr* 12, 200–6.
96. Ji, J., and Maren, S. (2007) Hippocampal involvement in contextual modulation of fear extinction. *Hippocampus* 17, 749–58.
97. Phelps, E.A., Delgado, M.R., Nearing, K.I., and LeDoux, J.E. (2004) Extinction learning in humans; role of the amygdala and vmPFC. *Neuron* 43, 897–905.
98. Walker, D.L., Ressler, K.J., Lu, K.T., and Davis, M. (2002) Facilitation of conditioned fear extinction by systemic administration or intra-amygdala infusions of d-cycloserine as assessed with fear-potentiated startle in rats. *J Neurosci* 22, 2343–51.
99. Ressler, K.J., Rothbaum, B.O., Tannenbaum, L., et al. (2004) Cognitive enhancers as adjuncts to psychotherapy: use of d-cycloserine in phobic individuals to facilitate extinction of fear. *Arch Gen Psychiatry* 61, 1136–44.
100. Davis, M. (2006) Neural systems involved in fear and anxiety measured with fear-potentiated startle. *Am Psychol* 61, 741–56.
101. Norberg, M.M., Krystal, J.H., and Tolin, D.F. (2008) A meta-analysis of d-cycloserine and the facilitation of fear extinction and exposure therapy. *Biol Psychiatry* 63, 1118–26.
102. Amorapanth, P., LeDoux, J.E., and Nader, K. (2000) Different lateral amygdala outputs mediate reactions and actions elicited by a fear-arousing stimulus. *Nat Neurosci* 3, 74–79.
103. Cain, C.K., and LeDoux, J.E. (2007) Escape from fear: a detailed behavioral analysis of two atypical responses reinforced by CS termination. *J Exp Psychol Anim Behav Process* 33, 451–63.
104. LeDoux, J.E., and Gorman, J.M. (2001) A call to action: overcoming anxiety through active coping. *Am J Psychiatry* 158, 1953–55.
105. van der Kolk, B.A., McFarlane, A.C., and Weisaeth, L. (eds.). (1996) *Traumatic Stress: The Effects of Overwhelming Experience on Mind, Body, and Society*. New York: Guilford Press.

106. Cloitre, M., Cohen, L.R. & Koenen, K.C. (2006) Treating Survivors of Childhood Abuse: Psychotherapy for the Interrupted Life. New York: Guilford Press.
107. LaBar, K.S., Gatenby, J.C., Gore, J.C., LeDoux, J.E., and Phelps, E.A. (1998) Human amygdala activation during conditioned fear acquisition and extinction: a mixed-trial fMRI study. *Neuron* 20, 937–45.
108. Morris, J.S., Ohman, A., and Dolan, R.J. (1998) Conscious and unconscious emotional learning in the human amygdala. *Nature* 393, 467–70.
109. Shin, L.M., Kosslyn, S.M., McNally, R.J., et al. (1997) Visual imagery and perception in posttraumatic stress disorder. A positron emission tomographic investigation. *Arch Gen Psychiatry* 54, 233–41.
110. Liberzon, I., Britton, J.C., and Phan, K.L. (2003) Neural correlates of traumatic recall in posttraumatic stress disorder. *Stress* 6, 151–56.
111. Shin, L.M., Orr, S.P., Carson, M.A., et al. (2004) Regional cerebral blood flow in the amygdala and medial prefrontal cortex during traumatic imagery in male and female Vietnam veterans with PTSD. *Arch Gen Psychiatry* 61, 168–76.
112. Schienle, A., Schafer, A., Walter, B., Stark, R., and Vaitl, D. (2005) Brain activation of spider phobics towards disorder-relevant, generally disgust- and fear-inducing pictures. *Neurosci Lett* 388, 1–6.
113. Phan, K.L., Fitzgerald, D.A., Nathan, P.J., and Tancer, M.E. (2006) Association between amygdala hyperactivity to harsh faces and severity of social anxiety in generalized social phobia. *Biol Psychiatry* 59, 424–29.



<http://www.springer.com/978-1-60327-328-2>

Post-Traumatic Stress Disorder

Basic Science and Clinical Practice

Shiromani, P.; Keane, T.; LeDoux, J.E. (Eds.)

2009, XIII, 418 p., Hardcover

ISBN: 978-1-60327-328-2

A product of Humana Press