

2

Epidemiology

Bong Jin Park, Han Kyu Kim, Burak Sade, and Joung H. Lee

Introduction

Meningiomas constitute 13–25% of primary intracranial neoplasms.^{1–3} In one population-based study, these tumors accounted for 40% of all primary intracranial neoplasms when tumors diagnosed incidentally at autopsy or by neuroimaging studies were included.⁴ Symptomatic tumors were encountered in 2.0/100,000 of the population and the asymptomatic ones in 5.7/100,000, with an overall incidence of 7.7/100,000. Asymptomatic meningiomas can be found in about 1–2.3% of all autopsies.⁵ However, the classical teaching of the past few decades, which may be erroneous, suggests the most common primary intracranial neoplasms are gliomas (50.3%), followed by meningiomas (20.9%), pituitary adenomas (15%) and nerve sheath tumors (8%).^{6,7} Such discrepancy in these incidence rates underscores the fact that the majority of meningiomas actually remain asymptomatic and undetected during life.⁸ With recent advances in neuroimaging, many asymptomatic meningiomas are being detected today, making the true incidence higher than those previously reported.⁹ It has been shown that the incidence for meningiomas increased 3- to 3.9-fold in the post-computed tomography (CT) decade.^{10,11}

Gender, Race, and Age

Meningiomas show a higher incidence among women as compared to men in most ethnic groups.^{12,13} The male-to-female ratio ranges from 1:1.4 to 2.6.^{14–16} In population-based studies, the mean annual crude incidence rates are reported to be 2–7/100,000 for women and 1–5/100,000 for men.^{1,3}

In a population-based study performed in Los Angeles County, African Americans showed a higher incidence (3.1/100,000) than Caucasians (2.3/100,000).¹² In this study, Asians living in Los Angeles had the lowest rates. In Caucasians, women are approximately twice as likely to develop meningiomas than men, whereas in African American populations the incidence is evenly distributed between males and females.¹ Two studies have also shown a lower incidence of meningiomas

in Asia as compared to western countries,^{17,18} whereas Chi et al. reported no significant difference in the incidence of meningiomas in relation to other intracranial tumors, namely, 20.8% in Korea, 18.5% in Japan, and 16.6% in China.¹⁹ Others have also observed no racial differences in incidence rates of meningiomas.¹³

The mean age at presentation is 56.4 years (range 10–85 years) in males and 55.9 years (range 26–86 years) in females, whereas in the subgroup of malignant and atypical meningiomas, the mean age shifts to 63.2 years (range 51–78 years) in males and 53.6 years (range 28–79 years) in females, a difference that is not statistically significant.²⁰ The incidence of meningiomas increases with age.²¹ In patients older than 70 years, they have been reported as the most common brain tumors, with an incidence of 50.6%. This represents an almost 3.5 times higher incidence in this age group as compared to those under the age of 70, and it applies to both sexes.^{4,22} The age-specific annual incidence rate increases in the eighth decade to 8.4/100,000.¹⁶

Histology and Location

The vast majority of meningiomas (92%) have a benign histology, whereas 8% show atypical or malignant features.²³ The most common histopathological subtype is the meningotheliomatous type (63%), followed by transitional (19%), fibrous (13%), and psammomatous (2%) meningiomas.²⁴ The majority of malignant meningiomas are located over the cerebral convexities. In approximately 73.3–75% of cases these tumors are located in the supratentorial compartment.⁸ The ratio of calvarial to basal skull meningiomas was reported as 2.3:1.²⁵ Extracranial metastases from meningiomas have been considered to be one of the strong indicators of malignancy and have been shown to occur in 11–23% of patients with malignant meningiomas.⁶

The most common locations include parasagittal/falcine 25%, convexity 19%, sphenoid ridge 17%, followed by suprasellar 9%, posterior fossa 8%, olfactory groove 8%, middle

fossa/Meckel's cave 4%, tentorial 3%, peri-torcular 3%, lateral ventricle 1–2%, foramen magnum 1–2%, and orbit/optic nerve sheath 1–2%.¹⁰ Among the parasagittal meningiomas, 49% are located over the anterior one third of the falx, with 29% in the middle third, and 22% along the posterior third.¹⁶ Medial sphenoid ridge meningiomas were more common than middle or lateral sphenoid ridge meningiomas. Multiple meningiomas or meningiomatosis is encountered in 2.5% of meningiomas. The incidence of ectopic location is 0.4% with the vast majority (73%) occurring inside the orbit, paranasal sinuses, eyelids, parotid gland, temporalis muscle, temporal bone, and zygoma. Distant sites have also been reported, such as the lungs, mediastinum, and the adrenal glands.^{6,26}

Recently, Lee et al. demonstrated an association between the histology of the tumor and its site of origin.²⁴ They showed a predominance of meningothelial meningiomas at the midline skull base and spinal locations. Based on this finding, as well as embryological and molecular features, they suggested that this particular subtype of meningiomas may indeed be unique, contrary to the traditional dogma that all benign meningiomas are identical or homogeneous tumors.

Meningiomas in Children

In children, meningiomas account for only 0.4–4.0% of primary intracranial neoplasms.^{1,27–29} The age-adjusted annual incidence was reported to be 1.32/1,000,000.³⁰ There is a male predominance, with a male-to-female ratio of 1.2 to 1.9:1.^{27,30}

The majority of meningiomas in the pediatric age group are located supratentorially (66%), whereas 19% occur in the posterior fossa and 17% present as intraventricular meningiomas.³⁰ They are usually seen in association with neurofibromatosis type 2 (NF-2) or following radiation therapy and show a significantly higher incidence of tumor calcification.^{27,28} In NF-2, it has been estimated that 50% of all patients develop meningiomas, and 30% of these patients have multiple meningiomas.⁶

In children, additional unique features include the significantly increased incidence of atypical (36.4%) and malignant (27.2%) subtypes. In infants, meningiomas are extremely rare and show a higher frequency in males and favor convexity location. On the other hand, there is a smaller incidence of seizure, and dural attachment is less frequently seen on preoperative imaging.¹

Spinal Meningiomas

In women, meningiomas are by far the most common primary spinal tumor, accounting for 58% of all spinal tumors, whereas in men they are third most common primary spinal tumor following gliomas and nerve sheath tumors.³¹ Spinal meningiomas are reported to be more frequent in western countries (25–46%) as compared to Asian countries, for

example, 14.1% in China and 8.6% in Thailand.^{17,19,32} Thoracic spine is the most common site (55–57.1%).^{17,19,32} The male-to-female ratio is 1:4 to 1:5.³³ In females, they are very common in the postmenopausal age group, with the majority (75–87%) occurring over the age of 40.^{34,35}

The tumor is located completely intradurally in 83–90%, extradurally in 5–14%, and both intradurally and extradurally in 5% of the cases.^{32,35} Extradural meningiomas are reported to be more common in children. In 50–68% of the cases, the tumor is located lateral to the spinal cord, in 18–31% posteriorly, and in 15–19% anteriorly.^{32,35}

Histologically, 43.9–56.9% of spinal meningiomas show psammomatous subtype, whereas 28–29.9% are meningothelial, 8–19% transitional, 2.3–5% are fibrous, and 0.6% are malignant meningiomas.^{32,35} The incidence of multiple spinal meningiomas is reported as 1–9%. In a recent review of our series, contrary to the above, meningothelial subtype was the most common (80%) in the spine.²⁴

Radiation

In a population-based study, the incidence of meningiomas among Hiroshima atomic bomb survivors was 8.7/100,000.³⁶ When this population was stratified according to the hypocenter to the explosion, the incidence of meningiomas was much higher in patients who had been closer to the site of explosion. Similarly, the incidence of meningiomas was reported as 9.5/100,000 population in the group who had undergone low-dose radiation treatment as children for tinea capitis in Israel.³⁷

In the medical or occupational setting, no significant associations were observed for diagnostic studies and increased meningioma incidence, but the use of radiation therapy to head and neck for neoplastic conditions has been shown to be associated with an increase in the incidence of meningiomas.³⁸

Radiation-induced meningiomas differ significantly from primary intracranial meningiomas in that their incidence of calvarial location, multiplicity, recurrence rate following complete resection, and malignant histology are higher.²⁵

Occupation

Association of increased risk of meningioma incidence has been suggested for various occupations in the literature. There is a huge diversity in the nature of proposed occupations such as dentists, teachers, managers, social workers, workers in the petroleum, rubber and plastics industry, auto body repairers, painters, chemists, carpenters, cooks, woodworkers, glass-makers, machine operators, as well as military workers, motor vehicle drivers, computer specialists, and so on.^{12,39,40}

Rajamaram et al. have stated that it might be practical to analyze these occupations in two groups.⁴⁰ Groups like teachers, managers, etc., tend to be relatively better educated and would be expected to have a higher awareness of their health status,

which would result in the increased detection of tumors in these groups, by earlier recognition of the symptoms, or by stronger willingness of the individual to seek medical care. On the other hand, exposure to some chemical agents or other environmental factors may be more influential in other groups. In this context, lead, tin, cadmium, benzene, and metal dusts and fumes have been suggested as possible contributors.^{12,39–41}

Cell Phone Use

Over the years, there has been much debate about the potential role of cell phones as a causative factor in the development of brain tumors because of the microwaves emitted by these devices. In a recent population-based case-control study conducted on 366 glioma and 381 meningioma patients with 1494 controls, no association was found between cell phone use and increased meningioma incidence. However, in this study, increased incidence for glioma occurrence was detected in individuals who had used cell phones for more than 10 years.⁴² A similar study in a relatively smaller scale also suggested a low risk for development of high-grade glioma for cell phone users, but the overall results did not show any association for either gliomas or meningiomas.⁴³ The study conducted by the Swedish Interphone Study Group also found no increase in incidence for gliomas or meningiomas located in the temporal or parietal lobes, regardless of tumor histology, phone type, or amount of phone use.⁴⁴

References

- Bondy M, Ligon BL. Epidemiology and etiology of intracranial meningiomas: A review. *J Neurooncol* 1996;29:197–205.
- Claus EB, Bondy ML, Schildkraut JM, et al. Epidemiology of intracranial meningioma. *Neurosurgery* 2005;57:1088–95.
- Longstreth WT Jr, Dennis LK, McGuire VM, et al. Epidemiology of intracranial meningioma. *Cancer* 1993;72:639–48.
- Radhakrishnan K, Mokri B, Parisi JE, et al. The trends in incidence of primary brain tumors in the population of Rochester, Minnesota. *Ann Neurol* 1995;37:67–73.
- Kurland LT, Schoenberg BS, Annegers JF, et al. The incidence of primary intracranial neoplasms in Rochester, Minnesota, 1935–1977. *Ann NY Acad Sci* 1982;381:6–16.
- Evans JJ, Lee JH, Suh J, et al. Meningiomas. In: Newell DW, Moore AJ, ed. *Neurosurgery: Principles and Practice*. London: Springer-Verlag, 2004;205–32.
- Hoffman SH, Propp JM, McCarthy BJ. Temporal trends in incidence of primary brain tumors in the United States, 1985–1999. *Neurooncology* 2006;8:27–37.
- Elia-Pasquet S, Provost D, Jaffre A, et al. Incidence of central nervous system tumors in Gironde, France. *Neuroepidemiology* 2004;23:110–7.
- Klaeboe L, Lonn S, Scheie D, et al. Incidence of intracranial meningiomas in Denmark, Finland, Norway and Sweden, 1986–1997. *Int J Cancer* 2005;20:996–1001.
- Christensen HC, Kosteljanetz M, Johansen C. Incidence of gliomas and meningiomas in Denmark, 1943 to 1997. *Neurosurgery* 2003;52:1327–34.
- Helseth A. The incidence of primary central nervous system neoplasms before and after computerized tomography availability. *J Neurosurg* 1995;83:999–1003.
- Preston-Martin S. Descriptive epidemiology of primary tumors of the brain, cranial nerves and cranial meninges in Los Angeles County. *Neuroepidemiology* 1989;8:283–95.
- Surawicz TS, McCarthy BJ, Kupelian V, et al. Descriptive epidemiology of primary brain and CNS tumors: results from the central brain tumor registry of the United States, 1990–1994. *Neurooncology* 1999;1:14–25.
- Alessandro GD, Giovanni MD, Iannizzi L, et al. Epidemiology of primary intracranial tumors in the Valle d'Aosta (Italy) during the 6-year period 1986–1991. *Neuroepidemiology* 1995;14:139–46.
- Lovaste MG, Ferrari G, Rossi G. Epidemiology of primary intracranial neoplasms: experiment in the Province of Trento (Italy), 1977–1984. *Neuroepidemiology* 1986;5:220–32.
- Rohringer M, Sutherland GR, Louw DF, et al. Incidence and clinicopathological features of meningioma. *J Neurosurg* 1989;71:665–72.
- Huang WQ, Zheng SJ, Tian QS, et al. Statistical analysis of central nervous system tumors in China. *J Neurosurg* 1982;56:555–64.
- Ng HK, Poon WS, South JR, et al. Tumors of the central nervous system in Chinese in Hong Kong: a histological review. *Aust NZ J Surg* 1988;58:573–8.
- Chi JG, Khang SK. Central nervous system tumors among Koreans. *J Kor Med Sci* 1989;4:77–90.
- Das A, Tang WY, Smith DR. Meningiomas in Singapore: demographic and biological characteristics. *J Neurooncol* 2000;47:153–60.
- Kuratsu JI, Ushio Y. Epidemiological study of primary intracranial tumors: a regional survey in Kumamoto prefecture in the southern part of Japan. *J Neurosurg* 1996;84:946–50.
- Kuratsu J, Ushio Y. Epidemiological study of primary intracranial tumors in elderly people. *J Neurol Neurosurg Psychiatry* 1997;63:116–8.
- Feun LG, Raub WA, Landy HJ, et al. Retrospective epidemiologic analysis of patients diagnosed with intracranial meningioma from 1977 to 1990 at the Jackson Memorial Hospital, Sylvester Comprehensive Cancer Center: the Jackson Memorial Hospital tumor registry experience. *Cancer Detect Prev* 1996;20:166–70.
- Lee JH, Sade B, Choi E, et al. Midline skull base and spinal meningiomas are predominantly of the meningotheelial histological subtype. *J Neurosurg* 2006;105:60–64.
- Sadamori N, Shibata S, Mine M, et al. Incidence of intracranial meningiomas in Nagasaki atomic-bomb survivors. *Int J Cancer* 1996;67:318–22.
- Staneczek W, Janisch W. Epidemiological data on meningiomas in East Germany 1961–1986: incidence, localization, age and sex distribution. *Clin Neuropathol* 1992;11:135–41.
- Baumgartner JE, Sorenson JM. Meningioma in the pediatric population. *J Neurooncol* 1996;29:223–8.
- Merten DF, Gooding CA, Newton TH, et al. Meningiomas of childhood and adolescence. *J Pediatrics* 1974;84:696–700.
- Schoenberg BS, Schoenberg DG, Christine BW, et al. The epidemiology of primary intracranial neoplasms of childhood: a population study. *Mayo Clin Proc* 1976;51:51–6.
- Kuratsu J, Ushio Y. Epidemiological study of primary intracranial tumors in childhood: a population-based survey in Kumamoto prefecture, Japan. *Pediatr Neurosurg* 1996;25:240–7.

31. Preston-Martin S. Descriptive epidemiology of primary tumors of the spinal cord and spinal meninges in Los Angeles County, 1972–1985. *Neuroepidemiology* 1990;9:106–11.
32. Cohen-Gadol AA, Zikel OM, Koch CA, et al. Spinal meningiomas in patients younger than 50 years of age: a 21-year experience. *J Neurosurg (Spine 3)* 2003;98:258–63.
33. Levy WJ Jr, Bay J, Dohn D. Spinal cord meningioma. *J Neurosurg* 1982;57:804–12.
34. Helseth A, Mork SJ. Primary intraspinal neoplasms in Norway, 1955 to 1986: a population-based survey of 467 patients. *J Neurosurg* 1989;71:842–5.
35. Solero CL, Fornari M, Giombini S, et al. Spinal meningiomas: review of 174 operated cases. *Neurosurgery* 1989;25:153–60.
36. Shintani T, Hayakawa N, Hoshi M, et al. High incidence of meningioma among Hiroshima atomic bomb survivors. *J Radiat Res* 1999;40:49–57.
37. Ron E, Modan B, Boice JD, et al. Tumors of the brain and nerve system after radiotherapy in childhood. *N Engl J Med* 1988;319:1033–9.
38. Phillips LE, Frankenfeld CL, Drangsholt M, et al. Intracranial meningioma and ionizing radiation in medical and occupational settings. *Neurology* 2005;64:350–2.
39. Navas-Acien A, Pollan M, Gustavsson P, et al. Occupation, exposure to chemicals and risk of gliomas and meningiomas in Sweden. *Am J Ind Med* 2002;42:214–27.
40. Rajaraman P, De Roos AJ, Stewart PA, et al. Occupation and risk of meningioma and acoustic neuroma in the United States. *Am J Ind Med* 2004;45:395–407.
41. Hu J, Little J, Xu T, et al. Risk factors for meningioma in adults: a case-control study in northeast China. *Int J Cancer* 1999;83:299–304.
42. Schuz J, Bohler E, Berg G, et al. Cellular phones, cordless phones and the risks of glioma and meningioma (Interphone Study Group, Germany). *Am J Epidemiol* 2006;163:512–20.
43. Christensen HC, Schuz J, Kosteljanetz M, et al. Cellular telephones and risk for brain tumors: a population based, incident case-control study. *Neurology* 2005;64:1189–95.
44. Lonn S, Ahlbom A, Hall P, et al. Long-term mobile phone use and brain tumor risk. *Am J Epidemiol* 2005;161:526–35.



<http://www.springer.com/978-1-84628-526-4>

Meningiomas

Diagnosis, Treatment, and Outcome

Lee, J.H. (Ed.)

2009, XXIII, 639 p., Hardcover

ISBN: 978-1-84628-526-4