

Contents

1	Introduction	1
	References	6
2	Splitting of the Main Problem into Four Sub-cases	9
2.1	Introduction	9
2.2	Two NITA Generated by a Non-real Element of Degree 3 not Derived from a Group and Lemmas	11
2.3	NITA Generated by b_3 and Satisfying $b_3^2 = b_4 + b_5$	23
2.4	General Information on NITA Generated by b_3 and Satisfying $b_3^2 = \bar{b}_3 + b_6$ and $b_3^2 = c_3 + b_6$	52
2.5	NITA Generated by b_3 Satisfying $b_3^2 = \bar{b}_3 + b_6$ and b_6 Nonreal and $b_{10} \in B$ is Real	55
2.6	NITA Generated by b_3 Satisfying $b_3^2 = c_3 + b_6$, $c_3 \neq b_3, \bar{b}_3, b_6$ Non-real, $(b_3b_8, b_3b_8) = 4$ and $c_3^2 = r_3 + s_6$	82
2.7	Structure of NITA Generated by b_3 and Satisfying $b_3^2 = c_3 + b_6$, $c_3 \neq b_3, \bar{b}_3$, $(b_3b_8, b_3b_8) = 3$ and c_3 Non-real	83
2.8	Structure of NITA Generated by b_3 and Satisfying $b_3^2 = c_3 + b_6$, $c_3 \neq b_3, \bar{b}_3$, $(b_3b_8, b_3b_8) = 3$ and c_3 Real	108
	References	150
3	A Proof of a Non-existence of Sub-case (2)	151
3.1	Introduction	151
3.2	Preliminary Results	152
3.3	Case $z = z_3$	158
3.4	Cases $z = z_4, z = z_5, z = z_6, z = z_7$, and $z = z_8$	174
3.5	Case $z = z_9$	180
4	Preliminary Classification of Sub-case (3)	187
4.1	Introduction	187
4.2	Preliminary Results	188
4.3	Case $R_{15} = x_5 + x_{10}$	190
4.4	Case $R_{15} = x_6 + x_9$	195

4.5	Case $R_{15} = x_7 + x_8$	209
4.6	Case $(b_3x_7, b_3x_7) = 3$	233
4.7	Case $b_3b_{10} = b_{15} + x_5 + y_5 + z_5$	238
5	Finishing the Proofs of the Main Results	243
5.1	Introduction	243
5.2	Proof of Theorem 5.1	247
5.3	Proof of Theorem 5.2	268
	References	271
Index		273

On Normalized Integral Table Algebras (Fusion Rings)
Generated by a Faithful Non-real Element of Degree 3

Arad, Z.; Bangteng, X.; Chen, G.; Cohen, E.; Haj Ihia

Hussam, A.; Muzychuk, M.

2011, X, 274 p., Hardcover

ISBN: 978-0-85729-849-2