

The burden of non-motor symptoms

Maria João Forjaz, Chandni Chandiramani and
Pablo Martinez-Martin

Although most of the symptoms of Parkinson's disease (PD) seem to be motor-based, the non-motor symptoms (NMSs) of PD, often overshadowed, also constitute a major clinical challenge. The NMSs of PD, including neuropsychiatric, cognitive, gastrointestinal and sensory, not only are observed in later stages but also predominate in the early stages of PD. Research suggests that NMSs are present at onset, while some can precede the motor symptoms, often by many years [1–3]. The identification of premotor stages is based on the early detection of the combination of NMSs with PD detection: for example, olfactory changes are the most common NMS presenting in the early stages of PD and occur in 90% of the patients [4,5]. Surveys reveal that 90% have at least one NMS, whereas 10% of PD patients exhibit five NMSs [6]. NMSs seem to be present before the diagnosis of PD and also inevitably emerge with progression of the disease, thus having an impact on patients' health-related quality of life (HRQoL) and placing an increased burden on caregivers.

The burden of non-motor symptoms on patients' health-related quality of life

Two important shifts

Interest in patients' HRQoL, and patient-reported outcomes including QoL measures, started when therapeutic interventions changed their focus from postponing death to improving HRQoL. As PD is a chronic debilitating condition, so far without cure, specialists became interested in measuring the impact of the disease and treatments in terms of HRQoL [7,8].

Recently, a second shift occurred, and there has been growing interest in measuring not only motor symptoms, but also NMSs. First, measures about specific NMSs, developed for the general population, were included in PD studies. Later on, there was an effort to develop domain-specific questionnaires for PD, such as the Scales for Outcomes in Parkinson's disease (SCOPA). Finally, specific scales, comprising a large number of NMSs in a single measure, such as the NMS Questionnaire (NMSQuest) and the NMS Scale (NMSS), were developed [9].

Studies using non-motor symptom measures

There is a high consistency of findings in studies using PD-specific NMS measures (Figure 2.1). Patients report, on average, 9–11 symptoms on the NMSQuest, the most frequent symptoms being nocturia, urgency, constipation and 'the blues'. The number of symptoms increases with Hoehn and Yahr (H&Y) stage [22] and disease duration. The mean severity score on the NMSS ranges from 31 to 91.

The most common NMSS domains are mood and sleep, and the NMSS scores are significantly associated with H&Y stage and motor symptoms. The relationship between NMSS and HRQoL is high, consistent and robust (Spearman $r = 0.70$ with PDQ-39 or -8) [15,18].

Motor versus non-motor symptoms: effect on patients' health-related quality of life

NMSs seem to have a greater impact on patients' HRQoL than motor symptoms [23–25]. The most frequent HRQoL determinants found in

a HRQoL review of PD were depression and H&Y stage, followed by disability, disease duration and gait disorders [7].

A study specifically analyzing the impact of motor symptoms and NMSs on patients' HRQoL found that, although motor variables contributed to 19% of the variance, inclusion of non-motor factors significantly improved the model (62%) [26]. The most important predictors of HRQoL were depression, sleep disorders and fatigue and, to a lesser extent, UPDRS motor score and H&Y stage.

Further research

NMSs have a great impact on patients' HRQoL. This relationship, when using the NMSS, is robust and of high magnitude. More research is needed to ascertain which domains of the NMSS mostly impact on which HRQoL aspects, and whether NMSs predict HRQoL in the medium or long term. Longitudinal studies are very important, because there is strong evidence that some NMSs may be preclinical characteristics of PD [27]. Finally, there are insufficient studies comparing the contribution of motor and non-motor symptoms on HRQoL, using specific NMS measures.

The burden of patient's non-motor symptoms on the caregiver

NMSs cause a significant burden to not only the patient's life, but also the caregiver's life. Figure 2.2 summarizes the main studies about the patient factors influencing the caregiver's QoL and burden.

The impact of the disease on the caregiver's wellbeing can be measured in terms of QoL or specific measures of burden. Questionnaires of the caregiver's QoL include the SF-36 [52], EQ-5D [53], WHOQOL-BREF [54] and the Scale of Quality of Life of Caregivers [55]. Older patients with lower HRQoL, higher disability and more severe motor and neuropsychological affectation are taken care of by caregivers with lower HRQoL [37,39,45,49].

There are several measures of caregivers' burden, the three most commonly used measures being: the Zarit Caregiver Burden Inventory [56]; the Caregiver Burden Inventory [57]; and the Subjective and Objective

Studies using specific non-motor symptom scales for Parkinson's disease

Reference	Sample	Symptoms, domains or score: M ± SD (range)	Most common symptoms or domains	Associated factors
NMSQuest	n = 100–545	Mean number of symptoms: 9 to 11	Nocturia, urgency, constipation, depression	H&Y stage, disease duration
Chaudhuri, et al (2006) [10]	123 PD patients (multinational)	Symptoms: 9.6	Symptoms: nocturia (67%), urgency (61%), remembering (51%)	H&Y stage ($r = 0.31$), disease duration ($r = 0.22$)
Martinez-Martin, et al (2007) [11]	545 PD patients (multinational)	Symptoms: 10.3 ± 5.4 (0–28)	Symptoms: nocturia (62%), urgency (56%), constipation (52%) Domains: digestive, sleep	Age of onset (before/after 50 years), disease duration, H&Y stage
Chaudhuri, et al (2010) [12]	242 PD patients (multinational)	Symptoms: 10.9 ± 5.6	Symptoms: nocturia (65%), urgency (60%), concentrating (50%) Domains: gastrointestinal, sleep	NMSs present more in akinesia PD type
Wang, et al (2010) [13]	117 PD patients (China)	Not reported	Symptoms: constipation (56%), sadness/blues (43%), remembering (41%)	Malnutrition
Cervantes-Arriaga, et al (2010) [14]	100 PD patients (Mexico)	Symptoms: 10.6 ± 5.5 (1–26)	Not reported	H&Y ($r < 0.25$, $p < 0.05$)
NMSS	n = 22–242	Mean of NMSS scores: 31–90	Domains: mood and sleep	H&Y stage, quality of life, motor symptoms
Chaudhuri, et al (2007) [15]	242 PD patients (multinational)	NMSS score: 56.5 ± 40.7 (0–243)	Not reported	PDQ-8 ($r = 0.70$); H&Y ($r = 0.33$); UPDRS section 4 ($r = 0.49$); UPDRS section 3 ($r = 0.35$)
Honig, et al (2009) [16]	22 patients with advanced PD receiving intrajejunal levodopa infusion (multinational)	NMSS score at baseline: 89.9 ± 56.5; at follow-up: 39.4 ± 33.9	Not reported	Change in NMSS score associated with: PDQ-8 ($r = 0.61$), UPDRS complications ($r = 0.55$); dyskinesia score ($r = 0.41$) and proportion of day in off ($r = 0.54$)

Kim, et al (2009) [17]	23 new PD patients (Korea)	Number of symptoms: 8.3 ± 4.0 (3–17). NMSS score: 37.0 ± 28.9	Symptoms: nocturia (65%), forget things (61%), restless legs (52%)	H&Y stage ($r = 0.43$, with number of symptoms)
Martinez-Martin, et al (2009) [18]	411 PD patients (multinational)	NMSS score: 57.1 ± 44.0 (0–233)	Domains with higher mean scores: mood/apathy (11.2), sleep/fatigue (9.6), urinary (9.0)	Group differences: disease duration, H&Y stage and CISI-PD severity level ($p < 0.001$). Correlations: PDQ-39 ($r = 0.70$), SCOPA-AUT ($r = 0.64$), EQ-5D index ($r = 0.57$). Other correlations ($r = 0.38$ – 0.51): H&Y; SCOPA-M, -PC and -COG; CISI-PD; PDSS
Wang, et al (2009) [19]	126 PD patients (China)	NMSS score: 31.06 ± 30.88 (0–177)	Domains with higher mean scores: mood (8.26), sleep/fatigue (6.33), miscellany (3.84)	Correlations with: sleep (PSQI, $r = 0.63$); anxiety (HAMA, $r = 0.52$); depression (GDS, $r = 0.45$)
Cervantes-Arriaga, et al (2010) [14]	100 PD patients (Mexico)	NMSS score: 69.7 ± 56.9 (0–237)	Not reported	H&Y ($r < 0.25$, $p = 0.05$)
Checklist of 48 NMSSs grouped into 12 domains	$n = 1072$–1130	Mean number of symptoms: 8	Domains: gastrointestinal	H&Y stage, quality of life
Barone, et al (2009) [20]	1072 PD patients (Italy)	Symptoms: 7.8 ± 4.9 (0–32) Domains: 5.5 ± 2.5 (0–12)	Symptoms: fatigue (58%), anxiety (56%), leg pain (38%) Domains: psychiatric (67%), sleep (64%), gastrointestinal (61%)	H&Y stage (all domains except cardiovascular); cognitive impairment (6 domains); PDQ-39 (all domains)
Colosimo, et al (2010) [21]	1130 PD patients (Italy)	Symptoms: 8 Domains: 5.5	Domains: gastrointestinal (62%), pain (61%), urinary (58%)	None analyzed

Figure 2.1 Studies using specific non-motor symptom scales for Parkinson's disease. CISI-PD, Clinical Impression of Severity Index for Parkinson's Disease; GDS, Geriatric Depression Scale; HAMA, Hamilton Rating Scale for Anxiety; H&Y, Hoehn and Yahr scale; NMS, non-motor symptom; NMSQuest, Non-Motor Symptoms Questionnaire; NMSS, Non-Motor Symptoms Scale; PD, Parkinson's disease; PDQ-39, Parkinson's disease Questionnaire – 39 items; PDQ-8, Parkinson's disease Questionnaire – 8 items; SCOPA-AUT, Scales for Outcomes in Parkinson's disease – Autonomic; SCOPA-COG, Scales for Outcomes in Parkinson's disease – Cognition; SCOPA-M, Scales for Outcomes in Parkinson's disease – Motor; SCOPA-PC, Scales for Outcomes in Parkinson's disease – Psychiatric Complications; UPDRS, Unified Parkinson's disease Rating Scale.

Studies about patient's factors influencing the caregiver's quality of life

Reference	Study type: sample, country, design	Patient factors: disease related, NMS, motor symptoms, others	Outcome measure: caregiver's burden; caregiver's QoL
Miller et al (1998) [28]	54 dyads (patient–caregiver), UK, multiple regression	NMS: depression	Caregiver's burden: GHQ-30, Depression (GDS, BDI); Machine Strain Scale
Carter et al (2004) [29]	380 caregivers, USA, group differences	Disease related: disease severity	Caregiver's burden: Family Care(giving) Inventory
Meara et al (1999) [30]	79 caregivers, UK, correlations	NMS: depression, cognition	Caregiver's burden: depression (GDS-15)
Habermann (2000) [31]	8 caregivers, USA, qualitative study	Disease related: disease severity NMS: Feelings of struggle and frustration	Caregiver's burden: interview about challenges that CG experience
Fernandez et al (2001) [32]	45 dyads, USA, multiple regression	Disease related: disease duration	Caregiver's burden: depression (HAM-D, BDI)
Caap-Ahlgren and Dehlin (2002) [33]	65 dyads, Sweden, multiple regression	NMS: depression Others: disability	Caregiver's burden: scale, depression (GDS)
Edwards and Scheetz (2002) [34]	41 caregivers, USA, multiple regression	Others: activities of daily living	Caregiver's burden: Zarit Caregiver Burden Inventory
Happe and Berger (2002) [35]	106 caregivers, Germany, multiple regression	Disease related: disease severity (with caregiver's sleep) NMS: depression (with caregiver's depression), sleep (with caregiver's sleep) NMS: cognition, depression	Caregiver's burden: Caregiver Burden Inventory, depression (CES-D); sleep
Thommessen et al (2002) [36]	58 couples, Norway, structural equation modelling		Caregiver's burden: Relative Stress Scale

Martinez-Martin et al (2004) [37]	86 dyads, Spain, correlations	Disease related: disease duration and severity NMS: Neuropsychological disorders Questionnaire (verbal communication, mental state, depression, sleep, bladder incontinence, perception alterations) Motor symptoms: UPDRS (I–IV) Others: disability NMS: depression, anxiety and sleep	Caregiver's QoL: Scale of Quality of Life of Caregivers
Pal et al (2004) [38]	40 PD patients and 23 caregivers, Canada, correlations		Caregiver's burden: caregiver's depression, anxiety and sleep
Martinez-Martin et al (2005) [39]	57 dyads, Spain, multiple regression	Motor symptoms: UPDRS (I–III) Others: disability (ISAPD–ADL); QoL (PDQ-8)	Caregiver's QoL: Scale of Quality of Life of Caregivers
Cifu et al (2006) [40]	49 dyads, USA, correlations	Disease related: disease severity NMS: cognition Motor symptoms: motor examination Others: disability	Caregiver's burden: Zarit Caregiver Burden Inventory; Caregiver Distress Scale
Schrag et al (2006) [41]	123 caregivers, UK, correlations	Disease related: disease duration and severity NMS: depression, hallucinations, confusion, falls Others: disability, QoL Others: patient's age	Caregiver's burden: Caregiver Burden Inventory, depression (BDI) Caregiver's QoL: Scale of Quality of life of Caregivers Caregiver's QoL: WHOQOL-BREF
Schestatsky et al (2006) [42]	21 dyads, Brazil, correlations		

Figure 2.2 Studies about patient's factors influencing the caregiver's quality of life (continues overleaf).

Studies about patient's factors influencing the caregiver's quality of life (continued)

Reference	Study type: sample, country, design	Patient factors: disease related, NMS, motor symptoms, others	Outcome measure: caregiver's burden; caregiver's QoL
Aarsland et al (2007) [43]	94 caregivers, Norway, multiple regression	NMS: depression, cognition (Relative Stress Scale), depression, neuropsychiatric symptoms, cognition (caregiver's depression and distress) NMS: agitation, psychosis, mood	Caregiver's burden: Relative Stress Scale; depression (BDI); distress (GHQ-12) Caregiver's burden: caregiver's distress (Neuropsychiatric Inventory)
Aarsland et al (2007) [43]	537, multinational, descriptive		
Kim et al (2007) [44]	76 caregivers, Korea, group differences	Disease related: disease duration (with subjective burden) and severity (with objective burden) Others: disability (UPDRS II with objective and subjective burden)	Caregiver's burden: subjective and objective Burden Scale
Martinez-Martin et al (2007) [45]	80 dyads, Spain, multiple regression	Disease related: disease severity NMS: mood (depression, anxiety) Others: QoL (EQ-5D), disability	Caregiver's burden: Zarit Caregiver Burden Inventory, HADS Caregiver's QoL: SF-36, EQ-5D
Carter et al (2008) [46]	219 caregivers, USA, multiple regression	Disease related: cognition (delayed recall), depression Others: disability (UPDRS-ADL)	Caregiver's burden: Family Care (giving) Inventory; depression (CES-D)
Lokk (2008) [47]	404 caregivers, Sweden, structured interview	Disease related: disease duration NMS: cognition, hallucinations, memory, bradyphrenia, sleep Motor symptoms: motor dysfunction, motor complications	Caregiver's burden: questions about perceived worry, stress, mood and sleep Caregiver's QoL: EQ-5D visual analogue scale (0–10)

Martinez-Martin et al (2008) [48]	286 dyads, Spain, multiple regression	<p>Disease related: disease severity (with caregiver's burden and depression)</p> <p>NMS: sleep, cognition, NMS (with caregiver's burden), NMS (with caregiver's QoL)</p> <p>Motor symptoms: motor complications (with caregiver's burden)</p> <p>Others: disability, QoL, age (with caregiver's burden), patient's QoL (with caregiver's QoL)</p> <p>NMS: behavioral problems (with burden); cognitive impairment (with QoL)</p> <p>Others: patient-caregiver relationship (with burden); perceived social support (with burden); frequency of breaks (with QoL)</p> <p>Disease related: disease severity and age of onset (dementia group)</p> <p>NMS: neuropsychiatric symptoms</p> <p>Others: UPDRS functional (dementia and depression patient groups)</p> <p>Others: motor fluctuations, safety</p>	<p>Caregiver's burden: Zarit Caregiver Burden Inventory, caregiver's anxiety and depression (HAD5)</p> <p>Caregiver's QoL: EQ-5D</p> <p>Caregiver's burden: Caregiver Burden Inventory</p> <p>Caregiver's QoL: Scale of Quality of Life of Caregivers</p> <p>Caregiver's burden: distress score of Neuropsychiatric Inventory</p> <p>Caregiver's burden: repertory grid</p>
D'Amelio et al (2009) [49]	Caregivers, Italy, path analysis		
Stella et al (2009) [50]	50 dyads, Brazil, correlations		
Roland et al (2010) [51]	5 spouses, Canada, semi-qualitative		

Figure 2.2 Studies about patient's factors influencing the caregiver's quality of life (continued). BDI, Beck Depression Inventory; CES-D, Center for Epidemiologic Studies Depression Scale; GDS, Geriatric Depression Scale; GHQ2, General Health Questionnaire-2; HADS, Hospital Anxiety and Depression Scale; HAM-D, Hamilton Rating Scale for Depression; H&Y, Hoehn and Yahr staging; ISAPD, Intermediate Scale for Assessment of Parkinson's Disease; NMS, non-motor symptoms; PD, Parkinson's disease; PDQ-8, Parkinson's disease Questionnaire – 8 items; QoL, quality of life; SF-36, Short Form Health Survey – 36 items; UPDRS, Unifies Parkinson's Disease Rating Scale; WHOQOL-BREF, World Health Organization Quality of Life – Brief.

Burden Scale [58]. Measures of specific symptoms or general distress have also been used to measure caregivers' burden.

The disease-related factors that most affect caregivers' burden are severity and duration. Almost half of the revised studies identified the patient's disability as an important determinant of the caregiver's burden. However, few studies found a significant association between motor dysfunction and the caregiver's burden [37,40,47,48]. Relative to NMSs, the caregiver's burden seems to be mostly affected by the patient's depression and impaired cognitive function, followed by sleep disturbances and hallucinations. A study specifically comparing the effect of motor and non-motor symptoms in the caregiver found that a patient's depression and cognitive impairment had a greater impact on a caregiver's depression and strain than a patient's disability [46]. Research about the impact of NMSs, using specific NMS measures, on caregivers' burdens and QoL is needed.

Economic costs

PD is the second most common neurodegenerative disorder; it increases with age and affects about 1 million people in the USA and about 5 million people worldwide [59]. Given the progressive nature of PD, such an increase will have a significant impact on the burden of the health-care system and caregivers, with considerable caregiving required in the later stages of PD [59].

NMSs range from autonomic and neuropsychiatric symptoms to sleep disturbances and sensory problems. Although cognitive impairment has a lesser impact compared with other NMSs, it continues to be one of the major reasons for admission to institutional care [60,61]. Autonomic dysfunction can occur early in the disease and may not necessarily be related to disease severity. One study reported that half the sample of hospitalized PD patients rated autonomic symptoms as 'a lot' or 'very much' [62].

In the UK, the annual direct costs for patients living in full-time institutional care was estimated at about £19 338, five times higher than in the home setting [63]. Costs for NMS drug therapy increase with clinical progression of symptoms, thus contributing to higher economic costs, especially for patients with a fixed income from pensions and govern-

ment benefits [61,64]. Economic costs are doubled by each H&Y stage, especially after stage 2.0 [65].

NMSs also account for the burden of hidden costs such as sick leave, early retirement and informal care not only for patients but also for caregivers in certain instances. The cost burden of NMSs is significantly high, especially in patients with advanced PD and increasingly severe symptoms, for whom there is a poorer QoL, reduced productivity and a greater need for health-care services, which in turn have an impact on direct and indirect costs.

Thus, identifying disease-modifying treatments early in the disease, before any functional or motor disability appears, is critical in reducing costs and preserving QoL. Evidence suggests that initial therapy with non-levodopa agents is cost-effective, prolongs time to levodopa initiation and delays the onset of dyskinesias [66,67]. Results of the ADAGIO trial hint that rasagiline can reduce the progression of NMSs, as assessed using Movement Disorder Society (MDS)-UPDRS [68–70].

Conclusions

There is a significant interrelationship of severity of disease, QoL, patient's and caregiver's burden, and cost of illness. NMSs contribute to the overall PD burden, which is a major determinant of QoL. An increasing awareness of 'at-risk' individuals, based on detection of some or a combination of NMSs, is essential for an early identification of PD patients. Identification of prodromal patients plays a key factor in preventing the burden of economic costs and improving QoL of patients and caregivers. Keeping this in mind, it becomes important to optimize the management of all aspects of NMSs in PD.

References

- 1 Shiba M, Bower JH, Maraganore DM, et al. Anxiety disorders and depressive disorders preceding Parkinson's disease: a case-control study. *Mov Disord* 2000;15:669-677.
- 2 Iranzo A, Molinuevo JL, Santamaría J, et al. Rapid-eye-movement sleep behavior disorder as an early marker for a neurodegenerative disorder: a descriptive study. *Lancet Neurol* 2006;5:572-577.
- 3 Antonini A. Non-motor symptoms in Parkinson's disease. *Eur Neurol Rev* 2009;4:25-27.
- 4 Mueller A, Abolmaali ND, Hakimi AR, et al. Olfactory bulb volumes in patients with idiopathic Parkinson's disease: a pilot study. *J Neural Transm* 2005;112:1363-1370.
- 5 Berendse HW. 1.4 Early diagnosis in PD: olfaction. *Parkinsonism Rel Disord* 2006;12:1-2.

- 6 Miyasaki JM, Shannon K, Voon V, et al. Practice Parameter: evaluation and treatment of depression, psychosis, and dementia in Parkinson disease (an evidence-based review): report of the Quality Standards Subcommittee of the American Academy of Neurology. *Neurology* 2006;66:996-1002.
- 7 Forjaz MJ, Frades-Payo B, Martinez-Martin P. [The current state of the art concerning quality of life in Parkinson's disease: II. Determining and associated factors.] *Rev Neurol* 2009;49:655-660.
- 8 Frades-Payo B, Forjaz MJ, Martinez-Martin P. [The current state of the art concerning quality of life in Parkinson's disease: I. Instruments, comparative studies and treatments]. *Rev Neurol* 2009;49:594-598.
- 9 Chaudhuri KR, Martinez-Martin P. Quantitation of non-motor symptoms in Parkinson's disease. *Eur J Neurol* 2008;15(suppl 2):2-7.
- 10 Chaudhuri KR, Martinez-Martin P, Schapira AH, et al. International multicenter pilot study of the first comprehensive self-completed nonmotor symptoms questionnaire for Parkinson's disease: the NMSQuest study. *Mov Disord* 2006;21:916-923.
- 11 Martinez-Martin P, Schapira AH, Stocchi F, et al. Prevalence of nonmotor symptoms in Parkinson's disease in an international setting; study using nonmotor symptoms questionnaire in 545 patients. *Mov Disord* 2007;22:1623-1629.
- 12 Chaudhuri KR, Prieto-Jurcynska C, Naidu Y, et al. The nondeclaration of nonmotor symptoms of Parkinson's disease to health care professionals: an international study using the nonmotor symptoms questionnaire. *Mov Disord* 2010;25:697-701.
- 13 Wang G, Wan Y, Cheng Q, et al. Malnutrition and associated factors in Chinese patients with Parkinson's disease: Results from a pilot investigation. *Parkinsonism Relat Disord* 2010;16:119-123.
- 14 Cervantes-Arriaga A, Rodriguez-Violante M, Villar-Velarde A, Lopez-Gomez M, Corona T. [Metric properties of clinimetric indexes for non-motor dysfunction of Parkinson's disease in Mexican population.] *Rev Invest Clin* 2010;62:8-14.
- 15 Chaudhuri KR, Martinez-Martin P, Brown RG, et al. The metric properties of a novel non-motor symptoms scale for Parkinson's disease: Results from an international pilot study. *Mov Disord* 2007;22:1901-1911.
- 16 Honig H, Antonini A, Martinez-Martin P, et al. Intrajejunal levodopa infusion in Parkinson's disease: a pilot multicenter study of effects on nonmotor symptoms and quality of life. *Mov Disord* 2009;24:1468-1474.
- 17 Kim HJ, Park SY, Cho YJ, et al. Nonmotor symptoms in de novo Parkinson disease before and after dopaminergic treatment. *J Neurol Sci* 2009;287:200-204.
- 18 Martinez-Martin P, Rodriguez-Blazquez C, Abe K, et al. International study on the psychometric attributes of the non-motor symptoms scale in Parkinson disease. *Neurology* 2009;73:1584-1591.
- 19 Wang G, Hong Z, Cheng Q, et al. Validation of the Chinese Non-Motor Symptoms Scale for Parkinson's disease: Results from a Chinese pilot study. *Clin Neurol Neurosurgery* 2009;111:523-526.
- 20 Barone P, Antonini A, Colosimo C, et al. The PRIAMO study: A multicenter assessment of nonmotor symptoms and their impact on quality of life in Parkinson's disease. *Mov Disord* 2009;24:1641-1649.
- 21 Colosimo C, Morgante L, Antonini A, et al. Non-motor symptoms in atypical and secondary parkinsonism: the PRIAMO study. *J Neurol* 2010;257:5-14.
- 22 Hoehn MM, Yahr MD. Parkinsonism: onset, progression, and mortality. *Neurology* 1967;17:427-442.
- 23 Global Parkinson's Disease Survey Steering Committee. Factors impacting on quality of life in Parkinson's disease: results from an international survey. *Mov Disord* 2002;17:60-67.
- 24 Lohle M, Storch A, Reichmann H. Beyond tremor and rigidity: non-motor features of Parkinson's disease. *J Neural Transm* 2009;116:1483-1492.
- 25 Wolters EC. Non-motor extranigral signs and symptoms in Parkinson's disease. *Parkinsonism Relat Disord* 2009;15(suppl 3):S6-S12.

- 26 Qin Z, Zhang L, Sun F, et al. Health related quality of life in early Parkinson's disease: Impact of motor and non-motor symptoms, results from Chinese levodopa exposed cohort. *Parkinsonism. Rel Disord* 2009;15:767-771.
- 27 Chaudhuri KR, Healy, DG, Schapira AH. Non-motor symptoms of Parkinson's disease: diagnosis and management. *Lancet Neurol* 2006;5:235-245.
- 28 Miller E, Berrios GE, Politynska BE. Caring for someone with Parkinson's disease: factors that contribute to distress. *Int J Geriatr Psychiatry* 1998;11:263-268.
- 29 Carter JH, Stewart BJ, Archbold PG, et al. Living with a person who has Parkinson's disease: the spouse's perspective by stage of disease. *Mov Disord* 2004;13:20-28.
- 30 Meara J, Mitchelmore E, Hobson P. Use of the GDS-15 geriatric depression scale as a screening instrument for depressive symptomatology in patients with Parkinson's disease and their carers in the community. *Age Aging* 1999;28:35-38.
- 31 Habermann B. Spousal perspective of Parkinson's disease in middle life. *J Adv Nurs* 2000;31:1409-1415.
- 32 Fernandez HH, Tabamo REJ, David RR, Friedman JH. Predictors of depressive symptoms among spouse caregivers in Parkinson's disease. *Mov Disord* 2001;16:1123-1125.
- 33 Caap-Ahlgren M, Dehlin O. Factors of importance to the caregiver burden experienced by family caregivers of Parkinson's disease patients. *Aging Clin Exp Res* 2002;14:371-377.
- 34 Edwards NE, Scheetz PS. Predictors of burden for caregivers of patients with Parkinson's disease. *J Neurosci Nurs* 2002;34:184-190.
- 35 Happe S, Berger K. The association between caregiver burden and sleep disturbances in partners of patients with Parkinson's disease. *Age Ageing* 2002;31:349-354.
- 36 Thommessen B, Aarsland D, Braekhus A, Oksengaard AR, Engedal K, Laake K. The psychosocial burden on spouses of the elderly with stroke, dementia and Parkinson's disease. *Int J Geriatr Psychiatry* 2002;17:78-84.
- 37 Martinez-Martin P, Guerrero-Diaz MT, Frades-Payo B. [Neuropsychological disorders in Parkinson's disease: evaluating them and their impact on the caregiver]. *Rev Neurol* 2004;39:639-645.
- 38 Pal PK, Thennarasu K, Fleming J, Schulzer M, Brown T, Calne SM. Nocturnal sleep disturbances and daytime dysfunction in patients with Parkinson's disease and in their caregivers. *Parkinsonism Relat Disord* 2004;10:157-168.
- 39 Martinez-Martin P, Benito-Leon J, Alonso F, et al. Quality of life of caregivers in Parkinson's disease. *Qual Life Res* 2005;14:463-472.
- 40 Cifu DX, Carne W, Brown R, et al. Caregiver distress in parkinsonism. *J Rehabil Res Dev* 2006;43:499-508.
- 41 Schrag A, Hovris A, Morley D, Quinn N, Jahanshahi M. Caregiver-burden in Parkinson's disease is closely associated with psychiatric symptoms, falls, and disability. *Parkinsonism Relat Disord* 2006;12:35-41.
- 42 Schestatsky P, Zanatto VC, Margis R, et al. Quality of life in a Brazilian sample of patients with Parkinson's disease and their caregivers. *Rev Bras Psiquiatr* 2006;28:209-211.
- 43 Aarsland D, Bronnick K, Ehrt U, et al. Neuropsychiatric symptoms in patients with Parkinson's disease and dementia: frequency, profile and associated care giver stress. *J Neurol Neurosurg Psychiatr* 2007;78:36.
- 44 Kim KS, Kim BJ, Kim KH, et al. Subjective and objective caregiver burden in Parkinson's disease. *Taehan Kanho Hakhoe Chi* 2007;37:242-248.
- 45 Martinez-Martin P, Forjaz MJ, Frades-Payo B, et al. Caregiver burden in Parkinson's disease. *Mov Disord* 2007a;22:924-931.
- 46 Carter JH, Stewart BJ, Lyons, KS, Archbold, PG. Do motor and nonmotor symptoms in PD patients predict caregiver strain and depression? *Mov Disord* 2008;23:1211-1216.
- 47 Lökk J. Caregiver strain in Parkinson's disease and the impact of disease duration. *Eur J Phys Rehabil Med* 2008;44:39-45.

- 48 Martinez-Martin P, Arroyo S, Rojo-Abuin JM, Rodriguez-Blazquez C, Frades B, de Pedro CJ. Burden, perceived health status, and mood among caregivers of Parkinson's disease patients. *Mov Disord* 2008;23:1673-1680.
- 49 D'Amelio M, Terruso V, Palmeri B, et al. Predictors of caregiver burden in partners of patients with Parkinson's disease. *Neurol Sci* 2009;30:171-174.
- 50 Stella F, Banzato CE, Quagliato EM, Viana MA, Christofoletti G. Psychopathological features in patients with Parkinson's disease and related caregivers' burden. *Int J Geriatr Psychiatry* 2009;24:1158-1165.
- 51 Roland KP, Jenkins ME, Johnson AM. An exploration of the burden experienced by spousal caregivers of individuals with Parkinson's disease. *Mov Disord* 2010;25:189-193.
- 52 Ware JE Jr, Sherbourne CD. The MOS 36-item short-form health survey (SF-36): I. Conceptual framework and item selection. *Med Care* 1992;30:473-483.
- 53 EuroQol Group. EuroQol-a new facility for the measurement of health-related quality of life. *Health Policy* 1990;16:199-208.
- 54 WHOQOL Group. Development of the World Health Organization WHOQOL-BREF quality of life assessment. *Psychol Med* 1998;28:551-558.
- 55 Glozman JM, Bicheva KG, Fedorova NV. Scale of quality of life of care-givers (SQLC). *J Neurol* 1998;245:39-41.
- 56 Zarit SH, Reever KE, Bach-Peterson J. Relatives of the impaired elderly: correlates of feelings of burden. *Gerontologist* 1980;20:649-655.
- 57 Novak M, Guest C. Application of a multidimensional caregiver burden inventory. *The Gerontologist* 1989;29:798-803.
- 58 Montgomery RJV, Gonyea JG, Hooyman NR. Caregiving and the experience of subjective and objective burden. *Family Relations* 1985;34:19-26.
- 59 Olanow CW, Stern MB, Sethi K. The scientific and clinical basis for the treatment of Parkinson disease. *Neurology* 2009;72(suppl 4):1-136.
- 60 Goetz CG, Stebbins GT. Risk factors for nursing home placement in advanced Parkinson's disease. *Neurology* 1993;43:2227-2229.
- 61 Guttman M, Slaughter PM, Theriault ME, DeBoer DP, Naylor CD. Burden of parkinsonism: a population-based study. *Mov Disord* 2002;18:313-319.
- 62 Magerkurth C, Schnitzer R, Braune S. Symptoms of autonomic failure in Parkinson's disease: prevalence and impact on daily life. *Clin Auton Res* 2005;15:76-82.
- 63 Findley L, Aujla M, Bain PG, et al. Direct economic impact of Parkinson's disease: a research survey in the United Kingdom. *Mov Disord* 2003;18:1139-1145.
- 64 Cubo E, Martinez-Martin P, Gonzalez M, Frades B, ELEG Group. Impact of motor and non-motor symptoms on the direct costs of Parkinson's disease [Spanish]. *Neurologia* 2009;24:15-23.
- 65 Dodel RC, Singer M, Kohn-Volland R, et al. The economic impact of Parkinson's disease: An estimation based on a 3-month prospective analysis. *Pharmacoeconomics* 1998;14:299-312.
- 66 Pålhagen S, Heinonen E, Hagglund J, Kaugesaar T, Maki-Ikola O, Palm R. Selegiline slows the progression of the symptoms of Parkinson disease. *Neurology* 2006;66:1200-1206.
- 67 Haycox A, Armand C, Murteira S, Cochran J, François C. Cost effectiveness of rasagiline and pramipexole as treatment strategies in early Parkinson's disease in the UK setting: An economic Markov model evaluation. *Drugs Aging* 2009;26:791-801.
- 68 Parkinson Study Group. A controlled, randomized, delayed-start study of rasagiline in early Parkinson disease. *Arch Neurol* 2004;61:561-566.
- 69 Olanow CW, Rascol O, Hauser R, et al. A double-blind, delayed-start trial of rasagiline in Parkinson's disease. *N Engl J Med* 2009a;361:1268-1278.
- 70 Poewe W, Hauser R, Lang AE. Rasagiline 1 mg/day provides benefits in the progression of non-motor symptoms in patients with early Parkinson's disease: Assessment with the revised MDS-UPDRS. *Mov Disord* 2009;24:S272-S273.

Handbook of Non-Motor Symptoms in Parkinson's
Disease

Chaudhuri, K.R.; Martinez-Martin, P.; Odin, P.; Antonini,
A.

2011, XIII, 97 p. 15 illus. in color., Softcover

ISBN: 978-1-907673-23-8