

Preface

Wild crop relatives have been playing enormously important roles both in the depiction of plant genomes and the genetic improvement of their cultivated counterparts. They have contributed immensely to resolving several fundamental questions, particularly those related to the origin, evolution, phylogenetic relationship, cytological status and inheritance of genes of an array of crop plants; provided several desirable donor genes for the genetic improvement of their domesticated counterparts; and facilitated the innovation of many novel concepts and technologies while working on them directly or while using their resources. More recently, they have even been used for the verification of their potential threats of gene flow from genetically modified plants and invasive habits. Above all, some of them are contributing enormously as model plant species to the elucidation and amelioration of the genomes of crop plant species.

As a matter of fact, as a student, a teacher, and a humble science worker I was, still am and surely will remain fascinated by the wild allies of crop plants for their invaluable wealth for genetics, genomics and breeding in crop plants and as such share a deep concern for their conservation and comprehensive characterization for future utilization. It is by now a well established fact that wild crop relatives deserve serious attention for domestication, especially for the utilization of their phytomedicines and nutraceuticals, bioenergy production, soil reclamation, and the phytoremediation of ecology and environment. While these vastly positive impacts of wild crop relatives on the development and deployment of new varieties for various purposes in the major crop plants of the world agriculture, along with a few negative potential concerns, are envisaged the need for reference books with comprehensive deliberations on the wild relatives of all the major field and plantation crops and fruit and forest trees is indeed imperative. This was the driving force behind the inception and publication of this series.

Unlike the previous six book projects I have edited alone or with co-editors, this time it was very difficult to formulate uniform outlines for the chapters of this book series for several obvious reasons. Firstly, the status of the crop relatives is highly diverse. Some of them are completely wild, some are sporadically cultivated and some are at the initial stage of domestication for specific breeding objectives recently deemed essential. Secondly, the status of their conservation varies widely: some have been conserved, characterized and utilized; some have been eroded completely except for their presence in their center(s) of origin; some are at-risk or endangered due to genetic erosion, and some of them have yet to be explored. The third constraint is the variation in their relative worth, e.g. as academic model, breeding resource, and/or potential as “new crops.”

The most perplexing problem for me was to assign the chapters each on a particular genus to different volumes dedicated to crop relatives of diverse crops grouped based on their utility. This can be exemplified with *Arabidopsis*, which has primarily benefited the Brassicaceae crops but also facilitated genetic analyses and improvement in crop plants in other distant families; or with many wild relatives of forage crops that paved the way for the genetic analyses and breeding of some major cereal and millet crops. The same is true for wild crop relatives such as *Medicago truncatula*, which has paved the way for in-depth research on two crop groups of diverse use: oilseed and pulse crops belonging to the Fabaceae family. The list is too long to enumerate. I had no other choice but to compromise and assign the genera of crop relatives in a volume on the crop group to which they are taxonomically the closest and to which they have relatively greater contributions. For example, I placed the chapter on genus *Arabidopsis* in the volume on oilseeds, which deals with the wild relatives of Brassicaceae crops amongst others.

However, we have tried to include deliberations pertinent to the individual genera of the wild crop relatives to which the chapters are devoted. Descriptions of the geographical locations of origin and genetic diversity, geographical distribution, karyotype and genome size, morphology, etc. have been included for most of them. Their current utility status – whether recognized as model species, weeds, invasive species or potentially cultivable taxa – is also delineated. The academic, agricultural, medicinal, ecological, environmental and industrial potential of both the cultivated and/or wild allied taxa are discussed.

The conservation of wild crop relatives is a much discussed yet equally neglected issue albeit the in situ and ex situ conservations of some luckier species were initiated earlier or are being initiated now. We have included discussions on what has happened and what is happening with regard to the conservation of the crop relatives, thanks to the national and international endeavors, in most of the chapters and also included what should happen for the wild relatives of the so-called new, minor, orphan or future crops.

The botanical origin, evolutionary pathway and phylogenetic relationship of crop plants have always attracted the attention of plant scientists. For these studies morphological attributes, cytological features and biochemical parameters were used individually or in combinations at different periods based on the availability of the required tools and techniques. Access to different molecular markers based on nuclear and especially cytoplasmic DNAs that emerged after 1980 refined the strategies required for precise and unequivocal conclusions regarding these aspects. Illustrations of these classical and recent tools have been included in the chapters.

Positioning genes and defining gene functions required in many cases different cytogenetic stocks, including substitution lines, addition lines, haploids, monoloids and aneuploids, particularly in polyploid crops. These aspects have been dealt in the relevant chapters. Employment of colchiploidy, fluorescent or genomic in situ hybridization and Southern hybridization have reinforced the theoretical and applied studies on these stocks. Chapters on relevant genera/species include details on these cytogenetic stocks.

Wild crop relatives, particularly wild allied species and subspecies, have been used since the birth of genetics in the twentieth century in several instances such as studies of inheritance, linkage, function, transmission and evolution of genes. They have been frequently used in genetic studies since the advent of molecular markers. Their involvement in molecular mapping has facilitated the development of mapping

populations with optimum polymorphism to construct saturated maps and also illuminating the organization, reorganization and functional aspects of genes and genomes. Many phenomena such as genomic duplication, genome reorganization, self-incompatibility, segregation distortion, transgressive segregation and defining genes and their phenotypes have in many cases been made possible due to the utilization of wild species or subspecies. Most of the chapters contain detailed elucidations on these aspects.

The richness of crop relatives with biotic and abiotic stress resistance genes was well recognized and documented with the transfer of several alien genes into their cultivated counterparts through wide or distant hybridization with or without employing embryo-rescue and mutagenesis. However, the amazing revelation that the wild relatives are also a source of yield-related genes is a development of the molecular era. Apomictic genes are another asset of many crop relatives that deserve mention. All of these past and the present factors have led to the realization that the so-called inferior species are highly superior in conserving desirable genes and can serve as a goldmine for breeding elite plant varieties. This is particularly true at a point when natural genetic variability has been depleted or exhausted in most of the major crop species, particularly due to growing and promoting only a handful of so-called high-yielding varieties while disregarding the traditional cultivars and landraces. In the era of molecular breeding, we can map desirable genes and polygenes, identify their donors and utilize tightly linked markers for gene introgression, mitigating the constraint of linkage drag, and even pyramid genes from multiple sources, cultivated or wild taxa. The evaluation of primary, secondary and tertiary gene pools and utilization of their novel genes is one of the leading strategies in present-day plant breeding. It is obvious that many wide hybridizations will never be easy and involve near-impossible constraints such as complete or partial sterility. In such cases gene cloning and gene discovery, complemented by intragenic breeding, will hopefully pave the way for success. The utilization of wild relatives through traditional and molecular breeding has been thoroughly enumerated over the chapters throughout this series.

Enormous genomic resources have been developed in the model crop relatives, for example *Arabidopsis thaliana* and *Medicago truncatula*. BAC, cDNA and EST libraries have also been developed in some other crop relatives. Transcriptomes and metabolomes have also been dissected in some of them. However, similar genomic resources are yet to be constructed in many crop relatives. Hence this section has been included only in chapters on the relevant genera.

In this book series, we have included a section on recommendations for future steps to create awareness about the wealth of wild crop relatives in society at large and also for concerns for their alarmingly rapid decrease due to genetic erosion. The authors of the chapters have also emphasized on the imperative requirement of their conservation, envisaging the importance of biodiversity. The importance of intellectual property rights and also farmers' rights as owners of local landraces, botanical varieties, wild species and subspecies has also been dealt in many of the chapters.

I feel satisfied that the authors of the chapters in this series have deliberated on all the crucial aspects relevant to a particular genus in their chapters.

I am also very pleased to present many chapters in this series authored by a large number of globally reputed leading scientists, many of whom have contributed to the development of novel concepts, strategies and tools of genetics, genomics and breeding and/or pioneered the elucidation and improvement of particular plant

genomes using both traditional and molecular tools. Many of them have already retired or will be retiring soon, leaving behind their legacies and philosophies for us to follow and practice. I am saddened that a few of them have passed away during preparation of the manuscripts for this series. At the same time, I feel blessed that all of these stalwarts shared equally with me the wealth of crop relatives and contributed to their recognition and promotion through this endeavor.

I would also like to be candid with regard to my own limitations. Initially I planned for about 150 chapters devoted to the essential genera of wild crop relatives. However, I had to exclude some of them either due to insignificant progress made on them during the preparation of this series, my failure to identify interested authors willing to produce acceptable manuscripts in time or authors' backing out in the last minute, leaving no time to find replacements. I console myself for this lapse with the rationale that it is simply too large a series to achieve complete satisfaction on the contents. Still I was able to arrange about 125 chapters in the ten volumes, contributed by nearly 400 authors from over 40 countries of the world. I extend my heartfelt thanks to all these scientists, who have cooperated with me since the inception of this series not only with their contributions, but also in some cases by suggesting suitable authors for chapters on other genera. As happens with a mega-series, a few authors had delays for personal or professional reasons, and in a few cases, for no reason at all. This caused delays in the publication of some of the volumes and forced the remaining authors to update their manuscripts and wait too long to see their manuscripts in published form. I do shoulder all the responsibilities for this myself and tender my sincere apologies.

Another unique feature of this series is that the authors of chapters dedicated to some genera have dedicated their chapters to scientists who pioneered the exploration, description and utilization of the wild species of those genera. We have duly honored their sincere decision with equal respect for the scientists they rightly reminded us to commemorate.

Editing this series was, to be honest, very taxing and painstaking, as my own expertise is limited to a few cereal, oilseed, pulse, vegetable, and fruit crops, and some medicinal and aromatic plants. I spent innumerable nights studying to attain the minimum eligibility to edit the manuscripts authored by experts with even life-time contributions on the concerned genera or species. However, this indirectly awakened the "student-for-life" within me and enriched my arsenal with so many new concepts, strategies, tools, techniques and even new terminologies! Above all, this helped me to realize that individually we know almost nothing about the plants on this planet! And this realization strikingly reminded me of the affectionate and sincere advice of Dr. Norman Borlaug to keep abreast with what is happening in the crop sciences, which he used to do himself even when he had been advised to strictly limit himself to bed rest. He was always enthusiastic about this series and inspired me to take up this huge task. This is one of the personal and professional reasons I dedicated this book series to him with a hope that the present and future generations of plant scientists will share the similar feelings of love and respect for all plants around us for the sake of meeting our never-ending needs for food, shelter, clothing, medicines, and all other items used for our basic requirements and comfort. I am also grateful to his granddaughter, Julie Borlaug, for kindly extending her permission to dedicate this series to him.

I started editing books with the 7-volume series on Genome Mapping and Molecular Breeding in Plants with Springer way back in 2005, and I have since

edited many other book series with Springer. I always feel proud and satisfied to be a member of the Springer family, particularly because of my warm and enriching working relationship with Dr. Sabine Schwarz and Dr. Jutta Lindenborn, with whom I have been working all along. My special thanks go out to them for publishing this “dream series” in an elegant form and also for appreciating my difficulties and accommodating many of my last-minute changes and updates.

I would be remiss in my duties if I failed to mention the contributions of Phullara – my wife, friend, philosopher and guide – who has always shared with me a love of the collection, conservation, evaluation, and utilization of wild crop relatives and has enormously supported me in the translation of these priorities in my own research endeavors – for her assistance in formulating the contents of this series, for monitoring its progress and above all for taking care of all the domestic and personal responsibilities I am supposed to shoulder. I feel myself alien to the digital world that is the sine qua non today for maintaining constant communication and ensuring the preparation of manuscripts in a desirable format. Our son Sourav and daughter Devleena made my life easier by balancing out my limitations and also by willingly sacrificing the spare amount of time I ought to spend with them. Editing of this series would not be possible without their unwavering support.

I take the responsibility for any lapses in content, format and approach of the series and individual volumes and also for any other errors, either scientific or linguistic, and will look forward to receiving readers’ corrections or suggestions for improvement.

As I mentioned earlier this series consists of ten volumes. These volumes are dedicated to wild relatives of Cereals, Millets and Grasses, Oilseeds, Legume Crops and Forages, Vegetables, Temperate Fruits, Tropical and Subtropical Fruits, Industrial Crops, Plantation and Ornamental Crops, and Forest Trees.

This volume “Wild Crop Relatives – Genomic and Breeding Resources: Oilseeds” includes 17 chapters dedicated to *Arabidopsis*, *Brassica*, *Capsella*, *Carthamus*, *Crambe*, *Cuphea*, *Diplotaxis*, *Eruca*, *Helianthus*, *Hirschfeldia*, *Linum*, *Moricandia*, *Orychophragmus*, *Pachycladon*, *Ricinus*, *Sesamum* and *Sinapsis*. The chapters of this volume were authored by 33 scientists from 11 countries of the world including Canada, China, France, Germany, India, Italy, New Zealand, Poland, Spain, UK and USA.

It is my sincere hope that this volume and the series as a whole will serve the requirements of students, scientists and industries involved in studies, teaching, research and the extension of oilseed crops with an intention of serving science and society.

Clemson, USA

Chittaranjan Kole

Wild Crop Relatives: Genomic and Breeding Resources
Oilseeds

Kole, C. (Ed.)

2011, XXV, 295 p., Hardcover

ISBN: 978-3-642-14870-5