

# Contents

|                                                   |                                                                                                                                                             |    |
|---------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------|----|
| <b>1</b>                                          | <b>Introduction</b>                                                                                                                                         | 1  |
|                                                   | References                                                                                                                                                  | 5  |
| <br><b>Part I Synthesis of Indole Derivatives</b> |                                                                                                                                                             |    |
| <b>2</b>                                          | <b>Construction of 2-(Aminomethyl)indoles through Copper-Catalyzed Domino Three-Component Coupling and Cyclization</b>                                      | 9  |
| 2.1                                               | Introduction                                                                                                                                                | 9  |
| 2.1.1                                             | Synthesis of 2-(Aminomethyl)indoles Using Several Amines and Aldehydes                                                                                      | 10 |
| 2.1.2                                             | Synthesis of Substituted 2-(Aminomethyl)indoles Using Various Ethynylanilines and Secondary Amines                                                          | 13 |
| 2.1.3                                             | Construction of Polycyclic Indoles by Palladium-Catalyzed C–H Functionalization                                                                             | 14 |
| 2.1.4                                             | Synthetic Application to Calindol, Benzo[ <i>e</i> ][1,2]thiazines, and Indene                                                                              | 14 |
| 2.2                                               | Experimental Section                                                                                                                                        | 19 |
| 2.2.1                                             | General Methods                                                                                                                                             | 19 |
| 2.2.2                                             | General Procedure for Synthesis of 2-(Aminomethyl)indole                                                                                                    | 24 |
| 2.2.3                                             | General Procedure for Synthesis of Tetrahydropyridine-Fused Indole                                                                                          | 33 |
| 2.2.4                                             | General Procedure for Synthesis of Benzo[ <i>e</i> ][1,2]thiazine-1,1-dioxide                                                                               | 41 |
|                                                   | References                                                                                                                                                  | 44 |
| <b>3</b>                                          | <b>Facile Synthesis of 1,2,3,4-Tetrahydro-<math>\beta</math>-Carbolines by One-Pot Domino Three-Component Indole Formation and Nucleophilic Cyclization</b> | 47 |
| 3.1                                               | Experimental Section                                                                                                                                        | 52 |
| 3.1.1                                             | General Methods                                                                                                                                             | 52 |

|        |                                                                                                                                                                                                                                                                                                                                                                                                              |    |
|--------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----|
| 3.1.2  | General Procedure for Synthesis of<br><i>N</i> -Arylsulfonyl-2-ethynylaniline: Synthesis of<br>2-Ethynyl- <i>N</i> -mesitylenesulfonylaniline (1c). . . . .                                                                                                                                                                                                                                                  | 53 |
| 3.1.3  | <i>N</i> -( <i>p</i> -Bromobenzenesulfonyl)-2-ethynylaniline (1e) . . . .                                                                                                                                                                                                                                                                                                                                    | 53 |
| 3.1.4  | <i>N</i> -( <i>p</i> -Chlorobenzenesulfonyl)-2-ethynylaniline (1f). . . .                                                                                                                                                                                                                                                                                                                                    | 54 |
| 3.1.5  | 2-Ethynyl- <i>N</i> -( <i>p</i> -fluorosulfonyl)aniline (1g) . . . . .                                                                                                                                                                                                                                                                                                                                       | 54 |
| 3.1.6  | General Procedure for Synthesis of<br>1,2,3,4-Tetrahydro- $\beta$ -carboline by Domino<br>Copper-Catalyzed Three-Component Indole<br>Formation and Cyclization with <i>t</i> -BuOK: Synthesis of<br>2-Methyl-1-propyl-2,3,4,9-tetrahydro-1 <i>H</i> -pyrido[3,4- <i>b</i> ]-<br>indole (6a) and 2-Methyl-1-propyl-1,2,3,4-<br>tetrahydropyrazino[1,2- <i>a</i> ]indole (8a)<br>(Table 1, Entry 11) . . . . . | 54 |
| 3.1.7  | 2-Methyl-1-[2-(trimethylsilyl)ethenyl]-2,3,4,9-<br>tetrahydro-1 <i>H</i> -pyrido[3,4- <i>b</i> ]indole (6b) and<br>2-Methyl-1-[2(trimethylsilyl)ethenyl]-1,2,3,4-<br>tetrahydropyrazino[1,2- <i>a</i> ]indole (8b) . . . . .                                                                                                                                                                                 | 55 |
| 3.1.8  | 1-(Benzyloxymethyl)-2-methyl-2,3,4,9-tetrahydro-<br>1 <i>H</i> -pyrido[3,4- <i>b</i> ]indole (6c) and 1-(Benzyloxymethyl)-<br>2-methyl-1,2,3,4-tetrahydropyrazino[1,2- <i>a</i> ]indole (8c). . .                                                                                                                                                                                                            | 56 |
| 3.1.9  | 2-Methyl-2,3,4,9-tetrahydro-1 <i>H</i> -pyrido[3,4- <i>b</i> ]-<br>indole (6d) . . . . .                                                                                                                                                                                                                                                                                                                     | 56 |
| 3.1.10 | General Procedure for Synthesis of<br>1,2,3,4-Tetrahydro- $\beta$ -carboline by Domino<br>Copper-Catalyzed Three-Component Indole Formation<br>and Cyclization with MsOH: Synthesis of<br>2-Methyl-2,3-dihydropyrido[3,4- <i>b</i> ]indol-4(9 <i>H</i> )-one (7a)<br>(Conditions A) . . . . .                                                                                                                | 57 |
| 3.1.11 | 2-Allyl-2,3-dihydro-1 <i>H</i> -pyrido[3,4- <i>b</i> ]indol-4(9 <i>H</i> )-<br>one (7b) . . . . .                                                                                                                                                                                                                                                                                                            | 57 |
| 3.1.12 | 2-Butyl-2,3-dihydro-1 <i>H</i> -pyrido[3,4- <i>b</i> ]indol-4(9 <i>H</i> )-<br>one (7c) . . . . .                                                                                                                                                                                                                                                                                                            | 58 |
| 3.1.13 | General Procedure for Synthesis of<br>1,2,3,4-Tetrahydro- $\beta$ -carboline by Domino<br>Copper-Catalyzed Three-Component Indole<br>Formation and Cyclization by MsOH: Synthesis of<br>2-Benzyl-2,3-dihydro-1 <i>H</i> -pyrido[3,4- <i>b</i> ]indol-4(9 <i>H</i> )-one<br>(7d) (Conditions B) . . . . .                                                                                                     | 58 |
| 3.1.14 | ( <i>R</i> )-2,3-Dimethyl-2,3-dihydro-1 <i>H</i> -pyrido[3,4- <i>b</i> ]indol-<br>4(9 <i>H</i> )-one (7e) (Conditions C) . . . . .                                                                                                                                                                                                                                                                           | 59 |
| 3.1.15 | ( <i>R</i> )-3-Isobutyl-2-methyl-2,3-dihydro-<br>1 <i>H</i> -pyrido[3,4- <i>b</i> ]indol-4(9 <i>H</i> )-one (7f). . . . .                                                                                                                                                                                                                                                                                    | 59 |

|          |                                                                                                                                                                                                                                                                     |           |
|----------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------|
| 3.1.16   | ( <i>R</i> )-3-Benzyl-2-methyl-2,3-dihydro-1 <i>H</i> -pyrido[3,4- <i>b</i> ]indol-4(9 <i>H</i> )-one ( <b>7g</b> ) . . . . .                                                                                                                                       | 59        |
| 3.1.17   | 5,6,8,9,10,11,11 <i>a</i> ,12-Octahydroindolo[3,2- <i>b</i> ]quinolizine ( <b>7h</b> ) . . . . .                                                                                                                                                                    | 60        |
|          | References . . . . .                                                                                                                                                                                                                                                | 60        |
| <b>4</b> | <b>Concise Synthesis of Indole-Fused 1,4-Diazepines through Copper(I)-Catalyzed Domino Three-Component Coupling–Cyclization–<i>N</i>-Arylation under Microwave Irradiation . . . . .</b>                                                                            | <b>63</b> |
| 4.1      | Experimental Section . . . . .                                                                                                                                                                                                                                      | 67        |
| 4.1.1    | General Methods . . . . .                                                                                                                                                                                                                                           | 67        |
| 4.1.2    | General Procedure for Synthesis of 2-Ethynyl- <i>N</i> -methanesulfonylaniline: Synthesis of 2-Ethynyl- <i>N</i> -methanesulfonyl-4-methoxycarbonylaniline ( <b>1b</b> ) . . . . .                                                                                  | 68        |
| 4.1.3    | 2-Ethynyl- <i>N</i> -methanesulfonyl-4-trifluoromethyl-carbonylaniline ( <b>1c</b> ) . . . . .                                                                                                                                                                      | 68        |
| 4.1.4    | 2-Ethynyl- <i>N</i> -methanesulfonyl-4-methylaniline ( <b>1d</b> ) . . . . .                                                                                                                                                                                        | 69        |
| 4.1.5    | 2-Ethynyl- <i>N</i> -methanesulfonyl-5-trifluoromethyl-carbonylaniline ( <b>1e</b> ) . . . . .                                                                                                                                                                      | 69        |
| 4.1.6    | General Procedure for Synthesis of Indole-Fused 1,4-Diazepine through Three-Component Indole Formation- <i>N</i> -Arylation: Synthesis of 7- <i>n</i> -Butyl-7,8-dihydro-6 <i>H</i> -benzo[ <i>f</i> ]indolo[1,2- <i>a</i> ]-[1,4]diazepine ( <b>3a</b> ) . . . . . | 69        |
| 4.1.7    | 7-Methyl-7,8-dihydro-6 <i>H</i> -benzo[ <i>f</i> ]indolo[1,2- <i>a</i> ]-[1,4]diazepine ( <b>3b</b> ) . . . . .                                                                                                                                                     | 70        |
| 4.1.8    | 7-Benzyl-7,8-dihydro-6 <i>H</i> -benzo[ <i>f</i> ]indolo[1,2- <i>a</i> ]-[1,4]diazepine ( <b>3c</b> ) . . . . .                                                                                                                                                     | 70        |
| 4.1.9    | 7-Allyl-7,8-dihydro-6 <i>H</i> -benzo[ <i>f</i> ]indolo[1,2- <i>a</i> ]-[1,4]diazepine ( <b>3d</b> ) . . . . .                                                                                                                                                      | 71        |
| 4.1.10   | 7-Allyl-3-methoxycarbonyl-7,8-dihydro-6 <i>H</i> -benzo[ <i>f</i> ]indolo[1,2- <i>a</i> ][1,4]diazepine ( <b>3e</b> ) . . . . .                                                                                                                                     | 71        |
| 4.1.11   | 7-Allyl-3-trifluoromethyl-7,8-dihydro-6 <i>H</i> -benzo[ <i>f</i> ]indolo[1,2- <i>a</i> ][1,4]di-azepine ( <b>3f</b> ) . . . . .                                                                                                                                    | 71        |
| 4.1.12   | 7-Allyl-3-methyl-7,8-dihydro-6 <i>H</i> -benzo[ <i>f</i> ]indolo[1,2- <i>a</i> ][1,4]diazepine ( <b>3g</b> ) . . . . .                                                                                                                                              | 72        |
| 4.1.13   | 7-Allyl-2-trifluoromethyl-7,8-dihydro-6 <i>H</i> -benzo[ <i>f</i> ]indolo[1,2- <i>a</i> ][1,4]diazepine ( <b>3h</b> ) . . . . .                                                                                                                                     | 72        |
| 4.1.14   | Synthesis of <i>N</i> -[(2-bromothiophen-3-yl)methyl]-butan-1-amine ( <b>4</b> ) . . . . .                                                                                                                                                                          | 73        |
| 4.1.15   | 7-Allyl-7,8-dihydro-6 <i>H</i> -pyrido[3,2- <i>f</i> ]indolo[1,2- <i>a</i> ][1,4]diazepine ( <b>5</b> ) . . . . .                                                                                                                                                   | 73        |
| 4.1.16   | Synthesis of <i>N</i> -((2-bromopyridin-3-yl)methyl)prop-2-en-1-amine ( <b>6</b> ) . . . . .                                                                                                                                                                        | 73        |

|                      |                                                                                                                        |    |
|----------------------|------------------------------------------------------------------------------------------------------------------------|----|
| 4.1.17               | 7-Allyl-7,8-dihydro-6 <i>H</i> -indolo[1,2- <i>a</i> ]thieno-<br>[2,3- <i>f</i> ][1,4]diazepine ( <b>7</b> ) . . . . . | 74 |
| References . . . . . |                                                                                                                        | 74 |

## Part II Synthesis of Isoquinoline Derivatives

|                      |                                                                                                                                                                                                                                                                                                                                     |           |
|----------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------|
| <b>5</b>             | <b>Facile Synthesis of 3-(Aminomethyl)isoquinolines by<br/>Copper-Catalyzed Domino Four-Component Coupling<br/>and Cyclization . . . . .</b>                                                                                                                                                                                        | <b>79</b> |
| 5.1                  | Introduction . . . . .                                                                                                                                                                                                                                                                                                              | 79        |
| 5.2                  | Experimental Section . . . . .                                                                                                                                                                                                                                                                                                      | 83        |
| 5.2.1                | General Methods . . . . .                                                                                                                                                                                                                                                                                                           | 83        |
| 5.2.2                | General Procedure for Four-Component Isoquinoline<br>Formation . . . . .                                                                                                                                                                                                                                                            | 85        |
| References . . . . . |                                                                                                                                                                                                                                                                                                                                     | 88        |
| <b>6</b>             | <b>Rapid Access to 3-(Aminomethyl)isoquinoline-Fused Polycyclic<br/>Compounds by Copper-Catalyzed Four-Component Coupling,<br/>Cascade Cyclization, and Oxidation . . . . .</b>                                                                                                                                                     | <b>89</b> |
| 6.1                  | Experimental Section . . . . .                                                                                                                                                                                                                                                                                                      | 94        |
| 6.1.1                | General Procedure for Synthesis of (Aminomethyl)-<br>isoquinoline-Fused Polycyclic Compounds by Domino<br>Mannich-Type Reaction and Cascade Cyclization:<br>Synthesis of 6-[( <i>N,N</i> -Diisopropylamino)methyl]-3,4-<br>dihydro-2 <i>H</i> -pyrimido[2,1- <i>a</i> ]isoquinoline ( <b>12a</b> )<br>(Table 1, Entry 10) . . . . . | 94        |
| 6.1.2                | 6-[( <i>N,N</i> -Diisopropylamino)methyl]-9-Fluoro-3,4-<br>Dihydro-2 <i>H</i> -Pyrimido[2,1- <i>a</i> ]isoquinoline ( <b>12b</b> ) . . . . .                                                                                                                                                                                        | 95        |
| 6.1.3                | 6-[( <i>N,N</i> -Diisopropylamino)methyl]-10-Fluoro-3,4-<br>Dihydro-2 <i>H</i> -Pyrimido[2,1- <i>a</i> ]isoquinoline ( <b>12c</b> ) . . . . .                                                                                                                                                                                       | 95        |
| 6.1.4                | 6-[( <i>N,N</i> -Diisopropylamino)methyl]-9-Methyl-3,4-<br>Dihydro-2 <i>H</i> -Pyrimido[2,1- <i>a</i> ]isoquinoline ( <b>12d</b> ) . . . . .                                                                                                                                                                                        | 96        |
| 6.1.5                | 6-[( <i>N,N</i> -Diisopropylamino)methyl]-10-Methoxy-3,4-<br>Dihydro-2 <i>H</i> -Pyrimido[2,1- <i>a</i> ]isoquinoline ( <b>12e</b> ) . . . . .                                                                                                                                                                                      | 96        |
| 6.1.6                | 6-(Piperidin-1-ylmethyl)-3,4-Dihydro-2 <i>H</i> -Pyrimido-<br>[2,1- <i>a</i> ]isoquinoline ( <b>12f</b> ) . . . . .                                                                                                                                                                                                                 | 96        |
| 6.1.7                | 6-[( <i>N,N</i> -Diallylamino)methyl]-3,4-Dihydro-2 <i>H</i> -<br>Pyrimido[2,1- <i>a</i> ]isoquinoline ( <b>12g</b> ) . . . . .                                                                                                                                                                                                     | 97        |
| 6.1.8                | 6-[[ <i>N,N</i> -Bis(( <i>R</i> )-1-phenylethyl)amino]methyl]-3,4-<br>Dihydro-2 <i>H</i> -Pyrimido[2,1- <i>a</i> ]isoquinoline ( <b>12h</b> ) . . . . .                                                                                                                                                                             | 97        |
| 6.1.9                | 5-[( <i>N,N</i> -Diisopropylamino)methyl]-2,3-Dihydroimidazo-<br>[2,1- <i>a</i> ]isoquinoline ( <b>13</b> ) . . . . .                                                                                                                                                                                                               | 98        |

|                      |                                                                                                                                            |     |
|----------------------|--------------------------------------------------------------------------------------------------------------------------------------------|-----|
| 6.1.10               | 7-[( <i>N,N</i> -Diisopropylamino)methyl]-2,3,4,5-Tetrahydro<br>[1,3]diazepino[2,1- <i>a</i> ]isoquinoline ( <b>14</b> ) . . . . .         | 98  |
| 6.1.11               | 8-[( <i>N,N</i> -Diisopropylamino)methyl]-3,4,5,6-<br>Tetrahydro-2 <i>H</i> -[1,3]diazocino[2,1- <i>a</i> ]isoquinoline ( <b>15</b> ). . . | 98  |
| 6.1.12               | 6-[( <i>N,N</i> -Diisopropylamino)methyl]benzimidazo-<br>[2,1- <i>a</i> ]isoquinoline ( <b>16</b> ) . . . . .                              | 99  |
| References . . . . . |                                                                                                                                            | 99  |
| <b>7</b>             | <b>Conclusions</b> . . . . .                                                                                                               | 101 |

Copper-Catalyzed Multi-Component Reactions  
Synthesis of Nitrogen-Containing Polycyclic Compounds

Ohta, Y.

2011, XVI, 104 p., Hardcover

ISBN: 978-3-642-15472-0