

# Contents

<b>1</b>	<b>Factors Limiting Motor Recovery After Facial Nerve Injury</b>	<b>1</b>
1.1	Altered Synaptic Input to the Axotomized Hyperexcitable Facial Motoneurons	1
1.2	Excessive Collateral Branching of Axons at the Lesion Site	2
1.3	Role of Cytoskeleton Reorganization During Axonal Regrowth	3
1.4	Exchange of Nerve Impulses Between Adjacent Axons	5
1.5	Vigorous Terminal Sprouting of Axons in the Denervated Muscles	6
1.6	Cellular Correlates of Muscle Reinnervation: the Role of Terminal Schwann Cells	6
1.7	Molecular Correlates of Muscle Reinnervation: Role of Sprouting-Inducing Stimuli	7
1.8	Questions Still Open	7
1.9	Methodological Approach	9
<b>2</b>	<b>Attempts to Improve Axonal Pathfinding and Quality of Target Reinnervation</b>	<b>11</b>
2.1	Efforts to Reduce Collateral Axonal Branching at the Lesion Site	12
2.1.1	Neutralization of Trophic Factors at the Lesion Site Reduced Collateral Axonal Branching, but Did Not Improve Recovery of Function	12
2.1.2	Local Stabilization of Microtubule Assembly Improved Recovery of Facial Nerve Function After Repair	25
2.2	Efforts to Reduce Axonal Sprouting in Denervated Muscles	33
2.2.1	Direct Modification of Microtubule Dynamics in Reinnervated Muscles Failed to Reduce Terminal Axonal Sprouting	33
2.2.2	Intraoperative Electrical Stimulation Prior to Reconstructive Surgery Did Not Improve Recovery of Function	35
2.2.3	Postoperative Electrical Stimulation of Paralyzed Vibrissal Muscles Did Not Improve Recovery of Function	40
2.2.4	Manual Stimulation of Paralyzed Vibrissal Muscles Following Facial Nerve Injury Promoted Full Recovery of Whisking	46
2.2.5	Manual Stimulation of Facial Muscles Improved Functional Recovery After Hypoglossal–Facial Anastomosis or Interpositional Nerve Grafting	53
2.2.6	Manual Stimulation of the Suprahyoid–Sublingual Region Diminished Polyinnervation of the Motor Endplates and Improved Recovery of Function After Hypoglossal Nerve Injury in Rats	63
2.2.7	Manual Stimulation of Forearm Muscles Did Not Improve Recovery of Motor Function After Injury to a Mixed Peripheral Nerve	78
2.2.8	Manually Stimulated Recovery of Motor Function After Facial Nerve Repair Requires Intact Sensory Input	84

<b>3</b>	<b>Discussion</b>	93
3.1	Significance of Axonal Branching at the Lesion Site	93
3.1.1	Reduced Collateral Branching Failed to Promote Recovery of Whisking Function	93
3.1.2	Effect of Perturbed Microtubule Assembly	96
3.2	Unsuccessful Ways to Reduce Intramuscular Axonal Sprouting in Denervated Muscles	100
3.2.1	Intraoperative Electrical Stimulation (IOES) Prior to Reconstructive Surgery	100
3.2.2	Postoperative Electrical Stimulation (POES) of Paralyzed Vibrissal Muscles	102
3.3	Successful Ways to Reduce Intramuscular Axonal Sprouting in Paralyzed Muscles	103
3.3.1	Manual Stimulation of Paralyzed Vibrissal Muscles After FFA	103
3.3.2	Manual Stimulation of Paralyzed Facial Muscles After HFA or IPNG	105
3.3.3	Manual Stimulation of Paralyzed Orbicularis Oculi Muscle After FFA	108
3.3.4	Manual Stimulation of Paralyzed Suprahyoid–Sublingual Muscles After HHA	109
3.4	Unsuccessful Manual Stimulation of Paralyzed Forearm Muscles After MMA	112
3.4.1	Clinical Relevance of Median Nerve Injury	112
3.4.2	The Effects of Manual Stimulation	113
3.4.3	Significance of the Intact Trigeminal Sensory Input	114
<b>4</b>	<b>Conclusions</b>	119
	<b>References</b>	121
	<b>Index</b>	141

Physical Rehabilitation of Paralysed Facial Muscles:  
Functional and Morphological Correlates

Angelov, D.N.

2011, XII, 144 p. 22 illus., 14 illus. in color., Softcover

ISBN: 978-3-642-18119-1