

Contents

7.1 Introduction	305
7.2 The Cochlea and the Cochlear Nerve	306
7.2.1 The Middle Ear and the Cochlea: Mechanical Transmission of Sound	306
7.2.2 Cochlear Hair Cells: Transduction and Amplification.....	308
7.2.3 Spiral Ganglion Cells and the Cochlear Nerve: Neural Transmission	308
7.2.4 The Auditory Periphery: Generation of Evoked Activity	308
7.2.5 Hearing Loss	309
Clinical Case 7.1 Cerebellopontine Angle Tumour	310
7.3 The Brain Stem Auditory System	312
7.3.1 The Cochlear Nuclei: Diversification of Cochlear Input	312
7.3.2 The Superior Olivary Complex: Recreation of Auditory Space	313
7.3.3 The Upper Brain Stem: Integration of Ascending Auditory Pathway.....	313
7.3.4 Brain Stem Topography: Generation of Evoked Potentials	314
Clinical Case 7.2 Impaired Sound Localization Following a Midline Pontine Lesion.....	314
7.4 The Forebrain Auditory System	315
7.4.1 The Auditory Thalamus	315
7.4.2 The Acoustic Radiation	315
7.4.3 The Auditory Cortex: Sequential Levels of Auditory Processing	316
7.4.4 Auditory Disorders Related to Stroke.....	319
Clinical Case 7.3 Auditory Agnosia Caused by Bilateral Lesions Restricted to the Auditory Radiations	321
Clinical Case 7.4 Neuropathology of Auditory Agnosia Following Bilateral Temporal Lobe Infarction	322
Clinical Case 7.5 Auditory Hallucinations Following a Metastasis in Heschl's Gyrus.....	324
7.5 The Descending Auditory System	325
References	326


H.J. ten Donkelaar (✉)
935 Department of Neurology, Radboud University Nijmegen Medical
Centre, P.O. Box 9101, 6500 HB Nijmegen, The Netherlands
e-mail: h.tendonkelaar@neuro.umcn.nl

7.1 Introduction

The ear or vestibulocochlear organ is composed of external, middle and inner parts (Fig. 7.1). The external ear consists of the auricle and the external acoustic meatus with the outer layer of the tympanic membrane. The middle ear is formed by the tympanic cavity, the auditory ossicles and the inner layer of the tympanic membrane. The inner ear comprises the labyrinth, a series of fluid-filled spaces in the petrous part of the temporal bone. The auditory part of the inner ear consists of the cochlea with the organ of Corti, which contains hair cells as auditory receptors. Receptors sensitive to high frequencies are located near the cochlear base and those sensitive to low frequencies near the apex of the cochlea. The hair cells are innervated by the peripheral processes of bipolar ganglion cells in the spiral ganglion. Their central processes form the cochlear division of the vestibulocochlear nerve and terminate in the cochlear nuclei. The principal auditory pathway passes from the cochlea, via the cochlear nuclei, the inferior colliculus and the medial geniculate body (MGB) to the contralateral auditory cortex on the dorsal surface of the superior temporal gyrus. Each MGB is bilaterally innervated, so that each hemisphere receives cochlear input bilaterally. All of the components of the auditory pathway are tonotopically organized.

At birth, humans have about 20,000 inner and outer hair cells in the organ of Corti, which often do not last a lifetime as they do not regenerate when lost (Stone et al. 1998). By the age of 65–75 years, many individuals have a bilateral, high-frequency progressive hearing loss known as presbycusis associated with hair cell attrition. Hair cell loss is the most common cochlear defect causing hearing impairment in presbycusis and noise-induced hearing loss. Hearing disorders due to brain stem lesions are rare because of the bilateral projections of the central auditory pathways. Midline pontine lesions may result in impaired sound localization due to interruption of the input of the superior olivary complex (see Sect. 7.3.2 and *Clinical case 7.2*). Disorders of

Fig. 7.1 Overview of the external, middle and internal ear (from Brödel 1946)



auditory perception may follow strokes in the territory of the internal carotid arteries or of the vertebrobasilar system. The central disorders of auditory perception may result from lesions of either the right and the left or both cerebral hemispheres, usually involving parietotemporal cortical areas as illustrated in *Clinical cases* (see Sect. 7.4.4).

7.2 The Cochlea and the Cochlear Nerve


7.2.1 The Middle Ear and the Cochlea: Mechanical Transmission of Sound

The **middle ear** comprises the tympanic cavity, the tympanic membrane, the three auditory ossicles, two middle ear muscles, air-filled cavities formed by the mastoid antrum and mastoid air cells, and the auditory tube. The **tympanic cavity** communicates with these air-filled cavities and through the auditory tube with the nasopharynx (Fig. 7.1). The three **auditory ossicles** are the hammer or **malleus**, the anvil or **incus** and the stirrup or **stapes**. The head of the malleus is anchored to the tympanic membrane, whereas the base of the stapes is connected to the fenestra vestibuli or oval window. Sound waves set the tympanic membrane into vibrating movements, which via the auditory ossicles are transmitted to the inner ear. The **inner ear** or cochlea is a fluid-filled tube that is coiled two and a half times. In cross-section, it has a broad base, a pointed apex and a central pillar called the modiolus. The osseous labyrinth

communicates with the tympanic cavity through two openings in its medial wall, the oval window or **fenestra vestibuli** and the round window or **fenestra cochleae**. The oval window is closed by the base of the stapes, so that vibrations of the auditory ossicles are transmitted to the perilymph of the inner ear.

Motion of the auditory ossicles is modified by two small **middle ear muscles**, the tensor tympani and the stapedius. The **tensor tympani** is the largest of the two. It is attached to the handle of the malleus and is innervated by the trigeminal nerve. The smaller **stapedius** attaches anteriorly to the head of the stapes and is innervated by the facial nerve. The stapedius and tensor tympani motoneurons form separate cell groups, situated close to the facial and motor trigeminal nuclei, respectively (Lyon 1978; Mizuno et al. 1982; Shaw and Baker 1983). The stapedius functions to protect the auditory receptors of the inner ear against excessive stimulation caused by too strong sound pressure. The sound pressure depends on the amplitude of the waves: the greater the amplitude, the higher the sound pressure. The stapedius contracts in response to sounds above 70 dB (the intensity of loud conversation), damping the movements of the auditory ossicle chain. The tensor tympani contracts to louder sounds, especially impulse noises. The **acoustic middle ear reflex** includes projections from the ventral cochlear nucleus via the superior olivary nuclear complex to the motor nuclei of the trigeminal and facial nerves (Borg 1973). With electro-acoustic impedance measurements, **stapedius muscle contraction** can be readily detected in response to ipsilateral or contralateral sound, giving objective information

Fig. 7.2 The foetal cochlear duct. At 16 weeks of development, the cochlear nerve (*cn*) fibres pass through a central pillar, the modiolus (*mod*), whereas their cells of origin form the spiral ganglia (*spg*). Below, details of the spiral organ are shown for 25 weeks of development. Abbreviations: *bm* basilar membrane; *cd* cochlear duct; *ihc*, *ohc* inner and outer hair cells; *st* scala tympani; *sva* stria vascularis; *sve* scala vestibuli; *tC* tunnel of Corti; *tm* tectorial membrane; *vm* vestibular (Reissner) membrane (from ten Donkelaar et al. 2006)



about the functional state of the middle and the inner ear, the auditory and facial nerves and the central auditory pathways in the lower brain stem. Ipsilateral and contralateral measurements can distinguish between right, left and midline lesions of the lower brain stem (Hayes and Jerger 1981).

The **cochlea** is composed of three chambers or scalae: the scala vestibuli, the scala media and the scala tympani, separated from each other by the vestibular membrane of Reissner and the basilar membrane (Fig. 7.2). The inner scala media is filled with endolymph, which is rich in potassium and has the character of intracellular fluid. The perilymph of the outer scalae vestibuli and tympani has approximately the same composition as the cerebrospinal fluid. The two perilymph compartments form one space, since they are continuous with each other at the apex of the cochlea (the helicotrema). The perilymph drains to the subarachnoid space. The **scala media** or **cochlear duct** contains the organ of Corti, which rests on the basilar membrane (Fig. 7.3). The superior wall of

the cochlear duct (the membrane of Reissner) angles downwards from lateral to medial, making the cochlear duct wedge shaped. The lateral wall is the stria vascularis. The thickened epithelium that constitutes the **organ of Corti** can be divided into hair cells and supporting cells. The hair cells are the sensory receptor cells of which there is a single row of inner hair cells and three rows of outer hair cells. The supporting cells include the inner and outer pillar cells, which are separated by the tunnel of Corti extending the length of the cochlea. Both the hair cells and the supporting cells are overlaid by the gelatinous tectorial membrane. In humans, there are 12,000 outer hair cells in three rows at the basal turn, increasing to four to five rows in the second and apical turns, and 3,500 inner hair cells in a single row (Retzius 1884; Bredberg 1968; Kimura 1975). On their apical side, the hair cells contain contractile proteins, including an actin cuticular plate, and about 100 stereocilia, graded in length, which extend to the overlying tectorial membrane. The stereocilia




Fig. 7.3 Photomicrograph of the human cochlea (from ten Donkelaar et al. 2006; courtesy Jo Curfs, Nijmegen)

are composed of the active contractile proteins actin and myosin (Flock 1980; Corwin and Warchol 1991). The afferent fibres from the hair cells pass from the organ of Corti through small openings in the osseous lamina into the modiolus. Their cell bodies are located in the modiolus in Rosenthal's canal as the spiral cochlear ganglion (Sect. 7.2.3). The inner ear is vascularized by the **internal auditory artery** which in some 80% is a branch of the anterior inferior cerebellar artery (Kim et al. 1990; Schuknecht 1993).

7.2.2 Cochlear Hair Cells: Transduction and Amplification

The human ear can detect sound waves with frequencies between 20 and 20,000 cycles per second or Hertz (Hz), i.e. approximately ten octaves of sound. The human ear has the greater sensitivity for sounds around 1,000 Hz. The greater the frequency, the higher the pitch. The sound vibrations that enter the scala vestibuli and the perilymph at the oval window produce displacement of the basilar membrane before they finally dissipate back to the middle ear by movements of the membrane covering the round window. **Transduction** of sound occurs in the sensory cells of the organ of Corti. Oscillations of the basilar membrane produce a shearing force on the stereocilia of the receptor cells, which are in firm contact with the non-oscillating tectorial membrane. The tilting of the stiff cilia is the adequate stimulus for the auditory receptor cells. The

inner and outer hair cells have different roles in the transduction of energy within the cochlea. Inner hair cells provide direct input to almost all of the axons in the cochlear nerve. Their activity is modified by local **amplification** of the motion of the basilar membrane produced by the outer hair cells and hair-cell related supporting cells (Flock et al. 1999; Moore and Linthicum 2004). Several molecules have been identified as having a vital role in hair-cell transduction. They are specifically expressed in and around the stereocilia and mutations in their genes lead to deafness (Steel and Kros 2001; ten Donkelaar et al. 2006).

7.2.3 Spiral Ganglion Cells and the Cochlear Nerve: Neural Transmission

The transition from hair cell activity to neural activity occurs within the cochlea. Activation of the stereocilia results in changes in the intracellular potential that lead to the release of a neurotransmitter from synaptic vesicle clusters at the base of the hair cells. Opposite such a cluster of synaptic vesicles, bulbous nerve terminals are found on the outer surface of the cell wall. Six to eight such terminals are present on the base of each inner hair cell, and a smaller number on each outer hair cell (Nadol 1990). These terminals continue as short unmyelinated processes, forming the “dendritic” segment of cochlear nerve fibres. They become myelinated when they enter the osseous spiral lamina. Here, they reach their cells of origin, the **spiral ganglion cells**. The **spiral ganglion** extends only half-way from the base of the cochlea to the apex. Therefore, the peripheral processes, containing hair cells in the apical and middle turns of the cochlea, extend down through the modiolus to reach the most apical ganglion cells. In humans, there are about 35,000 spiral ganglion cells (Hinojosa et al. 1985; Spoendlin 1985). Two types of ganglion cells are found (Spoendlin 1985). The majority (90–95%) are type I cells and contact inner hair cells. The unmyelinated peripheral processes of the remaining ganglion cells (5–10%), the type II cells, contact the outer hair cells. The central processes of both types of ganglion cells form the cochlear nerve (Spoendlin and Schrott 1989). The **cochlear nerve** enters the ventral cochlear nucleus on the ventrolateral side of the inferior cerebellar peduncle (see Fig. 7.6a). Upon entering the brain stem, primary auditory fibres bifurcate into equally sized ascending and descending branches (Moore and Osen 1979).

7.2.4 The Auditory Periphery: Generation of Evoked Activity

Neural activity is reflected in the **brain stem auditory-evoked potentials** or **responses** (BAEPS or BAERs), an externally recordable series of small amplitude and short latency wave-like potentials evoked by a transient stimulus




Fig. 7.4 Brain stem (ABR), middle latency (MLR) and cortical (ACR) auditory-evoked responses (from Pasman 1997; courtesy Jaco Pasman, Nijmegen)

such as a click (Jewett et al. 1970; Stockard et al. 1978, 1986). Auditory-evoked responses (AERs) can be subdivided according to their latency into brain stem (ABR), middle latency (MLR) and cortical (ACR) AERs (Fig. 7.4). In humans, the ABR or BAEP is characterized by six or sometimes seven deflections (I–VII) in the first 9 ms after the stimulus. Waves I, III and V are of greatest interest since they reflect volume-conducted activity from the levels of the acoustic nerve, pons and midbrain, respectively. The earliest BAEP waves (waves I and II) are generated by the cochlear nerve, prior to its entrance into the brain stem (Stockard et al. 1978, 1986; Moller and Jannetta 1982; Martin et al. 1995; Fig. 7.5). A potential corresponding to wave II of the scalp-recorded BAEP can be recorded intrasurgically from the surface of the human cochlear nerve as it passes through the internal auditory meatus and crosses the intradural space (Martin et al. 1995). This supports an earlier dipole localization study (Scherger and von Cramon 1985). Therefore, both waves I and II of the human BAEPs are generated by activity in axons of the cochlear nerve. BAEPs can distinguish between pathologies of the middle and inner ear, the auditory nerve and the brain stem. There are three major applications of BAEPs in adults: (1) the detection of tumours in the region of the posterior cranial fossa; (2) evolution of coma; and (3) assessment of patients with suspected demyelinating diseases such as MS. Acoustic neurinomas may cause complete loss of wave I on the side of the lesion or a significant increase in the I to III interpeak latency.

7.2.5 Hearing Loss

Two types of hearing loss can be distinguished: conductive and sensorineural. **Conductive hearing loss** is related to




Fig. 7.5 Relationship between components of the brain stem-auditory evoked response and the auditory projection pathway. Abbreviations: CI colliculus inferior; CN cochlear nuclei; II lateral lemniscus; MGB medial geniculate body; nVIII vestibulocochlear nerve; SO superior olive; I–V waves of BAEP (after Stockard et al. 1978)

defects in conductive mechanisms in the middle ear, resulting from conditions such as otitis media and otosclerosis. **Sensorineural hearing loss** is caused by disease in the cochlea or its central connection, the cochlear nerve. Hearing loss of cochlear origin is common and can result from a variety of conditions, including tumours, infections, temporal bone fractures or from exposure to excessive noise or ototoxic drugs (Schuknecht 1993). In **presbycusis**, the hearing loss of the aged, the loss begins with degeneration of outer hair cells at the basal end of the cochlea, but does not seriously affect hearing until the upper range of speech frequencies, around 3,000 Hz, is affected. Noise-induced hearing loss and severe blows to the head tend to affect the anterior basal turn of the cochlea, the region that processes 3,000–4,000 Hz (Moore and Linthicum 2004). **Tinnitus**, characterized by noise in the ears such as ringing, humming or whistling, is a common symptom in disorders of the inner ear, but it can also occur in disorders affecting the VIIIth nerve such as an **acoustic neurinoma** (see **Clinical case 7.1**) and with verteobasilar disease. Sudden onset of unilateral or bilateral deafness usually accompanied by dizziness or vertigo can be a sign of occlusion of the basilar artery (Huang et al. 1993; Levine and Häusler 2001).

Clinical Case 7.1 Cerebellopontine Angle Tumour

Electrocochleography and auditory brain stem response (ABR) are important electrophysiological tools for routine use in diagnosing vestibular schwannomas (Eggermont et al. 1980; Chandrasekhar et al. 1995). Kaga et al. (1997) reported a case of a vestibular schwannoma in which electrocochleography and ABR were correlated with temporal bone pathology (Fig. 7.6).

Case report: A 74-year-old female presented with a left hearing impairment. In 1975, she had undergone mastectomy of her left breast and in 1987, at the age of 73, she was treated with cobalt radiotherapy for a recurrence of the breast cancer. Pure tone audiometry revealed threshold elevation in the middle- and high-frequency range. ABR showed no response in the left ear but electrocochleography showed clear compound action potentials. CT scanning

and MRI demonstrated the presence of a medium-sized cerebellopontine angle tumour in the left ear (Fig. 7.7a, b). Three years later, she died of metastatic lung cancer and sepsis. At autopsy, metastases of the breast cancer were found in the right upper lobe of the lung and in the right temporal lobe of the brain. The temporal bone pathology consisted primarily of a large schwannoma, originating from the left inferior vestibular nerve and occupying the left internal auditory meatus (Fig. 7.7c, d). The organ of Corti was well preserved in each turn. In the modiolus, the numbers of spiral ganglion cells and cochlear nerve fibres in each turn were decreased. These histological findings suggest that clear compound action potentials were recorded from the distal part of the cochlear nerve in spite of the presence of the vestibular schwannoma. ABR could not be detected because of the blockade of the proximal portion of the cochlear nerve by the vestibular schwannoma.




Fig. 7.6 (a) The course of the vestibulocochlear nerve from the inner ear to the brain stem on the right and (b) the cerebellopontine angle. The black arrow in (b) points at an acoustic neurinoma and the way it extends (white arrows). Abbreviations: *nIII* oculomotor nerve; *nIV*

trochlear nerve; *nV1* ophthalmic nerve; *nV2* maxillary nerve; *nV3* mandibular nerve; *nVI* abducens nerve; *nIX* glossopharyngeal nerve; *nX* vagal nerve; *nXI* accessory nerve; *sps* superior petrosal sinus ((b) after ten Donkelaar et al. 2007)




Fig. 7.7 (a, b) CT and MRI demonstrating the presence of a medium-sized tumour in the left internal auditory canal and the cerebellopontine angle. (c) Mid-modiolar section of the left ear showing enlargement of the internal auditory canal occupied by a vestibular

schwannoma (HE stain). (d) Magnification of the vestibular schwannoma with mixed Antoni A and B cell types (HE stain; from Kaga et al. 1997)

Selected References

- Chandrasekhar SS, Brackmann DE, Kalpna K, Devgan KK (1995) Utility of auditory brainstem response audiometry in diagnosis of acoustic neuromas. *Am J Otol* 16:63–67
- Eggermont JJ, Don M, Brackmann DE (1980) Electrocochleography and auditory brainstem electric responses in patients with pontine angle tumours. *Ann Otol Rhinol Laryngol* 89:1–19

- Kaga K, Iwasaki S, Tamura A, Suzuki J-I, Haebara H (1997) Temporal bone pathology of acoustic neuroma correlating with presence of electrocochleography and absence of auditory brainstem response. *J Laryngol Otol* 111:967–972

7.3 The Brain Stem Auditory System

Upon entering the brain stem, the central processes of the spiral ganglion cells bifurcate and distribute to the cells of the dorsal and ventral cochlear nuclei (Sect. 7.3.1). The organization of the terminations was first described by Lorente de Nó (1933), based on his Golgi studies in a 4-day-old cat. In squirrel monkeys, fibres from the basal turn of the cochlea project to dorsal regions of the ventral cochlear nucleus, whereas apical fibres project to ventral regions (Moskowitz and Liu 1972). The primary cochlear nuclei contribute bilateral ascending projections to the superior olivary complex and to the lateral lemniscus (Sect. 7.3.2). The majority of the lateral lemniscal fibres ascend directly to the inferior colliculus (Sect. 7.3.3). Ascending projections from the inferior colliculus form the brachium of the inferior colliculus and reach the MGB (Sect. 7.4.1), which via the acoustic radiation (Sect. 7.4.2) projects to the auditory cortex (Sect. 7.4.3).


7.3.1 The Cochlear Nuclei: Diversification of Cochlear Input

The human cochlear nuclei consist of a large ventral nucleus and a smaller dorsal nucleus (Moore and Osen 1979; Terr and Edgerton 1985; Adams 1986). The **dorsal cochlear**

nucleus contains a large variety of cell types, and is situated dorsolateral to the inferior cerebellar peduncle. The **ventral cochlear nucleus** contains many different cell types and has anteroventral, ventral and posteroventral subnuclei, which borders are not well defined, however. The cochlear nuclei receive a rich **blood supply** from multiple sources, including branches of the anterior and posterior inferior cerebellar arteries (Oas and Baloh 1992).

The **secondary auditory projections** from the cochlear nuclei to the superior olivary complex and the inferior colliculus take various routes (Fig. 7.8). Ipsilaterally, a major projection from both ventral and dorsal cochlear nuclei reaches the superior olivary complex (Sect. 7.3.2). Contralaterally, there are three major ascending cochlear projections (Strominger 1973; Strominger et al. 1977): (1) the largest originates in the ventral part of the ventral cochlear nucleus and forms the **trapezoid body**; its axons may proceed directly to the contralateral lemniscus or terminate in the superior olivary complex; (2) fibres from the dorsal part of the ventral cochlear nucleus form the **intermediate acoustic stria**; they contribute to the lateral lemniscus; and (3) a contralateral projection from the dorsal cochlear nucleus, forming the **dorsal acoustic stria**. The dorsal and intermediate acoustic striae and the trapezoid body converge to form the **lateral lemniscus**. The auditory nuclei do not only serve as relay nuclei in the ascending auditory projection, but also as reflex centres. Efferents from the cochlear nuclei enter the reticular formation, where they contact neurons of the

Fig. 7.8 (a) Overview of the auditory projections in the human brain (after ten Donkelaar et al. 2007); (b–d) the position of the cochlear nuclei (in red), the lateral lemniscus (in light red) and the colliculus inferior (in red) in horizontal sections of the brain stem (after Duvernoy 1995). Abbreviations: *ar* acoustic radiation; *bci* brachium of colliculus inferior; *CI* colliculus inferior; *cn* cochlear nerve; *Cod*, *Cov* dorsal and ventral cochlear nuclei; *CS* colliculus superior; *ct* corpus trapezoideum; *gtt* gyrus temporalis transversus (Heschl's gyrus); *ll* lateral lemniscus; *MGB* medial geniculate body; *pt* planum temporale; *sad* stria acoustica dorsalis; *SO* superior olive






Fig. 7.8 (continued)

ascending reticular activating system (see Chap. 5), and give rise to the **auditory-evoked startle reflex**.

7.3.2 The Superior Olivary Complex: Recreation of Auditory Space

The superior olivary complex is the first site for **binaural convergence**. In primates, the cochlear nuclei project to the superior olivary complex on both sides of the brain stem (Strominger 1973; Strominger et al. 1977). The **superior olivary complex** is located in the caudal pons, lateral to the medial lemniscus and dorsal to the spinothalamic tract. The complex contains the medial superior olivary nucleus, the lateral superior olivary nucleus and the nucleus of the trapezoid body. The latter nucleus is indistinct in apes and vestigial in humans (Moore 2000). The superior olivary complex is important for the **localization of sounds** (Moore and Linthicum 2004). A sound is localized by two means depending on its frequency: (1) low-frequency sounds activate the two ears at somewhat different times (interaural time differences); (2) high-frequency sounds activate the two ears with somewhat different intensities (interaural intensity differences). Neurons in the medial superior olivary nucleus are tuned to low-frequency stimuli and are sensitive to interaural time differences. The projection from the ventral cochlear nucleus is thought to contribute to this sensitivity. In contrast, neurons in the lateral superior olivary nucleus are tuned to high-frequency stimuli and are sensitive to interaural intensity differences. The lateral superior olivary nucleus receives a monosynaptic excitatory connection from the ipsilateral

ventral cochlear nucleus and a disynaptic inhibitory connection from the contralateral ventral cochlear nucleus via the nucleus of the trapezoid body. Since the dorsal cochlear nucleus does not innervate the superior olivary complex, it is believed not to play a role in the localization of sounds.

Behavioural studies in cats have implicated the superior olivary complex in the **recreation of auditory space**. Cats with lesions above the level of the superior olivary complex, in the lateral lemniscus, the inferior colliculus, the MGB or the auditory cortex, are unable to locate a sound source in the spatial field contralateral to the lesion, whereas cats with lesions below the superior olivary complex have more diffuse deficits (Casseday and Neff 1975; Thompson and Masterton 1978; Jenkins and Masterton 1982). A comparable deficit has been observed in human subjects with extensive midline pontine lesions that eliminated crossed input to the superior olivary complex on both sides (Griffiths et al. 1997a; Furst et al. 2000; see *Clinical case 7.2*). These animal and human studies suggest that the auditory spatial field is recreated in the brain stem by transformations occurring at the level of the superior olivary complex.

7.3.3 The Upper Brain Stem: Integration of Ascending Auditory Pathway

The **lateral lemniscus** is clearly visible in the rostral pons and the midbrain. Most of its fibres terminate in the inferior colliculus. Many of these fibres send a collateral branch to the nuclei of the lateral lemniscus, which innervate the inferior colliculus and also directly the MGB. In most mammalian

species, the lateral lemniscus contains sizable ventral, intermediate and dorsal lemniscal nuclei (Moore 1987). In humans, only the dorsal lemniscal nucleus is well developed (Geniec and Morest 1971; Moore 1987). It gives rise to Probst's commissure to the contralateral inferior colliculus.

The **inferior colliculus** is composed of three nuclei: central, external and pericentral. The **central nucleus** is the principal nucleus of the inferior colliculus and receives input from: (1) the direct pathway from the dorsal and ventral cochlear nuclei; (2) projections arising from the ipsilateral and contralateral superior olivary complex and (3) fibres from the dorsal nucleus of the lateral lemniscus. These projections all pass via the lateral lemniscus. The central nucleus is laminated (Geniec and Morest 1971) and processes the physical characteristics of sounds for auditory perception. In this nucleus, neurons in a single layer are maximally sensitive to similar tonal frequencies. The function of the other two nuclei of the inferior colliculus is not entirely clear. Lesion studies in cats suggest that the **external** and **pericentral nuclei** play a role in acousticomotor function such as the orientation of the head and body to auditory stimuli. The inferior colliculus projects to the MGB via the brachium of

the inferior colliculus, which is macroscopically visible on the lateral surface of the midbrain. The inferior colliculi are interconnected via the commissure of the inferior colliculi.

7.3.4 Brain Stem Topography: Generation of Evoked Potentials

Waves I and II of the ABR are generated by the cochlear nerve. The subsequent waves III–VI are generated within the brain stem (see Fig. 7.5). Intracranial recordings made from the surface of the human brain stem and dipole studies suggest that wave III is generated by a volley of action potentials in axons emerging from the cochlear nuclei in the ventral acoustic stria (Stockard et al. 1978, 1986; Møller and Jannetta 1982; Scherg and von Cramon 1985). Waves IV and V are generated further rostrally in the brain stem: wave IV most likely at the level of the superior olivary complex contralateral to the stimulated ear, presumably by the bend in the axonal pathway occurring at that point, and wave V by synaptic activity in the inferior colliculus (Møller and Jannetta 1982; Moore et al. 1996).

Clinical Case 7.2 Impaired Sound Localization Following a Midline Pontine Lesion

In a 45-year-old female patient with an extensive **midline pontine lesion**, eliminating crossed input to the superior olivary complex on both sides, Griffiths et al. (1997a, b) observed that the patient had no difficulty in detecting frequency and amplitude modulation and no general deficit in detection of auditory temporal information, but she was *unable* to determine by sound alone the location and direction of motion of objects in the environment, such as ringing telephones and passing trains. Furst and co-workers analyzed sound localization in patients with multiple sclerosis and brain stem infarcts (Furst et al. 2000; 1995; Aharonson et al. 1998). Levine and Häusler (2001) reported another case (see **Case report**).

Case report: An 80-year-old male presented with sudden onset of vertigo and vomiting. On examination, he was found to have a left gaze palsy, dysphagia, dysarthria, and a right hemiplegia that included only the lower face. He had no auditory complaints, and his bedside hearing evaluation was unremarkable. MRI showed a left trapezoid body infarct, the location of which is indicated in Fig. 7.9a. A year later, he was evaluated with a battery of hearing tests. Despite an age-appropriate audiogram and normal BAERs, all fusion tests were abnormal for the three stimuli used (clicks, low-pass noise and high-pass noise) and for interaural time or level disparities (Fig. 7.9b). Just noticeable differences were highly abnormal, and regardless of the size or type of interaural disparity, the patient indicated that everything sounded as though it were coming from or near the centre of his head (Fig. 7.9c). Unlike normal subjects, nothing was heard coming from the far right or left.




Fig. 7.9 Impaired sound localization in a patient with a lower pontine lesion (arrow in **a**) involving the trapezoid body; (**b**) brain stem auditory evoked responses; (**c**) sound lateralization (after Levine and

Häusler 2001; see text for explanation). *Abbreviations:* Co cochlear nuclei; IC inferior colliculus; II lateral lemniscus; TB trapezoid body; vas ventral acoustic stria

Selected References

- Aharonson V, Furst M, Levine RA, Chaigrecht M, Korczyn AD (1998) Lateralization and binaural discrimination of patients with pontine lesions. *J Acoust Soc Am* 103:2624–2633
- Furst M, Levine RA, Korczyn AD, Fullerton BC, Tadmor R, Algom D (1995) Brainstem lesions and click lateralization in patients with multiple sclerosis. *Hear Res* 82:109–124
- Furst M, Aharonson V, Levine RA, Fullerton BC, Tadmor R, Pratt H, et al. (2000) Sound lateralization and interaural discrimination. Effects of brainstem infarcts and multiple sclerosis lesions. *Hear Res* 143:29–42

- Griffiths TD, Bates D, Rees A, Witton C, Golkar A, Green GGR (1997) Sound movement detection deficit due to a brainstem lesion. *J Neurol Neurosurg Psychiatry* 62:522–526
- Levine RA, Häusler R (2001) Auditory disorders in stroke. In: Bogousslavsky J, Caplan LR (eds) *Stroke syndromes*, 2nd ed. Cambridge University Press, Cambridge, pp 144–161

This case is based on a case report by Levine and Häusler (2001).

7.4 The Forebrain Auditory System

For decades, the dominant species for research on the auditory forebrain has been the cat, but the focus has now clearly shifted to non-human primates. Although the subcortical auditory systems of monkeys and cats are largely similar, there are important differences in cortical organization (Kaas and Hackett 2000).

7.4.1 The Auditory Thalamus

The **medial geniculate body (MGB)** or **nucleus** is clearly visible on the inferior surface of the inferior thalamus. The MGB contains several divisions, the principal auditory relay nucleus is the **ventral or principal medial geniculate nucleus** (Winer 1984). The ventral division of the MGB is laminated. It receives the major ascending auditory projection from the also laminated central nucleus of the inferior colliculus. For both

nuclei, lamination is a structural correlate of precise tonotopic organization. In contrast, the dorsal and medial divisions of the MGB are not laminated and receive much less dense input from the inferior colliculus. The ventral medial geniculate nucleus projects via the auditory radiation to the tonotopically organized primary auditory cortex. The dorsal and medial sub-nuclei project to higher-order auditory cortical areas in the planum temporale, areas that do not have such a precise tonotopic organization as the primary auditory cortex.

7.4.2 The Acoustic Radiation

In 1882, Constantin von Monakow first described the origin of the acoustic radiation from the MGB in rabbit experiments. The classic studies in the human brain located the proximal part of the acoustic radiation just caudal to the thalamus, where it originates from the MGB, then passes through the sublenticular, posterior part of the internal capsule to curve around the inferior sulcus of the insula before reaching Heschl's gyrus




Fig. 7.10 The acoustic and optic radiations in coronal (a) and sagittal (b) probabilistic maps (after Rademacher et al. 2002). *Abbreviations:* AR acoustic radiation; OR optic radiation

(Dejerine 1895; Flechsig 1920; Pfeifer 1920). In a more recent study, Rademacher et al. (2002) showed the stereotaxic localization, intersubject variability and interhemispheric differences of the human acoustic radiation (Fig. 7.10). They showed that the location of the acoustic radiation varies considerably between individuals and hemispheres.

7.4.3 The Auditory Cortex: Sequential Levels of Auditory Processing

The **primary auditory cortex (A1)** is located on the transverse temporal or Heschl's gyrus in the temporal lobe of the cerebral cortex and corresponds to area 41. It is surrounded by secondary auditory areas (A2): caudally the caudomedial area, also known as the planum temporale, and rostrally, the rostral area. Geschwind and Levitsky (1968) demonstrated that the **planum temporale** is larger on the left side in the majority of the postmortem brains they examined. **Asymmetry** of the planum temporale may form the substrate for left hemispheric dominance for language-related auditory processes (Geschwind and Galaburda 1985; Dorsaint-Pierre et al. 2006) and is correlated with handedness (Steinmetz et al. 1989, 1991).

Heschl's gyrus is located largely within the lateral sulcus (von Economo and Horn 1930; Fig. 7.11). The transverse temporal gyrus is often partially duplicated into a double, or occasionally triple convexity (Pfeifer 1920; Steinmetz et al. 1989; Penhune et al. 1996; Leonard et al. 1998; Morosan et al. 2001). The cytoarchitecture of the human auditory cortex has been described by Brodmann (1908, 1909), von Economo and Koskinas (1925), Galaburda and Sanides (1980) and, more recently by Hackett et al. (2001) and Hackett




Fig. 7.11 The human auditory cortex. The primary auditory cortex is composed of two fields, TD and TC. On the right side (R), these occupy a double transverse temporal gyrus (Heschl's gyrus); on the left side (L), they correspond to a single Heschl's gyrus and a part of the more caudally situated planum temporale (PT). TD and TC are composed of markedly granular subareas (dotted in red) and less granular areas. Note the distinct right-left asymmetries with a larger planum temporale on the left side. *Abbreviations:* A anterior; HG Heschl's gyrus; P posterior; TA superior temporal area; TB magnocellular supratemporal area; TC transverse supratemporal area; TD intercalate supratemporal area; TG temporopolar area (after Brodal 1981)

and Kaas (2004) and Morosan et al. (2001) and Rademacher et al. (2001a, b). The primary auditory cortex was designated area 41 by Brodmann, TC by von Economo and Koskinas and




Fig. 7.12 Regional parcellation of the right human superior temporal cortex (rostral is below) according to (a) Brodmann, (b) von Economo and Koskinas, (c) Galaburda and Sanides and (d) Morosan and co-workers (after Hackett and Kaas 2004 and Morosan et al. 2001). In (a–c), core areas are shown in red, belt areas in medium red and parabelt areas in light red. Abbreviations: *KAlt*, *KAm* lateral and medial auditory

koniocortex; *PaAc/d*, *paAe*, *PaAl/paAl*, *PaAr* caudal/dorsal, external, lateral and rostral auditory parakoniocortex; *ProA* proauditory cortex; *TA1*, *TA2*, *TB*, *TBma*, *TBmp*, *TC*, *TD* subdivisions by von Economo and Koskinas; *Te1.0*, *Te1.1*, *Te1.2*, *Te3* subdivisions by Morosan and co-workers; *TG1*, *TG2* temporopolar subdivisions of von Economo and Koskinas; *Tpt* temporoparietal area; 22–52 Brodmann areas

KAm and *KAlt* (medial and lateral auditory **koniocortex**) by Galaburda and Sanides (Fig. 7.12a–c). *KAm* is the most medial and the most granular area, whereas the more lateral *KAlt* is less granular. Morosan et al. (2001) suggested three areas with well-developed layers IV, *Te1.1*, *Te1.0* and *Te1.2*, to represent the primary auditory cortex (Fig. 7.12d). There is considerable variability in size of the auditory koniocortex and its extent does not coincide with gyral or sulcal anatomy (Galaburda and Sanides 1980; Rademacher et al. 1993, 2001a, b; Hackett et al. 2001; Morosan et al. 2001). The human auditory koniocortex (area 41/*TC*/*KA*/*Te*) is homologous to the core area of the monkey auditory cortex. Based on parvalbumin staining, Wallace et al. (2002) suggested that Heschl's gyrus contains two core fields, partially surrounded by at least six belt fields that lie mostly on the superior temporal gyrus. In an fMRI study, Wessinger et al. (2001) showed that pure tones primarily activate the core and that more complex sounds activate belt areas.

The **primate auditory core area** is located in the centre of the superior temporal plane (Hackett et al. 2001; Fig. 7.13). In primates, a centrally located core region containing two or three subdivisions including the primary auditory area (*A1*), a surrounding belt of cortex with some seven divisions, and a lateral parabelt region comprised of at least two fields, have been described. In monkeys, the **core region** can be identified




Fig. 7.13 Auditory and auditory-related cortices in macaque monkeys. Core areas (*A1*, *R*, *RT*) are shown in red, belt areas (*CL*, *CM*, *AL*, *RM*, *RTL*, *RTM*) in medium red and parabelt areas (*CPB*, *RPB*) in light red. Major sulci have been opened to show the extent of auditory-related cortex. Abbreviations: *A1* primary auditory area; *AL* anterolateral area; *as* arcuate sulcus; *CL*, *CM* caudolateral and caudomedial areas; *CPB* caudal parabelt area; *cs* central sulcus; *h*, *l* high and low frequencies; *IN* insula; *ips* intraparietal sulcus; *lus* lunate sulcus; *PL* posterolateral area; *ps* principal sulcus; *R* rostral area; *RM* rostromedial area; *RPB* rostral parabelt area; *RT* rostrot temporal primary auditory cortex; *RTL*, *RTM* lateral and medial rostromedial areas; *STG* superior temporal gyrus; *sts* superior temporal sulcus; *vbls* ventral bank of lateral sulcus (after Hackett et al. 2001)

on the basis of *specific* anatomical and physiological features. The region shows dense immunostaining for parvalbumin in layer IV, surrounded by a more lightly stained belt, which is flanked by a very sparsely stained parabelt (Jones et al. 1995; Kosaki et al. 1997). Parvalbumin staining also marks the human core auditory cortex in humans (Nakahara et al. 2000; Wallace et al. 2002; Chirry et al. 2003). In macaque, chimpanzee and human brains, Hackett et al. (2001) identified the auditory core from serial sets of adjacent sections processed for cytoarchitecture, myeloarchitecture, acetylcholinesterase and cytochrome oxidase. The position of the core region with respect to major sulci and gyri in the superior temporal region varied most in chimpanzee and human brains.

In monkeys, most neurons of the ventral division of the MGB project to the core cortex (Mesulam and Pandya 1973; Burton and Jones 1976; Luethke et al. 1989; Rauschecker et al. 1997). These **thalamocortical projections** terminate in layers IV and lower III in regular patches of higher density label, separated by areas of less dense labelling (Pandya and Rosene 1993; Hashikawa et al. 1995). In contrast, the medial and dorsal divisions of the MGB project to the core area diffusely. It seems likely that the human primary auditory cortex also receives dense thalamic input. This input explains the cochleotopic organization shown in this area by functional imaging, including magnetoencephalography (MEG) (Elberling et al. 1982; Hari et al. 1989; Pantev et al. 1995; Lutkenhoner and Steinstrater 1998), PET (Lauter et al. 1985; Ottaviani et al. 1997; Lockwood et al. 1999), fMRI (Wessinger et al. 1997; Scheich et al. 1998; Di Salle et al. 2001) and microelectrode mapping studies in epilepsy patients (Howard et al. 1996).

The human auditory koniocortex is surrounded rostrally, laterally and caudally by an area of **parakoniocortex** (Fig. 7.12). This region covers the lateral part of the transverse temporal gyrus, and extends rostrally and caudally over the superior temporal plane. The auditory parakoniocortex has been called area 42 by Brodmann and TB by von Economo and Koskinas. Galaburda and Sanides (1980) distinguished three regions: (1) a rostral auditory parakoniocortex (PaAr) on the rostral aspect of the superior temporal plane; (2) a lateral, internal auditory parakoniocortex (PaAl), lateral to A1; and (3) a caudal auditory parakoniocortex (PaAc), covering the caudal portion of the superior temporal plane and extending around the insula to the parietal operculum. In its turn, the parakoniocortex is surrounded by an extensive area of auditory cortex that covers the remaining of the superior temporal plane and the lateral surface of the superior temporal gyrus, except for its rostral pole. This region was described as area 22 by Brodmann (1909), as TA by von Economo and Koskinas (1925) and as external auditory parakoniocortex (PaAe) by Galaburda and Sanides (1980).

In primates (Fig. 7.13), anatomical and physiological studies defined a **belt area** surrounding the core rostrally, laterally and caudally (Pandya and Sanides 1973; Galaburda

and Pandya 1983; Morel and Kaas 1992; Morel et al. 1993; Hackett et al. 1998a). The area rostral and lateral to the belt is nowadays known as the **parabelt** (Morel et al. 1993; Hackett et al. 1998a). Both belt and parabelt areas differ from the core area in their pattern of thalamic input. The macaque belt area receives projections from only the medial and dorsal divisions of the MGB (Rauschecker et al. 1997), whereas the parabelt area is also innervated by these two divisions of the MGB but, moreover, by the medial division of the pulvinar (Trojanowski and Jacobson 1975; Burton and Jones 1976; Hackett et al. 1998b).

Ablation of the core of macaque auditory cortex eliminates responses to auditory stimuli in the adjacent belt region (Rauschecker et al. 1997), suggesting that input from the medial and dorsal geniculate nuclei is not sufficient to support auditory processing in the absence of direct projections from the ventral geniculate nucleus. Instead, information processing to the secondary auditory cortical areas appears to depend on transcortical projections that pass successively from core to belt to parabelt cortex (Jones and Powell 1970; Seltzer and Pandya 1978; FitzPatrick and Imig 1980; Luethke et al. 1989; Morel and Kaas 1992; Morel et al. 1993; Hackett et al. 1998a). Tardif and Clarke (2001) studied the intrinsic connectivity of human auditory areas with anterograde and retrograde labelling of the carbocyanine dye DiI. With DTI, the tracts connecting the Heschl's gyri via the corpus callosum have been studied (Hofer and Frahm 2006; Westerhausen et al. 2009). These interhemispheric connections are located more rostrally within the posterior callosal third than those connecting the posterior parts of both superior temporal gyri.

The idea of a two-stream, **what/where** organization of sensory cortex originated in the visual system (see Chap. 8). In rhesus monkeys, such a dichotomy has also been demonstrated for the cortical auditory projections (Rauschecker and Tian 2000). The **'where' (dorsal) pathway** is thought to link A1 via the caudomedial belt with the frontal eye field and parietal targets (Romanski et al. 1999; Fig. 7.14) that are implicated in *spatial processing*. The **'what' (ventral) pathway** is thought to represent a *pattern information* stream that originates in the anterior core and belt areas and influences targets within the temporal lobe. A similar **two-stream organization** may exist in the human auditory cortex (Griffiths et al. 2000; Alain et al. 2001; Maeder et al. 2001; Wessinger et al. 2001; Clarke et al. 2002). The right insula is activated by a moving sound image (Griffiths et al. 1994) and, conversely, a patient with a right hemispheric stroke causing atrophy of the right insula was unable to detect sound source movement by either phase or loudness cues (Griffiths et al. 1996). Subjects listening to stimulus movement stimulated by changes in binaural timing show maximal activity in the inferior parietal area, particularly on the right side (Griffiths et al. 1998; Weeks et al. 1999). These findings suggest: (1) that there is a transcortical passage of information from




Fig. 7.14 Topography of auditory-related projections. Caudal (CPB) and rostral (RPB) subdivisions of the parabelt and the superior temporal gyrus (STG) project topographically to segregated regions of superior temporal, posterior parietal and prefrontal cortices. Abbreviations: *as* arcuate sulcus; *cs* central sulcus; *ips* intraparietal sulcus; *ls* lateral sulcus; *lus* lunate sulcus; *ps* principal sulcus; *sts* superior temporal sulcus; *Tpt* temporoparietal area (after: Hackett and Kaas 2004)

auditory koniocortex mediocaudalwards across the insula into the parietal lobe, during processing of information on sound source position and motion (see Hackett and Kaas 2004); and (2) a dominant functional role of the right hemisphere in sound localization.

In **functional imaging studies**, simple auditory tasks such as passive listening to white noise bursts, tones or consonant-vowel speech syllables, activate restricted areas within the lateral fissure on the superior temporal plane (Zatorre et al. 1992, 1994; Binder et al. 1994, 1997, 2000; Zatorre and Binder 2000). The extent of the activation varies from subject to subject and may spread rostralwards and caudalwards on the superior temporal plane. The area of activation is generally within and around the transverse temporal gyrus. With exposure of subjects to more complex stimuli such as passive listening to tone patterns, single words, pseudowords or narrative text, activity is not only present in the cortex of the superior temporal gyrus, but now foci of activation appear on the lateral aspect of the superior temporal gyrus in area 22/TA/PaAc (Binder et al. 1994). The human primary auditory cortex is functionally organized in a **tonotopic manner**. In a combined fMRI and DTI study, Upadhyay et al. (2007) showed that the connectivity pattern in the human primary auditory cortex is similar to that described in tonotopic mapping studies on macaque monkeys (Morel et al. 1993) and cats (Lee et al. 2004; Lee and Winer 2005).

In general, activity is **bilaterally equal**. With complex stimuli, language in particular, the question arises whether there is a right–left asymmetry in the response. Since handedness influences hemispheric lateralization, imaging studies of speech processing are normally restricted to neurologically normal right-handers. In them, there is a tendency for

greater activation of the left hemisphere during tasks that depend on word meaning. Left lateralization of speech characterizes both males and females (Frost et al. 1999). The functional significance of greater left hemispheric activity is implied by imaging studies of stroke patients after infarctions of the left perisylvian area (Weiller et al. 1995; Heiss et al. 1997; Mummery et al. 1999). Subjects who showed good recovery of speech perception had increasing activation of the left temporal cortex surrounding the infarct. Some indications for an **opposite asymmetry** in processing **musical stimuli** come from cases of pathology:

1. A patient with a **right** thalamic tumour experienced **distorted perception of music** but not of voices (Roeser and Daly 1974).
2. Cortical activity has been demonstrated in the right superior temporal lobe during **musical hallucinations** (Kasai et al. 1999).
3. A case of **amusia**, a form of auditory agnosia, was seen after an infarct involving the right insula (Griffiths et al. 1997b).

7.4.4 Auditory Disorders Related to Stroke

Disorders of auditory perception may follow strokes in the territory of the internal carotid arteries or of the vertebrobasilar system (Levine and Häusler 2001; Lechevalier et al. 2007; Kaga 2009), and appear as:

- **Auditory agnosia**, the impossibility of recognizing environmental sounds, words and music, which the patient, however, is said to hear
- **Pure word deafness**, the impossibility to understand spoken language to repeat or to write under dictation in the absence of other signs of aphasia
- **Cortical deafness**, the feeling of being deaf contrasting with the integrity of the tonal audiogram
- **Amusia**, auditory agnosia specific for music

The central disorders of auditory perception may result from lesions of either the right, the left or both cerebral hemispheres, usually involving parietotemporal cortical areas. **Cortical deafness** is characterized by bilateral abolition of the middle and late latencies of auditory potentials, caused by bilateral lesions of the primary auditory cortices. Such patients have the feeling of being deaf to all types of auditory stimuli, but often say they are *not* deaf, rather that they do not understand what is said to them. The term **subcortical deafness** is used to indicate an auditory disorder clinically identical to cortical deafness, but due to lesions in subcortical areas of the brain. It was first described by Le Gros Clark and Russell (1938). The ischaemic lesions involved the two external capsules and extended sufficiently downwards to interrupt the acoustic radiations, while sparing the auditory cortices. Recent cases were reported by Woods (1996), Levine and Häusler (2001) and Kaga et al. (2005; see **Clinical case 7.3**).

Since the pioneering studies of Ferrier (1875) and Henschen (1920), there has been a longstanding debate as to whether bilateral destruction of either the primary auditory cortex or the acoustic radiation results in **auditory agnosia**. In macaque monkeys, bilateral lesions of the primary auditory cortex apparently do not cause permanent deafness (Heffner and Heffner 1990). Less recovery of function in the human brain, compatible with the clinical diagnosis of auditory agnosia, may or may not have been caused by the inclusion of the surrounding auditory association areas (Lechevalier et al. 2007).

Tanaka et al. (1991) differentiated three clinical syndromes of auditory agnosia: (1) **disconnection syndromes**, destroying the acoustic radiation and causing auditory agnosia (*prephonemic deficit*); (2) **cortical lesions** of the left superior temporal lobe may result in pure word deafness (*linguistic deficit*) and (3) unilateral or bilateral **temporoparietal** or **subcortical lesions** have been documented in patients with non-verbal auditory agnosia (*deficit to environmental sounds*). Lesions occurring peripherally to the MGB (*prethalamal*) may cause hearing loss and those bilaterally located centrally to the MGB (*postthalamal*) may result in auditory agnosia. Small lesions of the MGB may be related to auditory hallucinations (Fukutake and Hattori 1998). **Pure word deafness** may be the result of left or bilateral temporal lesions, possibly due to disconnection as suggested by Liepmann and Storch (1902). Recent cases were reported by Kaga et al. (2000; see **Clinical case 7.4**) and Levine and Häusler (2001).

Disorders of music perception following cerebral damage can be divided into two categories (Lechevalier et al. 2007):

1. **Multimode perceptive disorders** affecting more or selectively musical sounds, but with verbal and environmental sound difficulties
2. A **pure amusia**, where only music perception is affected (for **congenital amusia** see Ayotte et al. 2000)

In both monkeys and humans, neurons in **core areas** respond strongly to narrow-band sounds such as tones, whereas neurons in **belt areas** respond better to more complex sounds such as noise (Wessinger et al. 2001; Rauschecker and Tian 2004; Tian and Rauschecker 2004; Bendor and Wang 2006). Within the core areas, two mirror symmetric **tonotopic maps** sharing a low-frequency border have been identified, corresponding to A1 and the rostral field R (Formisano et al. 2003; Bendor and Wang 2006). In monkeys, a third core area (RT) has been found that lies rostral to R (Kaas and Hackett 2000; Hackett and Kaas 2004). Kaas and Hackett postulated that each core area is connected to medial and lateral neighbouring belt areas (see Fig. 7.14), with additional belt areas located on the rostral and caudal ends of the core. Three of these lateral belt areas (caudal-lateral, middle-lateral and antero-lateral) have been mapped

electrophysiologically and possess similar mirror tonotopic maps to those of their adjacent core (Rauschecker and Tian 2004; Tian and Rauschecker 2004). In an fMRI study, Patterson et al. (2002) identified a specific region in the lateral part of Heschl's gyrus that was preferentially activated by temporally regular sounds with a **pitch**. They determined that only lateral Heschl's gyrus, a non-primary auditory region rostralateral to the primary auditory cortex, responded to the temporal regularity of pitch of the acoustic stimuli. Other imaging studies (Penagos et al. 2004; Schneider et al. 2005) have confirmed these findings.

Musical perception is not a uniform competence in the general population. Some patients will have had musical training, others not. Peretz (2001) estimated that 5–10% of individuals are completely unable to distinguish the pitches of two notes of music or to memorize the smallest musical tone. Geschwind and Galaburda (1985) suggested that rightward deviation from the usual pattern of cerebral asymmetry may be associated with increased giftedness for talents for which the right hemisphere is assumed to be important. With MR morphometry, Schlaug et al. (1995) presented evidence for structural brain asymmetry in musicians. Musicians with **perfect pitch** revealed stronger leftward asymmetry of the planum temporale than non-musicians or musicians without perfect pitch. This suggests that outstanding musical ability is associated with increased leftward asymmetry of the cortex subserving music-related functions.

Neuropsychological studies in epileptic patients who underwent a unilateral temporal cortectomy have contributed to our knowledge of the localization of musical functions (Liégeois-Chauvel et al. 1998). A right temporal cortectomy was found to disturb melodic perception as well as the perception of pitch intervals, whereas a left-sided lobectomy did not disturb perception of the intervals. These data underline the key role of the superior temporal gyrus in discrimination of melodies. Cortectomy of the posterior part of T1, including the planum temporale, the lateral part of Heschl's gyrus and Brodmann area 22, is more striking for the processing of pitch and variations of rhythm than cortectomy of the rostral part of T1. Disorders of the perception of rhythm and metre (recognition of a cadence of march or waltz) can be dissociated. The right and left rostral parts of T1 would be implicated in the processing of metre. Griffiths et al. (1997b) reported a patient with lesions of the middle and posterior temporal areas and the insula of the right hemisphere. The patient complained of not being able to appreciate music. Neuropsychological testing showed a deficit of musical perception without disturbance of the perception of noises, environmental sound and speech sounds. His ability to detect continuous changes of sound frequency was preserved. However, a disturbance in the analysis of rapid sequences of notes seemed to be the basis of his musical perception deficit. Neuroimaging studies have revealed that rhythm perception

activates area 44 and that detection of pitch changes relies on the left cuneus and precuneus (Platel et al. 1997, 2003).

Auditory hallucinations are observed in brain stem (Ross et al. 1975; Cambier et al. 1987; Fisher and Tapia 1987; Griffiths 2000) and temporal lobe (Lechevalier et al. 1985) strokes. Cambier et al. (1987) reported five purely auditory observations of *hallucinosi* (hallucinations, regarded by the

patient as abnormal), four of which were attributed to paramedian strokes of the pons and one to an infarct of the dorsolateral mesencephalon. Auditory hallucinations following temporal lobe lesions are unusual and have specific characteristics (Liepmann and Storch 1902; Lechevalier et al. 1985; Augustin et al. 2001; Evers and Ellger 2004; Sacks 2007; see *Clinical case 7.5*).

Clinical Case 7.3 Auditory Agnosia Caused by Bilateral Lesions Restricted to the Auditory Radiations

Bilateral lesions of the auditory radiations are rare (Tanaka et al. 1991; Woods 1996; Kaga et al. 2005; see **Case report**).

Case report: Kaga et al. (2005) reported a patient with auditory agnosia due to bilateral lesions of the auditory radiations. A 43-year-old male patient experienced mild left temporal hemiplegia due to a right putaminal haemorrhage. He recovered completely but hypertension persisted. When he was 53 years old, he had a left putaminal haemorrhage and

went into a coma. After recovering from the coma and the right hemiplegia, he could hear but could not discriminate speech sounds. Brain CT and MRI demonstrated small bilateral lesions restricted to the auditory radiations (Fig. 7.15a, b). Pure-tone audiograms recorded 1 and 4 years after the second haemorrhage are shown in Fig. 7.15c, d. MEG demonstrated the disappearance of middle latency responses and AEP studies showed a very small Pa peak. In contrast, a positron emission tomographic study showed a marked bilateral increase in blood flow in the auditory cortex in response to both click and monosyllable stimuli. This may be due to activation of the auditory cortex via non-specific pathways.




Fig. 7.15 Auditory agnosia caused by bilateral lesions restricted to the auditory radiations. In the axial (a) and coronal (b) MRIs, the auditory radiations are bilaterally damaged (arrows) by small brain

infarcts. Pure-tone audiograms recorded one (c) and four (d) years after the second haemorrhage (from Kaga et al. 2005)




Fig. 7.15 (continued)

Selected References

Kaga K, Kurauchi T, Nakamura M, Shindo M, Ishii K (2005) Magnetoencephalography and positron emission tomography studies of a patient with auditory agnosia caused by bilateral lesions confined to the auditory radiations. *Acta Otolaryngol* 125:1351–1355

Tanaka Y, Kano T, Yoshida M, Yamadori A (1991) “So-called” cortical deafness. Clinical, neurophysiological and radiological observations. *Brain* 114:2385–2401

Woods RP (1996) Correlation of brain structure and function. In: Toga AW, Mazziotta JC (eds) *Brain mapping: the systems*. Academic, San Diego, CA, pp 365–402

Clinical Case 7.4 Neuropathology of Auditory Agnosia Following Bilateral Temporal Lobe Infarction

Severe auditory deficits due to bilateral lesions of the primary auditory cortex or the auditory radiations is very rare. The resulting hearing problem is referred to as **auditory agnosia** or **cortical deafness**. Kaga et al. (2000) reported a patient who came to autopsy (see **Case report**).

Case report: Kaga’s case of auditory agnosia due to bilateral lesions of the auditory cortex was first diagnosed in 1975 when the patient was 37 years old. He was admitted to hospital for examination following his second cerebrovascular accident. MRI of the lesions on admission is shown in Fig. 7.16a, b. A comprehensive follow-up examination of auditory function was periodically conducted until his sudden death 15 years later. His brain was studied neuropathologically. Initial pure-tone audiometry revealed moderate

sensorineural hearing loss in the right ear and mild sensorineural hearing loss in the left ear. Repeated pure-tone audiometry revealed that bilaterally thresholds became progressively poorer over time. Speech audiometry of both ears consistently revealed that the patient was unable to discriminate any monosyllabic words. In general, speech and hearing tests demonstrated that he could not comprehend spoken words but could comprehend written commands and gestures. Neuropathological examination of the brain revealed a total defect and neuronal loss of the superior temporal gyrus, including Heschl’s gyrus, and total gliosis of the MGB (MGB; Fig. 7.16c, d). In the right hemisphere, subcortical necrosis, gliosis in the centre of the superior temporal gyrus and partial gliosis of the MGB were found (Fig. 7.16c, e). These data support the clinical observations of imperception of speech sounds, music and environmental sounds, which may be due to progressive degeneration of both MGBs.




Fig. 7.16 Auditory agnosia following bilateral temporal lobe infarction. (a, b) Axial (here the left side is on the left) and coronal (here the left side is on the right) MRIs showing a large infarct in the left hemisphere and a small infarct in the right hemisphere including the auditory cortex. (c) Lateral views of the brain and two horizontal sections in which the auditory cortex is present. In the right hemisphere, a small

infarct is present in the upper part of the lateral sulcus, whereas in the left hemisphere extensive infarction can be seen in Broca's area, the superior temporal gyrus and the supramarginal gyrus. (d, e) HE-stained sections of the medial geniculate body (MGB). In the left MGB, neurons have been completely replaced by glial cells (d), whereas in the right MGB (e) there is partial neuronal preservation (from Kaga et al. 2000)




Fig. 7.16 (continued)

Selected Reference

Kaga K, Shindo M, Tanaka Y, Haebara H (2000) Neuropathology of auditory agnosia following bilateral temporal lobe lesions: a case study. *Acta Otolaryngol* 120:259–262

Clinical Case 7.5 Auditory Hallucinations Following a Metastasis in Heschl's Gyrus

Case report: A 64-year-old patient presented with word-finding difficulties. He suffered from coronary sclerosis with exercise-induced angina pectoris but he had no previous neurological complaints. On neurological examination, there were no focal signs but his speech was non-fluent with word-finding difficulties and suboptimal comprehension. On hospital admission, he repeatedly complained of **auditory hallucinations**, consisting of incomprehensible

words and sounds. On MRI, a contrast-enhancing lesion was found in the left gyrus of Heschl (Fig. 7.17) that appeared to be part of a more lobular contrast-enhancing in the left parietotemporal region with surrounding oedema. The auditory hallucinations disappeared on treatment with dexamethasone. A biopsy showed that the tumour was a gemistocytary astrocytoma for which he was treated with radiotherapy and temozolamide.

This case was kindly provided by Peter van Domburg (Department of Neurology, Orbis Medical Centre, Sittard, The Netherlands).





Fig. 7.17 T1-contrast MRIs of a metastasis in the left gyrus of Heschl that caused auditory hallucinations (courtesy Peter van Domburg, Sittard)

Fig. 7.18 Efferent control of the cochlea. Abbreviations: *Cod*, *Cov* dorsal and ventral cochlear nuclei; *ct* corpus trapezoidum; *ihc* inner hair cells; *ll* lateral lemniscus; *ocb* olivocochlear bundle; *ohc* outer hair cells; *PON* periolivary nuclei; *SO* superior olivary complex (after Nieuwenhuys 1984)



7.5 The Descending Auditory System

Parallel with the pathways from the organ of Corti to the auditory cortex, there is an uninterrupted chain of neurons conducting impulses in the opposite, descending direction. The final link in this **descending auditory system** is formed by the **olivocochlear bundle of Rasmussen**, which originates in the **peri-olivary nuclei** around the superior olivary

nucleus (Fig. 7.18). Most of the fibres of the olivocerebellar bundle decussate in the tegmentum. They enter the vestibular nerve and join the cochlear nerve via the vestibulocochlear anastomosis (Schuknecht 1993) to terminate in the inner and outer hair cells of the organ of Corti. The human olivocochlear system has been identified with acetylcholinesterase histochemistry (Schuknecht et al. 1959) and choline acetyltransferase immunohistochemistry (Moore et al. 1999; Moore and Linthicum 2004).

References

- Adams JC (1986) Neuronal morphology of the human cochlear nucleus. *Arch Otol Head Neck Surg* 112:1253–1261
- Alain C, Arnott SR, Hevenor S, Graham S, Grand CL (2001) 'What' and 'where' in the human auditory system. *Proc Natl Acad Sci USA* 98:12301–12306
- Augustin J, Guegan-Massardier E, Levillain D et al (2001) Musical hallucinosis following infarction of the right middle cerebral artery. *Rev Neurol (Paris)* 157:289–292
- Ayotte J, Peretz I, Hyde K (2000) Congenital amusia: a group study of adults afflicted with a music-specific disorder. *Brain* 125:238–251
- Bendor D, Wang X (2006) Cortical representations of pitch in monkeys and humans. *Curr Opin Neurobiol* 16:391–399
- Binder JR, Rao SM, Hammeke TA, Yetkin FZ, Jasmanowicz A, Bandettini PA et al (1994) Functional magnetic resonance imaging of human auditory cortex. *Ann Neurol* 35:662–672
- Binder JR, Frost JA, Hammeke TA, Cox RW, Rao SM, Prieto T (1997) Human brain language areas identified by functional magnetic resonance imaging. *J Neurosci* 17:353–362
- Binder JR, Frost JA, Hammeke TA, Bellrowan PSF, Springer JA, Kaufman JN, Possing ET (2000) Human temporal lobe activation by speech and nonspeech sounds. *Cereb Cortex* 10:512–528
- Borg E (1973) On the neuronal organization of the acoustic middle ear reflex. A physiological and anatomical study. *Brain Res* 49:101–123
- Bredberg G (1968) Cellular pattern and nerve supply of the human organ of Corti. *Acta Otolaryngol, Suppl* 236:1–135
- Brodal A (1981) *Neurological Anatomy, in relation to clinical medicine*, 3rd edn. Oxford University Press, New York
- Brödel M (1946) Three unpublished drawings of the anatomy of the human ear. Saunders, Philadelphia, PA
- Brodman K (1908) Beiträge zur histologischen Lokalisation der Grosshirnrinde. VI. Mitteilung: Die Cortexgliederung des Menschen. *J Psychol Neurol (Lpz)* 10:231–246
- Brodman K (1909) Vergleichende Lokalisationslehre der Grosshirnrinde in ihren Prinzipien dargestellt auf Grund des Zellenbaues. Barth, Leipzig, English translation by LJ Garey (1999) Brodman's Localisation in the Cerebral Cortex. Imperial College Press, London
- Burton H, Jones EG (1976) The posterior thalamic region and its cortical projection in New World and Old World monkeys. *J Comp Neurol* 168:249–301
- Cambier J, Decroix JP, Masson C (1987) Auditory hallucinations in lesions of the brain stem. *Rev Neurol (Paris)* 150:255–262
- Casseday JH, Neff WD (1975) Auditory localization: role of auditory pathways in the brain stem of the cat. *J Neurophysiol* 38:852–858
- Chirý O, Tardif E, Magistretti PJ, Clarke S (2003) Patterns of calcium-binding proteins support parallel and hierarchical organization of human auditory areas. *Eur J Neurosci* 17:397–410
- Clarke S, Bellmann-Thiran A, Maeder P, Adriani M, Vernet O, Regli L et al (2002) What and where inhuman audition: Selective deficits following focal hemispheric lesions. *Exp Brain Res* 147:8–15
- Corwin JT, Warchol ME (1991) Auditory hair cells: structure, function, development, and regeneration. *Annu Rev Neurosci* 14:301–333
- Dejerine J (1895) *Anatomie des centres nerveux*, Tome 1. Reuff, Paris
- Di Salle F, Formisano E, Seifritz E, Linden DE, Scheffler SC et al (2001) Functional fields in human auditory cortex revealed by time-resolved fMRI without interference of EPI noise. *Neuroimage* 13:328–338
- Dorsaint-Pierre R, Penhune VB, Watkins KE, Neelin P, Lerch JP, Bouffard M et al (2006) Asymmetries of the planum temporale and Heschl's gyrus: relationship to language lateralization. *Brain* 129:1164–1176
- Duvernoy HM (1995) *The human brain stem and cerebellum: surface, structure, vascularization and three-dimensional sectional anatomy with MRI*. Springer, New York
- Elberling C, Bak C, Kofoed B, Lebech J, Saermark K (1982) Auditory magnetic fields. Source localization and tonotopic organization in the right hemisphere of human brain. *Scand Audiol* 10:203–207
- Evers S, Ellger T (2004) The clinical spectrum of musical hallucinations. *J Neurol Sci* 227:55–65
- Ferrier D (1875) Experiments on the brain of monkeys. *Proc R Soc Lond B23*:409–432
- Fisher CM, Tapia J (1987) Lateral medullary infarction extending to the lower pons. *J Neurol Neurosurg Psychiatry* 50:620–624
- FitzPatrick KS, Imig IJ (1980) Auditory cortico-cortical connections in the owl monkey. *J Comp Neurol* 192:589–610
- Flechsig P (1920) *Anatomie des menschlichen Gehirns und Rückenmark des Menschen aufgrund entwicklungsgeschichtlicher Untersuchungen*. Engelmann, Leipzig
- Flock Å (1980) Contractile proteins in hair cells. *Hear Res* 2:411–412
- Flock Å, Flock B, Fridberger A, Scarfone E, Ulfendahl M (1999) Supporting cells contribute to control of hearing sensitivity. *J Neurosci* 19:4498–4507
- Formisano E, Kim DS, Di Salle F, van de Moortele PF, Uğurbil K, Goebel R (2003) Mirror-symmetric tonotopic maps in human primary auditory cortex. *Neuron* 40:859–869
- Frost JA, Binder JR, Springer JA, Hammeke TA, Bellrowan PS, Rao SM, Cox RW (1999) Language processing is strongly left lateralized in both sexes. Evidence from functional MRI. *Brain* 122:190–208
- Fukutake T, Hattori T (1998) Auditory illusions caused by a small lesion in the right medial geniculate body. *Neurology* 51:1469–1471
- Furst M, Aharonson V, Levine RA, Fullerton BC, Tadmor R, Pratt H et al (2000) Sound lateralization and interareal discrimination. Effects of brainstem infarcts and multiple sclerosis lesions. *Hear Res* 143:29–42
- Galaburda A, Pandya DN (1983) The intrinsic architectonic and connective organization of the superior temporal region of the rhesus monkey. *J Comp Neurol* 221:169–184
- Galaburda A, Sanides F (1980) Cytoarchitectonic organization of the human auditory cortex. *J Comp Neurol* 190:597–610
- Geniec P, Morest DK (1971) The neuronal architecture of the human posterior colliculus. *Acta Otolaryngol, Suppl* 295:1–33
- Geschwind N, Galaburda AM (1985) Cerebral localization: Biological mechanisms, associations, and pathology I. A hypothesis and a program for research. *Arch Neurol* 42:428–459
- Geschwind N, Levitsky W (1968) Human brain: left-right asymmetries in the temporal speech region. *Science* 161:186–187
- Griffiths TD (2000) Musical hallucinosis in acquired deafness: phenomenology and substrate. *Brain* 123:2065–2076
- Griffiths TD, Bench CJ, Frackowiak RSJ (1994) Human cortical areas selectively activated by apparent sound movement. *Curr Biol* 4:892–895
- Griffiths TD, Rees A, Witton C, Shakir RA, Henning GB, Green GGR (1996) Evidence for a sound movement area in the human cerebral cortex. *Nature* 342:425–427
- Griffiths TD, Bates D, Rees A, Witton C, Gholkar A, Green GGR (1997a) Sound movement detection deficit due to a brainstem lesion. *J Neurol Neurosurg Psychiatry* 62:522–526
- Griffiths TD, Rees A, Witton C, Cross PM, Shakir RA, Green GGR (1997b) Spatial and temporal auditory processing deficits following right hemisphere infarction. A psychophysical study. *Brain* 120:785–794
- Griffiths TD, Rees G, Rees A, Green GGR, Witton C, Rowe D et al (1998) Right parietal cortex is involved in the perception of sound movement in humans. *Nat Neurosci* 1:74–79
- Griffiths TD, Green GGR, Rees A, Rees G (2000) Human brain areas involved in the analysis of auditory movement. *Hum Brain Mapp* 9:72–80
- Hackett TA, Kaas JH (2004) Auditory cortex in primates: Functional subdivisions and processing streams. In: Galaburda A (ed) *The cognitive neuroscience III*. MIT, Cambridge, MA, pp 215–232

- Hackett TA, Stepniewska I, Kaas JH (1998a) Subdivisions of auditory cortex and ipsilateral cortical connections of the parabelt auditory cortex in macaque monkeys. *J Comp Neurol* 394:475–495
- Hackett TA, Stepniewska I, Kaas JH (1998b) Thalamocortical connections of the parabelt auditory cortex in macaque monkeys. *J Comp Neurol* 400:271–286
- Hackett TA, Preuss TM, Kaas JH (2001) Architectonic identification of the core region in auditory cortex of macaques, chimpanzees, and humans. *J Comp Neurol* 441:197–222
- Hari R, Hämäläinen M, Kaukoranta E, Mäkelä J, Joutsiniemi S-L, Tiikonen J (1989) Selective listening modifies activity of the human auditory cortex. *Exp Brain Res* 74:463–470
- Hashikawa T, Molinari M, Rausell E, Jones EG (1995) Patchy and laminar terminations of medial geniculate axons in monkey auditory cortex. *J Comp Neurol* 362:195–208
- Hayes D, Jerger J (1981) Patterns of acoustic reflex and auditory brainstem response abnormality. *Acta Otolaryngol* 92:199–209
- Heffner HE, Heffner RS (1990) Effect of bilateral auditory cortex lesions on absolute thresholds in Japanese macaques. *J Neurophysiol* 64:191–205
- Heiss W-D, Karbe H, Weber-Luxemburger G, Herholz K, Kessler J, Pietrzyk U, Pawlik G (1997) Speech-induced cerebral metabolic activation reflects recovery from aphasia. *J Neurol Sci* 145:213–217
- Henschen SE (1920) Über die Hörsphäre. *J Psychol Neurol (Lpz)* 22:319–473
- Hinojosa R, Seligsohn R, Lerner SA (1985) Ganglion cell counts in the cochleae of patients with normal audiograms. *Acta Otolaryngol* 99:8–13
- Hofer S, Frahm J (2006) Topography of the human corpus callosum revisited – comprehensive fiber tractography using diffusion tensor magnetic resonance imaging. *Neuroimage* 32:989–994
- Howard MA, Vokov IO, Abbas PJ, Damasio H, Ollendiek MC, Granner MA (1996) A chronic microelectrode investigation of the tonotopic organization of human auditory cortex. *Brain Res* 724:260–264
- Huang MH, Huang CC, Ryu SJ, Chu NS (1993) Sudden bilateral hearing impairment in vertebrobasilar occlusive disease. *Stroke* 24:132–137
- Jenkins WM, Masterton RB (1982) Sound localization: effects of unilateral lesions in the central auditory system. *J Neurophysiol* 47:987–1016
- Jewett RL, Romano MN, Williston JS (1970) Human auditory evoked potentials: Possible brain stem components detected on the scalp. *Science* 167:1517–1518
- Jones EG, Powell TPS (1970) An anatomical study of converging sensory pathways within the cerebral cortex of the monkey. *Brain* 93:793–820
- Jones EG, Dell’Anna ME, Molinari M, Rausell E, Hashikawa T (1995) Subdivisions of macaque monkey auditory cortex revealed by calcium-binding protein immunoreactivity. *J Comp Neurol* 362:153–170
- Kaas JH, Hackett TA (2000) Subdivisions of auditory cortex and processing streams in primates. *Proc Natl Acad Sci USA* 97:11793–11799
- Kaga K (2009) Central auditory pathway disorders. Springer, Tokyo
- Kaga K, Iwasaki S, Tamura A, Suzuki J-I, Haebara H (1997) Temporal bone pathology of acoustic neuroma correlating with presence of electrocochleography and absence of auditory brainstem response. *J Laryngol Otol* 111:967–972
- Kaga K, Shindo M, Tanaka Y, Haebara H (2000) Neuropathology of auditory agnosia following bilateral temporal lobe lesions: A case study. *Acta Otolaryngol* 120:259–262
- Kaga K, Kurauchi T, Nakamura M, Shindo M, Ishii K (2005) Magnetoencephalography and positron emission tomography studies of a patient with auditory agnosia caused by bilateral lesions confined to the auditory radiations. *Acta Otolaryngol* 125:1351–1355
- Kasai H, Asada T, Yumoto M, Jakeya J, Matsuda H (1999) Evidence for functional abnormality in the right auditory cortex during musical hallucinations. *Lancet* 354:1703–1705
- Kim H-N, Kim YH, Park IY, Kim GR, Chung IH (1990) Variability of the surgical anatomy of the neurovascular complex of the cerebellopontine angle. *Ann Otol Rhinol Laryngol* 99:288–295
- Kimura RS (1975) The ultrastructure of the organ of Corti. *Int Rev Cytol* 42:173–222
- Kosaki H, Hashikawa T, He J, Jones EG (1997) Tonotopic organization of auditory cortical fields delineated by parvalbumin immunoreactivity in macaque monkeys. *J Comp Neurol* 386:304–316
- Lauter JL, Herscovitch P, Formby C, Raichle ME (1985) Tonotopic organization in human auditory cortex revealed by positron emission tomography. *Hear Res* 20:199–205
- Lechevalier B, Eustache F, Rossa Y (1985) Les troubles de la perception de la musique d’origine neurologique. Masson, Paris
- Lechevalier B, Lambert J, Moreau S, Platel H, Viader F (2007) Auditory disorders related to strokes. In: Godefroy O, Bogousslavsky J (eds) *The behavioral and cognitive neurology of stroke*. Cambridge University Press, Cambridge, pp 348–368
- Lee CC, Winer JA (2005) Principles governing auditory cortex connections. *Cereb Cortex* 15:1804–1814
- Lee CC, Schreiner CE, Imaizumi K, Winer JA (2004) Tonotopic and heterotopic projection systems in physiologically defined auditory cortex. *Neuroscience* 128:871–887
- Le Gros Clark WE, Russell WR (1938) Cortical deafness without aphasia. *Brain* 61:375–383
- Leonard CM, Puranik C, Kuldau JM, Lombardino LJ (1998) Normal variation in the frequency and location of human auditory cortex landmarks. Heschl’s gyrus: Where is it? *Cereb Cortex* 8:397–406
- Levine RA, Häusler R (2001) Auditory disorders in stroke. In: Bogousslavsky J, Caplan LR (eds) *Stroke Syndromes*, 2nd edn. Cambridge University Press, Cambridge, pp 144–161
- Liégeois-Chauvel C, Peretz I, Babai M, Laguiton V, Chauvel P (1998) Contribution of different cortical areas in the temporal lobes to music processing. *Brain* 121:1853–1867
- Liepmann H, Storch E (1902) Der mikroskopische Gehirnbefund bei dem Fall Gorstelle. *Monatsschr Psychiatr Neurol* 11:115–120
- Lockwood AH, Salvi RJ, Coad ML, Arnold SA, Wack DS, Murphy BW, Burkard RF (1999) The functional anatomy of the normal human auditory system: Responses to 0.5 and 4.0 kHz tones at varied intensities. *Cereb Cortex* 9:65–76
- Lorente de Nó R (1933) Anatomy of the eighth nerve. The central projections of the nerve endings of the internal ear. *Laryngoscope* 43:1–38
- Luethke LE, Krubitzer LA, Kaas JH (1989) Connections of primary auditory cortex in the New World monkey, *Saguinus*. *J Comp Neurol* 285:487–513
- Lutkenhoner B, Steinstrater O (1998) High-precision neuromagnetic study of the functional organization of the human auditory cortex. *Audiol Neuro-Otol* 3:191–213
- Lyon MJ (1978) The central localization of the motor neurons to the stapedius muscle in the cat. *Brain Res* 143:437–444
- Maeder PP, Meuli RA, Adriani M, Bellmann A, Fornari E, Thiran JP et al (2001) Distinct pathways involved in sound recognition and localization: a human fMRI study. *Neuroimage* 14:802–816
- Martin WH, Pratt H, Schweigler JW (1995) The origin of the human auditory brain stem response wave II. *Electroencephalogr Clin Electrophysiol* 96:357–370
- Mesulam M-M, Pandya DN (1973) The projections of the medial geniculate complex within the Sylvian fissure of the rhesus monkey. *Brain Res* 60:315–333
- Mizuno N, Nomura S, Konishi A, Uemura-Sumi M, Takahashi O, Yasui Y et al (1982) Localization of motoneurons innervating the tensor tympani muscle: a horseradish peroxidase study in the guinea pig and cat. *Neurosci Lett* 31:205–208
- Moller AR, Jannetta PJ (1982) Auditory evoked potentials recorded intracranially from the brainstem in man. *Exp Neurol* 78:144–157
- Moore JK (1987) The human auditory brain stem. A comparative view. *Hear Res* 29:1–32

- Moore JK (2000) Organization of the human superior olivary complex. *Microsc Res Tech* 51:403–412
- Moore JK, Linthicum FH Jr (2004) Auditory system. In: Paxinos G, Mai JK (eds) *The human nervous system*, 2nd edn. Elsevier, Amsterdam, pp 1241–1279
- Moore JK, Osen KK (1979) The cochlear nuclei in man. *Am J Anat* 154:393–418
- Moore JK, Ponton CW, Eggermont JJ, Wu BJ-C, Huang JQ (1996) Perinatal maturation of the auditory brainstem response: Changes in path length and conduction velocity. *Ear Hear* 17:411–418
- Moore JK, Simmons DD, Guan Y-L (1999) The human olivocochlear system: organization and development. *Audiol Neuro-Otol* 4: 311–325
- Morel A, Kaas JH (1992) Subdivision and connections of auditory cortex in owl monkeys. *J Comp Neurol* 318:27–63
- Morel A, Garraghty PE, Kaas JH (1993) Tonotopic organization, architectonic fields, and connections of auditory cortex in macaque monkeys. *J Comp Neurol* 335:437–459
- Morosan P, Rademacher J, Schleicher A, Amunts K, Schormann T, Zilles K (2001) Human primary auditory cortex: cytoarchitectonic subdivisions and mapping into a spatial reference system. *Neuroimage* 13:684–701
- Moskowitz N, Liu JC (1972) Central projections of the spiral ganglion of the squirrel monkey. *J Comp Neurol* 144:335–344
- Mummery CJ, Ashburner J, Scott SK, Wise RJS (1999) Functional neuroimaging of speech perception in six normal and two aphasic subjects. *J Acoust Soc Am* 106:449–457
- Nadol JB Jr (1990) Synaptic morphology of inner and outer hair cells of the human organ of Corti. *J Electron Microscop Tech* 15:187–196
- Nakahara H, Yamada S, Mizutani T, Murayama S (2000) Identification of the primary auditory field in archival human brain tissue via immunocytochemistry of parvalbumin. *Neurosci Lett* 286:29–32
- Nieuwenhuys R (1984) Anatomy of the auditory pathways, with emphasis on the brain stem. *Adv Oto-Rhino-Laryngol* 34:25–38
- Oas JG, Baloh RW (1992) Vertigo and the anterior inferior cerebellar artery syndrome. *Neurology* 42:2274–2279
- Ottaviani F, Di Girolamo S, Briglia G, De Rossi G, Di Giuda D, Di Nardo W (1997) Tonotopic organization of human auditory cortex analyzed by SPET. *Audiology* 36:241–248
- Pandya DN, Rosene DL (1993) Laminar termination patterns of thalamic, callosal, and association afferents in the primary auditory area of the rhesus monkey. *Exp Neurol* 119:220–234
- Pandya DN, Sanides F (1973) Architectonic parcellation of the temporal operculum in rhesus monkey and its projection pattern. *Z Anat Entwickl-Gesch* 139:127–161
- Pantev C, Bertrand O, Eulitz C, Verkindt C, Hampson S, Schuierer G, Elbert T (1995) Specific tonotopic organization of different areas of the human auditory cortex revealed by simultaneous magnetic and electric recordings. *Electroencephalogr Clin Neurophysiol* 94:26–40
- Pasman JW (1997) Auditory Evoked Responses in Preterm Infants. Thesis, University of Nijmegen
- Patterson RD, Uppenkamp S, Johnsrude IS, Griffiths TD (2002) The processing of temporal pitch and melody information in auditory cortex. *Neuron* 36:767–776
- Penagos H, Melcher JR, Oxenham AJ (2004) A neural representation of pitch salience in non-primary auditory cortex revealed with functional magnetic resonance imaging. *J Neurosci* 24:6810–6815
- Penhune VB, Zatorre RJ, MacDonald JD, Evans AC (1996) Interhemispheric anatomical differences in human primary auditory cortex: probabilistic mapping and volume measurement from magnetic resonance scans. *Cereb Cortex* 6:661–672
- Peretz I (2001) Brain specialization for music: new evidence from congenital amusia. *Ann N Y Acad Sci* 930:153–165
- Pfeifer RA (1920) Myelogenetisch-anatomische Untersuchungen über das kortikale Ende der Hörleitung. *Abh math phys Kl Sächs Akad Wiss* 37:1–54
- Platel H, Price C, Baron JC, Wise R, Lambert J, Frackowiak RSJ et al (1997) The structural components of music perception: a functional anatomical study. *Brain* 120:229–243
- Platel H, Baron JC, Desgranges B, Bernard F, Eustache F (2003) Semantic and episodic memory of music are subserved by distinct neural networks. *Neuroimage* 20:244–256
- Rademacher J, Caviness VS Jr, Steinmetz H, Galaburda AM (1993) Topographical variation of the human primary cortices: Implications for neuroimaging, brain mapping, and neurobiology. *Cereb Cortex* 3:313–329
- Rademacher J, Morosan P, Schormann T, Schleicher A, Werner C, Freund H-J, Zilles K (2001a) Probabilistic mapping and volume measurement of human primary auditory cortex. *Neuroimage* 13: 669–683
- Rademacher J, Morosan P, Schleicher A, Freund H-J, Zilles K (2001b) Human primary auditory cortex in women and men. *Neuroreport* 12:1561–1566
- Rademacher J, Bürgel U, Zilles K (2002) Stereotaxic localization, intersubject variability, and interhemispheric differences of the human auditory thalamocortical system. *Neuroimage* 17:142–160
- Rauschecker JP, Tian B (2000) Mechanisms and streams for processing of “what” and “where” in auditory cortex. *Proc Natl Acad Sci USA* 97:11800–11806
- Rauschecker JP, Tian B (2004) Processing of band-passed noise in the lateral auditory belt cortex of the rhesus monkey. *J Neurophysiol* 91:2578–2589
- Rauschecker JP, Tian B, Pons T, Mishkin M (1997) Serial and parallel processing in rhesus monkey auditory cortex. *J Comp Neurol* 382:89–103
- Retzius G (1884) *Das Gehörorgan der Wirbelthiere*, vol II. Samson and Wallin, Stockholm
- Roeser RJ, Daly DD (1974) Auditory cortex disconnection associated with thalamic tumor. A case report. *Neurology* 24:555–559
- Romanski LM, Bates JF, Goldman-Rakic PS (1999) Auditory belt and parabelt projections to the prefrontal cortex in the rhesus monkey. *J Comp Neurol* 403:141–157
- Ross ED, Jossman PB, Bell B, Sabin T, Geschwind N (1975) Musical hallucinations in deafness. *J Am Med Assoc* 231:620–622
- Sacks O (2007) *Musophilia. Tales of music and the brain*. Knopf, New York
- Scheich H, Baumgart F, Gaschler-Markefski B, Tegeler C, Tempelmann C, Heinze HJ et al (1998) Functional magnetic resonance imaging of a human auditory cortex area involved in foreground-background decomposition. *Eur J Neurosci* 10:803–809
- Scherg M, von Cramon D (1985) A new interpretation of the generators of BAEP waves I-V: Results of spatiotemporal dipole modeling. *Electroencephalogr Clin Neurophysiol* 62:290–299
- Schlaug G, Jäncke L, Huang Y, Steinmetz H (1995) In vivo evidence of structural brain asymmetry in musicians. *Science* 267:699–701
- Schneider P, Sluming V, Roberts N, Scherg M, Goebel R, Specht HJ et al (2005) Structure and functional asymmetry of lateral Heschl’s gyrus reflects pitch perception preference. *Nat Neurosci* 8: 1241–1247
- Schuknecht HF (1993) *Pathology of the ear*, 2nd edn. Lea and Febiger, Philadelphia, PA
- Schuknecht HF, Churchill JA, Doran R (1959) The localization of acetylcholinesterase in the cochlea. *Arch Otolaryngol* 69:549–559
- Seltzer B, Pandya DN (1978) Afferent cortical connections and architectonics of the superior temporal sulcus and surrounding cortex in the rhesus monkey. *Brain Res* 149:1–24
- Shaw M, Baker R (1983) The locations of stapedius and tensor tympani motoneurons in the cat. *J Comp Neurol* 216:10–19
- Spoendlin H (1985) Anatomy of cochlear innervation. *Am J Otolaryngol* 6:453–467
- Spoendlin H, Schrott A (1989) Analysis of the human auditory nerve. *Hear Res* 143:25–38

- Steel KP, Kros CJ (2001) A genetic approach to understanding auditory function. *Nat Genet* 27:143–149
- Steinmetz H, Rademacher J, Huang Y, Hefter H, Zilles K, Thron A, Freund H-J (1989) Cerebral asymmetry: MR planimetry of the human planum temporale. *J Comput Assist Tomogr* 13:996–1005
- Steinmetz H, Volkman J, Jäncke L, Freund H-J (1991) Anatomical left-right asymmetry of language-related temporal cortex is different in left- and right-handers. *Ann Neurol* 29:315–319
- Stockard JJ, Stockard JE, Sherbrough FW (1978) Nonpathologic factors influencing brainstem auditory evoked potentials. *Am J EEG Technol* 18:177
- Stockard JJ, Stockard JE, Sherbrough FW (1986) Brainstem auditory evoked potentials in neurology: Methodology, interpretation, and clinical application. In: Aminoff MJ (ed) *Electrodiagnosis in clinical neurology*, 2nd edn. Churchill Livingstone, New York
- Stone JS, Oesterle EC, Rubel EW (1998) Recent insights into regeneration of auditory and vestibular hair cells. *Curr Opin Neurol* 11:17–24
- Strominger NL (1973) The origin, course and distribution of the dorsal and intermediate acoustic striae in the rhesus monkey. *J Comp Neurol* 147:209–234
- Strominger NL, Nelson LR, Dougherty WJ (1977) Second-order auditory pathways in the chimpanzee. *J Comp Neurol* 172:349–365
- Tanaka Y, Kano T, Yoshida M, Yamadori A (1991) “So-called” cortical deafness. Clinical, neurophysiological and radiological observations. *Brain* 114:2385–2401
- Tardif E, Clarke S (2001) Intrinsic connectivity of human auditory areas: A tracing study with DiI. *Eur J Neurosci* 13:1045–1050
- ten Donkelaar HJ, Lammens M, Cruysberg JRM, Cremers CWJR (2006) Development and developmental disorders of the brain stem. In: ten Donkelaar HJ, Lammens M, Hori A (eds) *Clinical neuroembryology: Development and developmental disorders of the human central nervous system*. Springer, Heidelberg, pp 269–308
- ten Donkelaar HJ, Lohman AHM, Keyser A, van der Vliet AM (2007) Het centrale zenuwstelsel. In: ten Donkelaar HJ, Lohman AHM, Moorman AFM (eds) *Klinische Anatomie en Embryologie*, 3rd edn. Maarssen, Elsevier, pp 981–1141 (in Dutch)
- Terr LI, Edgerton BJ (1985) Three-dimensional reconstruction of the cochlear nuclear complex in humans. *Arch Otolaryngol* 111:495–501
- Thompson GC, Masterton RB (1978) Brainstem auditory pathways involved in reflexive head orientation to sound. *J Neurophysiol* 45:1183–1202
- Tian B, Rauschecker JP (2004) Processing of frequency-modulated sounds in the lateral auditory belt cortex of the rhesus monkey. *J Neurophysiol* 92:2993–3013
- Trojanowski JQ, Jacobson S (1975) Areal and laminar distribution of some pulvinar cortical efferents in rhesus monkeys. *J Comp Neurol* 169:371–392
- Upadhyay J, Ducros M, Knaus TA, Lindgren KA, Silver A, Tager-Flusberg H, Kim D-S (2007) Function and connectivity in human primary auditory cortex: A combined fMRI and DTI study at 3 Tesla. *Cereb Cortex* 17:2420–2432
- von Economo C, Horn L (1930) Über Windungsrelief, Maße und Rindenarchitektur der Supratemporalfläche, ihre individuellen und ihre Seitenunterschiede. *Z Neurol Psychiatr* 130:678–757
- von Economo C, Koskinas GN (1925) *Die Cytoarchitektur der Hirnrinde des erwachsenen Menschen*. Springer, Heidelberg, English translation by LC Triarhou (2008) *Atlas of cytoarchitectonics of the adult human cerebral cortex*. Karger, Basel
- Wallace MN, Johnson PW, Palmer AR (2002) Histochemical identification of cortical areas in the auditory region of the human brain. *Exp Brain Res* 143:499–508
- Weeks RA, Aziz-Sultan A, Bushara KO, Tian B, Wessinger CM, Dang N et al (1999) A PET study of human auditory spatial processing. *Neurosci Lett* 262:155–158
- Weiller C, Isensee C, Rijntjes M, Huber W, Müller S, Bier D et al (1995) Recovery from Wernicke’s aphasia: a positron emission tomographic study. *Ann Neurol* 37:723–732
- Wessinger CM, Buonocore MH, Kussmaul CL, Mangun R (1997) Tonotopy in human auditory cortex examined with functional magnetic resonance imaging. *Hum Brain Mapp* 5:18–25
- Wessinger CM, VanMeter J, Tian B, Van Lare J, Pekar J, Rauschecker JP (2001) Hierarchical organization of the human auditory cortex revealed by functional magnetic resonance imaging. *J Cogn Neurosci* 13:1–7
- Westerhausen R, Grüner R, Specht K, Hugdahl K (2009) Functional relevance of interindividual differences in temporal lobe callosal pathways: a DTI tractography study. *Cereb Cortex* 19:1322–1329
- Winer JA (1984) The human medial geniculate body. *Hear Res* 15:225–247
- Woods RP (1996) Correlation of brain structure and function. In: Toga AW, Mazziotta JC (eds) *Brain mapping: the methods*. Academic, San Diego, CA, pp 313–341
- Zatorre RJ, Binder JR (2000) Functional and structural imaging of the human auditory system. In: Toga AW, Mazziotta JC (eds) *Brain mapping: the systems*. Academic, San Diego, CA, pp 365–402
- Zatorre RJ, Evans AC, Meyer E, Giedde A (1992) Lateralization of phonetic and pitch discrimination in speech processing. *Science* 256:846–849
- Zatorre RJ, Evans AC, Meyer E (1994) Neural mechanisms underlying melodic perception and memory for pitch. *J Neurosci* 14:1908–1919



<http://www.springer.com/978-3-642-19133-6>

Clinical Neuroanatomy
Brain Circuitry and Its Disorders
ten Donkelaar, D.H.J.
2011, XXIV, 834 p., Hardcover
ISBN: 978-3-642-19133-6