

Chapter 2

Recombinant Antibodies and In Vitro Selection Technologies

C. Ronald Geyer, John McCafferty, Stefan Dübel,
Andrew R.M. Bradbury, and Sachdev S. Sidhu

Abstract

Over the past decade, the accumulation of detailed knowledge of antibody structure and function has enabled antibody phage display to emerge as a powerful in vitro alternative to hybridoma methods for creating antibodies. Many antibodies produced using phage display technology have unique properties that are not obtainable using traditional hybridoma technologies. In phage display, selections are performed under controlled, in vitro conditions that are tailored to suit demands of the antigen and the sequence encoding the antibody is immediately available. These features obviate many of the limitations of hybridoma methodology, and because the entire process relies on scalable molecular biology techniques, phage display is also suitable for high-throughput applications. Thus, antibody phage display technology is well suited for genome-scale biotechnology and therapeutic applications. This review describes the antibody phage display technology and highlights examples of antibodies with unique properties that cannot easily be obtained by other technologies.

Key words: In vitro selection, Phage display, Antibodies

1. Introduction

Methods for generating antibodies were initially developed more than a century ago with the production of polyclonal antibody preparations from animal immunizations (1). The advent of hybridoma technology in 1975 enabled the production of monoclonal antibodies through the fusion of myeloma cells with antibody producing B-cells (2). Hybridoma technology advanced our capacity for research and diagnostics by providing homogenous, purified antibody preparations that improved tracking, detection, and quantitation of target molecules in cells and serum. The hybridoma technology, however, is not without its limitations. With regard to generating human therapeutics, hybridoma antibodies are typically from murine sources, which limits their therapeutic

applications due to human anti-mouse antibody reaction (3, 4). A variety of strategies have been developed to address this problem, including chimerization and humanization strategies (5–9), and transgenic animals with human immunoglobulin loci (10–14). Despite these efforts, the generation of antibodies by hybridoma technology is still costly and time-consuming. Further, since these antibodies are produced in animals, it is difficult to generate them against toxic and highly conserved antigens (15) as well as antigens that are not stable in animal systems.

Alongside the hybridoma technology, methods have been established to generate antibodies using in vitro display technologies. The first such method was antibody phage display, introduced 20 years ago (16–18), followed by yeast, ribosome, puromycin-based plasmid, and bacterial display systems (19–22). Selection platforms (23–25) and the design of antibody fragments (26, 27) for making libraries have been widely dealt with in previous reviews. The purpose of this review is to illustrate how in vitro selection, especially phage display, has yielded antibodies with remarkable properties that are difficult to obtain using traditional immunization methods.

Principles behind methods used in all in vitro display systems are similar whatever the display platform, and center on the coupling of genotype (gene) to phenotype (binding protein). In practice, this comprises the creation of DNA libraries encoding binding molecules such as antibodies, the display of the encoded proteins, the application of selective pressure based on the binding properties of the encoded proteins, followed by growth and screening of individual clones. Since in vitro display methods are performed in bacterial or yeast systems, the turnaround time for antibody generation is less, and the potential for high-throughput generation of binders is greater (28). The power of in vitro antibody selection is further enhanced by the ability to precisely control selection conditions. In contrast to animal immunization, where there is little control over the nature of antibodies produced, manipulation of selection conditions can be carried out in vitro, for example, by presentation of specific conformations of the target antigen or by including competitors to direct selection towards targets or epitopes of interest. In vitro selection methods also overcome the problem of tolerance, which limits the potential for making anti-self antibodies. As tolerance is applied to specific variable heavy (VH) and light (VL) domain combinations that recognize self-antigens, when in vitro libraries are created from natural sources new combinations with the capacity to recognize self-antigens can be created. This has been proven for example by the selection of hundreds of human antibodies from naive libraries to human targets (29–32). For libraries that are constructed using synthetic diversity, the concept of tolerance does not apply. This enables the selection of antibodies against highly conserved targets such as ubiquitin (33, 34), histones (35), hemoglobins (36), and posttranslational modifications (37–39).

Another advantage of in vitro selected antibodies is that the gene encoding the antibody is cloned simultaneously with selection. This is perhaps the most crucial difference between hybridoma and in vitro selection technologies and provides many advantages for engineering selected antibodies. For example, affinity maturation of selected antibodies can be easily performed using in vitro selection technologies. Maturation of antibody affinity to the picomolar range (40–44) has become relatively routine and, at least in one case, femtomolar affinity has been achieved (45). These affinities are far higher than those that can be obtained by immunization, which are limited to ~100 pM by the physiological mechanisms of B-cell activation (46–48). In addition, antibody specificities can be broadened or narrowed by appropriate selection conditions. As a result, in vitro selection has yielded antibodies with remarkable properties that are either a direct result of the flexibility and control that can be applied to all aspects of the selection processes, or novel properties developed as a direct result of the recombinant nature of selected proteins. Finally, the availability of the antibody gene allows the creation of a large variety of antibody derivatives with added functions by simple subcloning.

2. Phage-Displayed Antibody Libraries

Antibody libraries for in vitro selections can be generated from immunized repertoires, natural naïve repertoires, or designed “synthetic” repertoires. Strategies to generate antibody libraries have been described extensively (49, 50) and are discussed briefly here. Immune antibody libraries are generated by cloning antibody fragments, either single-chain variable fragments (scFvs) or antigen binding fragments (Fabs), from IgG mRNAs obtained from activated B-cells (49, 50). These libraries are biased for members that bind a specific antigen and require that a new library be generated for each antigen of interest. Immune antibody libraries have been generated against a number of different species (51, 52). Human immune libraries have been constructed from virus-infected patients to generate neutralizing antibodies and from cancer patients to isolate tumor specific antibodies (53–57).

Although natural naïve antibody libraries have been generated using IgM or IgG mRNAs from resting B-cells, initial results indicated that libraries based on IgM mRNA yield more binders (17), probably because the IgG fraction is biased towards recent immune responses. Naturally rearranged variable region genes have been used to construct large antibody fragment libraries (32, 58–61). In contrast to immune libraries, naïve libraries can be used to generate antibodies against a variety of antigens; however, they generally bind with lower affinity and may need to be affinity matured.

In synthetic antibody libraries, antibody diversity is designed and synthesized in a controlled manner. In these libraries, the composition of complementarity determining regions (CDRs) can be precisely defined. A number of approaches have been used to design synthetic libraries and they vary in the number of variable framework regions used, the design of CDR diversity, and the library construction method. Synthetic libraries have been constructed using a variety of different variable framework genes (62–65), with diversity introduced into the CDRs, primarily in CDRH3 and CDRL3. Results from selections with these libraries have shown that larger libraries yield antibodies with higher affinity and greater specificity (50), and that specific variable framework regions are over represented in the selected antibody fragments. This observation led to the development of libraries using a single VH and VL combination (66–68). A number of different libraries have been devised that differ in the variable gene and the diversification strategy. Most libraries use a common VH domain (VH3–23) as it is stable, expressed well in bacteria and on phage, and pairs with most VL domains. Synthetic libraries are constructed by cloning oligonucleotides into the CDRs of defined antibody fragments, and thus, these libraries are not limited to the diversity present in natural repertoires. However, libraries have been created by grafting natural CDRs into single frameworks (69), as well as by using designed oligonucleotides that mimic the CDR diversity observed in natural repertoires (68, 70). Libraries have also been designed with restricted diversity in the CDRs, based on the observation that tyrosine and serine are enriched in the antigen-binding sites of antibodies (71–73).

In addition to variable domain and CDR design, antibody fragments must be fused to the phage coat protein in order to establish the genotype/phenotype connection. Phage display requires that antibody fragments be displayed rather than IgGs. The two most popular antibody fragments used to display the variable antigen binding domains are the Fab and the scFv. The Fab is a heterodimer consisting of the variable and first constant domains of heavy and light chains. The scFv consists of the variable domains from the light and heavy chains joined by a peptide linker. These antibody fragments are most commonly displayed on phage by fusing them to pIII or pVIII coat proteins. The pVIII coat protein can potentially enable the display of multiple polypeptides on the phage surface, as approximately 2,500 copies of pVIII are present on each phage particle (74). However, large proteins are not well tolerated as pVIII fusions, which limits their use for displaying antibody fragments (75), and in direct comparisons, pIII display appears to be more efficient than pVIII (76). Thus, antibody fragments are more commonly fused to the pIII coat protein. There are approximately five pIII coat proteins on one tip of the phage particle. Fusion to the pIII coat protein results in low-level display of antibody fragments using phagemid systems (49), and display

levels can easily be switched between monovalent and oligovalent display (77), which allows selections from large libraries to be optimized for obtaining higher affinity antibody fragments by avoiding avidity effects.

2.1. Diverse Applications of Antibody Phage Display

Since the invention of antibody phage display, intellectual property issues have delayed its broad use and have limited the number of synthetic antibodies in the clinic during the 1990s. Nonetheless, as of 2010, phage display technology has been used to generate at least 35 human antibodies that are in clinical development (78). The FDA has approved two of these antibodies (adalimumab and belimumab) and one is under review (raxibacumab). The success of phage-derived antibodies in clinical trials is similar to monoclonal antibodies derived from other technologies (78). The number of antibodies generated using phage display is rapidly expanding and a comprehensive catalog is beyond the scope of this review. Below, we highlight some examples of phage-derived antibodies generated against extracellular targets as well as some of the unique features of antibodies that can be generated using phage display technologies.

2.2. Antibodies Against Extracellular Targets

Phage display technologies are extremely powerful for generating functional antibodies that disrupt normal or pathological extracellular signaling. Phage display selects for antibodies that bind their target with high affinity, however this does not guarantee that they will have the desired function. Phage display, however, can produce many antibodies that bind a given target, increasing the chance that some of the antibodies will possess the desired properties. One such example was the use of phage display to generate more than 1,200 antibodies against the B-lymphocyte stimulator (Blys) (79), a potent cytokine for B-cell proliferation and differentiation. Biochemical and cellular assays were used to subsequently identify antibodies, many with subnanomolar affinities, which blocked B-cell activation by inhibiting the interaction between Blys and its receptor. One of these antibodies, which showed specificity for secreted Blys, was affinity matured and shown to be a potent inhibitor of Blys signaling (80). This antibody, belimumab, has been approved by the FDA in March 2011 for use in treatment of systemic lupus erythematosus.

A second example of an antibody isolated by phage display against a cytokine target is the tumor necrosis factor alpha (TNF α) blocking antibody, adalimumab. TNF α is a proinflammatory mediator implicated in autoimmune conditions. Adalimumab has been approved for the treatment of several conditions including rheumatoid arthritis, ankylosing spondylitis, chronic plaque psoriasis, and Crohn's disease, which was the first fully human antibody approved by the FDA in 2002. A number of other antibodies against soluble ligands have been generated by phage display and are in advanced clinical trials (81).

Another strategy to block receptor signaling is to target receptor sites that prevent ligand binding. A recent series of studies highlights the use of phage-derived antibodies to block Insulin-like Growth Factor 1 Receptor (IGF-1R) signaling (82–84). Phage display was used to generate antibodies against two unique epitopes of IGF-1R. Both antibodies blocked binding of Insulin-like Growth Factor 1 (IGF-1) and Insulin-like Growth Factor 2 (IGF-2), but they did so by either directly competing for ligand binding or by an allosteric mechanism, which decreased the affinity of ligand binding (84). Interestingly, cotreatment with both antibodies improved both the potency and extent of IGF-1 and IGF-2 blockade compared to treatment with either antibody alone. Similar results have been observed with Her2 (Human Epidermal growth factor Receptor 2), where combinations of antibodies that bind unique epitopes have greater activity than either antibody alone (85).

Antibodies have also been generated to block ligand-induced conformational changes in Notch receptors (86). The ectodomain of the Notch receptor contains multiple epidermal growth factor (EGF) repeats and ligand binding induces a conformational change at the juxtamembrane negative regulatory region, which causes a protease cleavage site to be exposed. Subsequent proteolysis causes the intracellular domain to be translocated to the nucleus. Phage display was used to generate antibodies that target the juxtamembrane negative regulatory regions of Notch-1 and Notch-2 (86). These antibodies bind and stabilize the “closed” conformation of the Notch receptor, preventing proteolytic cleavage.

Ligands often act either by causing dimerization of their target receptors or by inducing conformational changes in preexisting dimers. Antibodies targeting the ligand-binding domain can, in some instances, mimic the effect of the natural ligand and cause receptor activation rather than inhibition. For example, phage display was used to generate antibodies that bind to Muscle Specific Kinase (MuSK) (87) or CD40 (88) and function as agonists for receptor activation (87). In another recent study, over 500 distinct antibodies were generated against TRAIL receptor-1 (TRAIL-R1) and TRAIL receptor-2 (TRAIL-R2) (89). TRAIL is a homotrimeric ligand that causes multimerization of TRAIL receptors, which in turn leads to apoptosis, particularly in tumor cells overexpressing the receptors. Ten agonistic antibodies specific for TRAIL-R1 and six antibodies acting only on TRAIL-R2 were identified. As expected, these antibodies competed for binding with TRAIL, but surprisingly, they were active agonists as monovalent antibodies in either scFv or Fab formats, and activity was not enhanced upon conversion to IgG. The mechanism of action for this unusual agonistic activity is still unclear.

In addition to selecting antibodies that bind to purified proteins, phage display can be used to select antibodies that recognize targets expressed on the surfaces of cells. A number of selection

protocols have been developed to select antibodies that bind cell surface proteins. These include strategies that incorporate negative selections or preabsorption steps (90–94), strategies to remove unbound phage (95, 96), and the pathfinder approach (97, 98). In vitro selection schemes have also been devised to select for antibodies that mediate receptor internalization (99–101). In these selections, phage libraries are incubated with target cells and then phage that bind the cell surface are removed and phage antibodies inside the cell are isolated. This strategy is useful for generating antibodies to deliver drugs to specific cells (102, 103). The ability to perform selections directly on cells with negative selections has proven to be a powerful trait of phage display technology.

2.3. Antibodies Against Infectious Disease Targets

Phage display has been used to select antibodies against a variety of infectious agents. For example, antibodies have been isolated that discriminate between strains of Hanta (104), Dengue (105, 106), Influenza (107, 108), Ebola (109), and Venezuelan equine encephalitis virus (110). Further, phage display selections do not require purified virus. For the Venezuelan equine encephalitis virus selection, the use of competitive binding conditions allowed antibodies to be generated against impure virus preparations (110). In these selections, cell extracts from uninfected cells were added to the binding buffer, which eliminated the isolation of antibodies against components of the host cell and allowed antibodies to be generated against the viral envelope. Human antibodies have also been selected against a number of bacterial bio-threat targets, including *Brucella melitensis* (111), *Burkholderia mallei*, *Burkholderia pseudomallei* (112), and anthrax toxins (113–117) and spores (118).

In one study, antibodies were used to block protein interactions associated with influenza entry into target cells. Phage display was used to generate antibodies that recognize the H5 hemagglutinin influenza ectodomain (119, 120). Structural characterization of one of these antibodies bound to H5 showed that it binds to hemagglutinin by inserting its heavy chain into a highly conserved pocket in the stem region, which prevents structural reorganizations required for membrane fusion. This conserved epitope is found in many different influenza viruses and this antibody was shown to neutralize H5N1, H1N1, H2N2, H6N1, H6N2, H8N4, and H9N2 viruses. Although antibodies have not been generated against this epitope by traditional immunization, and antibodies with this specificity do not normally arise during infection, antibodies with similar VH gene usage and neutralizing activity have been selected from phage antibody libraries created from human IgM+ memory B-cells from recently infected individuals (121).

2.4. Antibodies with Ultra-High Specificity

With protein targets, antibodies have been selected that display high specificity for a chosen target. For example, antibodies have been generated that differentiate between chicken and quail

lysozyme, which differ by only four amino acids (122). Another example is the isolation of antibodies that distinguish between the SH2 domains of ABL1 and ABL2 tyrosine kinase (29, 123), which differ by only 11%, and for which it has not been possible to obtain specific antibodies by immunization. In such studies, negative selection steps have been incorporated into phage display selections to generate antibodies with desired specificity. For example, antibodies were generated that recognize only fetal and not adult hemoglobin (124). In this study, antibodies that recognize adult hemoglobin were depleted by preincubating the antibody phage library with adult hemoglobin prior to each round of selection against fetal hemoglobin. Phage display has also been used to generate antibodies that recognize specific protein complexes, for example the generation of antibodies that recognize unique peptides in the context of specific MHC molecules (125–128).

The technology has also been applied to selectively target alternatively spliced fibronectin variants associated with tumor neovasculature. Extra-domain A (EDA) and B (EDB) are fibronectin variants each of which contain an additional domain, both being highly conserved between human and mouse. Using a synthetic antibody library, it was possible to select human/mouse cross-reactive scFvs against each of the recombinantly purified extra domains (129, 130), and these antibodies were effective for immunohistochemical analysis *in vitro* and for biodistribution studies *in vivo*. In the case of the anti-EDB antibody (130), the modular nature of the scFv was exploited to engineer numerous fusion proteins with potential for cancer therapy (131), and three of these derivatives are now in clinical trials.

2.5. Antibodies Against Specific Protein Conformations

Phage display has been used to generate antibodies that recognize specific protein conformations. Many signaling proteins exist in specific conformational states that mediate distinct cellular responses. Antibodies that recognize specific protein conformations provide a unique resource for characterizing signaling pathways. These types of antibodies are difficult to generate via immunization strategies, as protein conformations are often unstable in an immunized animal. In contrast, *in vitro* selection technologies are ideally suited for these applications because selection conditions can be precisely controlled to favor particular conformations. Negative selections can be used to deplete nonspecific binders and affinity maturation strategies can be employed to fine-tune specificity.

Phage display has been used to generate antibody fragments that specifically recognize the GTP-bound form of Rab6 (132) and active and inactive forms of Caspase-1 (133). scFvs specific to the GTP-bound form of the small guanosine triphosphatase (GTPase) Rab6 were generated by performing selections against a

GTP-locked mutant (132). Fabs specific to on and off states of caspase-1 were generated by selecting libraries against caspase-1 complexed to small molecules that lock it in the on or off state (133). Fabs were converted into full-length IgGs to produce highly sensitive affinity reagents that could be used to probe the localization of active caspase-1 in cells (133). Conformation-specific antibodies have also been generated against active cell membrane receptors by performing phage selections on whole cells (134).

2.6. Antibodies Against Integral Membrane Proteins

Integral membrane proteins absolutely require a membrane or detergent environment to maintain their native conformation. Consequently, the generation of conformation-specific antibodies against membrane proteins by animal immunization is severely limited by the denaturing effects of the serum environment. In contrast, the ability to control selection conditions makes in vitro techniques much more amenable to this task. By performing selections in the presence of detergent, high affinity Fabs were isolated against the citrate transporter CitS from *Klebsiella pneumoniae* (135), as well as against the potassium channel KcsA from *Streptomyces lividans* (136). In the latter case, the Fabs were used as crystallization chaperones that enabled the elucidation of the crystal structure of the full-length potassium channel.

2.7. Antibodies Against RNA

Phage display has been used to generate antibodies that recognize structured RNA molecules (137), which have proven to be essentially nonimmunogenic for hybridoma methods. Using a nuclease free selection buffer, high affinity Fabs were isolated against a structured domain from the *Tetrahymena* group I intron. The structure of the Fab/RNA complex was solved to high resolution, highlighting the use of antibody fragments as chaperones for RNA crystallization. Fabs were also obtained against a class I ligase ribozyme and were used as chaperones to obtain the crystal structure (138). One Fab recognized a small, discrete sequence in the ribozyme and retained binding capacity when this sequence was transferred to other RNA structures, providing a novel RNA crystallization chaperone system.

2.8. Antibodies Against Posttranslational Modifications

Phage display has been useful for detecting posttranslational modifications that have proven intractable to immunization. For example, sulfotyrosine is a posttranslational modification predicted to occur in 30% of all secretory and membrane proteins (139). Perhaps because of its ubiquitous nature, traditional immunizations have consistently failed to produce anti-sulfotyrosine antibodies. However, using phage display, antibodies were readily generated to recognize proteins containing sulfotyrosine (but not tyrosine or tyrosine phosphate) independently of protein context or sequence (39, 140).

3. Exploiting the Recombinant Nature of In Vitro Antibodies

Since phage display is an in vitro selection method, it offers many advantages for engineering antibodies. The phage display system provides the antibody gene and sequence following selection against a particular target. This allows antibodies to be easily further evolved and engineered to improve binding, to narrow or broaden specificity, or to improve expression as IgGs or as fusions to functional moieties.

Antibody fragments isolated from an initial phage display selection can be used directly as an affinity reagent, or they can be used as leads for further improving binding. Since the sequence of isolated antibody fragments are rapidly determined by sequencing, it is straightforward to make second-generation libraries by introducing mutations into antibody fragments. Affinity maturation strategies have been used to generate antibodies with affinities that exceed those of natural antibodies, which are limited to a ceiling of $K_d > 0.1$ nM by the nature of the B cell response (46–48). There are many different approaches for introducing diversity into antibody fragments to improve the affinities obtained from combinatorial libraries. With in vitro affinity maturation selections, randomized antibody fragments undergo selection with increased pressure to identify variants with enhanced affinity (141). In general, there are two approaches for generating diversification: targeted and nontargeted. There are many examples of in vitro affinity maturation, and here we highlight some key studies that demonstrate the power of the process.

Error prone PCR is the most common method for introducing nontargeted mutations (142). In this method, sequence diversity is randomly introduced into the antibody fragment gene by mutagenic PCR strategies (50). The down side of this approach is that deleterious mutations can be introduced into the conserved framework region, which reduces the number of functional antibody fragments in the library. DNA shuffling is another method for introducing nontargeted mutations (143). In this method, a group of closely related sequence are randomly fragmented and then reassembled by PCR, which leads to a shuffling of DNA fragments. The approach can be combined with PCR mutagenesis to further enhance diversity. This method was used to increase the affinity of an scFv for fluorescein by 1,000-fold, resulting in subpicomolar affinity (45).

As an alternative to random PCR mutagenesis, knowledge of the antibody sequence enables precise targeting of mutations for affinity maturation. Targeted mutation strategies have the advantage of focusing mutations to CDR loops, which are most likely to enhance affinity without introducing deleterious mutations in regions that may affect protein folding and stability. Targeted mutagenesis can be

performed using degenerate oligonucleotides, which allows for precise control over the locations where diversity is introduced. Further, CDRs can be targeted in either a parallel or sequential fashion. By targeting CDR loops in phage-displayed antibody libraries, an anti-HIV-1 antibody (44) and an anti-c-erbB-2 (43) antibody were affinity matured to the low picomolar range.

In addition to affinity, the specificity of antibodies can be altered by phage display. While absolute specificity for a single antigen is generally the goal of antibody design, cross-reactivity is desirable for certain applications. For example, in the case of antibody therapeutics, species cross-reactivity enables assessment of therapeutic efficacy and toxicity in animal models. Cross-reactive antibodies are often difficult to obtain by hybridoma methods because of the conservation of functional sites on proteins across species. In contrast, in vitro phage antibody libraries are not affected by immune tolerance, and generation of antibodies that target conserved sites across species orthologues has proven to be the rule rather than the exception. For example, antibodies that cross-react with human and mouse VEGF were obtained directly from phage libraries without further selections to broaden specificity (144, 145). For BAFF/BLyS receptor 3 (BR3), antibodies generated against human BR3 showed weak cross-reactivity with mouse BR3. In this case, phage display was used to select cross-reactive antibodies from secondary libraries (146). This strategy has also been used to generate antibodies with cross-reactivity towards CXCL10 and CXCL9 homologues (147). In an extreme example, this approach has been used to broaden the specificity of trastuzumab so that it cross-reacts with Erb-B2 and VEGF, two proteins that share no sequence or structural homology (148). In this case, secondary libraries were created by diversifying the light chain, which plays a minor role in Erb-B2 recognition. Extensive affinity maturation produced antibodies with low nanomolar affinity for both ErbB2 and VEGF (148).

The ability to improve affinity and broaden specificity also has major implications for the development of antibodies against infectious disease agents. For the effective inhibition of viral infection and bacterial toxins, antibodies must be of very high affinity, and at the same time, they should be cross-reactive with a variety of antigen subtypes to afford broad protection against pathogen variants. A powerful example of using affinity and specificity selection cycles was demonstrated for an antibody with broad specificity for different subtypes of Botulinum toxins. Remarkably, this antibody is able to recognize Botulinum toxins A, B, E, and F, all the serotypes that afflict humans (149, 150).

The ability to rapidly obtain the gene for a selected antibody fragment allows the antibody to be easily engineered by simple subcloning strategies. Antibody fragments produced from phage display selections can be subcloned into IgG expression systems to

produce antibodies in mammalian tissue culture systems (151). Antibody fragments have also been engineered with other functions by fusing them to peptides and proteins that induce dimerization (152) and multimerization (153–155) to facilitate detection and purification, or that provide them with fluorescent (156–158) or enzymatic (159) properties. In vivo peptide biotinylation tags have been fused to the C-termini of antibody fragments to enable antibodies to be immobilized or multimerized (153, 160–163). Antibody fragments have been fused to the dimeric enzyme alkaline phosphatase, which provides both dimerization and alkaline phosphatase activity that greatly enhances functionality and simplifies screening (30, 159). scFvs have also been fused to Fc domains, converting them into antibody-like molecules with properties similar to IgGs (164–167).

Recombinant technologies have enabled the generation of a large variety of bispecific antibodies that recognize two different targets (ref. 168 and see also Chapter 16). This can be accomplished by engineering two different Fc domains to allow heterologous pairing (169, 170). Alternatively, scFvs can be fused recombinantly to IgGs to impart bifunctionality (171). This strategy has recently been used to generate bispecific antibodies against IGF-1R, which blocked ligand binding better than either monospecific IgG (172). The bispecific antibodies also showed an improved ability to reduce the growth of multiple tumor cell lines, to inhibit ligand-induced IGF-1R signaling in tumor cells, and to block in vivo tumor growth (172). Bispecific antibody fragments have also been generated by varying the peptide linker length that connects VH and VL domains in scFvs. This strategy has been used to generate dimers (173–175), trimers (174, 176, 177), and tetramers (178). Various other bispecific antibody designs have also been created (see ref. 179 for a review).

Within the context of improved antibody therapeutics, fusion proteins that can extend the capabilities of natural IgGs have been constructed. By exchanging or engineering the Fc region, antibodies with designed pharmacokinetics and improved effector functions have been obtained (for reviews see refs. 180, 181).

4. High-Throughput Antibody Selections and Next-Generation Sequencing

The ease and speed with which antibody fragments can be selected using phage display, usually between 1 and 2 weeks, allows the technology to be implemented in a high-throughput manner (29, 30, 123, 182, 183) and see also Chapters 3–6. Initial experiments using a limited number of targets showed that antibodies could be generated from semiautomated selections using phage antibody

libraries (182–184). More recent studies have shown that phage display selections can be scaled up to target a larger number of antigens (29, 30, 123, 182, 183). For example, phage antibody selections were carried out on over 400 different antigens representing 292 proteins. In total, 25% of antibodies screened were positive, and 80% of these were specific when screened against irrelevant antigens (30). The practicality of generating antibodies against a broad array of different targets was further demonstrated in a recent multinational study in which antibodies against 20 different SH2 domains were generated via immunization and hybridoma technology or by phage display technology (31). The phage display selections were successful against all 20 targets and yielded at least ten unique binders for each target. Both phage display and hybridoma technologies produced many binders with low-nanomolar affinities. Antibodies were validated using a number of different assays, including microarrays, immunoblots, immunofluorescence, and immunoprecipitation. Overall, this study shows that antibodies with high affinity and specificity can be efficiently generated using high-throughput phage technologies.

Next-generation sequencing (NGS) technologies have been used to improve the characterization of mouse immunizations and phage display selections and to speed up the identification of antibodies. Several studies have utilized NGS for high-resolution analysis of natural (185) and synthetic (186) antibody repertoires. The 454 sequencing platform, which provides sequencing reads between 250 and 400 bases, was used to provide information on the diversity of CDRs and on VH and VL pairings (185, 187). The Illumina sequencing platform, which provides a higher number of shorter reads (~100 bases), was used to analyze the CDRH3 region of an scFv library (186). NGS platforms provide information on V-gene family frequency, CDR length and diversity, and a comparison of the theoretical and actual properties of the library (188). NGS has also been used to characterize how the immunoglobulin repertoire changes after immunization, where NGS was used to monitor enrichment in antigen specific V-genes (189).

For phage display selections, NGS was used to monitor enrichment of antibody sequences during successive rounds of selection (186). The information generated by NGS can be used to bypass antibody screening, which is time-consuming and expensive. For antibodies generated by mouse immunization, NGS was used to identify heavy and light chains and pairings between them were inferred based on their frequencies in the repertoire (189). A similar strategy was used to identify antibodies from phage display selections (186). High frequency antibody fragments were identified following rounds of selection and desired antibody fragments were recovered by PCR (186). Identification of antibody sequences following selection eliminates the characterization of redundant clones. Further, it reduces the amount of target protein

required to perform and characterize antibodies from immunizations or phage selections. Lastly, NGS was used to characterize an antibody phage display selection against a protein target IL-6, expressed on the surface of *E. coli* (190). This study highlighted the potential for using NGS to characterize phage display selections against complex targets.

5. Conclusions and Future Perspectives

It is now well accepted in the scientific community that there is an urgent need to improve antibody quality in general, as an alarmingly high proportion of commercial antibodies either show poor specificity, or fail to even recognize their targets (191–194). At the same time, high-throughput genomics and proteomics technologies have vastly expanded the scope of proteins and pathways that now await detailed analysis at the cell biology level. To deal with the thousands of new proteins revealed by genomics and proteomics projects, there is an urgent need for high quality antibodies, and it is clear that the current hybridoma methods are not suitable for this task.

In this landscape, the emergence of high quality in vitro antibody libraries is both timely and opportune. Numerous studies have reported in vitro repertoires that routinely yield antibodies that rival or surpass hybridoma antibodies in terms of functionality. Moreover, while further improvements in hybridoma technology are likely to be slight, in vitro repertoires and selection methods continue to improve. Universal in vitro libraries that can provide antibodies against virtually any antigen are now a reality, and it is hoped that the technology can be broadly disseminated in the near future.

The recombinant nature of in vitro repertoires is a fundamental advantage that extends the technology beyond the scope of hybridoma technology. With synthetic antibodies in particular, frameworks can be chosen for favorable traits such as low immunogenicity or high stability, initial clones can be rapidly affinity matured and reformatted, and antibodies can be shared and distributed in the form of synthetic DNA. Furthermore, precise control over selection conditions allows for high precision engineering of specificity and affinity. Further standardization of libraries and selection methods will enable the adaptation of the technology to high-throughput pipelines to enable antibody generation on a proteome scale, and the ability to select directly against cells and tissues will further expand the scope of the technology. Clearly, in vitro antibody libraries are ideally suited for addressing the challenges of cell biology in the genomics era, and the technology is poised to play an ever-expanding role in the future of biological research.

References

1. von Behring E, Kitasato S (1890) Über das zustandekommen der diphtherie-immunität und der tetanus-immunität bei thieren. Deut Med Wochenzeitschr 16:1113–1114
2. Köhler G, Milstein C (1975) Continuous cultures of fused cells secreting antibody of predefined specificity. Nature 256:495–497
3. Courtenay-Luck NS, Epenetos AA, Moore R et al (1986) Development of primary and secondary immune responses to mouse monoclonal antibodies used in the diagnosis and therapy of malignant neoplasms. Cancer Res 46:6489–6493
4. Tjandra JJ, Ramadi L, McKenzie IF (1990) Development of human anti-murine antibody (HAMA) response in patients. Immunol Cell Biol 68:367–376
5. Almagro JC, Fransson J (2008) Humanization of antibodies. Front Biosci 13:1619–1633
6. Hwang WYK, Foote J (2005) Immunogenicity of engineered antibodies. Methods 36:3–10
7. Kashmiri SVS, De Pascalis R, Gonzales NR, Schlom J (2005) SDR grafting—a new approach to antibody humanization. Methods 36:25–34
8. Studnicka GM, Soares S, Better M et al (1994) Human engineered monoclonal antibodies retain full specific binding activity by preserving non-CDR complementarity-modulating residues. Protein Eng 7:805–814
9. Osbourn J, Groves M, Vaughan T (2005) From rodent reagents to human therapeutics using antibody guided selection. Methods 36: 61–68
10. Fishwild DM, O'Donnell SL, Bengoechea T et al (1996) High-avidity human IgG kappa monoclonal antibodies from a novel strain of minilocus transgenic mice. Nat Biotechnol 14:845–851
11. Jakobovits A (1995) Production of fully human antibodies by transgenic mice. Curr Opin Biotechnol 6:561–566
12. Kuroiwa Y, Kasinathan P, Sathiyaseelan T et al (2009) Antigen specific human polyclonal antibodies from hyperimmunized cattle. Nat Biotechnol 27:173–181
13. Lonberg N, Huszar D (1995) Human antibodies from transgenic mice. Int Rev Immunol 13:65–93
14. Weiner LM (2006) Fully human therapeutic monoclonal antibodies. J Immunother 29:1–9
15. Winter G, Milstein C (1991) Man-made antibodies. Nature 349:293–299
16. McCafferty J, Griffiths AD, Winter G et al (1990) Phage antibodies: filamentous phage displaying antibody variable domains. Nature 348:552–554
17. Marks JD, Hoogenboom HR, Bonnert TP et al (1991) By-passing immunization. Human antibodies from V-gene libraries displayed on phage. J Mol Bio 222:581–597
18. Breitling F, Dübel S, Seehaus T (1991) A surface expression vector for antibody screening. Gene 104:147–153
19. Boder ET, Wittrup KD (1997) Yeast surface display for screening combinatorial polypeptide libraries. Nat Biotechnol 15:553–557
20. Feldhaus MJ, Siegel RW, Opresko LK et al (2003) Flow-cytometric isolation of human antibodies from a nonimmune *Saccharomyces cerevisiae* surface display library. Nat Biotechnol 21:163–170
21. He M, Taussig MJ (2007) Rapid discovery of protein interactions by cell-free protein technologies. Biochem Soc Trans 35:962–965
22. Jostock T, Dübel S (2005) Screening of molecular repertoires by microbial surface display. Comb Chem High Throughput Screen 8:127–133
23. Paschke M (2006) Phage display systems and their applications. Appl Microbiol Biotechnol 70:2–11
24. Zahnd C, Amstutz P, Pluckthun A (2007) Ribosome display: selecting and evolving proteins in vitro that specifically bind to a target. Nat Methods 4:269–279
25. Chao G, Lau WL, Hackel BJ et al (2006) Isolating and engineering human antibodies using yeast surface display. Nat Protoc 1: 755–768
26. Benhar I (2007) Design of synthetic antibody libraries. Expert Opin Biol Ther 7:763–779
27. Bradbury AR, Marks JD (2004) Antibodies from phage antibody libraries. J Immunol Methods 290:29–49
28. Dübel S, Stoevesandt O, Taussig MJ et al (2010) Generating recombinant antibodies to the complete human proteome. Trends Biotechnol 28:333–339
29. Mersmann M, Meier D, Mersmann J et al (2010) Towards proteome scale antibody selections using phage display. New Biotechnol 27:118–128
30. Schofield DJ et al (2007) Application of phage display to high throughput antibody generation and characterization. Genome Biol 8:R254
31. Colwill K, Gräslund S, Persson MA et al (2011) A roadmap to generate renewable protein binders to the human proteome. Nat Methods 8:551–561

32. Hust M, Meyer T, Voedisch B et al (2011) A human scFv antibody generation pipeline for proteome research. *J Biotechnol* 152: 159–170
33. Koide A, Bailey CW, Huang X et al (1998) The fibronectin type III domain as a scaffold for novel binding proteins. *J Mol Biol* 284: 1141–1151
34. Angenendt P, Wilde J, Kijanka G et al (2004) Seeing better through a MIST: evaluation of monoclonal recombinant antibody fragments on microarrays. *Anal Chem* 76:2916–2921
35. Philibert P, Stoessel A, Wang W (2007) A focused antibody library for selecting scFvs expressed at high levels in the cytoplasm. *BMC Biotechnol* 7:1472–6750
36. Parsons HL, Earnshaw JC, Wilton J et al (1996) Directing phage selections towards specific epitopes. *Protein Eng* 9:1043–1049
37. Lassen KS, Bradbury AR, Rehfeld JF et al (2008) Microscale characterization of the binding specificity and affinity of a monoclonal anti-sulfotyrosyl IgG antibody. *Electrophoresis* 29:2557–2564
38. Kehoe JW, Velappan N, Walbol M et al (2006) Using phage display to select antibodies recognizing post-translational modifications independently of sequence context. *Mol Cell Proteomics* 5:2350–2363
39. Hoffhines AJ, Damoc E, Bridges KG et al (2006) Detection and purification of tyrosine-sulfated proteins using a novel anti-sulfotyrosine monoclonal antibody. *J Biol Chem* 281:37877–37887
40. Raza A, Garcia-Rodriguez C, Lou J et al (2005) Molecular evolution of antibody affinity for sensitive detection of botulinum neurotoxin type A. *J Mol Biol* 351:158–169
41. Lee CV, Liang WC, Dennis MS et al (2004) High-affinity human antibodies from phage-displayed synthetic Fab libraries with a single framework scaffold. *J Mol Biol* 340: 1073–1093
42. Hanes J, Schaffitzel C, Knappik A et al (2000) Picomolar affinity antibodies from a fully synthetic naive library selected and evolved by ribosome display. *Nat Biotechnol* 18: 1287–1292
43. Schier R, McCall A, Adams GP et al (1996) Isolation of picomolar affinity anti-c-erbB-2 single-chain Fv by molecular evolution of the complementarity determining regions in the center of the antibody binding site. *J Mol Biol* 263:551–567
44. Yang WP, Green K, Pinz-Sweeney S et al (1995) CDR walking mutagenesis for the affinity maturation of a potent human anti-HIV-1 antibody into the picomolar range. *J Mol Biol* 254:392–403
45. Boder ET, Midelfort KS, Wittrup KD (2000) Directed evolution of antibody fragments with monovalent femtomolar antigen-binding affinity. *Proc Natl Acad Sci USA* 97: 10701–10705
46. Foote J, Eisen HN (1995) Kinetic and affinity limits on antibodies produced during immune responses. *Proc Natl Acad Sci USA* 92: 1254–1256
47. Foote J, Eisen HN (2000) Breaking the affinity ceiling for antibodies and T cell receptors. *Proc Natl Acad Sci USA* 97:10679–10681
48. Batista FD, Neuberger MS (1998) Affinity dependence of the B cell response to antigen: a threshold, a ceiling, and the importance of off-rate. *Immunity* 8:751–759
49. Sidhu SS (2005) Phage display in biotechnology and drug discovery. CRC Press, Boca Raton, FL
50. Ponsel D, Neugebauer J, Ladetzki-Baehs K et al (2011) High affinity, developability and functional size: the holy grail of combinatorial antibody library generation. *Molecules* 16:3675–3700
51. Azzazy HM, Highsmith WE Jr (2002) Phage display technology: clinical applications and recent innovations. *Clin Biochem* 35: 425–445
52. Benhar I (2007) Design of synthetic antibody libraries. *Expert Opin Biol Ther* 7:763–779
53. Burton DR, Barbas CF III, Persson MA et al (1991) A large array of human monoclonal antibodies to type I human immunodeficiency virus from combinatorial libraries of asymptomatic seropositive individuals. *Proc Natl Acad Sci USA* 88:10134–10137
54. Kramer RA, Marissen WE, Goudsmit J et al (2005) The human antibody repertoire specific for rabies virus glycoprotein as selected from immune libraries. *Eur J Immunol* 35:2131–2145
55. de Carvalho NC, Williamson RA, Parren PW et al (2002) Neutralizing human Fab fragments against measles virus recovered by phage display. *J Virol* 76:251–258
56. Zebedee SL, Barbas CF III, Hom YL et al (1992) Human combinatorial antibody libraries to hepatitis B surface antigen. *Proc Natl Acad Sci USA* 89:3175–3179
57. Cai X, Garen A (1995) Anti-melanoma antibodies from melanoma patients immunized with genetically modified autologous tumor cells: selection of specific antibodies from single-chain Fv fusion phage libraries. *Proc Natl Acad Sci USA* 92:6537–6541

58. Vaughan TJ, Williams AJ, Pritchard K et al (1996) Human antibodies with sub-nanomolar affinities isolated from a large non-immunized phage display library. *Nat Biotechnol* 14:309–314
59. Lloyd C, Lowe D, Edwards B et al (2009) Modelling the human immune response: performance of a 1011 human antibody repertoire against a broad panel of therapeutically relevant antigens. *Protein Eng Des Sel* 22:159–168
60. Sheets MD, Amersdorfer P, Finnern R et al (1998) Efficient construction of a large non-immune phage antibody library: the production of high-affinity human single-chain antibodies to protein antigens. *Proc Natl Acad Sci USA* 95:6157–6162
61. De Haard HJ (2002) Construction of large naive Fab libraries. *Methods Mol Biol* 178:87–100
62. Nissim A, Hoogenboom HR, Tomlinson IM et al (1994) Antibody fragments from a ‘single pot’ phage display library as immunochemical reagents. *EMBO J* 13:692–698
63. de Kruif J, Boel E, Logtenberg T (1995) Selection and application of human single chain Fv antibody fragments from a semi-synthetic phage antibody display library with designed CDR3 regions. *J Mol Biol* 248: 97–105
64. Griffiths AD, Williams SC, Hartley O et al (1994) Isolation of high affinity human antibodies directly from large synthetic repertoires. *EMBO J* 13:3245–3260
65. Hoogenboom HR, Winter G (1992) By-passing immunisation. Human antibodies from synthetic repertoires of germline VH gene segments rearranged in vitro. *J Mol Biol* 227:381–388
66. Pini A, Viti F, Santucci A et al (1998) Design and use of a phage display library. Human antibodies with subnanomolar affinity against a marker of angiogenesis eluted from a two-dimensional gel. *J Biol Chem* 273: 21769–21776
67. Hoet RM, Cohen EH, Kent RB et al (2005) Generation of high-affinity human antibodies by combining donor-derived and synthetic complementarity-determining-region diversity. *Nat Biotechnol* 23:344–348
68. Lee CV, Liang WC, Dennis MS et al (2004) High-affinity human antibodies from phage-displayed synthetic Fab libraries with a single framework scaffold. *J Mol Biol* 340: 1073–1093
69. Söderlind E, Strandberg L, Jirholt P et al (2000) Recombining germline-derived CDR sequences for creating diverse single-framework antibody libraries. *Nat Biotechnol* 18:852–856
70. Rothe C, Urlinger S, Lohning C et al (2008) The human combinatorial antibody library HuCAL GOLD combines diversification of all six CDRs according to the natural immune system with a novel display method for efficient selection of high-affinity antibodies. *J Mol Biol* 376:1182–1200
71. Sidhu SS, Li B, Chen Y, Fellouse FA et al (2004) Phage-displayed antibody libraries of synthetic heavy chain complementarity determining regions. *J Mol Biol* 338:299–310
72. Fellouse FA, Wiesmann C, Sidhu SS (2004) Synthetic antibodies from a four-amino-acid code: a dominant role for tyrosine in antigen recognition. *Proc Natl Acad Sci USA* 101:12467–12472
73. Fellouse FA, Li B, Compaan DM, Peden AA et al (2005) Molecular recognition by a binary code. *J Mol Biol* 348:1153–1162
74. Russel M, Linderöth NA, Sali A (1997) Filamentous phage assembly: variation on a protein export theme. *Gene* 192:23–32
75. Iannolo G, Minenkova O, Petruzzelli R et al (1995) Modifying filamentous phage capsid: limits in the size of the major capsid protein. *J Mol Biol* 248:835–844
76. Kretzschmar T, Geiser M (1995) Evaluation of antibodies fused to minor coat protein III and major coat protein VIII of bacteriophage M13. *Gene* 155:61–65
77. Rondot S, Koch J, Breitling F et al (2001) A helper phage to improve single chain antibody presentation in phage display. *Nat Biotechnol* 19:75–78
78. Nelson AL, Dhimolea E, Reichert JM (2010) Development trends for human monoclonal antibody therapeutics. *Nat Rev Drug Discov* 9:767–774
79. Edwards BM, Barash SC, Main SH et al (2003) The remarkable flexibility of the human antibody repertoire; isolation of over one thousand different antibodies to a single protein, BLYS. *J Mol Biol* 334:103–118
80. Baker KP, Edwards BM, Main SH et al (2003) Generation and characterization of LymphoStat-B, a human monoclonal antibody that antagonizes the bioactivities of B lymphocyte stimulator. *Arthritis Rheum* 48:3253–3265
81. Lloyd C, Lowe D, Edwards B et al (2009) Modelling the human immune response: performance of a 1011 human antibody repertoire against a broad panel of therapeutically relevant antigens. *Protein Eng Des Sel* 22:159–168

82. Dong J, Demarest SJ, Sereno A et al (2010) Combination of two insulin-like growth factor-I receptor inhibitory antibodies targeting distinct epitopes leads to an enhanced antitumor response. *Mol Cancer Ther* 9: 2593–2604
83. Doern A, Cao X, Sereno A et al (2009) Characterization of inhibitory anti-insulin-like growth factor receptor antibodies with different epitope specificity and ligand-blocking properties: implications for mechanism of action in vivo. *J Biol Chem* 284:10254–10267
84. Dong J, Sereno A, Snyder WB et al (2011) Stable IgG-like bispecific antibodies directed toward the type I insulin-like growth factor receptor demonstrate enhanced ligand blockade and anti-tumor activity. *J Biol Chem* 286: 4703–4717
85. Baselga J, Gelmon KA, Verma S et al (2010) Phase II trial of pertuzumab and trastuzumab in patients with human epidermal growth factor receptor 2-positive metastatic breast cancer that progressed during prior trastuzumab therapy. *J Clin Oncol* 28:1138–1144
86. Wu Y, Cain-Hom C, Choy L et al (2010) Therapeutic antibody targeting of individual Notch receptors. *Nature* 464:1052–1057
87. Xie MH, Yuan J, Adams C et al (1997) Direct demonstration of MuSK involvement in acetylcholine receptor clustering through identification of agonist ScFv. *Nat Biotechnol* 15:768–771
88. Ellmark P, Andersson H, Abayneh S et al (2008) Identification of a strongly activating human anti-CD40 antibody that suppresses HIV type 1 infection. *AIDS Res Hum Retroviruses* 24:367–373
89. Dobson CL, Main S, Newton P et al (2009) Human monomeric antibody fragments to TRAIL-R1 and TRAIL-R2 that display potent in vitro agonism. *MAbs* 1:552–562
90. Eisenhardt SU, Schwarz M, Bassler N et al (2007) Subtractive single-chain antibody (scFv) phage-display: tailoring phage-display for high specificity against function-specific conformations of cell membrane molecules. *Nat Protoc* 2:3063–3073
91. Huie MA, Cheung MC, Muench MO et al (2001) Antibodies to human fetal erythroid cells from a nonimmune phage antibody library. *Proc Natl Acad Sci USA* 98:2682–2687
92. Noronha EJ, Wang X, Desai SA et al (1998) Limited diversity of human scFv fragments isolated by panning a synthetic phage-display scFv library with cultured human melanoma cells. *J Immunol* 161:2968–2976
93. Ridgway JB, Ng E, Kern JA et al (1999) Identification of a human anti-CD55 single-chain Fv by subtractive panning of a phage library using tumor and nontumor cell lines. *Cancer Res* 59:2718–2723
94. Van Ewijk W, de Kruif J, Germeraad WT et al (1997) Subtractive isolation of phage-displayed single-chain antibodies to thymic stromal cells by using intact thymic fragments. *Proc Natl Acad Sci USA* 94:3903–3908
95. Giordano RJ, Cardo-Vila M, Lahdenranta J et al (2001) Biopanning and rapid analysis of selective interactive ligands. *Nat Med* 7: 1249–1253
96. Williams BR, Sharon J (2002) Polyclonal anti-colorectal cancer Fab phage display library selected in one round using density gradient centrifugation to separate antigen-bound and free phage. *Immunol Lett* 81:141–148
97. Osbourn JK, Derbyshire EJ, Vaughan TJ et al (1998) Pathfinder selection: in situ isolation of novel antibodies. *Immunotechnology* 3:293–302
98. Osbourn JK, Earnshaw JC, Johnson KS et al (1998) Directed selection of MIP-1 alpha neutralizing CCR5 antibodies from a phage display human antibody library. *Nat Biotechnol* 16:778–781
99. Poul MA, Becerril B, Nielsen UB et al (2000) Selection of tumor-specific internalizing human antibodies from phage libraries. *J Mol Biol* 301:1149–1161
100. Zhou Y, Marks JD (2009) Identification of target and function specific antibodies for effective drug delivery. *Methods Mol Biol* 525:145–160
101. Crépin R, Goenaga AL, Jullienne B et al (2010) Development of human single-chain antibodies to the transferrin receptor t, that effectively antagonize the growth of leukemias and lymphomas. *Cancer Res* 70: 5497–5506
102. Park JW, Kirpotin DB, Hong K et al (2001) Tumor targeting using anti-her2 immunoliposomes. *J Control Release* 74:95–113
103. Nielsen UB, Kirpotin DB, Pickering EM et al (2002) Therapeutic efficacy of anti-ErbB2 immunoliposomes targeted by a phage antibody selected for cellular endocytosis. *Biochim Biophys Acta* 1591:109–118
104. Velappan N, Martinez JS, Valero R et al (2007) Selection and characterization of scFv antibodies against the Sin Nombre hantavirus nucleocapsid protein. *J Immunol Methods* 321:60–69

105. Cabezas S, Rojas G, Pavon A et al (2009) Phage-displayed antibody fragments recognizing dengue 3 and dengue 4 viruses as tools for viral serotyping in sera from infected individuals. *Arch Virol* 154:1035–1045
106. Cabezas S, Rojas G, Pavon A et al (2008) Selection of phage-displayed human antibody fragments on Dengue virus particles captured by a monoclonal antibody: application to the four serotypes. *J Virol Methods* 147: 235–243
107. Lim AP, Chan CE, Wong SK et al (2008) Neutralizing human monoclonal antibody against H5N1 influenza HA selected from a Fab-phage display library. *Virol J* 5:130
108. Okada J, Ohshima N, Kubota-Koketsu R et al (2010) Monoclonal antibodies in man that neutralized H3N2 influenza viruses were classified into three groups with distinct strain specificity: 1968–1973, 1977–1993 and 1997–2003. *Virology* 397:322–330
109. Meissner F, Maruyama T, Frentsch M et al (2002) Detection of antibodies against the four subtypes of ebola virus in sera from any species using a novel antibody-phage indicator assay. *Virology* 300:236–243
110. Kirsch MI, Hülseweh B, Nacke C et al (2008) Development of human antibody fragments using antibody phage display for the detection and diagnosis of Venezuelan equine encephalitis virus (VEEV). *BMC Biotechnol* 8:66
111. Hayhurst A, Happe S, Mabry R et al (2003) Isolation and expression of recombinant antibody fragments to the biological warfare pathogen *Brucella melitensis*. *J Immunol Methods* 276:185–196
112. Zou N, Newsome T, Li B et al (2007) Human single-chain Fv antibodies against *Burkholderia mallei* and *Burkholderia pseudomallei*. *Exp Biol Med* (Maywood) 232:550–556
113. Maynard JA, Maassen CB, Leppla SH et al (2002) Protection against anthrax toxin by recombinant antibody fragments correlates with antigen affinity. *Nat Biotechnol* 20:597–601
114. Wild MA, Xin H, Maruyama T et al (2003) Human antibodies from immunized donors are protective against anthrax toxin in vivo. *Nat Biotechnol* 21:1305–1306
115. Steiniger SC, Altobelli LJ 3rd, Zhou B, Janda KD (2007) Selection of human antibodies against cell surface-associated oligomeric anthrax protective antigen. *Mol Immunol* 44:2749–2755
116. Pelat T, Hust M, Laffly E et al (2007) High-affinity, human antibody-like antibody fragment (single-chain variable fragment) neutralizing the lethal factor (LF) of *Bacillus anthracis* by inhibiting protective antigen-LF complex formation. *Antimicrob Agents Chemother* 51:2758–2764
117. Cirino NM, Sblattero D, Allen D et al (1999) Disruption of anthrax toxin binding with the use of human antibodies and competitive inhibitors. *Infect Immun* 67:2957–2963
118. Zhou B, Wirsching P, Janda KD (2002) Human antibodies against spores of the genus *Bacillus*: a model study for detection of and protection against anthrax and the bioterrorist threat. *Proc Natl Acad Sci USA* 99:5241–5246
119. Sui J, Hwang WC, Perez S et al (2009) Structural and functional bases for broad-spectrum neutralization of avian and human influenza A viruses. *Nat Struct Mol Biol* 16:265–273
120. Sun L, Lu X, Li C et al (2009) Generation, characterization and epitope mapping of two neutralizing and protective human recombinant antibodies against influenza A H5N1 viruses. *PLoS One* 4:e5476
121. Throsby M, van den Brink E, Jongeneelen M et al (2008) Heterosubtypic neutralizing monoclonal antibodies cross-protective against H5N1 and H1N1 recovered from human IgM+ memory B cells. *PLoS One* 3:e3942
122. Ayris J, Woods T, Bradbury A, Pavlik P (2007) High-throughput screening of single-chain antibodies using multiplexed flow cytometry. *J Proteome Res* 6:1072–1082
123. Pershad K, Pavlovic JD, Gräslund S et al (2010) Generating a panel of highly specific antibodies to 20 human SH2 domains by phage display. *Protein Eng Des Sel* 23:279–288
124. Parsons HL, Earnshaw JC, Wilton J et al (1996) Directing phage selections towards specific epitopes. *Protein Eng* 9:1043–1049
125. Mutuberria R, Satijn S, Huijbers A et al (2004) Isolation of human antibodies to tumor-associated endothelial cell markers by in vitro human endothelial cell selection with phage display libraries. *J Immunol Methods* 287:31–47
126. Cohen CJ, Denkberg G, Lev A et al (2003) Recombinant antibodies with MHC-restricted, peptide-specific, T-cell receptor-like specificity: new tools to study antigen presentation and TCR-peptide-MHC interactions. *J Mol Recognit* 16:324–332
127. Engberg J, Krogsgaard M, Fugger L (1999) Recombinant antibodies with the

- antigen-specific, MHC restricted specificity of T cells: novel reagents for basic and clinical investigations and immunotherapy. *Immunotechnology* 4:273–278
128. Stryhn A, Andersen PS, Pedersen LO et al (1996) Shared fine specificity between T-cell receptors and an antibody recognizing a peptide/major histocompatibility class I complex. *Proc Natl Acad Sci USA* 93:10338–10342
 129. Villa A, Trachsel E, Kaspar M et al (2008) A high-affinity human monoclonal antibody specific to the alternatively spliced ED α domain of fibronectin efficiently targets tumor neo-vasculature in vivo. *Int J Cancer* 122:2405–2413
 130. Pini A, Viti F, Santucci A et al (1998) Design and use of a phage display library. Human antibodies with subnanomolar affinity against a marker of angiogenesis eluted from a two-dimensional gel. *J Biol Chem* 273:21769–21776
 131. Schliemann C, Neri D (2010) Antibody-based vascular tumor targeting. *Recent Results Cancer Res* 180:201–216
 132. Nizak C, Monier S, del Nery E et al (2003) Recombinant antibodies to the small GTPase Rab6 as conformation sensors. *Science* 300:984–987
 133. Gao J, Sidhu SS, Wells JA (2009) Two-state selection of conformation-specific antibodies. *Proc Natl Acad Sci USA* 106:3071–3076
 134. Eisenhardt SU, Schwarz M, Bassler N, Peter K (2007) Subtractive single-chain antibody (scFv) phage-display: tailoring phage-display for high specificity against function-specific conformations of cell membrane molecules. *Nat Protoc* 2:3063–3073
 135. Rothlisberger D, Pos KM, Pluckthun A (2004) An antibody library for stabilizing and crystallizing membrane proteins - selecting binders to the citrate carrier CitS. *FEBS Lett* 564:340–348
 136. Uysal S, Vásquez V, Tereshko V et al (2009) Crystal structure of full-length KcsA in its closed conformation. *Proc Natl Acad Sci USA* 106:6644–6649
 137. Ye JD, Tereshko V, Frederiksen JK et al (2008) Synthetic antibodies for specific recognition and crystallization of structured RNA. *Proc Natl Acad Sci USA* 105:82–87
 138. Koldobskaya Y, Duguid EM, Shechner DM et al (2010) A portable RNA sequence whose recognition by a synthetic antibody facilitates structural determination. *Nat Struct Mol Biol* 18:100–106
 139. Monigatti F, Gasteiger E, Bairoch A, Jung E (2002) The Sulfinator: predicting tyrosine sulfation sites in protein sequences. *Bioinformatics* 18:769–770
 140. Kehoe JW, Velappan N, Walbolt M et al (2006) Using phage display to select antibodies recognizing post-translational modifications independently of sequence context. *Mol Cell Proteomics* 5:2350–2363
 141. Thie H, Voedisch B, Dübel S et al (2009) Affinity maturation by phage display. *Methods Mol Biol* 525:309–322
 142. Cadwell RC, Joyce GF (1994) Mutagenic PCR. *PCR Methods Appl* 3:S136–S140
 143. Stemmer WP (1994) DNA shuffling by random fragmentation and reassembly: *in vitro* recombination for molecular evolution. *Proc Natl Acad Sci USA* 91:10747–10751
 144. Fellouse FA, Wiesmann C, Sidhu SS (2004) Synthetic antibodies from a four-amino-acid code: a dominant role for tyrosine in antigen recognition. *Proc Natl Acad Sci USA* 101:12467–12472
 145. Liang WC, Wu X, Peale FV et al (2006) Cross-species vascular endothelial growth factor (VEGF)-blocking antibodies completely inhibit the growth of human tumor xenografts and measure the contribution of stromal VEGF. *J Biol Chem* 281:951–961
 146. Lee CV, Hymowitz SG, Wallweber HJ et al (2006) Synthetic anti-BR3 antibodies that mimic BAFF binding and target both human and murine B cells. *Blood* 108:3103–3111
 147. Fagète S, Ravn U, Gueneau F et al (2009) Specificity tuning of antibody fragments to neutralize two human chemokines with a single agent. *MAbs* 1:288–296
 148. Bostrom J, Yu SF, Kan D et al (2009) Variants of the antibody herceptin that interact with HER2 and VEGF at the antigen binding site. *Science* 323:1610–1614
 149. Volk WA, Bizzini B, Snyder RM et al (1984) Neutralization of tetanus toxin by distinct monoclonal antibodies binding to multiple epitopes on the toxin molecule. *Infect Immun* 45:604–609
 150. Zwick MB, Labrijn AF, Wang M et al (2001) Broadly neutralizing antibodies targeted to the membrane-proximal external region of human immunodeficiency virus type 1 glycoprotein gp41. *J Virol* 75:10892–10905
 151. Cheson BD, Leonard JP (2008) Monoclonal antibody therapy for B-cell non-Hodgkin's lymphoma. *N Eng J Med* 359:613–626
 152. de Kruif J, Logtenberg T (1996) Leucine zipper dimerized bivalent and bispecific scFv antibodies from a semi-synthetic antibody phage display library. *J Biol Chem* 271:7630–7634

153. Thie H, Binius S, Schirrmann T et al (2009) Multimerization domains for antibody phage display and antibody production. *New Biotechnol* 26:314–321
154. Hudson PJ, Kortt AA (1999) High avidity scFv multimers; diabodies and triabodies. *J Immunol Methods* 231:177–189
155. Dübel S, Breitling F, Kontermann R et al (1995) Bifunctional and multimeric complexes of streptavidin fused to single chain antibodies (scFv). *J Immunol Methods* 178:201–209
156. Huang D, Shusta EV (2006) A yeast platform for the production of single-chain antibody-green fluorescent protein fusions. *Appl Environ Microbiol* 72:7748–7759
157. Hink MA, Griep RA, Borst JW et al (2000) Structural dynamics of green fluorescent protein alone and fused with a single chain Fv protein. *J Biol Chem* 275:17556–17560
158. Casey JL, Coley AM, Tilley LM et al (2000) Green fluorescent antibodies: novel in vitro tools. *Protein Eng* 13:445–452
159. Griep RA, van Twisk C, Kerschbaumer RJ et al (1999) pSKAP/S: an expression vector for the production of single-chain Fv alkaline phosphatase fusion proteins. *Protein Expr Purif* 16:63–69
160. Al-Mrabeh A, Ziegler A, Cowan G et al (2009) A fully recombinant ELISA using in vivo biotinylated antibody fragments for the detection of potato leafroll virus. *J Virol Methods* 159:200–205
161. Predonzani A, Arnoldi F, Lopez-Requena A et al (2008) In vivo site-specific biotinylation of proteins within the secretory pathway using a single vector system. *BMC Biotechnol* 8:41
162. Warren DJ, Bjerner J, Paus E et al (2005) Use of an in vivo biotinylated single-chain antibody as capture reagent in an immunometric assay to decrease the incidence of interference from heterophilic antibodies. *Clin Chem* 51: 830–838
163. Cloutier SM, Couty S, Tersikh A et al (2000) Streptabody, a high avidity molecule made by tetramerization of in vivo biotinylated, phage display-selected scFv fragments on streptavidin. *Mol Immunol* 37:1067–1077
164. Moutel S et al (2009) A multi-Fc-species system for recombinant antibody production. *BMC Biotechnol* 9:14
165. Hu S, Shively L, Raubitschek A et al (1996) Minibody: a novel engineered anti-carcino-embryonic antigen antibody fragment (single-chain Fv-CH3) which exhibits rapid, high-level targeting of xenografts. *Cancer Res* 56:3055–3061
166. Gilliland LK, Norris NA, Marquardt H et al (1996) Rapid and reliable cloning of antibody variable regions and generation of recombinant single chain antibody fragments. *Tissue Antigens* 47:1–20
167. Shan D, Press OW, Tsu TT et al (1999) Characterization of scFv-Ig constructs generated from the anti-CD20 mAb 1F5 using linker peptides of varying lengths. *J Immunol* 162:6589–6595
168. Kontermann RE (2010) Alternative antibody formats. *Curr Opin Mol Ther* 12:176–183
169. Merchant AM, Zhu Z, Yuan JQ et al (1998) An efficient route to human bispecific IgG. *Nat Biotechnol* 16:677–681
170. Ridgway JB, Presta LG et al (1996) ‘Knobs-into-holes’ engineering of antibody CH3 domains for heavy chain heterodimerization. *Protein Eng* 9:617–621
171. Coloma MJ, Morrison SL (1997) Design and production of novel tetravalent bispecific antibodies. *Nat Biotechnol* 15:159–163
172. Dong J, Sereno A, Snyder WB et al (2011) Stable IgG-like bispecific antibodies directed toward the type I insulin-like growth factor receptor demonstrate enhanced ligand blockade and anti-tumor activity. *J Biol Chem* 286:4703–4717
173. Kortt AA, Dolezal O, Power BE et al (2001) Dimeric and trimeric antibodies: high avidity scFvs for cancer targeting. *Biomol Eng* 18: 95–108
174. Lawrence LJ, Kortt AA, Iliades P et al (1998) Orientation of antigen binding sites in dimeric and trimeric single chain Fv antibody fragments. *FEBS Lett* 425:479–484
175. Perisic O, Webb PA, Holliger P et al (1994) Crystal structure of a diabody, a bivalent antibody fragment. *Structure* 2:1217–1226
176. Atwell JL, Breheney KA, Lawrence LJ et al (1999) scFv multimers of the anti-neuraminidase antibody NC10: length of the linker between VH and VL domains dictates precisely the transition between diabodies and triabodies. *Protein Eng* 12: 597–604
177. Pei XY, Holliger P, Murzin AG et al (1997) The 2.0-Å resolution crystal structure of a trimeric antibody fragment with noncognate VH-VL domain pairs shows a rearrangement of VH CDR3. *Proc Natl Acad Sci USA* 94:9637–9642
178. Le Gall F, Kiprianov SM, Moldenhauer G et al (1999) Di-, tri- and tetrameric single chain Fv antibody fragments against human CD19: effect of valency on cell binding. *FEBS Lett* 453:164–168

179. Muller D, Kontermann RE (2010) Bispecific antibodies for cancer immunotherapy: current perspectives. *BioDrugs* 24:89–98
180. Beck A, Wagner-Rousset E, Bussat MC et al (2008) Trends in glycosylation, glycoanalysis and glycoengineering of therapeutic antibodies and Fc-fusion proteins. *Curr Pharm Biotechnol* 9:482–501
181. Presta LG (2008) Molecular engineering and design of therapeutic antibodies. *Curr Opin Immunol* 20:460–470
182. Hallborn J, Carlsson R (2002) Automated screening procedure for high-throughput generation of antibody fragments. *Biotechniques Suppl*, 30–37
183. Lou J, Marzari R, Verzillo V et al (2001) Antibodies in haystacks: how selection strategy influences the outcome of selection from molecular diversity libraries. *J Immunol Methods* 253:233–242
184. Kawe M, Forrer P, Amstutz P et al (2006) Isolation of intracellular proteinase inhibitors derived from designed ankyrin repeat proteins by genetic screening. *J Biol Chem* 281: 40252–40263
185. Glanville J, Zhai W, Berka J et al (2009) Precise determination of the diversity of a combinatorial antibody library gives insight into the human immunoglobulin repertoire. *Proc Natl Acad Sci USA* 106:20216–20221
186. Ravn U, Gueneau F, Baerlocher L et al (2010) By-passing in vitro screening-next generation sequencing technologies applied to antibody display and in silico candidate selection. *Nucleic Acids Res* 38:e193
187. Ge X, Mazor Y, Hunicke-Smith SP et al (2010) Rapid construction and characterization of synthetic antibody libraries without DNA amplification. *Biotechnol Bioeng* 106: 347–357
188. Fischer N (2011) Sequencing antibody repertoires: the next generation. *MAbs* 3: 17–20
189. Reddy ST, Ge X, Miklos AE et al (2010) Monoclonal antibodies isolated without screening by analyzing the variable-gene repertoire of plasma cells. *Nat Biotechnol* 28: 965–969
190. Zhang H, Torkamani A, Jones TM et al (2011) Phenotype-information-phenotype cycle for deconvolution of combinatorial antibody libraries selected against complex systems. *Proc Natl Acad Sci USA* 108: 13456–13461
191. Bordeaux J, Welsh A, Agarwal S et al (2010) Antibody validation. *Biotechniques* 48: 197–209
192. Pozner-Moulis S, Cregger M, Camp RL et al (2007) Antibody validation by quantitative analysis of protein expression using expression of Met in breast cancer as a model. *Lab Invest* 87:251–260
193. Grimsey NL, Goodfellow CE, Scotter EL et al (2008) Specific detection of CB1 receptors; cannabinoid CB1 receptor antibodies are not all created equal! *J Neurosci Methods* 171:78–86
194. Saper CB (2005) An open letter to our readers on the use of antibodies. *J Comp Neurol* 493:477–478



<http://www.springer.com/978-1-61779-930-3>

Antibody Methods and Protocols
Proetzel, G.; Ebersbach, H. (Eds.)
2012, X, 325 p., Hardcover
ISBN: 978-1-61779-930-3
A product of Humana Press