

Contents

1	Introduction	1
	References	6

Part I Total Synthesis of Pachastrissamine (Jaspine B)

2	Total Synthesis through Palladium-Catalyzed Bis-Cyclization of Bromoallenes	11
2.1	Experimental Section	17
2.1.1	General Methods	17
2.1.2	<i>tert</i> -Butyl (<i>R</i>)-4-[(<i>R</i>)-3-Bromopropa-1,2-dienyl]-2,2-dimethyloxazolidine-3-carboxylate (12a).	18
2.1.3	<i>N</i> -[(2 <i>R</i> ,4 <i>R</i>)-5-Bromo-1-hydroxypenta-3,4-dien-2-yl]-benzamide (9a)	18
2.1.4	(3 <i>aS</i> ,6 <i>aS</i>)-6-Methylene-2-phenyl-3 <i>a</i> ,4,6,6 <i>a</i> -tetrahydrofuro[3,4- <i>d</i>]oxazole (8)	19
2.1.5	(<i>R</i>)- <i>N</i> -[5-(Methoxymethyl)-2,3-dihydrofuran-3-yl]-benzamide (13a)	19
2.1.6	(<i>R</i>)- <i>N</i> -{5-[(2,2,2-Trifluoroethoxy)methyl]-2,3-dihydrofuran-3-yl}benzamide (13b)	19
2.1.7	<i>N</i> -(5-Methylfuran-3-yl)benzamide (14)	20
2.1.8	<i>tert</i> -Butyl (<i>R</i>)-4-[(<i>S</i>)-3-Bromopropa-1,2-dienyl]-2,2-dimethyloxazolidine-3-carboxylate (12b).	20
2.1.9	<i>N</i> -[(2 <i>R</i> ,4 <i>S</i>)-5-Bromo-1-hydroxypenta-3,4-dien-2-yl]-benzamide (9b)	20
2.1.10	[(3 <i>aS</i> ,6 <i>S</i> ,6 <i>aS</i>)-2-Phenyl-3 <i>a</i> ,4,6,6 <i>a</i> -tetrahydrofuro[3,4- <i>d</i>]oxazol-6-yl]methanol (15)	21
2.1.11	(3 <i>aS</i> ,6 <i>S</i> ,6 <i>aS</i>)-2-Phenyl-6-tetradecyl-3 <i>a</i> ,4,6,6 <i>a</i> -tetrahydrofuro[3,4- <i>d</i>]oxazole (16)	21

2.1.12	(3a <i>S</i> ,6 <i>S</i> ,6a <i>S</i>)-2-Phenyl-6-(3-phenylpropyl)-3a,4,6,6a-tetrahydrofuro[3,4- <i>d</i>]oxazole (17a)	22
2.1.13	(3a <i>S</i> ,6 <i>S</i> ,6a <i>S</i>)-6-Ethyl-2-phenyl-3a,4,6,6a-tetrahydrofuro[3,4- <i>d</i>]oxazole (17b).	23
2.1.14	(3a <i>S</i> ,6 <i>S</i> ,6a <i>S</i>)-6-Isobutyl-2-phenyl-3a,4,6,6a-tetrahydrofuro[3,4- <i>d</i>]oxazole (17c).	23
2.1.15	(3a <i>S</i> ,6 <i>S</i> ,6a <i>S</i>)-6-(2-Methylallyl)-2-phenyl-3a,4,6,6a-tetrahydrofuro[3,4- <i>d</i>]oxazole (17d).	24
2.1.16	(1 <i>S</i> ,4 <i>S</i> ,7 <i>S</i>)-5-(4-Phenylhepta-1,6-dien-4-yl)-2-oxa-5-azabicyclo[2.2.1]heptan-7-ol (18).	24
2.1.17	(3a <i>S</i> ,6 <i>S</i> ,6a <i>S</i>)-6-(But-3-enyl)-2-phenyl-3a,4,6,6a-tetrahydrofuro[3,4- <i>d</i>]oxazole 42 (17e)	25
References	26

3 Total Synthesis through Palladium-Catalyzed Bis-Cyclization of Propargyl Chlorides and Carbonates

3.1	Experimental Section	38
3.1.1	General Methods	38
3.1.2	<i>tert</i> -Butyl (<i>S</i>)-4-[(<i>S</i>)-1-Hydroxyhexadec-2-yn-1-yl]-2,2-dimethyloxazolidine-3-carboxylate (<i>syn</i> -6)	39
3.1.3	<i>tert</i> -Butyl (<i>S</i>)-4-[(<i>R</i>)-1-Hydroxyhexadec-2-yn-1-yl]-2,2-dimethyloxazolidine-3-carboxylate (<i>anti</i> -6)	39
3.1.4	<i>tert</i> -Butyl (<i>S</i>)-4-[(<i>S</i>)-1-(Methoxycarbonyloxy)hexadec-2-yn-1-yl]-2,2-dimethyloxazolidine-3-carboxylate (<i>syn</i> -7)	40
3.1.5	<i>tert</i> -Butyl (<i>S</i>)-4-[(<i>R</i>)-1-(Methoxycarbonyloxy)hexadec-2-yn-1-yl]-2,2-dimethyloxazolidine-3-carboxylate (<i>anti</i> -7).	40
3.1.6	(2 <i>S</i> ,3 <i>S</i>)-2-Benzamido-1-hydroxyoctadec-4-yn-3-yl Methyl Carbonate (<i>syn</i> -8)	41
3.1.7	(2 <i>S</i> ,3 <i>R</i>)-2-Benzamido-1-hydroxyoctadec-4-yn-3-yl Methyl Carbonate (<i>anti</i> -8)	41
3.1.8	<i>tert</i> -Butyl (<i>S</i>)-4-[(<i>R</i>)-1-Chlorohexadec-2-yn-1-yl]-2,2-dimethyloxazolidine-3-carboxylate (<i>syn</i> -9)	42
3.1.9	<i>tert</i> -Butyl (<i>S</i>)-4-[(<i>S</i>)-1-Chlorohexadec-2-yn-1-yl]-2,2-dimethyloxazolidine-3-carboxylate (<i>anti</i> -9)	42
3.1.10	<i>N</i> -[(2 <i>S</i> ,3 <i>R</i>)-3-Chloro-1-hydroxyoctadec-4-yn-2-yl]benzamide (<i>syn</i> -12)	43
3.1.11	<i>N</i> -[(2 <i>S</i> ,3 <i>S</i>)-3-Chloro-1-hydroxyoctadec-4-yn-2-yl]benzamide (<i>anti</i> -12).	43
3.1.12	[(4 <i>S</i> ,5 <i>R</i>)-5-(Pentadec-1-yn-1-yl)-2-phenyl-4,5-dihydrooxazol-4-yl]methanol (<i>cis</i> -13) and [(2 <i>R</i> ,3 <i>R</i>)-2-(Hydroxymethyl)-3-(pentadec-1-yn-1-yl)aziridin-1-yl](phenyl)-methanone (14)	44

3.1.13	[(4 <i>S</i> ,5 <i>S</i>)-5-(Pentadec-1-ynyl)-2-phenyl-4,5-dihydrooxazol-4-yl]methanol (<i>trans</i> -13)	44
3.1.14	Synthesis of the Authentic Sample of <i>cis</i> -13 from the Known Compound <i>syn</i> -6.	45
3.1.15	Synthesis of the Authentic Sample of <i>trans</i> -13 from the Known Compound <i>anti</i> -6	45
3.1.16	General Procedure for Palladium-Catalyzed Cascade Cyclization of Propargyl Chlorides: Synthesis of (3 <i>aS</i> ,6 <i>aS</i> , <i>E</i>)-2-Phenyl-6-tetradecylidene-3 <i>a</i> ,4,6,6 <i>a</i> -tetrahydrofuro[3,4- <i>d</i>]oxazole ((<i>E</i>)-2) from <i>syn</i> -12.	46
3.1.17	General Procedure for Palladium-Catalyzed Cascade Cyclization of Propargyl Carbonates: Synthesis of (3 <i>aS</i> ,6 <i>aS</i> , <i>E</i>)-2-Phenyl-6-tetradecylidene-3 <i>a</i> ,4,6,6 <i>a</i> -tetrahydrofuro[3,4- <i>d</i>]oxazole ((<i>E</i>)-2) from <i>syn</i> -8.	46
3.1.18	(3 <i>aS</i> ,6 <i>aS</i> , <i>Z</i>)-2-Phenyl-6-tetradecylidene-3 <i>a</i> ,4,6,6 <i>a</i> -tetrahydrofuro[3,4- <i>d</i>]oxazole ((<i>Z</i>)-2).	46
3.1.19	(3 <i>aS</i> ,6 <i>S</i> ,6 <i>aS</i>)-2-Phenyl-6-tetradecyl-3 <i>a</i> ,4,6,6 <i>a</i> -tetrahydrofuro[3,4- <i>d</i>]oxazole (15)	47
3.1.20	(2 <i>S</i> ,3 <i>S</i> ,4 <i>S</i>)-4-Amino-2-tetradecyltetrahydrofuran-3-ol [Pachastrissamine (1)]	47
3.1.21	(2 <i>S</i> ,3 <i>S</i> ,4 <i>S</i>)-4-(Benzylamino)-2-tetradecyltetrahydrofuran-3-ol (<i>N</i> -Benzylpachastrissamine) (16).	48
3.1.22	(2 <i>S</i> ,3 <i>S</i> ,4 <i>S</i>)-4-Amino-2-tetradecyltetrahydrofuran-3-ol (Pachastrissamine) (1) from 16	48
	References	48

Part II Total Synthesis of Lysergic Acid, Lysergol, and Isolysergol

4 Palladium-Catalyzed Domino Cyclization of Amino Allenes Bearing a Bromoindolyl Group and Its Application to Total

	Synthesis of Ergot Alkaloids.	53
4.1	Experimental Section	60
4.1.1	General Methods	60
4.1.2	4-Bromo-1-tosyl-1 <i>H</i> -indole-3-carbaldehyde (9)	60
4.1.3	4-Bromo-3-(bromomethyl)-1-tosyl-1 <i>H</i> -indole (10)	61
4.1.4	4-Bromo-1-tosyl-3-[2-(trimethylsilylethynyl)-1,3-dithian-2-yl]methyl-1 <i>H</i> -indole (12).	61
4.1.5	(±)-1-(4-Bromo-1-tosyl-1 <i>H</i> -indol-3-yl)-4-(trimethylsilyl)but-3-yn-2-ol (13).	62
4.1.6	Methyl (±)-(<i>E</i>)-3-[1-(4-Bromo-1-tosyl-1 <i>H</i> -indol-3-yl)-but-3-yn-2-yloxy]acrylate (14).	63
4.1.7	(±)-(<i>E</i>)-4-Bromo-1-tosyl-3-{2-[3-(triisopropylsilyloxy)prop-1-en-yloxy]but-3-ynyl}-1 <i>H</i> -indole (15).	63

4.1.8	(±)-(2 <i>S</i> , <i>aR</i>)-6-(4-Bromo-1-tosyl-1 <i>H</i> -indol-3-yl)-2-(triisopropylsilyloxymethyl)hexa-3,4-dien-1-ol (16a) and (±)-(2 <i>R</i> , <i>aR</i>)-Isomer (16b)	64
4.1.9	Determination of Relative Configuration of 16a: Synthesis of (±)-4-Bromo-1-tosyl-3-[[(2 <i>R</i> ,5 <i>S</i>)-5-(triisopropylsilyloxymethyl)-5,6-dihydro-2 <i>H</i> -pyran-2-yl]methyl]-1 <i>H</i> -indole (17)	65
4.1.10	<i>N</i> -[(2 <i>S</i> , <i>aR</i>)-6-(4-Bromo-1-tosyl-1 <i>H</i> -indol-3-yl)-2-(triisopropylsilyloxymethyl)hexa-3,4-dienyl]-2-nitrobenzenesulfonamide (18a) and Its (±)-(2 <i>R</i> , <i>aR</i>)-Isomer (18b)	66
4.1.11	(±)- <i>N</i> -[(2 <i>S</i> , <i>aR</i>)-6-(4-Bromo-1-tosyl-1 <i>H</i> -indol-3-yl)-2-(triisopropylsilyloxymethyl)-hexa-3,4-dienyl]-4-methylbenzenesulfonamide (19a) and Its (±)-(2 <i>R</i> , <i>aR</i>)-Isomer (19b)	67
4.1.12	(±)-(6 <i>aR</i> ,9 <i>S</i>)-7-(2-Nitrophenylsulfonyl)-4-tosyl-9-(triisopropylsilyloxymethyl)-4,6,6 <i>a</i> ,7,8,9-hexahydroindolo[4,3- <i>fg</i>]quinoline (20a) and Its (±)-(6 <i>aS</i> ,9 <i>S</i>)-Isomer (20b) (Table 2, Entry 3)	68
4.1.13	(±)-(6 <i>aR</i> ,9 <i>S</i>)-4,7-Ditosyl-9-(triisopropylsilyloxymethyl)-4,6,6 <i>a</i> ,7,8,9-hexahydroindolo[4,3- <i>fg</i>]quinoline (21a) and Its (±)-(6 <i>aS</i> ,9 <i>S</i>)-Isomer (21b) (Table 4.2, Entry 12).	69
4.1.14	(±)-(6 <i>aR</i> ,9 <i>S</i>)-7-Methyl-4-tosyl-9-(triisopropylsilyloxymethyl)-4,6,6 <i>a</i> ,7,8,9-hexahydroindolo[4,3- <i>fg</i>]quinoline (22a) and Its (±)-(6 <i>aS</i> ,9 <i>S</i>)-Isomer (22b)	70
4.1.15	Methyl (±)-(6 <i>aR</i> ,9 <i>S</i>)-4,7-ditosyl-4,6,6 <i>a</i> ,7,8,9-hexahydroindolo[4,3- <i>fg</i>]quinoline-9-carboxylate (23a).	72
4.1.16	(±)-Methyl Isolysergate (24a) and (±)-Methyl Lysergate (24b)	73
	References	74
5	Total Synthesis of (+)-Lysergic Acid, (+)-Lysergol, and (+)-Isolysergol	77
5.1	Experimental Section	88
5.1.1	General Methods	88
5.1.2	(<i>S</i>)- <i>N</i> -[2-(Hydroxymethyl)but-3-ynyl]-4-methylbenzenesulfonamide (11)	89
5.1.3	(2 <i>R</i> ,5 <i>S</i>)-5-Ethynyl-2-phenyl-3-tosyl-1,3-oxazinane (12)	89
5.1.4	(2 <i>R</i> ,5 <i>S</i>)-5-(Iodoethynyl)-2-phenyl-3-tosyl-1,3-oxazinane (13)	90

5.1.5	<i>S</i> -Ethyl 2-(4-Bromo-1 <i>H</i> -indol-3-yl)-ethanethioate (15)	90
5.1.6	<i>S</i> -Ethyl 2-(4-Bromo-1-tosyl-1 <i>H</i> -indol-3-yl)-ethanethioate (7)	91
5.1.7	3-Allyl-4-bromo-1 <i>H</i> -indole (18)	91
5.1.8	3-Allyl-4-bromo-1-tosyl-1 <i>H</i> -indole (19)	92
5.1.9	2-(4-Bromo-1-tosyl-1 <i>H</i> -indol-3-yl)acetaldehyde (8)	92
5.1.10	(<i>R</i>)-1-(4-Bromo-1-tosyl-1 <i>H</i> -indol-3-yl)-4-[(2 <i>R</i> ,5 <i>S</i>)-2-phenyl-3-tosyl-1,3-oxazinan-5-yl]but-3-yn-2-ol (20a) and Its (<i>S</i>)-Isomer (20b)	93
5.1.11	1-(4-Bromo-1-tosyl-1 <i>H</i> -indol-3-yl)-4-[(2 <i>R</i> ,5 <i>S</i>)-2-phenyl-3-tosyl-1,3-oxazinan-5-yl]-but-3-yn-2-one (16)	93
5.1.12	(<i>R</i>)-1-(4-Bromo-1-tosyl-1 <i>H</i> -indol-3-yl)-4-[(2 <i>R</i> ,5 <i>S</i>)-2-phenyl-3-tosyl-1,3-oxazinan-5-yl]but-3-yn-2-ol (20a) . . .	94
5.1.13	(2 <i>R</i> ,5 <i>S</i>)-5-[(<i>R</i>)-4-(4-Bromo-1-tosyl-1 <i>H</i> -indol-3-yl)buta-1,2-dienyl]-2-phenyl-3-tosyl-1,3-oxazinanane (21a)	94
5.1.14	<i>N</i> -[(2 <i>S</i> ,4 <i>R</i>)-6-(4-Bromo-1-tosyl-1 <i>H</i> -indol-3-yl)-2-(hydroxymethyl)hexa-3,4-dienyl]-4-methyl benzenesulfon-amide (5a)	95
5.1.15	Determination of Relative Configuration of the Allenamide 5a: Synthesis of the Authentic Sample (±)-5a by Desilylation of the Known Allenamide (±)-22a	96
5.1.16	(<i>S</i>)-1-(4-Bromo-1-tosyl-1 <i>H</i> -indol-3-yl)-4-[(2 <i>R</i> ,5 <i>S</i>)-2-phenyl-3-tosyl-1,3-oxazinan-5-yl]but-3-yn-2-ol (20b) . . .	96
5.1.17	(2 <i>R</i> ,5 <i>S</i>)-5-[(<i>S</i>)-4-(4-Bromo-1-tosyl-1 <i>H</i> -indol-3-yl)buta-1,2-dienyl]-2-phenyl-3-tosyl-1,3-oxazinanane (21b) . .	97
5.1.18	<i>N</i> -[(2 <i>S</i> ,4 <i>S</i>)-6-(4-Bromo-1-tosyl-1 <i>H</i> -indol-3-yl)-2-(hydroxymethyl)hexa-3,4-dienyl]-4-methyl benzenesulfon-amide (5b)	97
5.1.19	[(6 <i>aR</i> ,9 <i>S</i>)-4,7-Ditosyl-4,6,6 <i>a</i> ,7,8,9-hexahydroindolo [4,3- <i>fg</i>]quinolin-9-yl]methanol (4a)	98
5.1.20	Determination of Relative Configuration of the Alcohol 4a: Synthesis of the Authentic Sample (±)-4a by Desilylation of the Known Tetracyclic Indole (±)-23a	98
5.1.21	[(6 <i>aS</i> ,9 <i>S</i>)-4,7-Ditosyl-4,6,6 <i>a</i> ,7,8,9-hexahydroindolo [4,3- <i>fg</i>]quinolin-9-yl]methanol (4b)	99
5.1.22	(+)-Isolysergol (3)	99
5.1.23	Methyl (6 <i>aR</i> ,9 <i>S</i>)-4,7-Ditosyl-4,6,6 <i>a</i> ,7,8,9-hexahydroindolo[4,3- <i>fg</i>]quinoline-9-carboxylate (24a)	100

5.1.24	(+)-Methyl Isolysergate (25a) and (+)-Methyl Lysergate (25b)	100
5.1.25	(+)-Lysergol (2) and (+)-Isolysergol (3)	101
5.1.26	(+)-Lysergic Acid (1)	101
5.1.27	Determination of Optical Purity of Lysergic Acid (1)	102
References		102
6	Conclusions	105

<http://www.springer.com/978-4-431-54042-7>

Total Synthesis of Bioactive Natural Products by
Palladium-Catalyzed Domino Cyclization of Allenes and
Related Compounds

Inuki, S.

2012, XVI, 106 p., Hardcover

ISBN: 978-4-431-54042-7