

Chapter 2

PIP Kinases from the Cell Membrane to the Nucleus

Mark Schramp, Andrew Hedman, Weimin Li, Xiaojun Tan
and Richard Anderson

Abstract Phosphatidylinositol 4,5-bisphosphate (PIP₂) is a membrane bound lipid molecule with capabilities to affect a wide array of signaling pathways to regulate very different cellular processes. PIP₂ is used as a precursor to generate the second messengers PIP₃, DAG and IP₃, indispensable molecules for signaling events generated by membrane receptors. However, PIP₂ can also directly regulate a vast array of proteins and is emerging as a crucial messenger with the potential to distinctly modulate biological processes critical for both normal and pathogenic cell physiology. PIP₂ directly associates with effector proteins via unique phosphoinositide binding domains, altering their localization and/or enzymatic activity. The spatial and temporal generation of PIP₂ synthesized by the phosphatidylinositol phosphate kinases (PIPKs) tightly regulates the activation of receptor signaling pathways, endocytosis and vesicle trafficking, cell polarity, focal adhesion dynamics, actin assembly and 3' mRNA processing. Here we discuss our current understanding of PIPKs in the regulation of cellular processes from the plasma membrane to the nucleus.

Keywords Cell Migration · mRNA processing · Phosphatidylinositol phosphate kinase (PIPK) · Phosphatidylinositol 4,5-bisphosphate (PIP₂) · Vesicle trafficking

2.1 Introduction

Studies by Hokin and Hokin on exocrine tissue in the 1950s brought to light how changes in phospholipids could regulate cellular processes (Hokin and Hokin 1953). Later, discoveries in the 1980s advanced our understanding of how phosphatidyli-

R. Anderson (✉) · M. Schramp · W. Li
Department of Pharmacology, School of Medicine and Public Health,
3710 Medical Sciences Center, University of Wisconsin Medical School,
1300 University Ave., Madison, WI 53706
e-mail: raanders@wisc.edu

A. Hedman · X. Tan
Program in Molecular and Cellular Pharmacology
University of Wisconsin-Madison
1300 University Ave., Madison, WI 53706

inositol 4,5-bisphosphate (PIP₂) can be utilized by phospholipase C to generate IP₃ (Streb et al. 1983), a molecule that mobilizes Ca²⁺ stores from the endoplasmic reticulum, and by PI3K to generate PIP₃, a signaling molecule initially discovered downstream of the Src oncoprotein critical for cell proliferation and transformation (Whitman et al. 1988). In 1985, Lassing and Lindberg discovered that PIP₂ could directly associate with profilactin and profilin (Lassing and Lindberg 1985). These studies provided the backbone of PIP₂ biology and illuminate the importance of PIP₂ as a signaling nexus that not only is modified to generate PIP₃, DAG and IP₃, but can also directly modulate the activities of an ever growing array of proteins to regulate virtually every cellular process. The past few years have seen a resurgent interest in the regulation of PIP₂ synthesis. Unlike PIP₃, PIP₂ levels are relatively high and undergo only small changes in total cellular content upon stimulation (Heck et al. 2007). However, the spatial and temporal targeting of phosphatidyl-inositol phosphate kinases (PIPKs), the molecules that generate the vast majority of cellular PIP₂, via the actions of its functional protein associates can result in a dramatic and localized surge in PIP₂ levels to coordinate the activation of its regulated signaling pathways (Doughman et al. 2003b; Honda et al. 1999; Kisseleva et al. 2005; Sasaki et al. 2005). The findings that PIPK activity is enhanced by Rho and Arf family GTPases provided regulatory mechanisms and clues to the processes that PIP₂ generation might affect (Krauss et al. 2003; Weernink et al. 2004). Perhaps even more surprising, roles for PIPKs and PIP₂ within the nucleus are defining how these molecules can regulate mRNA processing and gene expression (Mellman et al. 2008). Thus how PIPKs are regulated has emerged as crucial yet relatively unexplored frontier in molecular biology that could dramatically impact how we view normal and pathogenic behavior. This chapter will focus on the role that PIPKs and PIP₂ generation play in modulating signaling pathways and cellular process from the plasma membrane to the nucleus.

2.2 The Enzymes. Sequence, Structure and Enzymology

2.2.1 *Sequence and Structure of Phosphatidylinositol Phosphate Kinases*

There are three known families of PIP kinases, type I, II and III, which catalyze the formation of phosphatidylinositol bisphosphates. Type I PIP kinases contain three separate genes, α , β and γ , and primarily catalyze phosphorylation of PI4P to PI4,5P₂ (Anderson et al. 1999; Doughman et al. 2003a; Heck et al. 2007; Ishihara et al. 1996; Jenkins et al. 1991; Loijens et al. 1996; Schill and Anderson 2009b). Type II PIP kinases, also have three separate genes, α , β and γ , and generate PI4,5P₂ by phosphorylating the 4' OH position of PI5P (Boronenkov and Anderson 1995; Rameh et al. 1997). First identified in yeast for their function in vacuole morphology, the most recently discovered type III PIP kinases generate PI3,5P₂ by phosphorylating PI3P (Cooke et al. 1998; Gary et al. 1998; McEwen et al. 1999; Yamamoto et al. 1995). Within this family of kinases are the yeast Fab1p and the human PikFYVE

(Cabezas et al. 2006; Sbrissa et al. 1999). At ~ 2200 amino acids the type III PIP Kinases are much larger in size than type I and II kinases and contain an N-terminal FYVE domain that allows for association with endosomal compartments (Gary et al. 1998; Odorizzi et al. 1998; Sbrissa et al. 1999; Yamamoto et al. 1995). The conserved feature of all PIP kinases is ~ 280 amino acid kinase domain (Ishihara et al. 1996).

2.2.1.1 Sequences of PIP Kinases

There is little conserved sequence among PIPK subtypes, with 60% identity among PIPKI, 60–77% identity among PIPKII, while PikFYVE and Fab1p only share 20% identity. Compared to the yeast MSS4p, human type I PIPKs share only 20% identity even though they catalyze the same reaction and can complement Mss4p deficient cells (Homma et al. 1998). Little identity is conserved between PIPK subtypes, PIPKI and PIPKII share less than 30% identity, while type PIPKIII shares less than 20% identity with human PIPKI and PIPKII. In the C-terminal region of type I, II and III PIPKs is the kinase homology domain, which is responsible for catalytic activity and is the only region that shares significant homology between family members (Anderson et al. 1999; Boronenkov and Anderson 1995; Ishihara et al. 1998; Loijens et al. 1996). PIPKI α , β and γ are approximately 75–80% identical, PIPKII α , β and γ are 66–78% identical, and 41% identity is shared between the PikFYVE and Fab1p kinase domains. There is $\sim 30\%$ shared identity in the kinase domain between family members.

While there is little overall sequence conservation among PIPKs, there are highly conserved regions within the kinase domain that are important for catalytic activity and substrate recognition. These essential components include the G-loop, catalytic residues and the activation loop. Deletion of regions N-terminal and C-terminal to the 280 amino acid kinase domain abrogate activity, indicating the importance of these regions in maintaining structural integrity (Ishihara et al. 1998). Starting from the N-terminus, PIPK kinase domains (highlighted in yellow) (Fig. 2.1) contain the consensus sequence of GxSGS in PIPKI and a conserved IIK, corresponding to a region found in protein kinases known as the G-loop, which consists of a glycine patch followed by a downstream lysine that mediates nucleotide binding (Anderson et al. 1999; Hanks et al. 1988; Heck et al. 2007; Ishihara et al. 1998; Rao et al. 1998; Saraste et al. 1990). Mutations in this region can inhibit kinase activity. Mutation of the murine PIPKI α Glycine 124 in the GxSGS consensus to Valine reduced kinase activity by one third, while mutating the downstream Lysine, K138, to Alanine completely abolished kinase activity (Ishihara et al. 1998). This Lysine in the IIK sequence corresponds with Lysine 72 in Protein Kinase A, which together with Mg²⁺ coordinates the α and β phosphates in ATP. Mutation of K72 also abolishes kinase activity of PKA and mutation of the corresponding residue abolishes activity of all PIP kinases (Anderson et al. 1999; Heck et al. 2007; Iyer et al. 2005; Knighton et al. 1991a, 1991b; Rao et al. 1998; Taylor et al. 1992, 1993).

C-terminal to the G-loop are two highly conserved regions responsible for PIPK catalytic activity (highlighted in magenta) (Fig. 2.1). These are the DLKGS and




Fig. 2.1 ClustalW multiple sequence alignment for human and yeast PIPK kinase domains. Conserved regions are highlighted—G-loop (yellow) for ATP binding, catalytic kinase domain sequences (Magenta), activation loop (cyan). Conserved residues (Red) PIPKIIβ K150, D216, D278 and D369 are required for PIP kinase activity

MDYSL motifs. The **DLKGS** motif shows similarity to the **HRDLK** catalytic sequence in PKA, but is not equivalent based on structure (Anderson et al. 1999; Rao et al. 1998; Yamamoto et al. 1995). However the conserved Aspartate in the **DLKGS** motif has been shown to be required for kinase activity in PIPKIIβ and PIPKIIγ (Anderson et al. 1999; Narkis et al. 2007). Furthermore, human D253 N mutations in PIPKIIγ that result in an **NLKGS** sequence lead to lethal contractual syndrome type 3 disease (Narkis et al. 2007). The PKA Aspartate 166 in the **HRDLK** motif functions as a catalytic base, corresponding to Aspartate 278 within the **MDYSL** motif in PIPKIIβ that superimposes on its structure (Krupa et al. 2004; Rao et al. 1998). Additionally, the catalytic Aspartate 369 in PIPKIIβ, within the (G/A)**IIDIL** motif, corresponds to Aspartate 184 in PKA found within a conserved **DFG** motif. This region coordinates Mg^{2+} ions and is required for kinase activity (Heck et al. 2007; Rao et al. 1998; Zheng et al. 1991). These four conserved residues (K138, D216, D278 and D369 in PIPKIIβ) in PIPKs are essential for kinase activity.

Finally, at the C-terminus of the kinase homology domain is a region of 20–30 amino acids (highlighted in cyan) (Fig. 2.1), known as the activation loop, responsible for substrate specificity and PIPK intracellular targeting (Anderson et al. 1999; Burden et al. 1999; Kunz et al. 2000, 2002; Rao et al. 1998). The activation loop is common amongst protein kinases (Cox and Taylor 1994; Hanks and Hunter 1995; Johnson et al. 1996; Rao et al. 1998) and its modification is one means to regulate these proteins. The activation loop of PIP kinases begins with the consensus **GIIDIL** with the aspartate conserved among other protein kinases, and ending with

an RF that is conserved in PIP kinases. In experiments where the activation loop of PIPKI β and PIPKII β were swapped, the specific activity of these chimeric PIP kinases matched its activation loop, for example, PIPKI β containing the II β activation loop phosphorylated PI5P, while chimeric PIPKII β containing the I β activation loop phosphorylated PI4P (Kunz et al. 2000). Additionally, these domain swaps altered PIPKI targeting (Kunz et al. 2000). Furthermore, specific residues conserved within PIPKI and PIPKII are essential for specificity. PIPKII β A381 to E point mutations alter substrate specificity from PI5P to PI4P, while the reverse mutation, E362A, in PIPKI β allowed for PIP₂ synthesis from PI4P and PI5P, demonstrating the importance of this Glutamate in PI4P specificity (Kunz et al. 2002). The role of the activation loop was further shown through complementation studies in yeast deficient in Mss4p. Yeast cells lacking Mss4p can be rescued by the PIPKII β chimera containing the I β specificity loop, thereby restoring type I activity, but not the PIPKI β chimera with the II β loop (Kunz et al. 2000). There is also a conserved KK motif in the activation loop found among PIPKs. Mutation of this KK to NN reduced substrate affinity, while a conservative KK to RR mutation did not alter function (Kunz et al. 2000). This suggests that this KK motif is important for lipid substrate interaction, but not for recognition of specific PIP isomers.

2.2.1.2 PIP Kinase Structure

Structures have been solved for human PIPKII β (PDB ID: 1BO1) and PIPKII γ (PDB ID: 2GK9), with publications focusing on PIPKII β (Burden et al. 1999; Kunz et al. 2000; Rao et al. 1998). The PIPKII β structure reveals a homodimer, with two PIPKII β monomers flush against one another. A set of antiparallel beta sheets and a single alpha helix form a flat face that allows for membrane association (Burden et al. 1999). Ten basic residues result in a net charge of +14 on this face; mutation of three lysines on this face to glutamates (K72/76/78) inhibited lipid binding and partially inhibited phosphoinositide binding, suggesting the basic face plays a role in membrane association but is not the only requirement for substrate binding (Burden et al. 1999). Figure 2.2a, b show that the planar face that interacts with the plasma membrane consists primarily of a beta sheet and a single helix. Individual monomers are shown in blue and green, with the basic residues found on the planar face, shown in red. Figure 2.2c shows the molecules from the side, with the planar face that interacts with the membrane facing downwards.

The conserved regions of the kinase domain described above are highlighted in Fig. 2.3. The G-loop (yellow) links two beta strands. The kinase catalytic DLKGS and MDYSL motifs (shown in magenta) are also in this pocket and the activation loop (shown in cyan) faces the membrane. The conserved catalytic residues within PIPKII β (K150, D216, D278 and D369) are highlighted in red. The structures of ATP and the PIPKII β substrate, PI5P, were modeled onto the structure (Fig. 2.3b) (Rao et al. 1998). This reveals a pocket, where the γ phosphate of ATP is oriented such that it faces the membrane, while PI5P fits such that the four hydroxyl of the myo-inositol ring faces this phosphate, consistent with PIPKII β function (Heck et al.




Fig. 2.2 PIPKII β structure. **a**, **b** and **c** Pymol rendering of the PIPKII β structure (PDB: 1BO1). PIPKII β forms a dimer (individual monomers in *blue* and *green*). **a** and **b** show the planar face that binds to the membrane, with key basic residues highlighted in *red*. **c** shows the side view of PIPKII β dimer, with membrane binding plane shown facing down

2007; Rao et al. 1998). Proposed residues necessary for ATP (133G, 136G, 139F, 148V, 150K, 201T, 203N, 204V, 205F, 278D, 282L, 368I, 369D) (Blue) and PI5P binding (134R, 218K, 224R, 239K, 278D, 372T) (Red) in the model are shown in Fig. 2.3b (Rao et al. 1998). The planar face of the dimer allows for the pocket to interact with the phosphoinositide head group and allow for catalytic activity without interfering with membrane structure.

2.2.2 Enzymology of PIPKs

The basic reaction catalyzed by PIP kinases is the ATP-dependent phosphorylation of phosphatidylinositol monophosphates to produce phosphatidylinositol bisphosphates (Heck et al. 2007). The substrate specificity of PIPKI, II and III was discovered through kinase activity on purified phosphatidylinositol phosphates from biological sources and synthetic phosphoinositides. These studies identified the primary substrate activity of PI4P for PIPKI, PI5P for PIPKII and PI3P for PIPKIII (Bazenet et al. 1990; Cabezas et al. 2006; Rameh et al. 1997; Sbrissa et al. 1999; Zhang et al. 1997). However, the PIP kinases can also utilize other PIP substrates with lower activity. For example, PIPKIs can phosphorylate PI3P to produce PI3,4P₂, and subsequently PI3,4,5P₃, or PI3,5P₂ (Tolias et al. 1998; Zhang et al. 1997). While the PIPKI affinity for PI3P is similar to PI4P, the reaction rate is higher for PI4P (Zhang et al. 1997). Similarly, PIPKII can also catalyze the phosphorylation of PI3P to PI3,4P₂ but is not the preferred substrate (Rameh et al. 1997). Finally, PIPKIII has been reported to generate PI5P from PI (Sbrissa et al. 1999; Shisheva et al. 1999). Though the enzymes can make multiple products, the *in vivo* importance of enzyme specificity can be observed through yeast complementation studies. Yeast deficient




Fig. 2.3 PIPKII β structure. **a** and **b** Pymol rendering of PIPKII β highlighting conserved regions—G loop (yellow), catalytic kinase domain sequences (magenta) and activation loop (cyan). Conserved residues required for PIP kinase activity are shown in red (PIPKII β K150, D216, D278 and D369). Highlighting residues believed to mediate ATP (blue) and PI5P (red) binding on PIPKII β structure

in MSS4p can be rescued using human PIPKI but not PIPKII (Homma et al. 1998), while yeast deficient in Fab1p can only be rescued by human PIPKIII (McEwen et al. 1999). However, the function of substrate diversity for each kinase *in vivo* has not been examined closely.

The activity of PIP kinases is regulated by a variety of stimuli, including lipids, proteins and post-translational modifications. Following their initial purification, phosphatidic acid was shown to enhance PIPKI kinase activity (Jenkins et al. 1994; Moritz et al. 1992a, b). More recently, phosphorylation has been shown to regulate PIPKs. Activation of Protein Kinase A (PKA) resulted in phosphorylation of PIPKI α on Serine 214 in the kinase domain, reducing activity, and activation of PKC via LPA treatment reduced PIPKI α phosphorylation and enhanced kinase activity (Park et al. 2001). Also, PKD can phosphorylate the activation loop of PIPKII α at Thr 376, and mutation of this site reduced kinase activity (Hinchliffe and Irvine 2006). Phosphorylation of PIPKs does not always affect kinase activity as Casein Kinase (CK) II phosphorylation of PIPKII α at Ser304 affects localization, but not kinase activity (Hinchliffe et al. 1999a, b). PIPKI and PIPKIII also have protein kinase activity, as these enzymes can auto-phosphorylate, reducing enzyme activity (Itoh et al. 2000; Sbrissa et al. 2000). Protein-protein interactions are also critical regulators




Fig. 2.4 PIPKs are recruited to intracellular compartments by PIP₂-effector proteins. *1* PIPKs interact with proteins that mediate their localization within the cell. *2* Once at a specific site, PIPKs generate PIP₂. *3* The interacting proteins that targeted PIPKs to specific sites are then modulated by the newly synthesized PIP₂. *4* The activated PIPK effectors regulate various cellular processes

of PIP kinase activity. Interactions with GTPases and specific proteins such as Talin and adaptor protein (AP) complexes have been shown to enhance PIPK activity (Di Paolo et al. 2002; Krauss et al. 2006; Weernink et al. 2004). The roles of phosphorylation and protein-protein interactors in regulating PIPK activity will be described in detail later.

The unique sequences in PIPKs allow for specific interactions with proteins that serve to target PIPKs to specific compartments. Often, these same interacting proteins are PIP₂ effectors (Heck et al. 2007). These protein-protein interactions target PIP kinases to specific compartments to generate a local pool of PIP₂, which can then regulate the activity of its effectors. This hypothesis is illustrated in Fig. 2.4. Multiple examples of regulation of PIP kinase targeted PIP₂ production have been demonstrated with diverse roles in cell-adhesion, migration, protein trafficking and nuclear signaling.

2.3 Membrane Associated PIPKs Drive Cell Migration and Vesicle Trafficking

2.3.1 PIPKs Help Regulate Directional Migration

Cell migration is a critical process for normal development, wound healing, cell survival and immunological responses, but can also have deleterious effects as evidenced by its role seen in metastatic tumor formation. Cell migration requires a tight

coordination of many molecular biological processes, including the establishment of cell polarity, the organized formation and turnover of adhesive structures and a regulated control of dynamic cytoskeletal rearrangements (Fig. 2.5a). Regulation of these processes produces the well-characterized migration pattern defined by a continuous cycling between membrane protrusions at the leading cell front, anchoring of the newly formed front, the rolling of the cell body forward, and the release of posterior adhesions to retract the trailing cell membrane. Many signal transduction pathways can influence these processes in large part due to their ability to regulate the spatio-temporal generation of the key lipid messenger, PI4,5P₂, herein referred to as PIP₂.

2.3.1.1 PIPKs Regulate PIP₂ Synthesis at the Leading Edge to Drive Membrane Protrusion

In 1985 Lassing and Lindberg discovered that PIP₂ specifically interacts with profilin, thus starting a long and multifaceted look at the role of phosphoinositides in migration and invasion (Lassing and Lindberg 1985). Since then, PIP₂ has been shown to directly bind many actin-associated proteins that both directly and indirectly regulate the cellular cytoskeletal machinery (Yin and Janmey 2003). Increased PIP₂ levels promote actin polymerization, whereas a decrease results in actin disassembly. PIP₂ controls actin polymerization at many levels, including inactivating proteins involved in actin severing or depolymerization as well as enhancing actin polymerization, branching and bundling (Sechi and Wehland 2000; Takenawa and Itoh 2001). During cell migration, rampant F-actin synthesis at the migrating front pushes the cell membrane forward (Fig. 2.5) resulting in the formation of subcellular structures including membrane ruffles, lamellipodia, microvilli, motile actin comets, filopodia, microspikes and dorsal ruffles (Janmey and Lindberg 2004; Nicholson-Dykstra et al. 2005). All of these membranous protrusions have been observed in different cell types overexpressing type I PIPKs, a family of kinases responsible for the vast majority of PIP₂ produced in a cell (Honda et al. 1999; Matsui et al. 1999; Rozelle et al. 2000; Shibasaki et al. 1997; Yamamoto et al. 2001). Using GFP-tagged PH domains or antibodies that specifically recognize PIP₂, this lipid messenger was found to concentrate in dynamic, actin-rich regions of the cell (Tall et al. 2000).

PIP₂ plays a multifaceted role in regulating F-actin dynamics at these protrusive cell fronts. PIP₂ directly binds gelsolin, a capping protein found at the barbed end of F-actin filaments, blocking its association with F-actin thereby inducing a rapid and local actin polymerization (Niggli 2005). The association of PIP₂ with profilin frees up actin monomers so that they can be incorporated into the growing filaments (Lambrechts et al. 2002; Skare and Karlsson 2002). Using *in vitro* actin co-sedimentation assays, PIP₂ was shown to attenuate the association between F-actin and cofilin, an actin-severing and depolymerizing factor (Gorbatyuk et al. 2006). However, cofilin activity enhances migration (Sidani et al. 2007). The hydrolysis of PIP₂ by Phospholipase C (PLC) releases and activates a distinct membrane-bound pool of cofilin, triggering the formation of membrane protrusions and the establishment of a leading edge (van Rheenen et al. 2007). A localized activation of cofilin enhances the




Fig. 2.5 PIPKs are involved in multiple processes associated with cell migration. **a** PIPKs are integral components at each step in the migration process. Cell migration involves a continuous cycle of membrane protrusion, adhesion formation and maturation at the leading edge, adhesion disassembly (mediated by PIPKI β) at the trailing edge and contractile forces to pull the cell body forward. **b** At the leading edge, PIPKI α regulates Rac and Cdc42 to initiate F-actin synthesis. PIPKI γ isoforms are involved in the trafficking of proteins to the leading edge, the assembly of proteins that mediate adhesions and the maturation of these adhesions to induce actomyosin contractions

number of F-actin barbed ends, setting a site for membrane protrusion, however, curtailing that activity is also required to promote enhanced filament formation and actin bundling required to push the membrane forward. The dynamic interplay between actin severing and F-actin synthesis is further observed in the association of PIP₂ with villin. PIP₂ binding to villin differentially regulates the protein's activities, inhibiting its actin severing and capping activity while enhancing its ability to bundle actin (Kumar and Khurana 2004; Kumar et al. 2004).

While uncovering roles for PIP₂ in regulating actin-associated proteins has been widely successful, defining the specific proteins involved in its synthesis during cell migration has been less concrete. PIPKI α is recruited to membrane ruffles and increases PIP₂ levels in response to EGF and PDGF (Fig. 2.5b) (Doughman et al. 2003b; Honda et al. 1999). Depletion of PIPKI α in mouse embryo fibroblasts blocks cell migration, and PIPKI α -deficient mast cells display atypical actin organization (Kisseleva et al. 2005; Sasaki et al. 2005). Both PIPKI α and PIPKI β have been shown to directly bind Rac1, a member of the Rho family of small GTPases that help regulate actin-cytoskeletal dynamics (Hall and Nobes 2000), and are thus thought to be Rac1 effectors (Weernink et al. 2004). PIPKI α is targeted to lamellipodia and its presence within signaling cascades that include Rho-GTPases is widely known (Doughman et al. 2003b). In addition, the uncapping of actin filaments in cells expressing a constitutively active mutant of Rac1 could be blocked by the overexpression of a dominant negative mutant of PIPKI β but not a kinase-dead mutant of PIPKI α (Tolias et al. 1998, 2000). However, conflicting evidence fails to define whether PIPKI α is upstream or downstream of these proteins. Mutations in the PIPKI α binding region on Rac1 prevents translocation to the plasma membrane and fails to activate downstream signaling responses to integrin-mediated adhesion (Del Pozo et al. 2002). A more recent report shows that PIPKI α regulates Rac1 activation and downstream F-actin assembly, focal adhesion formation and directed migration (Chao et al.). Other reports identify PIPKI α as an effector of Rho-GTPases downstream of G protein-coupled and growth factor receptor activation (Chatah and Abrams 2001). GTPase activation is facilitated by guanine-nucleotide exchange factors (GEFs) that are often activated by phosphoinositides including PIP₂ (Di Paolo and De Camilli 2006). However, effectors of GTPases are often the PIPKs that generate phosphoinositides (Di Paolo and De Camilli 2006). This positive feedback loop results in the generation of membrane domains enriched for specific phosphoinositides. Oude Weernink et al. have shown that all type I PIPK isoforms can associate with RhoA and Rac1 but not Cdc42, another Rho family GTPase (Weernink et al. 2004). However, Cdc42 can stimulate PIP5K activity (Weernink et al. 2004). The concomitant stimulation of neuronal Wiskott-Aldrich syndrome protein (N-WASP) by Cdc42 and PIP₂ increases actin filament nucleation and F-actin synthesis via activation of the actin-related protein 2/3 complex (Arp2/3) (Prehoda et al. 2000).

2.3.1.2 PIPKI γ Regulates the Formation and Maturation of Integrin-mediated Contacts

Following membrane protrusion at the migrating front, the cell anchors this region to the underlying matrix via the formation of specialized adhesive structures, including

focal adhesions, focal complexes, podosomes, invadopodia and hemidesmosomes (Fig. 2.5) (Broussard et al. 2008; Webb et al. 2002). Cell migration requires a tight coordination of the formation/turnover of these structures (Webb et al. 2002). Focal adhesions are large, multi-protein complexes that physically link the cellular cytoskeleton to the extra-cellular matrix (ECM). Their formation generally involves integrin molecules and is dependent on PIP₂ (Ling et al. 2002). Many of the proteins required for focal contact formation, such as α -actinin, vinculin, talin and ezrin, are positively regulated by PIP₂ (Ling et al. 2006). PIPKI γ 2 is required for EGF-stimulated directional migration and is phosphorylated by the EGFR and Src within its C-terminus (Sun et al. 2007). PIPKI γ 2 phosphorylation modulates an association with talin, facilitating its assembly into dynamic adhesive complexes at the leading edge (Ling et al. 2002, 2003). Talin is a scaffolding protein that can bind both β integrin, resulting in its activation and engagement to the ECM, and F-actin or vinculin at nascent adhesions (Nayal et al. 2004). Furthermore, talin binds PIP₂ and this interaction is required for targeting talin to nascent contacts and enhancing its association with β integrin (Ling et al. 2002; Martel et al. 2001). Expression of a PIP₂-specific PH domain blocked the membrane localization of talin essential for focal adhesion maintenance (Martel et al. 2001). In addition, a local enrichment of PIP₂ enhances vinculin assembly into newly formed focal contacts (Chandrasekar et al. 2005). PIP₂ binding to vinculin disrupts its auto-inhibition, exposing protein binding domains that mediate actin and talin association (Bakolitsa et al. 2004). Additionally, PIP₂ binding to vinculin exposes a proline rich domain resulting in a vinculin/VASP association that recruits profilin/G-actin or Arp2/3, facilitating their assembly into nascent contacts and enhancing F-actin nucleation and synthesis (De-Mali et al. 2002; Millard et al. 2004). This suggests that PIPKI γ 2 is recruited to newly forming contacts and the generation of PIP₂ aides their maturation. The result is a rigid adhesive complex that strengthens the cell infrastructure, a critical step in migration as contractile forces use these anchors to pull a cell in the direction of migration.

2.3.1.3 Trafficking of Integrin-containing Vesicles to the Leading Edge Is Mediated by PIPKI γ and PIP₂

Coordinated cell migration requires the trafficking of newly synthesized and endocytosed proteins to the leading edge (Caswell and Norman 2006; Prigozhina and Waterman-Storer 2004; Schmoranz et al. 2003). Impairment of the endocytosis, trafficking and exocytosis of integrins drastically impairs the establishment of polarity and directionality of cell migration (Tayeb et al. 2005; Zovein et al.). Furthermore, PIP₂ generation is required for vesicular trafficking (Wenk and De Camilli 2004).

PIPKI γ knockout studies have revealed vesicle trafficking defects both at the plasma membrane and in endosome-like structures in neuronal cells (Di Paolo et al. 2004; Wenk et al. 2001). Furthermore, several studies define a pivotal role for the exocyst protein complex in the polarized trafficking of transmembrane proteins during cell polarization and migration (Letinic et al. 2009). The exocyst complex consists of eight subunits and facilitates the exocytosis of post-Golgi and endocytic recycling

endosomes (including integrin-containing vesicles) to regions of rapid membrane expansion, such as occurs in migrating cells (Yeaman et al. 2001). Depletion of exocyst complex components inhibits wound healing and migration (Rosse et al. 2006). Rab11 and Arf6, two small GTPases that regulate PIPKs, also regulate the exocyst complex and integrin trafficking (Oztan et al. 2007). Furthermore, at least two exocyst subunits, Sec3 and Exo70, bind PIP₂ via conserved basic residues in their C-termini (Liu et al. 2007). PIPKI γ 2 is able to associate with the exocyst complex and regulate the trafficking of integrin molecules in migrating cells (unpublished data). PIPKI γ could play a role in the assembly of the exocyst complex onto vesicles (Fig. 2.5). In epithelial cells, PIPKI γ 2 is required for the basolateral sorting and endocytosis of E-cadherin via the AP1B and AP2 adaptor protein (AP) complexes respectively (Ling et al. 2007). This occurs by a novel mechanism where PIPKI γ interacts directly with E-cadherin at regions within the PIP kinase domain and then recruits specific AP complex subunits through interactions within its unique C-terminus. This may be a mechanism for regulation of both cell adhesion and migration. In addition, a localized generation of PIP₂ may be important for vesicle fusion at the plasma membrane (Fig. 2.5). PIP₂ and type PIPKI activity have been shown to be critical during Ca²⁺-activated secretion of large dense-core vesicles (Hay et al. 1995). A critical player in this process, synaptotagmin, binds PIP₂ via C2 domains and mediates membrane fusion (Bai et al. 2004). Furthermore, Gong et al. showed that chromaffin cells taken from the adrenal gland of PIPKI γ $-/-$ mice displayed a defect in the readily releasable pool of vesicles containing catecholamine and a delay in fusion pore expansion (Gong et al. 2005). Both Sec3 and Exo70 were shown to bind PIP₂ at the plasma membrane in yeast, a critical interaction for normal exocytosis and maintenance of cell morphology (He et al. 2007; Zhang et al. 2008).

2.3.1.4 Rear Retraction of the Cell Requires the Dissociation and Internalization of Integrin and Acto-myosin Contractility

During cell migration, new adhesions are formed along the protruding edge. Their maturation develops a focal point used to generate of forces required to push the membrane in the direction of migration. In addition, the maturation of focal adhesions stabilizes forces used to pull the rear of the cell forward. Overexpression of type I PIPKs has been shown to induce the formation of F-actin stress fibers, though the formation of actin-based protrusions is inhibited at the cell rear (Shibasaki et al. 1997; Yamamoto et al. 2001). Instead, rear retraction occurs at least in part through RhoA-regulated acto-myosin contractility (Yoshinaga-Ohara et al. 2002). In fibroblasts, integrin-mediated adhesion to fibronectin results in a rapid increase in PIP₂ synthesis and previous work has shown that PIPKI γ interacts with RhoA resulting in enhanced PIP₂ production (Fig. 2.5) (Weernink et al. 2004). Lokuta et al. show that PIPKI γ 2 is enriched in the uropod of directionally migrating neutrophils (Lokuta et al. 2007). Overexpression of a kinase-inactive mutant of PIPKI γ 2 compromised uropod formation and retraction of the cell rear (Lokuta et al. 2007). When bound to GTP, RhoA can signal through ROCK to initiate acto-myosin contractility via its ability to enhance the phosphorylation of the myosin light chain both directly and

indirectly through MLCK (Schramp et al. 2008). ROCK also plays a critical role in the RhoA-mediated activation of PIP5 K to synthesize PIP₂, a signaling pathway utilized during neurite remodeling (van Horck et al. 2002; Yamazaki et al. 2002).

In addition, focal contacts at the rear of the cell need to be disassembled. A prevailing theory of cell migration adheres to the idea that disassembled adhesions are internalized via endocytosis and recycled to the protruding edge where they once again are utilized in the formation of nascent adhesions (Pierini et al. 2000). The protease calpain, FAK, phosphatases and kinases that regulate FAK, microtubules and dynamin 2 have all been shown to regulate focal adhesion disassembly (Brousard et al. 2008; Burridge et al. 2006; Ezratty et al. 2005; Franco et al. 2004). All three isoforms of type I PIPKs have been implicated in clathrin-mediated endocytosis (Franco and Huttenlocher 2005). Recently, Chao et al. showed that PIPKI β is required for β 1 integrin uptake and adhesion turnover during cell migration (Chao et al.). Localized PIPK recruitment and PIP₂ formation can recruit molecules that directly regulate endocytosis to the plasma membrane and modulate proteins involved in the inactivation of focal adhesions (Fig. 2.5). PIP₂ can directly bind many endocytic clathrin adaptors, including AP-2, AP180 and epsin (Qualmann et al. 2000). Clathrin adaptor components and dynamin become enriched at focal adhesion sites prior to internalization (Chao and Kunz 2009; Ezratty et al. 2005). Loss of PIPKI β blocked the assembly of clathrin adaptors at adhesion plaques and prevented an association between dynamin and FAK, an interaction required for endocytosis (Chao et al.). Interestingly, in addition to its role in promoting F-actin synthesis at nascent contacts, a role for vinculin in promoting focal adhesion disassembly is also emerging. The introduction of a vinculin mutant unable to bind PIP₂ into mouse melanoma cells repressed its translocation from the membrane to the cytosol and blocked the disassembly of focal adhesions, cell spreading and migration (Chandrasekar et al. 2005). Calpain is another PIP₂ regulated protein that directly mediates focal adhesion disassembly (Franco and Huttenlocher 2005; Franco et al. 2004). Calpain is a protease that can cleave adhesion proteins including talin, vinculin, paxillin and β integrin and is required for focal adhesion turnover and cell migration (Franco and Huttenlocher 2005; Franco et al. 2004). However, the activation cues to signal these processes remain to be fully uncovered.

PIPKs are proving to be integral components in helping regulate the many processes required for cell migration. The existence of multiple PIPK isoforms may enable cells to coordinate PIP₂ synthesis at specific sites to regulate the forces that induce membrane protrusion, the trafficking events that localize proteins at the leading edge, the assembly, maturation and endocytosis of adhesions and the forces generated to retract the trailing rear of the cell.

2.3.2 PIP Kinases in Adaptor Protein Complex Assembly and Protein Trafficking

In the past decade, considerable evidence has demonstrated that PIPKs and their lipid biproducts, PIP₂, are actively involved in various protein trafficking processes.

Table 2.1 Adaptor complexes consist of different subunits

| AP complex | Subunits |
|------------|--------------------------------------|
| AP-1A | $\beta 1, \gamma, \sigma 1, \mu 1A$ |
| AP-1B* | $\beta 1, \gamma, \sigma 1, \mu 1B$ |
| AP-2 | $\beta 2, \alpha, \sigma 2, \mu 2$ |
| AP-3A | $\beta 3A, \delta, \sigma 3, \mu 3A$ |
| AP-3B* | $\beta 3B, \delta, \sigma 3, \mu 3B$ |
| AP-4 | $\beta 4, \epsilon, \sigma 4, \mu 4$ |

*Tissue specific AP complexes

Among these, the most widely characterized is the role of PIPKs in clathrin-mediated endocytosis and endosomal recycling, both of which are dependent on AP complexes. PIPKs directly bind AP complexes and their ability to synthesize a localized pool of PIP₂ results in AP complex assembly and activation. Here we will summarize how PIPKs regulate different AP complexes to modulate distinct protein trafficking processes.

2.3.2.1 AP Complexes

AP complexes are indispensable components of clathrin-coated vesicles in the endocytic and post-Golgi trafficking pathways. AP complexes not only link clathrin to the membrane, but also determine the specificity of cargo selection at various membrane compartments (Nakatsu and Ohno 2003; Ohno 2006). To date, four ubiquitously expressed AP complexes (AP-1A, AP-2, AP-3A, and AP-4) have been identified (Dell’Angelica et al. 1999; Nakatsu and Ohno 2003), with two additional complexes, AP-1B and AP-3B, found specifically in epithelial and neuronal cells respectively (Fölsch et al. 1999; Nakatsu and Ohno 2003; Ohno 2006). Each AP complex consists of two large adaptin subunits, one derived from the β -class ($\beta 1-4$) and one of γ, α, δ or ϵ , together with one medium subunit ($\mu 1-4$) and one small subunit ($\sigma 1-4$) (Table 2.1) (Jackson 1998; Ohno 2006). In each AP complex, the large adaptin subunits are responsible for membrane association through PIP₂ (Gaidarov et al. 1999; Ohno 2006; Rohde et al. 2002). The μ subunit determines specific cargo selection, while the β subunit is required for clathrin recruitment (Ohno 2006). Therefore, different AP complexes direct distinct cargo proteins at the membrane surface of vesicles (Fig. 2.6).

2.3.2.2 PIP Kinases in Regulation of AP Complex Assembly

Recent studies have demonstrated that PIP kinases are involved in clathrin-mediated vesicle trafficking at the plasma membrane. PIPKs have been found to directly bind AP complexes, regulating AP complex assembly and the formation of clathrin-coated pits. Here we will discuss how PIPKs regulate protein trafficking at the plasma membrane through specific interactions with AP complexes.




Fig. 2.6 AP complexes mediate endocytic and post-Golgi trafficking. Different AP complexes are involved in clathrin-mediated protein trafficking between different compartments. PIPKI γ 2 interacts with AP-1B and AP-2 complexes, modulating basolateral transport and endocytosis

E-cadherin maintains epithelial cell morphology by forming adherens junctions (AJs) with adjacent cells. The amount of E-cadherin on the plasma membrane directly determines the strength of AJs (Yap et al. 2007). In addition to E-cadherin gene expression, post-translational regulations such as exocytosis, endocytosis, recycling and lysosomal degradation have also been implicated as important factors affecting the stability of AJs (Schill and Anderson 2009a). Furthermore, PIP₂ has been shown to be an essential regulator of different E-cadherin trafficking processes (Schill and Anderson 2009a). PIPKI γ directly binds all type I classical cadherins and colocalizes with E-cadherin in epithelial cells (Akiyama et al. 2005; Ling et al. 2007). Specifically, PIPKI γ 2 has been found to regulate AJ assembly by modulating E-cadherin trafficking between endosomes and the plasma membrane (Ling et al. 2007). Expression of wild-type PIPKI γ 2 promotes both internalization and recycling of E-cadherin, while a kinase dead mutant of PIPKI γ 2 inhibits both of these processes (Ling et al. 2007), indicating that phosphoinositide generation is required for E-cadherin trafficking.

PIPKI γ 2 can directly associate with both E-cadherin and the μ 1B subunit of AP-1B (Ling et al. 2007), suggesting that it could act as a scaffold to link these proteins (Fig. 2.7). Within the unique PIPKI γ 2 C-terminus is the sequence YSPL, Yxx ϕ tyrosine-sorting motif (x represents any and ϕ represents a bulky hydrophobic amino




Fig. 2.7 PIPKI γ 2/AP-1B interaction regulates E-cadherin trafficking. PIPKI γ 2 interacts with the μ 1B subunit of AP-1B through the Yxx ϕ motif in the C-terminus and recruits E-cadherin via the kinase domain. Additionally, PIPKI γ 2-generated PIP₂ assembles PIP₂-interacting proteins of the AP-1B and clathrin vesicle complexes at specific sites

acid) recognized by the μ 1B subunit of AP-1B (Bairstow et al. 2006; Bonifacino and Traub 2003; Ling et al. 2007; Sugimoto et al. 2002). It has been demonstrated that the PIPKI γ 2 YSPL sequence is responsible for the μ 1B interaction, as a mutation of tyrosine to a phenylalanine disrupted their interaction and subsequently decreased the association between E-cadherin and μ 1B (Ling et al. 2007). Interestingly, PIPKI γ 2 also binds to the β 2 subunit of AP-2 (Nakano-Kobayashi et al. 2007). Whether the PIPKI γ 2/AP-2 interaction is also involved in E-cadherin trafficking is not clear.

Current evidence supports a model where PIPKI γ 2 acts as both a scaffold and direct regulator of complex assembly through the generation of PIP₂ during E-cadherin trafficking to the plasma membrane (Fig. 2.7). The μ subunit of the AP complex is the predominate molecule mediating selection through a direct interaction with cargo proteins (Ohno 2006; Sugimoto et al. 2002). However, PIPKI γ 2 could function as a scaffold protein that links E-cadherin to the μ 1B subunit of the AP-1B complex. At the same time, PIPKI γ 2 also functions by temporally and specifically producing PIP₂, which in turn recruits and regulates AP complex proteins and other components of the clathrin-coated vesicle. Certain patients suffering from hereditary diffuse gastric cancer have been found to contain a mutation within the PIPKI γ 2-binding site in the C terminus of E-cadherin (Yabuta et al. 2002; Ling et al. 2007). The mutated E-cadherin shows inhibited interaction with PIPKI γ 2, and localizes mostly to cytoplasmic compartments rather than the plasma membrane, losing the ability to form AJs (Ling et al. 2007; Suriano et al. 2003; Yabuta et al. 2002). These results underscore the role of PIPKI γ 2 in E-cadherin trafficking to the plasma membrane.




Fig. 2.8 PIPKIγ2/AP2 interaction modulates endocytosis. **a** PIPKIγ2 directly interacts with the AP-2 complex through multiple sites. **b** PIPKIγ2 facilitates clathrin-AP-2-mediated endocytosis of YxxΦ-motif-containing cargos. Activated Arf6 recruits and activates PIPKIγ2, leading to PI4,5P₂ generation, which in turn recruits AP-2 (1). Dephosphorylation of PIPKIγ2 by calcineurin or another upstream molecule (2) induces PIPKIγ2-binding to the Ear domain of the β2 subunit of AP-2, further activating PIPKIγ2 (3). Conformational change in μ2 caused by Thr156 phosphorylation (4) promotes cargo binding (5) to μ2 through YxxΦ motif with the help of PI4,5P₂. This results in μ2 binding to and activating the PIPKIγ2 kinase domain (6). Clathrin-engagement to AP-2 disrupts PIPKIγ2 association with the β2 subunit (7). PI4,5P₂ accumulation in step 1, 3, and 6 recruits proteins involved in clathrin vesicle formation. PIPKIγ2 is removed from AP-2 by phosphorylation at Ser645 or other unidentified mechanisms before maturation of the clathrin-coated vesicle (8). **c** PIPKIγ2 may enhance clathrin-AP-2-mediated endocytosis of YxxΦ-independent cargos. Conformational change in μ2 caused by Thr156 phosphorylation (4) promotes PIPKIγ2 association with μ2 through one of the two YxxΦ motifs in the C-terminus of PIPKIγ2 (5), which may in turn enhance the interaction between PIPKIγ2 kinase domain and the μ2 subunit (6). YxxΦ engagement to μ2 may allow cargo selection through YxxΦ-independent mechanisms (7). Clathrin-binding to β2 (8) might facilitate these processes

During clathrin-mediated endocytosis, PIPKIγ2 regulates AP-2 complex assembly by directly binding multiple subunits in the AP-2 complex (Fig. 2.8a). The kinase domain of PIPKIγ2 binds to the μ2 subunit, an association that does not block cargo engagement to μ2 (Krauss et al. 2006). In this situation, cargo proteins bind to μ2 through tyrosine or dileucine sorting motifs. This interaction can then potentially activate PIPKIγ2 activity (Kahlfeldt et al. 2010; Krauss et al. 2006). In addition to this, the tyrosine-based sorting motif (⁶⁴¹SWVYSPL⁶⁴⁷) within the PIPKIγ2 specific C-terminal tail has also been found to directly bind to the μ2 subunit of AP-2 (Bairstow et al. 2006). A recent study discovered another tyrosine-based sorting motif (⁴⁹⁵RSYPTLED⁵⁰²) within the C terminus of PIPKIγ2 that can bind to the μ2 subunit (Kahlfeldt et al. 2010). Moreover, the C-terminal tail of PIPKIγ2 can also interact with the β2 subunit of AP-2, enhancing PIPKIγ2 activity (Kahlfeldt et al. 2010;

Nakano-Kobayashi et al. 2007; Thieman et al. 2009). Where PIPKI γ 2 interacts with the AP-2 complex is believed to mediate the formation of different clathrin-coated vesicles that regulate the trafficking of various cargo proteins (Kahlfeldt et al. 2010; Kwiatkowska 2010).

The PIPKI γ 2/AP-2 interaction is subject to regulation by phosphorylation of the PIPKI γ 2 C-terminus. Src-mediated phosphorylation of PIPKI γ 2 at Tyr644 (Tyr 649 in the human isoform) inhibits its interaction with the μ 2 subunit (Bairstow et al. 2006; Ling et al. 2003) and based on crystallographic data could impair PIPKI γ 2 binding to the β 2 subunit (Kahlfeldt et al. 2010). In addition, phosphorylation of PIPKI γ 2 at Ser645 (Ser650 in the human isoform) by cyclin-dependent kinase-5 (Cdk5) diminishes PIPKI γ 2 interaction with the β 2 subunit (Nakano-Kobayashi et al. 2007; Thieman et al. 2009), while dephosphorylation of PIPKI γ 2, possibly by calcineurin upon plasma membrane depolarization, enhances this binding affinity in clathrin-mediated endocytosis at the presynapse (Nakano-Kobayashi et al. 2007). Interestingly, it has been shown that phosphorylation of either Tyr644 or Ser645 inhibits the phosphorylation of the other (Lee et al. 2005). Therefore, dephosphorylation of both Tyr644 and Ser645 promotes PIPKI γ 2 interaction with both β 2 and μ 2, while Tyr644 phosphorylation alone inhibits this interaction and Ser645 phosphorylation alone might increase PIPKI γ 2 binding to the μ 2 subunit. As clathrin competes with PIPKI γ 2 for the same binding site in the β 2 subunit (Thieman et al. 2009), a PIPKI γ 2 binding switch from β 2 to μ 2 via phosphorylation changes of Tyr644 and Ser645 might occur during AP-2 complex assembly (Kwiatkowska 2010).

PIPKI γ 2-mediated PIP₂ generation recruits AP-2 complex components (Fig. 2.8b) (Höning et al. 2005). Dephosphorylation of PIPKI γ 2 by upstream signals triggers its binding to the Ear domain of the β 2 subunit, which activates PIPKI γ 2 to produce PIP₂. In the presence of PIP₂, phosphorylation of μ 2 subunit at Thr156 causes a conformational change that favors cargo engagement through tyrosine sorting motifs (Höning et al. 2005; Olusanya et al. 2001). Although cargo association with μ 2 subunit via acidic dileucine motifs is not enhanced by the Thr156-phosphorylation-mediated conformational change, the μ 2 subunit has the highest affinity towards both cargo signals when PIP₂ is present (Höning et al. 2005). Cargo engagement induces μ 2 to interact with the kinase core domain of PIPKI γ 2 enhancing its activity. Dynamic PIP₂ production recruits other proteins, such as AP180, epsin and dynamin-2, to facilitate the formation of the clathrin-coated vesicles during endocytosis. It is noteworthy that although PIPKI γ 2 plays important roles in clathrin-AP-2 complex assembly, it is not enriched in clathrin-coated vesicles (Thieman et al. 2009; Wenk et al. 2001), suggesting that PIPKI γ 2 functions in the dynamic assembly of AP-2 complex and clathrin-coated pits, but it is not a component of mature vesicles. Clathrin binding to the β 2 subunit has been found to eliminate PIPKI γ 2 association with β 2 through the C-terminus (Thieman et al. 2009), yet whether this removes PIPKI γ 2 from the AP complex or only causes a binding switch of PIPKI γ 2 from β 2 to μ 2 still needs to be determined.

An alternative hypothesis is that PIPKI γ 2 Yxx Φ motifs associates with μ 2 like a cargo protein (Kahlfeldt et al. 2010; Kwiatkowska 2010) (Fig. 2.8c). This alternative Yxx Φ / μ 2 interaction might provide a mechanism for Yxx Φ sorting motif-independent cargo selections. Yxx Φ / μ 2 interactions have been suggested to facilitate μ 2 interactions with the kinase core domain of PIPKI γ 2 and activate PIPKI γ 2 (Kahlfeldt et al.

2010; Kwiatkowska 2010). Kahlfeldt et al. showed that PIPKI γ 2 activity could be increased when purified μ 2 and the C-tail of PIPKI γ 2 were incubated with cell lysates containing overexpressed PIPKI γ 2. However, it is possible that the PIPKI γ 2 C-terminus mimicks the effects of a cargo protein like EGFR, as they did not see any increase in PIPKI γ 2 activity when the lysates were incubated with μ 2 alone (Kahlfeldt et al. 2010). Although evidence shows that PIPKI γ 2 can interact with μ 2 through either the kinase core domain or through one of the two Yxx ϕ motifs (Bairstow et al. 2006; Kahlfeldt et al. 2010), whether the μ 2-PIPKI γ 2 complex alone can lead to PIP kinase activation *in vivo* or whether inhibiting the μ 2 association with the PIPKI γ 2 C-terminus impairs PIPKI γ 2 functions in AP-2 complex assembly is not clear. Future studies are needed to understand how the interaction between μ 2 and the Yxx ϕ motifs in the C-terminus of PIPKI γ 2 correlates with Yxx ϕ sorting motif-independent cargo selections.

2.4 Nuclear Localized PIPKs Regulate Gene Expression and mRNA Processing

2.4.1 Nuclear PIP Kinases and Phosphoinositides

Much as how the discovery of the phosphatidylinositol (PI) cycle became fundamental to our understanding of signal transduction and overall cell biology (Gurr et al. 1963; Hokin and Hokin 1953; Kleinig 1970), the discovery of nuclear phospholipid signaling has revolutionized our view of the processes regulated by phospholipids (Cocco et al. 1987, 1988). Still, nuclear phospholipid signaling is fulfilled by lipid second messengers and requires an array of coordinated activities from lipid kinases, phosphatases, and effectors. However, the lipid kinases within the nucleus appear to be more selective, in that not all identical lipid kinases exist in the nucleus and the phosphoinositides they generate differ both spatially and temporally. Interestingly, a substantial pool of nuclear phosphoinositides does not appear to be associated with the nuclear membrane (Payraastre et al. 1992). These observations are helping to unravel another facet of phospholipid signaling found in cells.

2.4.1.1 PIPKs and PIP₂ at the Nuclear Envelope

Continuous with the ER, the nuclear envelope is a bilayered membranous structure. Pioneering studies on isolated rat liver nuclei revealed the existence of PI and PIP pools in the nuclear envelope (Smith and Wells 1983a). A recent study in sea urchin sperm identified an atypical composition of polyphosphoinositides in the nuclear membrane, as much as 51% of the total phospholipids, an enrichment conserved in mammalian sperm (Garnier-Lhomme et al. 2009). Incubation of isolated nuclear envelopes with [γ -³²P]ATP resulted in rapid labeling of phospholipid

products, later identified as phosphatidylinositol 4-phosphate (PI4P), phosphatidylinositol 4, 5-bisphosphate (PI4,5P₂), and phosphatidic acid (PA), suggesting nuclear envelopes contain PIK, PIPK, and diacylglycerol kinase (DGK) activity (Smith and Wells 1983b). Additionally, substantial evidence demonstrates the requirement of PIP₂, diacylglycerol (DAG), and phospholipase C (PLC γ) in the membrane fusion events leading to nuclear envelope formation during mitosis (Dumas et al. 2010). In conjunction with this, PKC β II was identified as a mitotic lamin kinase and its phosphorylation of lamin B is critical for nuclear envelope disassembly, providing evidence that phosphoinositide signaling extended from the nuclear membrane into the nucleus (Goss et al. 1994; Thompson and Fields 1996). The hydrolysis of PIP by phosphomonoesterases and a PIP-dependant ATPase are also associated with the nuclear envelope (Smith and Wells 1984a, b). Besides the canonical PIPKs, phosphatidylinositol 3-kinase (PI3K) may also produce PI3,4P₂ at the nuclear surface and therefore act as a “PIPK” (Yokogawa et al. 2000). These studies suggest that a crucial phosphatidylinositide metabolic cycle is present at the nuclear membrane and is required for normal cellular processes. Interestingly, PIPKs and PIP₂ were not found to be associated with the invaginations of the nuclear envelope (Boronenkov et al. 1998), indicating that PIPK activity and PIP₂ production are temporally and spatially regulated. At present, it is unclear which phosphoinositides (other than PI4P, PI3,4P₂, and PI4,5P₂) and which PIPKs are involved in the synthesis of phosphoinositides at the nuclear envelope and even less is known about how nuclear PIPKs are regulated.

2.4.1.2 The Intra-nuclear PIPKs and PIP₂

Studies by Cocco et al. in the late 1980s revealed that not all PIPKs in the nucleus appeared to be associated with the nuclear membrane. Using highly purified nuclei from mouse erythroleukemia (MEL) and Swiss 3T3 cells, nuclear membranes were stripped away using detergents and the remaining matrix was able to produce both PI4P and PIP₂ (Cocco et al. 1987, 1988). The existence of phosphoinositides in non-membranous structures within the nucleus raised the question of where these second messengers and the enzymes that generate them localize. Plausible hypotheses include the nuclear matrix/nucleoskeleton, chromatin, and protein complexes that are associated with these nuclear microenvironments (Fig. 2.9). Payraastre and colleagues showed nuclear matrix-associated PI4 K, PIPKIs, DGK, and PLC in mouse NIH 3T3-fibroblasts and rat liver cells (Payraastre et al. 1992). Using laser scanning confocal microscopy and immunofluorescence staining, PIP, PIPKI α (PIPKI β in mouse), PIPKII α (PIP4K α), PIPKII β (PIP4K β), PI(4,5)P₂, and PI(3,4)P₂ were identified at nuclear speckles (interchromatin granule clusters) (Boronenkov et al. 1998). However, the amount of PI4P is about 20-fold higher than that of PI5P in nucleus, and [γ -³²P]ATP incubation with isolated rat liver nuclei determined the relative labeling ratio for PI4,5P₂ generation at the 5 vs. 4 OH position was ~ 1.8 , suggesting that PIPKIs are the major kinase for PIP₂ synthesis within the nucleus (Keune et al. 2010; Vann et al. 1997).




Fig. 2.9 The nuclear PIPKs and PIP₂ pools. The nuclear envelope continues from the ER and contains PIP₂ as well as associated PIPK activities in both outer and inner bilayers. The interchromatin granule clusters (nuclear speckles) nest another PIPK and PIP₂ pool. Some protein complexes docked on the nuclear matrix may also serve to recruit PIPK and PIP₂ or that the PIP₂/PIPK can regulate the assembly of such matrices

So far, nuclear PIPKI α has been identified in complex with a speckle targeted PIPKI α regulated-poly(A) polymerase (Star-PAP), a PIP₂-sensitive protein kinase, CKI α , and mRNA processing factors (Mellman et al. 2008). The yeast Mss4p (mammalian PIPK homologue) contains a functional nuclear localization signal (NLS) and can shuttle between the cytoplasm and the nucleus (Audhya and Emr 2003). However, it remains to be determined whether PIPKI α can shuttle between the cytoplasm and nucleus or is capable of generating the majority of the nuclear PIP₂. Recently, another PIPKI family member, PIPKI γ 4, was identified in the nucleus (Schill and Anderson 2009b). PIPKI γ 4 is the only PIPKI γ subfamily member to be detected in the nucleus though its function remains to be characterized. PIPKI β predominantly localizes in the nucleus in certain cell lines owing to a specific 17-amino-acid α -helix insertion (Ciruela et al. 2000). PIPKI α also partially localizes

Table 2.2 The identified nuclear PIP kinases

| PIPK | Substrate | Product | Localization | Reference |
|-----------------------|-----------|------------------|-----------------|---|
| PIPKI α | PI4P | PIP ₂ | Nuclear speckle | Boronenkov et al. 1998; Mellman et al. 2008 |
| PIPKI $\gamma/4$ | PI4P | PIP ₂ | Nucleus | Schill and Anderson 2009 |
| PIPKII β/α | PI5P | PIP ₂ | Nucleus | Boronenkov et al. 1998; Ciruella et al. 2000; Bultsma et al. 2010 |

to the nucleus via PIPKII β an interaction that is indispensable for protecting PIPKII β from degradation via the nuclear ubiquitin ligase, Cul3-SPOP (cullin 3-speckle-type POZ domain protein) (Bultsma et al. 2010; Bunce et al. 2008). It appears that the principle role of nuclear PIPKII β is to regulate PI5P levels rather than generate PIP₂ (Bultsma et al. 2010; Keune et al. 2010). The different PIPK isoforms in the nucleus and potentially different localizations of these kinases (Table 2.2) suggest the production of PIP₂ is spatially and temporally regulated, which could result in differential control of the cellular signaling and biological functions.

2.4.2 Regulation and Functions of the Nuclear PIPKs

Compared to the vast amount of data detailing the functions of PIPK and PIP₂ in the cytoplasm, the nuclear functions of these molecules are less understood. Emerging data indicate that nuclear PIP₂ regulates diverse processes including stress response, cell cycle control and mitosis, transcription, mRNA processing and export, DNA repair, chromatin remodeling, and gene expression (Barlow et al. 2010; Bunce et al. 2006b; D'Santos et al. 1998; Gonzales and Anderson 2006; Irvine 2003; Keune et al. 2010). It is becoming clearer that specific nuclear PIPKs produce localized PIP₂ in response to different external and internal signals. Nuclear PIPKs could be associated with scaffolds containing other proteins including effector proteins or even phospholipids, which are organized into the nuclear matrix. The resources of PIPs can be supplied locally or delivered by phosphoinositide-carrier proteins. Upon PIPK stimulation and PIP₂ production, conformational/stoichiometric changes in effector proteins could lead to their activation and the induction of nuclear biological processes (Bunce et al. 2006a).

2.4.2.1 Regulation of the Nuclear PIPKI α

PIPKI α has been shown to be targeted to nuclear speckles where it complexes with the non-canonical poly(A) polymerase Star-PAP, CKI α , RNA polymerase II, and splicing factors (Gonzales et al. 2008; Mellman et al. 2008). The PIPKI α C-terminus directly interacts with Star-PAP. It regulates the expression of select genes induced by oxidative stress through the generation of PIP₂ and modulation of Star-PAP activities at the 3'-end of pre-mRNAs (Mellman et al. 2008). However, the upstream


regulator of this PIPKI α intra-nuclear activity has not yet been identified. CKI α is one of the potential site-specific regulators of PIPKI α , but it needs another protein kinase for priming phosphorylation of PIPKI α . Another possibility could be PIP₂ downstream signaling molecules, which can mediate PIPKI α activities by a feedback mechanism. PIPKI α also associates with pre-mRNA splicing factors at nuclear speckles (Boronenkov et al. 1998; Mellman et al. 2008). Speckles are storage sites for these factors whose structure dynamically changes during the cell cycle concomitantly with changes in nuclear PIP₂ levels (Clarke et al. 2001; Lamond and Spector 2003). Interestingly, PIP₂ and PI3K all assemble in nuclear speckles (Boronenkov et al. 1998; Didichenko and Thelen 2001; Osborne et al. 2001). These observations suggest multiple roles for nuclear PIPKI α and PIP₂ in addition to mediating Star-PAP activity.

The tumor suppressor retinoblastoma protein RB (pRB) is another PIPKI α regulator. pRB interacts with and highly activates PIPKI α in a large T antigen-regulated manner (Divecha et al. 2002). In line with this, PIP₂ is sufficient to target the chromatin remodeling complex BAF, which can then be recruited to gene transcription sites by pRB (Zhao et al. 1998), and may further promote an interaction between chromatin and the nuclear matrix via the BAF complex subunit BRG-1 (Rando et al. 2002). The results implicate a role of PIPKI α in transcriptional control and in the regulation of proteins via their ability to bind PIP₂. PIP₂ can also bind histones H1 and H3 alleviating the suppression of transcription (Yu et al. 1998). These studies suggest PIPKI α and its product PIP₂ are involved in the regulation of different phases of gene transcription.

2.4.2.2 Regulation of the Nuclear PIPKII β /II α

PIPKII β in combination with PIPKII α (as addressed formerly) potentially contribute to the generation of a second nuclear PIP₂ pool. Studies in MEL cells demonstrate that the function of PIPKII in the nucleus is mainly to regulate nuclear PI5P levels, because depletion of PIPKII β by RNAi increased PI5P levels while overexpression of PIPKII β decreased the nuclear PI5P levels (Jones et al. 2006). Like nuclear PIP₂, nuclear PI5P levels also change during the cell cycle (Clarke et al. 2001), implying a role of PIPKII β and PI5P in cell cycle regulation. PIPKII β can be activated by phosphorylation at Ser326 by p38-MAPK in response to UV irradiation resulting in nuclear accumulation of PI5P (Jones et al. 2006). The Cul3-SPOP ubiquitin ligase complex also regulates PIPKII β levels and PI5P production (Bunce et al. 2008). In concert with PIPKII β , the type I PI4,5P₂ 4-phosphatase converts nuclear PIP₂ to PI5P and therefore modulates PI5P levels (Barlow et al. 2010; Bunce et al. 2008; Zou et al. 2007). PI5P can induce the activation of the p38-MAPK pathway and stimulate Cul3-SPOP activity for PIPKII β ubiquitylation (Bunce et al. 2008). As a PIPKII β interacting partner, PIPKII α nuclear targeting seems to be required for suppressing the Cul3-SPOP-dependent PIPKII β ubiquitylation, because expression of the kinase-dead PIPKII α and a nuclear targeting-defective PIPKII α were not

Fig. 2.10 PIPKs generate PIP_2 in the nucleus to control many signaling events. Nuclear PIPKs and PIP_2 regulate different yet specific cellular molecules and processes through interactions with select nuclear proteins or by producing other second messengers. chrom=chromatin; nmtx=nuclear matrix



able to suppress ubiquitylation of PIPKII β , whereas overexpression of the wild-type PIPKII β did (Bultsma et al. 2010). An extension of PIPKII β regulated signaling lies in that PI5P can interact with inhibitor of growth protein 2 (ING2) and regulate the localization of ING2 to chromatin. ING2, in turn, modulates the acetylation of the tumor suppressor protein p53, linking PIPKII β activities to transcriptional regulation and gene expression (Bultsma et al. 2010; Jones et al. 2006).

2.4.3 Downstream Signaling of the Nuclear PIPK and PIP_2

As discussed above, PIP_2 can directly regulate interacting proteins or it can be processed to generate other second messengers including PIP_3 (Fig. 2.10). Class I and class II PI3Ks have been found in nucleus with the class IA PI3K being responsible for generating the majority of nuclear PIP_3 [reviewed in (D'Santos et al. 1998; Gonzales and Anderson 2006)]. Another bifurcated signaling pathway from nuclear PIP_2 is mediated by the nuclear phosphoinositide-specific phospholipase C (PI-PLC). PI-PLCs break down PIP_2 into DAG and inositol-1,4,5-triphosphate (IP_3). DAG is a direct activator of nuclear PKCs, which are known regulators of gene transcription, DNA synthesis, stress response, mitosis and cell cycle progression (D'Santos et al. 1998). IP_3 receptors have been identified on both the outer and inner nuclear membrane (Humbert et al. 1996). The nuclear IP_3 is recognized to be involved in Ca^{2+} import into nucleus and nuclear Ca^{2+} homeostasis, processes important for gene expression, DNA synthesis and repair, chromatin condensation, protein import, and apoptosis (Bading et al. 1997; D'Santos et al. 1998; Hardingham et al. 1997; Malviya and Rogue 1998) (Fig. 2.10).

The PIPK and PIP_2 downstream signaling cascades are far more intricate than we have thought so far. Identification of more PIPK regulators and effectors as well as

other roles for PIPKs, perhaps even independent of their kinase activity, will enhance our understanding of phosphoinositide signaling in the nucleus.

Acknowledgments We apologize to those whose work could not be cited due to space limitations. We thank Suyong Choi, Yue Sun and Rakesh Singh for their scientific discussions and comments on the manuscript prior to submission. Structures rendered in The PyMOL Molecular Graphics System, Version 1.3, Schrödinger, LLC. Research in the authors' lab is supported by NIH grants GM057549, GM051968 and CA104708.

References

- Akiyama C, Shinozaki-Narikawa N, Kitazawa T, Hamakubo T, Kodama T, Shibasaki Y (2005) Phosphatidylinositol-4-phosphate 5-kinase gamma is associated with cell-cell junction in A431 epithelial cells. *Cell Biol Int* 29:514–520
- Anderson RA, Boronenkov IV, Doughman SD, Kunz J, Loijens JC (1999) Phosphatidylinositol phosphate kinases, a multifaceted family of signaling enzymes. *J Biol Chem* 274:9907–9910
- Audhya A, Emr SD (2003) Regulation of PI4,5P₂ synthesis by nuclear-cytoplasmic shuttling of the Mss4 lipid kinase. *EMBO J* 22:4223–4236
- Bading H, Hardingham GE, Johnson CM, Chawla S (1997) Gene regulation by nuclear and cytoplasmic calcium signals. *Biochem Biophys Res Commun* 236:541–543
- Bai J, Tucker WC, Chapman ER (2004) PIP₂ increases the speed of response of synaptotagmin and steers its membrane-penetration activity toward the plasma membrane. *Nat Struct Mol Biol* 11:36–44
- Bairstow SF, Ling K, Su X, Firestone AJ, Carbonara C, Anderson RA (2006) Type Igamma661 phosphatidylinositol phosphate kinase directly interacts with AP2 and regulates endocytosis. *J Biol Chem* 281:20632–20642
- Bakolitsa C, Cohen DM, Bankston LA, Bobkov AA, Cadwell GW, Jennings L, Critchley DR, Craig SW, Liddington RC (2004) Structural basis for vinculin activation at sites of cell adhesion. *Nature* 430:583–586
- Barlow CA, Laishram RS, Anderson RA (2010) Nuclear phosphoinositides: a signaling enigma wrapped in a compartmental conundrum. *Trends Cell Biol* 20:25–35
- Bazenet CE, Ruano AR, Brockman JL, Anderson RA (1990) The human erythrocyte contains two forms of phosphatidylinositol-4-phosphate 5-kinase which are differentially active toward membranes. *J Biol Chem* 265:18012–18022
- Bonifacino J, Traub L (2003) Signals for sorting of transmembrane proteins to endosomes and lysosomes. *Annu Rev Biochem* 72:395–447
- Boronenkov IV, Anderson RA (1995) The sequence of phosphatidylinositol-4-phosphate 5-kinase defines a novel family of lipid kinases. *J Biol Chem* 270:2881–2884
- Boronenkov IV, Loijens JC, Umeda M, Anderson RA (1998) Phosphoinositide signaling pathways in nuclei are associated with nuclear speckles containing pre-mRNA processing factors. *Mol Biol Cell* 9:3547–3560
- Broussard JA, Webb DJ, Kaverina I (2008) Asymmetric focal adhesion disassembly in motile cells. *Curr Opin Cell Biol* 20:85–90
- Bultsma Y, Keune WJ, Divecha N (2010) PIP4Kbeta interacts with and modulates nuclear localization of the high-activity PtdIns5P-4-kinase isoform PIP4Kalpha. *Biochem J* 430:223–235
- Bunce MW, Bergendahl K, Anderson RA (2006a) Nuclear PI(4,5)P₂: a new place for an old signal. *Biochim Biophys Acta* 1761:560–569
- Bunce MW, Gonzales ML, Anderson RA (2006b) Stress-ING out: phosphoinositides mediate the cellular stress response. *Sci STKE* 2006:pe46

- Bunce MW, Boronenkov IV, Anderson RA (2008) Coordinated activation of the nuclear ubiquitin ligase Cul3-SPOP by the generation of phosphatidylinositol 5-phosphate. *J Biol Chem* 283:8678–8686
- Burden LM, Rao VD, Murray D, Ghirlando R, Doughman SD, Anderson RA, Hurley JH (1999) The flattened face of type II beta phosphatidylinositol phosphate kinase binds acidic phospholipid membranes. *Biochemistry* 38:15141–15149
- Burridge K, Sastry SK, Sallee JL (2006) Regulation of cell adhesion by protein-tyrosine phosphatases. I. Cell-matrix adhesion. *J Biol Chem* 281:15593–15596
- Cabezas A, Pattni K, Stenmark H (2006) Cloning and subcellular localization of a human phosphatidylinositol 3-phosphate 5-kinase, PIKfyve/FabI. *Gene* 371:34–41
- Caswell PT, Norman JC (2006) Integrin trafficking and the control of cell migration. *Traffic* 7:14–21
- Chandrasekar I, Stradal TE, Holt MR, Entschladen F, Jockusch BM, Ziegler WH (2005) Vinculin acts as a sensor in lipid regulation of adhesion-site turnover. *J Cell Sci* 118:1461–1472
- Chao WT, Kunz J (2009) Focal adhesion disassembly requires clathrin-dependent endocytosis of integrins. *FEBS Lett* 583:1337–1343
- Chao WT, Ashcroft F, Daquinag AC, Vadakkan T, Wei Z, Zhang P, Dickinson ME, Kunz, J (2010) Type I phosphatidylinositol phosphate kinase beta regulates focal adhesion disassembly by promoting beta1 integrin endocytosis. *Mol Cell Biol* 30:4463–4479
- Chao WT, Daquinag AC, Ashcroft F, Kunz, J (2010) Type I PIPK-alpha regulates directed cell migration by modulating Rac1 plasma membrane targeting and activation. *J Cell Biol* 190:247–262
- Chatah NE, Abrams CS (2001) G-protein-coupled receptor activation induces the membrane translocation and activation of phosphatidylinositol-4-phosphate 5-kinase I alpha by a Rac- and Rho-dependent pathway. *J Biol Chem* 276:34059–34065
- Ciruela A, Hinchliffe KA, Divecha N, Irvine RF (2000) Nuclear targeting of the beta isoform of type II phosphatidylinositol phosphate kinase (phosphatidylinositol 5-phosphate 4-kinase) by its alpha-helix 7. *Biochem J* 346(Pt 3):587–591
- Clarke JH, Letcher AJ, D'Santos C S, Halstead JR, Irvine RF, Divecha N (2001) Inositol lipids are regulated during cell cycle progression in the nuclei of murine erythroleukaemia cells. *Biochem J* 357:905–910
- Cocco L, Gilmour RS, Ognibene A, Letcher AJ, Manzoli FA, Irvine RF (1987) Synthesis of polyphosphoinositides in nuclei of Friend cells. Evidence for polyphosphoinositide metabolism inside the nucleus which changes with cell differentiation. *Biochem J* 248:765–770
- Cocco L, Martelli AM, Gilmour RS, Ognibene A, Manzoli FA, Irvine RF (1988) Rapid changes in phospholipid metabolism in the nuclei of Swiss 3T3 cells induced by treatment of the cells with insulin-like growth factor I. *Biochem Biophys Res Commun* 154:1266–1272
- Cooke FT, Dove SK, McEwen RK, Painter G, Holmes AB, Hall MN, Michell RH, Parker PJ (1998) The stress-activated phosphatidylinositol 3-phosphate 5-kinase FabI α is essential for vacuole function in *S. cerevisiae*. *Curr Biol* 8:1219–1222
- Cox S, Taylor SS (1994) Holoenzyme interaction sites in the cAMP-dependent protein kinase. Histidine 87 in the catalytic subunit complements serine 99 in the type I regulatory subunit. *J Biol Chem* 269:22614–22622
- D'Santos CS, Clarke JH, Divecha N (1998) Phospholipid signalling in the nucleus. Een DAG uit het leven van de inositide signalering in de nucleus. *Biochim Biophys Acta* 1436:201–232
- Del Pozo MA, Kiosses WB, Alderson NB, Meller N, Hahn KM, Schwartz MA (2002) Integrins regulate GTP-Rac localized effector interactions through dissociation of Rho-GDI. *Nat Cell Biol* 4:232–239
- Dell'Angelica E, Mullins C, Bonifacino J (1999) AP-4, a novel protein complex related to clathrin adaptors. *J Biol Chem* 274:7278–7285
- DeMali KA, Barlow CA, Burridge K (2002) Recruitment of the Arp2/3 complex to vinculin: coupling membrane protrusion to matrix adhesion. *J Cell Biol* 159:881–891
- Di Paolo G, De Camilli P (2006) Phosphoinositides in cell regulation and membrane dynamics. *Nature* 443:651–657

- Di Paolo G, Pellegrini L, Letinic K, Cestra G, Zoncu R, Voronov S, Chang S, Guo J, Wenk MR, De Camilli P (2002) Recruitment and regulation of phosphatidylinositol phosphate kinase type I gamma by the FERM domain of talin. *Nature* 420:85–89
- Di Paolo G, Moskowitz HS, Gipson K, Wenk MR, Voronov S, Obayashi M, Flavell R, Fitzsimonds RM, Ryan TA, De Camilli P (2004) Impaired PtdIns(4,5)P₂ synthesis in nerve terminals produces defects in synaptic vesicle trafficking. *Nature* 431:415–422
- Didichenko SA, Thelen M (2001) Phosphatidylinositol 3-kinase c2alpha contains a nuclear localization sequence and associates with nuclear speckles. *J Biol Chem* 276:48135–48142
- Divecha N, Roefs M, Los A, Halstead J, Bannister A, D'Santos C (2002) Type I PIPkinases interact with and are regulated by the retinoblastoma susceptibility gene product-pRB. *Curr Biol* 12:582–587
- Doughman RL, Firestone AJ, Anderson RA (2003a) Phosphatidylinositol phosphate kinases put PI4,5P(2) in its place. *J Membr Biol* 194:77–89
- Doughman RL, Firestone AJ, Wojtasiak ML, Bunce MW, Anderson RA (2003b) Membrane ruffling requires coordination between type Ialpha phosphatidylinositol phosphate kinase and Rac signaling. *J Biol Chem* 278:23036–23045
- Dumas F, Byrne RD, Vincent B, Hobday TM, Poccia DL, Larijani B (2010) Spatial regulation of membrane fusion controlled by modification of phosphoinositides. *PLoS One* 5:e12208
- Ezraty EJ, Partridge MA, Gundersen GG (2005) Microtubule-induced focal adhesion disassembly is mediated by dynamin and focal adhesion kinase. *Nat Cell Biol* 7:581–590
- Fölsch H, Ohno H, Bonifacino J, Mellman I (1999) A novel clathrin adaptor complex mediates basolateral targeting in polarized epithelial cells. *Cell* 99:189–198
- Franco SJ, Huttenlocher A (2005) Regulating cell migration: calpains make the cut. *J Cell Sci* 118:3829–3838
- Franco SJ, Rodgers MA, Perrin BJ, Han J, Bennin DA, Critchley DR, Huttenlocher A (2004) Calpain-mediated proteolysis of talin regulates adhesion dynamics. *Nat Cell Biol* 6:977–983
- Gaidarov I, Krupnick JG, Falck JR, Benovic JL, Keen JH (1999) Arrestin function in G protein-coupled receptor endocytosis requires phosphoinositide binding. *Embo J* 18:871–881
- Garnier-Lhomme M, Byrne RD, Hobday TM, Gschmeissner S, Woscholski R, Poccia DL, Dufourc EJ, Larijani B (2009) Nuclear envelope remnants: fluid membranes enriched in sterols and polyphosphoinositides. *PLoS One* 4:e4255
- Gary JD, Wurmser AE, Bonangelino CJ, Weisman LS, Emr SD (1998) Fab1p is essential for PtdIns(3)P 5-kinase activity and the maintenance of vacuolar size and membrane homeostasis. *J Cell Biol* 143:65–79
- Gong LW, Di Paolo G, Diaz E, Cestra G, Diaz ME, Lindau M, De Camilli P, Toomre D (2005) Phosphatidylinositol phosphate kinase type I gamma regulates dynamics of large dense-core vesicle fusion. *Proc Natl Acad Sci U S A* 102:5204–5209
- Gonzales ML, Anderson RA (2006) Nuclear phosphoinositide kinases and inositol phospholipids. *J Cell Biochem* 97:252–260
- Gonzales ML, Mellman DL, Anderson RA (2008) CKIalpha is associated with and phosphorylates star-PAP and is also required for expression of select star-PAP target messenger RNAs. *J Biol Chem* 283:12665–12673
- Gorbatyuk VY, Nosworthy NJ, Robson SA, Bains NP, Maciejewski MW, Dos Remedios CG, King GF (2006) Mapping the phosphoinositide-binding site on chick cofilin explains how PIP₂ regulates the cofilin-actin interaction. *Mol Cell* 24:511–522
- Goss VL, Hocesvar BA, Thompson LJ, Stratton CA, Burns DJ, Fields AP (1994) Identification of nuclear beta II protein kinase C as a mitotic lamin kinase. *J Biol Chem* 269:19074–19080
- Gurr MI, Finean JB, Hawthorne JN (1963) The phospholipids of liver-cell fractions. I. The phospholipid composition of the liver-cell nucleus. *Biochim Biophys Acta* 70:406–416
- Hall A, Nobes CD (2000) Rho GTPases: molecular switches that control the organization and dynamics of the actin cytoskeleton. *Philos Trans R Soc Lond B Biol Sci* 355:965–970
- Hanks SK, Hunter T (1995) Protein kinases 6. The eukaryotic protein kinase superfamily: kinase (catalytic) domain structure and classification. *FASEB J* 9:576–596
- Hanks SK, Quinn AM, Hunter T (1988) The protein kinase family: conserved features and deduced phylogeny of the catalytic domains. *Science* 241:42–52

- Hardingham GE, Chawla S, Johnson CM, Bading H (1997) Distinct functions of nuclear and cytoplasmic calcium in the control of gene expression. *Nature* 385:260–265
- Hay JC, Fisette PL, Jenkins GH, Fukami K, Takenawa T, Anderson RA, Martin TF (1995) ATP-dependent inositide phosphorylation required for Ca^{2+} -activated secretion. *Nature* 374:173–177
- He B, Xi F, Zhang X, Zhang J, Guo W (2007) Exo70 interacts with phospholipids and mediates the targeting of the exocyst to the plasma membrane. *EMBO J* 26:4053–4065
- Heck JN, Mellman DL, Ling K, Sun Y, Wagoner MP, Schill NJ, Anderson RA (2007) A conspicuous connection: structure defines function for the phosphatidylinositol-phosphate kinase family. *Crit Rev Biochem Mol Biol* 42:15–39
- Hinchliffe KA, Irvine RF (2006) Regulation of type II PIP kinase by PKD phosphorylation. *Cell Signal* 18:1906–1913
- Hinchliffe KA, Ciruela A, Letcher AJ, Divecha N, Irvine RF (1999a) Regulation of type II α -phosphatidylinositol phosphate kinase localisation by the protein kinase CK2. *Curr Biol* 9:983–986
- Hinchliffe KA, Ciruela A, Morris JA, Divecha N, Irvine RF (1999b) The type II PIPKins (PtdIns5P 4-kinases): enzymes in search of a function? *Biochem Soc Trans* 27:657–661
- Hokin MR, Hokin LE (1953) Enzyme secretion and the incorporation of P32 into phospholipides of pancreas slices. *J Biol Chem* 203:967–977
- Homma K, Terui S, Minemura M, Qadota H, Anraku Y, Kanaho Y, Ohya Y (1998) Phosphatidylinositol-4-phosphate 5-kinase localized on the plasma membrane is essential for yeast cell morphogenesis. *J Biol Chem* 273:15779–15786
- Honda A, Nogami M, Yokozeki T, Yamazaki M, Nakamura H, Watanabe H, Kawamoto K, Nakayama K, Morris AJ, Frohman MA, Kanaho Y (1999) Phosphatidylinositol 4-phosphate 5-kinase α is a downstream effector of the small G protein ARF6 in membrane ruffle formation. *Cell* 99:521–532
- Höning S, Ricotta D, Krauss M, Späte K, Spolaore B, Motley A, Robinson M, Robinson C, Haucke V, Owen D (2005) Phosphatidylinositol-(4,5)-bisphosphate regulates sorting signal recognition by the clathrin-associated adaptor complex AP2. *Mol Cell* 18:519–531
- Humbert JP, Matter N, Artault JC, Koppler P, Malviya AN (1996) Inositol 1,4,5-trisphosphate receptor is located to the inner nuclear membrane vindicating regulation of nuclear calcium signaling by inositol 1,4,5-trisphosphate. Discrete distribution of inositol phosphate receptors to inner and outer nuclear membranes. *J Biol Chem* 271:478–485
- Irvine RF (2003) Nuclear lipid signalling. *Nat Rev Mol Cell Biol* 4:349–360
- Ishihara H, Shibasaki Y, Kizuki N, Katagiri H, Yazaki Y, Asano T, Oka Y (1996) Cloning of cDNAs encoding two isoforms of 68-kDa type I phosphatidylinositol-4-phosphate 5-kinase. *J Biol Chem* 271:23611–23614
- Ishihara H, Shibasaki Y, Kizuki N, Wada T, Yazaki Y, Asano T, Oka Y (1998) Type I phosphatidylinositol-4-phosphate 5-kinases. Cloning of the third isoform and deletion/substitution analysis of members of this novel lipid kinase family. *J Biol Chem* 273:8741–8748
- Itoh T, Ishihara H, Shibasaki Y, Oka Y, Takenawa T (2000) Autophosphorylation of type I phosphatidylinositol phosphate kinase regulates its lipid kinase activity. *J Biol Chem* 275:19389–19394
- Iyer GH, Moore MJ, Taylor SS (2005) Consequences of lysine 72 mutation on the phosphorylation and activation state of cAMP-dependent kinase. *J Biol Chem* 280:8800–8807
- Jackson T (1998) Transport vesicles: coats of many colours. *Curr Biol* 8:R609–R612
- Janmey PA, Lindberg U (2004) Cytoskeletal regulation: rich in lipids. *Nat Rev Mol Cell Biol* 5:658–666
- Jenkins GH, Subrahmanyam G, Anderson RA (1991) Purification and reconstitution of phosphatidylinositol 4-kinase from human erythrocytes. *Biochim Biophys Acta* 1080:11–18
- Jenkins GH, Fisette PL, Anderson RA (1994) Type I phosphatidylinositol 4-phosphate 5-kinase isoforms are specifically stimulated by phosphatidic acid. *J Biol Chem* 269:11547–11554
- Johnson LN, Noble ME, Owen DJ (1996) Active and inactive protein kinases: structural basis for regulation. *Cell* 85:149–158

- Jones DR, Bultsma Y, Keune WJ, Halstead JR, Elouarrat D, Mohammed S, Heck AJ, D'Santos CS, Divecha N (2006) Nuclear PtdIns5P as a transducer of stress signaling: an in vivo role for PIP4Kbeta. *Mol Cell* 23:685–695
- Kahlfeldt N, Vahedi-Faridi A, Koo S, Schäfer J, Krainer G, Keller S, Saenger W, Krauss M, Haucke V (2010) Molecular basis for association of PIPKI gamma-p90 with clathrin adaptor AP-2. *J Biol Chem* 285:2734–2749
- Keune W, Bultsma Y, Sommer L, Jones D, Divecha N (2010) Phosphoinositide signalling in the nucleus. *Adv Enzyme Regul* 51(1):91–99
- Kisseleva M, Feng Y, Ward M, Song C, Anderson RA, Longmore GD (2005) The LIM protein Ajuba regulates phosphatidylinositol 4,5-bisphosphate levels in migrating cells through an interaction with and activation of PIPKI alpha. *Mol Cell Biol* 25:3956–3966
- Kleinig H (1970) Nuclear membranes from mammalian liver. II. Lipid composition. *J Cell Biol* 46:396–402
- Knighton DR, Zheng JH, Ten Eyck LF, Ashford VA, Xuong NH, Taylor SS, Sowadski JM (1991a) Crystal structure of the catalytic subunit of cyclic adenosine monophosphate-dependent protein kinase. *Science* 253:407–414
- Knighton DR, Zheng JH, Ten Eyck LF, Xuong NH, Taylor SS, Sowadski JM (1991b) Structure of a peptide inhibitor bound to the catalytic subunit of cyclic adenosine monophosphate-dependent protein kinase. *Science* 253:414–420
- Krauss M, Kinuta M, Wenk MR, De Camilli P, Takei K, Haucke V (2003) ARF6 stimulates clathrin/AP-2 recruitment to synaptic membranes by activating phosphatidylinositol phosphate kinase type Igamma. *J Cell Biol* 162:113–124
- Krauss M, Kukhtina V, Pechstein A, Haucke V (2006) Stimulation of phosphatidylinositol kinase type I-mediated phosphatidylinositol (4,5)-bisphosphate synthesis by AP-2mu-cargo complexes. *Proc Natl Acad Sci U S A* 103:11934–11939
- Krupa A, Preethi G, Srinivasan N (2004) Structural modes of stabilization of permissive phosphorylation sites in protein kinases: distinct strategies in Ser/Thr and Tyr kinases. *J Mol Biol* 339:1025–1039
- Kumar N, Khurana S (2004) Identification of a functional switch for actin severing by cytoskeletal proteins. *J Biol Chem* 279:24915–24918
- Kumar N, Zhao P, Tomar A, Galea CA, Khurana S (2004) Association of villin with phosphatidylinositol 4,5-bisphosphate regulates the actin cytoskeleton. *J Biol Chem* 279:3096–3110
- Kunz J, Wilson MP, Kisseleva M, Hurley JH, Majerus PW, Anderson RA (2000) The activation loop of phosphatidylinositol phosphate kinases determines signaling specificity. *Mol Cell* 5:1–11
- Kunz J, Fuelling A, Kolbe L, Anderson RA (2002) Stereo-specific substrate recognition by phosphatidylinositol phosphate kinases is swapped by changing a single amino acid residue. *J Biol Chem* 277:5611–5619
- Kwiatkowska K (2010) One lipid, multiple functions: how various pools of PI(4,5)P(2) are created in the plasma membrane. *Cell Mol Life Sci* 67:3927–3946
- Lambrechts A, Jonckheere V, Dewitte D, Vandekerckhove J, Ampe C (2002) Mutational analysis of human profilin I reveals a second PI(4,5)-P2 binding site neighbouring the poly(L-proline) binding site. *BMC Biochem* 3:12
- Lamond AI, Spector DL (2003) Nuclear speckles: a model for nuclear organelles. *Nat Rev Mol Cell Biol* 4:605–612
- Lassing I, Lindberg U (1985) Specific interaction between phosphatidylinositol 4,5-bisphosphate and profilactin. *Nature* 314:472–474
- Lee S, Voronov S, Letinic K, Nairn A, Di Paolo G, De Camilli P (2005) Regulation of the interaction between PIPKI gamma and talin by proline-directed protein kinases. *J Cell Biol* 168:789–799
- Letinic K, Sebastian R, Toomre D, Rakic P (2009) Exocyst is involved in polarized cell migration and cerebral cortical development. *Proc Natl Acad Sci U S A* 106:11342–11347
- Ling K, Doughman RL, Firestone AJ, Bunce MW, Anderson RA (2002) Type I gamma phosphatidylinositol phosphate kinase targets and regulates focal adhesions. *Nature* 420:89–93
- Ling K, Doughman RL, Iyer VV, Firestone AJ, Bairstow SF, Mosher DF, Schaller MD, Anderson RA (2003) Tyrosine phosphorylation of type Igamma phosphatidylinositol phosphate kinase by Src regulates an integrin-talin switch. *J Cell Biol* 163:1339–1349

- Ling K, Schill NJ, Wagoner MP, Sun Y, Anderson RA (2006) Movin' on up: the role of PtdIns(4,5)P(2) in cell migration. *Trends Cell Biol* 16:276–284
- Ling K, Bairstow S, Carbonara C, Turbin D, Huntsman D, Anderson R (2007) Type I gamma phosphatidylinositol phosphate kinase modulates adherens junction and E-cadherin trafficking via a direct interaction with mu 1B adaptin. *J Cell Biol* 176:343–353
- Liu J, Zuo X, Yue P, Guo W (2007) Phosphatidylinositol 4,5-bisphosphate mediates the targeting of the exocyst to the plasma membrane for exocytosis in mammalian cells. *Mol Biol Cell* 18:4483–4492
- Loijens JC, Boronenkov IV, Parker GJ, Anderson RA (1996) The phosphatidylinositol 4-phosphate 5-kinase family. *Adv Enzyme Regul* 36:115–140
- Lokuta MA, Senetar MA, Bennin DA, Nuzzi PA, Chan KT, Ott VL, Huttenlocher A (2007) Type I gamma PIP kinase is a novel uropod component that regulates rear retraction during neutrophil chemotaxis. *Mol Biol Cell* 18:5069–5080
- Malviya AN, Rogue PJ (1998) “Tell me where is calcium bred”: clarifying the roles of nuclear calcium. *Cell* 92:17–23
- Martel V, Racaud-Sultan C, Dupe S, Marie C, Paulhe F, Galmiche A, Block MR, Albiges-Rizo C (2001) Conformation, localization, and integrin binding of talin depend on its interaction with phosphoinositides. *J Biol Chem* 276:21217–21227
- Matsui T, Yonemura S, Tsukita S, Tsukita S (1999) Activation of ERM proteins in vivo by Rho involves phosphatidyl-inositol 4-phosphate 5-kinase and not ROCK kinases. *Curr Biol* 9:1259–1262
- McEwen RK, Dove SK, Cooke FT, Painter GF, Holmes AB, Shisheva A, Ohya Y, Parker PJ, Michell RH (1999) Complementation analysis in PtdInsP kinase-deficient yeast mutants demonstrates that *Schizosaccharomyces pombe* and murine Fab1p homologues are phosphatidylinositol 3-phosphate 5-kinases. *J Biol Chem* 274:33905–33912
- Mellman DL, Gonzales ML, Song C, Barlow CA, Wang P, Kendzierski C, Anderson RA (2008) A PtdIns4,5P2-regulated nuclear poly(A) polymerase controls expression of select mRNAs. *Nature* 451:1013–1017
- Millard TH, Sharp SJ, Machesky LM (2004) Signalling to actin assembly via the WASP (Wiskott-Aldrich syndrome protein)-family proteins and the Arp2/3 complex. *Biochem J* 380:1–17
- Moritz A, De Graan PN, Gispén WH, Wirtz KW (1992a) Phosphatidic acid is a specific activator of phosphatidylinositol-4-phosphate kinase. *J Biol Chem* 267:7207–7210
- Moritz A, Westerman J, De Graan PN, Wirtz KW (1992b) Phosphatidylinositol 4-kinase and phosphatidylinositol-4-phosphate 5-kinase from bovine brain membranes. *Methods Enzymol* 209:202–211
- Nakano-Kobayashi A, Yamazaki M, Unoki T, Hongu T, Murata C, Taguchi R, Katada T, Frohman M, Yokozeki T, Kanaho Y (2007) Role of activation of PIP5Kgamma661 by AP-2 complex in synaptic vesicle endocytosis. *EMBO J* 26:1105–1116
- Nakatsu F, Ohno H (2003) Adaptor protein complexes as the key regulators of protein sorting in the post-Golgi network. *Cell Struct Funct* 28:419–429
- Narkis G, Ofir R, Manor E, Landau D, Elbedour K, Birk OS (2007) Lethal congenital contractural syndrome type 2 (LCCS2) is caused by a mutation in ERBB3 (Her3), a modulator of the phosphatidylinositol-3-kinase/Akt pathway. *Am J Hum Genet* 81:589–595
- Nayal A, Webb DJ, Horwitz AF (2004) Talin: an emerging focal point of adhesion dynamics. *Curr Opin Cell Biol* 16:94–98
- Nicholson-Dykstra S, Higgs HN, Harris ES (2005) Actin dynamics: growth from dendritic branches. *Curr Biol* 15:R346–R357
- Niggli V (2005) Regulation of protein activities by phosphoinositide phosphates. *Annu Rev Cell Dev Biol* 21:57–79
- Odorizzi G, Babst M, Emr SD (1998) Fab1p PtdIns(3)P 5-kinase function essential for protein sorting in the multivesicular body. *Cell* 95:847–858
- Ohno H (2006) Physiological roles of clathrin adaptor AP complexes: lessons from mutant animals. *J Biochem* 139:943–948

- Olusanya O, Andrews P, Swedlow J, Smythe E (2001) Phosphorylation of threonine 156 of the $\mu 2$ subunit of the AP2 complex is essential for endocytosis in vitro and in vivo. *Curr Biol* 11:896–900
- Osborne SL, Thomas CL, Gschmeissner S, Schiavo G (2001) Nuclear PtdIns(4,5)P₂ assembles in a mitotically regulated particle involved in pre-mRNA splicing. *J Cell Sci* 114:2501–2511
- Oztan A, Silvis M, Weisz OA, Bradbury NA, Hsu SC, Goldenring JR, Yeaman C, Apodaca G (2007) Exocyst requirement for endocytic traffic directed toward the apical and basolateral poles of polarized MDCK cells. *Mol Biol Cell* 18:3978–3992
- Park SJ, Itoh T, Takenawa T (2001) Phosphatidylinositol 4-phosphate 5-kinase type I is regulated through phosphorylation response by extracellular stimuli. *J Biol Chem* 276:4781–4787
- Payraastre B, Nievers M, Boonstra J, Breton M, Verkleij AJ, Van Bergen en Henegouwen PM (1992) A differential location of phosphoinositide kinases, diacylglycerol kinase, and phospholipase C in the nuclear matrix. *J Biol Chem* 267:5078–5084
- Pierini LM, Lawson MA, Eddy RJ, Hendey B, Maxfield FR (2000) Oriented endocytic recycling of $\alpha 5\beta 1$ in motile neutrophils. *Blood* 95:2471–2480
- Prehoda KE, Scott JA, Mullins RD, Lim WA (2000) Integration of multiple signals through cooperative regulation of the N-WASP-Arp2/3 complex. *Science* 290:801–806
- Prigozhina NL, Waterman-Storer CM (2004) Protein kinase D-mediated anterograde membrane trafficking is required for fibroblast motility. *Curr Biol* 14:88–98
- Qualmann B, Kessels MM, Kelly RB (2000) Molecular links between endocytosis and the actin cytoskeleton. *J Cell Biol* 150:F111–F116
- Rameh LE, Tolias KF, Duckworth BC, Cantley LC (1997) A new pathway for synthesis of phosphatidylinositol-4,5-bisphosphate. *Nature* 390:192–196
- Rando OJ, Zhao K, Janney P, Crabtree GR (2002) Phosphatidylinositol-dependent actin filament binding by the SWI/SNF-like BAF chromatin remodeling complex. *Proc Natl Acad Sci U S A* 99:2824–2829
- Rao VD, Misra S, Boronenkov IV, Anderson RA, Hurley JH (1998) Structure of type II β phosphatidylinositol phosphate kinase: a protein kinase fold flattened for interfacial phosphorylation. *Cell* 94:829–839
- Rohde G, Wenzel D, Haucke V (2002) A phosphatidylinositol (4,5)-bisphosphate binding site within $\mu 2$ -adaptin regulates clathrin-mediated endocytosis. *J Cell Biol* 158:209–214
- Rosse C, Hatzoglou A, Parrini MC, White MA, Chavrier P, Camonis J (2006) RalB mobilizes the exocyst to drive cell migration. *Mol Cell Biol* 26:727–734
- Rozelle AL, Machesky LM, Yamamoto M, Driessens MH, Insall RH, Roth MG, Luby-Phelps K, Marriott G, Hall A, Yin HL (2000) Phosphatidylinositol 4,5-bisphosphate induces actin-based movement of raft-enriched vesicles through WASP-Arp2/3. *Curr Biol* 10:311–320
- Saraste M, Sibbald PR, Wittinghofer A (1990) The P-loop—a common motif in ATP- and GTP-binding proteins. *Trends Biochem Sci* 15:430–434
- Sasaki J, Sasaki T, Yamazaki M, Matsuoka K, Taya C, Shitara H, Takasuga S, Nishio M, Mizuno K, Wada T et al (2005) Regulation of anaphylactic responses by phosphatidylinositol phosphate kinase type I $\{\alpha\}$. *J Exp Med* 201:859–870
- Sbrissa D, Ikonomov OC, Shisheva A (1999) PIKfyve, a mammalian ortholog of yeast Fab1p lipid kinase, synthesizes 5-phosphoinositides. Effect of insulin. *J Biol Chem* 274:21589–21597
- Sbrissa D, Ikonomov OC, Shisheva A (2000) PIKfyve lipid kinase is a protein kinase: down-regulation of 5'-phosphoinositide product formation by autophosphorylation. *Biochemistry* 39:15980–15989
- Schill N, Anderson R (2009a) Out, in and back again: PtdIns(4,5)P₂ regulates cadherin trafficking in epithelial morphogenesis. *Biochem J* 418:247–260
- Schill NJ, Anderson RA (2009b) Two novel phosphatidylinositol-4-phosphate 5-kinase type I γ splice variants expressed in human cells display distinctive cellular targeting. *Biochem J* 422(3):473–482
- Schmoranzler J, Kreitzer G, Simon SM (2003) Migrating fibroblasts perform polarized, microtubule-dependent exocytosis towards the leading edge. *J Cell Sci* 116:4513–4519

- Schramm M, Ying O, Kim TY, Martin GS (2008) ERK5 promotes Src-induced podosome formation by limiting Rho activation. *J Cell Biol* 181:1195–1210
- Sechi AS, Wehland J (2000) The actin cytoskeleton and plasma membrane connection: PtdIns(4,5)P(2) influences cytoskeletal protein activity at the plasma membrane. *J Cell Sci* 113(Pt 21):3685–3695
- Shibasaki Y, Ishihara H, Kizuki N, Asano T, Oka Y, Yazaki Y (1997) Massive actin polymerization induced by phosphatidylinositol-4-phosphate 5-kinase in vivo. *J Biol Chem* 272:7578–7581
- Shisheva A, Sbrissa D, Ikononov O (1999) Cloning, characterization, and expression of a novel Zn²⁺-binding FYVE finger-containing phosphoinositide kinase in insulin-sensitive cells. *Mol Cell Biol* 19:623–634
- Sidani M, Wessels D, Mouneimne G, Ghosh M, Goswami S, Sarmiento C, Wang W, Kuhl S, El-Sibai M, Backer JM et al (2007) Cofilin determines the migration behavior and turning frequency of metastatic cancer cells. *J Cell Biol* 179:777–791
- Skare P, Karlsson R (2002) Evidence for two interaction regions for phosphatidylinositol(4,5)-bisphosphate on mammalian profilin I. *FEBS Lett* 522:119–124
- Smith CD, Wells WW (1983a) Phosphorylation of rat liver nuclear envelopes. I. Characterization of in vitro protein phosphorylation. *J Biol Chem* 258:9360–9367
- Smith CD, Wells WW (1983b) Phosphorylation of rat liver nuclear envelopes. II. Characterization of in vitro lipid phosphorylation. *J Biol Chem* 258:9368–9373
- Smith CD, Wells WW (1984a) Characterization of a phosphatidylinositol 4-phosphate-specific phosphomonoesterase in rat liver nuclear envelopes. *Arch Biochem Biophys* 235:529–537
- Smith CD, Wells WW (1984b) Solubilization and reconstitution of a nuclear envelope-associated ATPase. Synergistic activation by RNA and polyphosphoinositides. *J Biol Chem* 259:11890–11894
- Streb H, Irvine RF, Berridge MJ, Schulz I (1983) Release of Ca²⁺ from a nonmitochondrial intracellular store in pancreatic acinar cells by inositol-1,4,5-trisphosphate. *Nature* 306:67–69
- Sugimoto H, Sugahara M, Fölsch H, Koide Y, Nakatsu F, Tanaka N, Nishimura T, Furukawa M, Mullins C, Nakamura N et al (2002) Differential recognition of tyrosine-based basolateral signals by AP-1B subunit mu1B in polarized epithelial cells. *Mol Biol Cell* 13:2374–2382
- Sun Y, Ling K, Wagoner MP, Anderson RA (2007) Type I gamma phosphatidylinositol phosphate kinase is required for EGF-stimulated directional cell migration. *J Cell Biol* 178:297–308
- Suriano G, Mulholland D, De Wever O, Ferreira P, Mateus A, Bruyneel E, Nelson C, Mareel M, Yokota J, Huntsman D, Seruca R (2003) The intracellular E-cadherin germline mutation V832 M lacks the ability to mediate cell-cell adhesion and to suppress invasion. *Oncogene* 22:5716–5719
- Takenawa T, Itoh T (2001) Phosphoinositides, key molecules for regulation of actin cytoskeletal organization and membrane traffic from the plasma membrane. *Biochim Biophys Acta* 1533:190–206
- Tall EG, Spector I, Pentyala SN, Bitter I, Rebecchi MJ (2000) Dynamics of phosphatidylinositol 4,5-bisphosphate in actin-rich structures. *Curr Biol* 10:743–746
- Tayeb MA, Skalski M, Cha MC, Kean MJ, Scaife M, Coppolino MG (2005) Inhibition of SNARE-mediated membrane traffic impairs cell migration. *Exp Cell Res* 305:63–73
- Taylor SS, Knighton DR, Zheng J, Ten Eyck LF, Sowadski JM (1992) Structural framework for the protein kinase family. *Annu Rev Cell Biol* 8:429–462
- Taylor SS, Radzio-Andzelm E, Knighton DR, Ten Eyck LF, Sowadski JM, Herberg FW, Yonemoto W, Zheng J (1993) Crystal structures of the catalytic subunit of cAMP-dependent protein kinase reveal general features of the protein kinase family. *Receptor* 3:165–172
- Thieman JR, Mishra SK, Ling K, Doray B, Anderson RA, Traub LM (2009) Clathrin regulates the association of PIPKgamma661 with the AP-2 adaptor beta2 appendage. *J Biol Chem* 284:13924–13939
- Thompson LJ, Fields AP (1996) betaII protein kinase C is required for the G2/M phase transition of cell cycle. *J Biol Chem* 271:15045–15053
- Tolias KF, Couvillon AD, Cantley LC, Carpenter CL (1998) Characterization of a Rac1- and RhoGDI-associated lipid kinase signaling complex. *Mol Cell Biol* 18:762–770

- Tolias KF, Hartwig JH, Ishihara H, Shibasaki Y, Cantley LC, Carpenter CL (2000) Type I α phosphatidylinositol-4-phosphate 5-kinase mediates Rac-dependent actin assembly. *Curr Biol* 10:153–156
- Van Horck FP, Lavazais E, Eickholt BJ, Moolenaar WH, Divecha N (2002) Essential role of type I α phosphatidylinositol 4-phosphate 5-kinase in neurite remodeling. *Curr Biol* 12:241–245
- Van Rheeën J, Song X, Van Roosmalen W, Cammer M, Chen X, Desmarais V, Yip SC, Backer JM, Eddy RJ, Condeelis JS (2007) EGF-induced PIP₂ hydrolysis releases and activates cofilin locally in carcinoma cells. *J Cell Biol* 179:1247–1259
- Vann LR, Wooding FB, Irvine RF, Divecha N (1997) Metabolism and possible compartmentalization of inositol lipids in isolated rat-liver nuclei. *Biochem J* 327(Pt 2):569–576
- Webb DJ, Parsons JT, Horwitz AF (2002) Adhesion assembly, disassembly and turnover in migrating cells—over and over and over again. *Nat Cell Biol* 4:E97–E100
- Weernink PA, Meletiadis K, Hommeltenberg S, Hinz M, Ishihara H, Schmidt M, Jakobs KH (2004) Activation of type I phosphatidylinositol 4-phosphate 5-kinase isoforms by the Rho GTPases, RhoA, Rac1, and Cdc42. *J Biol Chem* 279:7840–7849
- Wenk MR, De Camilli P (2004) Protein-lipid interactions and phosphoinositide metabolism in membrane traffic: insights from vesicle recycling in nerve terminals. *Proc Natl Acad Sci U S A* 101:8262–8269
- Wenk MR, Pellegrini L, Klenchin VA, Di Paolo G, Chang S, Daniell L, Arioka M, Martin TF, De Camilli P (2001) PIP kinase Igamm is the major PI(4,5)P₂ synthesizing enzyme at the synapse. *Neuron* 32:79–88
- Whitman M, Downes CP, Keeler M, Keller T, Cantley L (1988) Type I phosphatidylinositol kinase makes a novel inositol phospholipid, phosphatidylinositol-3-phosphate. *Nature* 332:644–646
- Yabuta T, Shinmura K, Tani M, Yamaguchi S, Yoshimura K, Katai H, Nakajima T, Mochiki E, Tsujinaka T, Takami M et al (2002) E-cadherin gene variants in gastric cancer families whose probands are diagnosed with diffuse gastric cancer. *Int J Cancer* 101:434–441
- Yamamoto A, DeWald DB, Boronenkov IV, Anderson RA, Emr SD, Koshland D (1995) Novel PI(4)P 5-kinase homologue, Fab1p, essential for normal vacuole function and morphology in yeast. *Mol Biol Cell* 6:525–539
- Yamamoto M, Hilgemann DH, Feng S, Bito H, Ishihara H, Shibasaki Y, Yin HL (2001) Phosphatidylinositol 4,5-bisphosphate induces actin stress-fiber formation and inhibits membrane ruffling in CV1 cells. *J Cell Biol* 152:867–876
- Yamazaki M, Miyazaki H, Watanabe H, Sasaki T, Maehama T, Frohman MA, Kanaho Y (2002) Phosphatidylinositol 4-phosphate 5-kinase is essential for ROCK-mediated neurite remodeling. *J Biol Chem* 277:17226–17230
- Yap A, Crampton M, Hardin J (2007) Making and breaking contacts: the cellular biology of cadherin regulation. *Curr Opin Cell Biol* 19:508–514
- Yeaman C, Grindstaff KK, Wright JR, Nelson WJ (2001) Sec6/8 complexes on trans-Golgi network and plasma membrane regulate late stages of exocytosis in mammalian cells. *J Cell Biol* 155:593–604
- Yin HL, Janney PA (2003) Phosphoinositide regulation of the actin cytoskeleton. *Annu Rev Physiol* 65:761–789
- Yokogawa T, Nagata S, Nishio Y, Tsutsumi T, Ihara S, Shirai R, Morita K, Umeda M, Shirai Y, Saitoh N, Fukui Y (2000) Evidence that 3'-phosphorylated polyphosphoinositides are generated at the nuclear surface: use of immunostaining technique with monoclonal antibodies specific for PI 3,4-P₂. *FEBS Lett* 473:222–226
- Yoshinaga-Ohara N, Takahashi A, Uchiyama T, Sasada M (2002) Spatiotemporal regulation of moesin phosphorylation and rear release by Rho and serine/threonine phosphatase during neutrophil migration. *Exp Cell Res* 278:112–122
- Yu H, Fukami K, Watanabe Y, Ozaki C, Takenawa T (1998) Phosphatidylinositol 4,5-bisphosphate reverses the inhibition of RNA transcription caused by histone H1. *Eur J Biochem* 251:281–287
- Zhang X, Loijens JC, Boronenkov IV, Parker GJ, Norris FA, Chen J, Thum O, Prestwich GD, Majerus PW, Anderson RA (1997) Phosphatidylinositol-4-phosphate 5-kinase isozymes catalyze the

- synthesis of 3-phosphate-containing phosphatidylinositol signaling molecules. *J Biol Chem* 272:17756–17761
- Zhang X, Orlando K, He B, Xi F, Zhang J, Zajac A, Guo W (2008) Membrane association and functional regulation of Sec3 by phospholipids and Cdc42. *J Cell Biol* 180:145–158
- Zhao K, Wang W, Rando OJ, Xue Y, Swiderek K, Kuo A, Crabtree GR (1998) Rapid and phosphoinositol-dependent binding of the SWI/SNF-like BAF complex to chromatin after T lymphocyte receptor signaling. *Cell* 95:625–636
- Zheng JH, Knighton DR, Parello J, Taylor SS, Sowadski JM (1991) Crystallization of catalytic subunit of adenosine cyclic monophosphate-dependent protein kinase. *Methods Enzymol* 200:508–521
- Zou J, Marjanovic J, Kisseleva MV, Wilson M, Majerus PW (2007) Type I phosphatidylinositol-4,5-bisphosphate 4-phosphatase regulates stress-induced apoptosis. *Proc Natl Acad Sci U S A* 104:16834–16839
- Zovein AC, Luque A, Turlo KA, Hofmann JJ, Yee KM, Becker MS, Fassler R, Mellman I, Lane TF, Iruela-Arispe ML (2010) Beta1 integrin establishes endothelial cell polarity and arteriolar lumen formation via a Par3-dependent mechanism. *Dev Cell* 18:39–51

Phosphoinositides I: Enzymes of Synthesis and
Degradation

Balla, T.; Wymann, M.; York, J.D. (Eds.)

2012, XVI, 356 p., Hardcover

ISBN: 978-94-007-3011-3