

Preface

When I was approached to shape a book about phosphoinositide signaling, I first felt honored and humbled. On second thought, this appeared to be an impossible task. Phosphoinositides have grown from being just a curious lipid fraction isolated from bovine brain, showing increased radioactive metabolic labeling during intense stimulation protocols, to become the focus of immense interest as key regulatory molecules that penetrate every aspect of eukaryotic biology. The expansion of this field in the last three decades has been enormous: it turned from a basic science exercise of a devoted few to highly translatable science relevant to a large number of human diseases (isn't this the nature of good basic science?). These include cancer, metabolic-, immuno- and neurodegenerative disorders, to name just a few. Reviewing the large number of enzymes that convert phosphoinositides would fill a book—let alone the diverse biological processes in which phosphoinositides play key regulatory roles. Given the interest, a collection of up-to-date reviews compiled in a book is clearly warranted, which was enough to sway me to accept this assignment. As one editor is unable to handle this enormous task, I was delighted when Matthias Wymann and John York were kind enough to join me in this ambitious effort.

When thinking about potential authors, the obvious choice would have been to approach the people whose contributions have been crucial to push and elevate this field to the level it is today. Bob Michell, prophetically placed phosphoinositides in the center of signal transduction in a 1975 *Biochem. Biophys. Acta* review (Michell 1975), Michael Berridge had a key role in linking phosphoinositides and Ca^{2+} signaling and whose fascinating reviews have inspired many of us (Berridge and Irvine 1984). Robin Irvine, whose group found that InsP_3 was a mixture of two isomers, the active $\text{Ins}(1,4,5)\text{P}_3$ and an inactive $\text{Ins}(1,3,4)\text{P}_3$, and who described the tetrakisphosphate pathway (Irvine et al. 1986), and who always challenges us with most provocative ideas. Philip Majerus, who has insisted on the importance of inositide phosphatases (Majerus et al. 1999) very early on. The group of Lewis Cantley, with the discovery of PI 3-kinase activities and the mapping of downstream effectors (Whitman et al. 1988; Franke et al. 1997), or the Waterfield lab where the first PI 3-kinase catalytic subunit was isolated and cloned (Otsu et al. 1991; Hiles et al. 1992). Peter Downes, who recognized the translational value of phosphoinositide research. Jeremy Thorner and Scott Emr, whose work in baker's yeast still forms the

foundation of our understanding of the role of inositol lipids in trafficking (Strahl and Thorner 2007) or Pietro De Camilli, whose group documented the central role of inositides in brain and synaptic biology (Cremona et al. 1999). There are many others who made valuable or even greater contributions to phosphoinositide research. The above list reflects my bias, as these researchers had the largest impact on my thinking and the directions of my work. Research is, however, a constantly evolving process and we (now Matthias and John being involved) wanted to involve contributions of scientists who represent a second or third wave of researchers infected with the interest in phosphoinositides. We made an effort to recruit authors who have been trainees of these founding laboratories. With this selection our goal was to sample the view of the current and future generation. By selecting their trainees, we feel that we pay tribute to the “Founding Fathers”, and show that the research they put in motion is alive and continues with fresh ideas, new ambitions and a translational and therapeutic value.

Phosphoinositide research in the 1980s went hand in hand with research on Ca^{2+} signaling pursued in “non-excitable” cells and was also marked with the discovery of the family of protein kinase C enzymes, regulated by diacylglycerol, one of the products of phosphoinositide-specific phospholipase C enzymes. These areas of research developed and expanded to form their own fields, and could not be discussed here in detail—even though they are linked historically to the development of phosphoinositide signaling. The enormous work of the groups of Yasutomi Nishizuka on protein kinase C, and Katsuhiko Mikoshiba on cloning and characterizing the $\text{Ins}(1,4,5)\text{P}_3$ receptors are prime examples of these achievements. Although we could not cover all these areas, we included a chapter on Ca^{2+} signaling via the $\text{Ins}(1,4,5)\text{P}_3$ receptor by Colin Taylor, a trainee of the Michael Berridge’s lab, where important links between Ca^{2+} release and $\text{Ins}(1,4,5)\text{P}_3$ receptor signaling were discovered. We also decided to allocate some space to inositol phosphates, the soluble counterparts of some of the phosphoinositides. These molecules for long had been viewed only as the metabolic products of the second messenger $\text{Ins}(1,4,5)\text{P}_3$ but recently gained significant prominence as regulators of important physiological processes. With the discovery of the highly phosphorylated and pyrophosphorylated inositols and the enzymes that produce them, it became clear that this system represents a whole new regulatory paradigm with exciting new developments.

Finally, it was a difficult dilemma whether to include a Chapter on the early history of phosphoinositides. We decided against it for a number of reasons. First, the really interesting history is traced back to studies that preceded the landmark 1975 Bob Michell review and included the work of the Hokins (1987), Bernard Agranoff (2009) and other pioneers of phosphoinositide research. Nobody could tell these early developments better than Bob Michell in his several recollections (Michell 1995) or Robin Irvine who commemorated the 20 years of $\text{Ins}(1,4,5)\text{P}_3$ and the period leading to its discovery (Irvine 2003). We encourage the young readers to go back and read these recollections, as they show several examples of how seemingly uninspiring observations formed the beginning of something that became huge as it unfolded. What came after these landmark discoveries is so overwhelming that each one of us has own views and subjective memories and stories to tell on some aspects of

it. As Editors we felt that our views should not be elevated above others on these historical aspects, and leave it to the authors of the individual Chapters to elucidate the diversity in this respect. The only exception is a Chapter on the history of PI 3-kinases by Alex Toker that we felt deserves special emphasis as it had the most transforming impact on the field since the late 1980s.

One needs to understand that selection of authors is a subjective process and does not always reflect on who contributed the most in a selected field. However, we are confident that proper credit is given in the individual Chapters to each groups and individuals whose work has moved this field forward. It should also be understood that a field that generates over 10,000 entries in PubMed with each keyword that relates to phosphoinositides cannot be covered without missing some aspects that could be important. However, we trust that this collection will be found useful for both the experts and the novices.

References

- Agranoff BW (2009) Turtles all the way: Reflections on myo-Inositol. *J Biol Chem* 284(32):21121–21126
- Berridge MJ, Irvine RF (1984) Inositol trisphosphate, a novel second messenger in cellular signal transduction. *Nature* 312:315–321
- Cremona O, et al (1999) Essential role of phosphoinositide metabolism in synaptic vesicle recycling. *Cell* 99:179–188
- Franke TF, Kaplan DR, Cantley LC, Toker A (1997) Direct regulation of the Akt protooncogene product by PI3,4P2. *Science* 275:665–668
- Hiles ID et al (1992) Phosphatidylinositol 3-kinase: structure and expression of the 110 kDa catalytic subunit. *Cell* 70:419–429
- Hokin LE (1987) The road to the phosphoinositide-generated second messengers. *Trends Pharmacol Sci* 8:53–56
- Irvine RF (2003) 20 years of Ins(1,4,5)P3, and 40 years before. *Nat Rev Mol Cell Biol* 4(7):586–590
- Irvine RF, Letcher AJ, Heslop JP, Berridge MJ (1986) The inositol tris/tetrakis phosphate pathway—demonstration of inositol (1,4,5)trisphosphate-3-kinase activity in mammalian tissues. *Nature* 320:631–634
- Majerus PW, Kisseleva MV, Norris FA (1999) The role of phosphatases in inositol signaling reactions. *J. Biol Chem* 274:10669–10672
- Michell B (1995) Early steps along the road to inositol-lipid-based signalling. *Trends Biochem Sci* 20(8):326–329
- Michell RH (1975) Inositol phospholipids and cell surface receptor function. *Biochim Biophys Acta* 415:81–147
- Otsu M et al (1991) Characterization of two 85 kDa proteins that associate with receptor tyrosine kinases, middle-T/pp60c-src complexes, and PI3-kinase. *Cell* 65:91–104
- Strahl T, Thorner J (2007) Synthesis and function of membrane phosphoinositides in budding yeast, *Saccharomyces cerevisiae*. *Biochim Biophys Acta* 1771(3):353–404
- Whitman M, Downes CP, Keeler M, Keller T, Cantley L (1988) Type-I phosphatidylinositol kinase makes a novel inositol phospholipid, phosphatidylinositol-3-phosphate. *Nature* 332:644–646

Phosphoinositides II: The Diverse Biological Functions

Balla, T.; Wymann, M.; York, J.D. (Eds.)

2012, XVI, 460 p., Hardcover

ISBN: 978-94-007-3014-4