

Chapter 2

Revisiting Complexation Between DNA and Polyethylenimine: The Effect of Uncomplexed Chains Free in the Solution Mixture on Gene Transfection

2.1 Introduction

In comparison with viral vectors, more efforts have recently been spent on the development of non-viral vectors because of few fatal accidents in clinical trials of viral carriers [1–3]. It has been well recognized that non-viral vectors have their own advantages, such as low immune toxicity, construction flexibility and facile fabrication, in the gene transfection, especially for clinical applications [4–6]. However, they are still much less efficient than their viral counterparts. Among thousands of experimentally tested non-viral vectors, PEI is still considered as one of the most efficient candidates to deliver genes and often served as a “golden standard” [7–9]. Previously, PEI has been chemically modified in different ways so that additional functions were introduced for a better gene delivery, including the incorporation of intracellular biodegradable linkers [10, 11], the PEGylation to improve the serum stability during circulation [12, 13] and the attachment of some functional molecules to target specific cells or tissues [12–15]. Less attention, however, has been paid to why PEI remains one of the best non-viral carriers and how it facilitates the intracellular trafficking [16–27].

The first step in the development of non-viral vectors is how to package long anionic DNA chains into a small particle, i.e., the DNA complexation and condensation. Previous study showed that the physical and colloidal characteristics of resultant polymer/DNA polyplexes are important for an effective delivery of gene [28–32], but results were controversial and not conclusive. Much of the past effort has been devoted to the synthesis of state-of-the-art polymers [33–37] and subsequent polyplex formation [38–41], rather than the fundamental understanding of how a non-viral vector promotes the intracellular trafficking of DNA besides its roles in DNA encapsulation and protection. The proposed proton-sponge effect has been well accepted and taken as granted by those who entered this field later [7, 42].

Using a combination of different methods to characterize the size, molar mass and surface charge of the DNA/PEI polyplexes formed under different conditions, we confirmed some of the previous literature results, but revealed that the average size and chain density of the resultant polyplexes are not that important in the gene

transfection. In order to achieve an optimal transfection efficiency, the polyplexes in the solution mixture should have a slightly positively charged surface with a N:P ratio higher than 10. It is understandable that a positively charged surface facilitates the attachment of polyplexes to the negatively charged cellular membrane so that the endocytosis is promoted. However, it is still not clear why a high N:P ratio is required. Few previous studies showed that when the N:P ratio is high, a large amount of PEI chains are free in the solution mixture of DNA and PEI [43, 44]. It was also revealed that these excessive polycationic chains are much more toxic than those bound to DNA inside the polyplexes [44, 45]. Moreover, Wagner et al. found that the removal of free PEI chains by size exclusion chromatography significantly reduced the gene transfection efficiency [44]. To our knowledge, only few previous studies have noted such an effect of free polycationic chains on the gene transfection [43–47]. Most of the researchers in this field have overlooked such a finding. The current study was designed to elucidate the impact of polymer chains free in the solution mixture on gene delivery.

2.2 Materials and Methods

2.2.1 Materials and Cell Lines

Branched PEI with a weight-averaged molar mass of 25,000 g/mol (*b*PEI-25 K, Sigma-Aldrich) was used without further purification. Initial plasmid DNA pGL3-control vector (5,256 bp) encoding modified firefly luciferase and pEGFP-N1 (4,700 bp) expressing enhanced green fluorescent protein (EGFP) were purchased from Promega (USA) and Clontech (Germany), respectively. A large amount of these plasmids were prepared using a Qiagen Plasmid Maxi Kit (Qiagen, Germany). POPO-3 iodide, fluorescein-5-isothiocyanate (FITC) and di-4-ANEPPDHQ were purchased from Invitrogen (USA). Fetal bovine serum (FBS), Dulbecco's modified Eagle's medium (DMEM) and penicillin–streptomycin were purchased from GIBCO (USA). 293T cells were grown at 37 °C, 5 % CO₂ in DMEM supplemented with 10 % FBS, penicillin at 100 units/mL and streptomycin at 100 µg/mL.

2.2.2 Formation of DNA/PEI Polyplexes

Plasmid DNA was complexed with PEI in either distilled water or phosphate buffered saline (PBS) to form the DNA/PEI polyplexes as follows. Different amounts of PEI solution ($C = 1 \times 10^{-4} - 1 \times 10^{-3}$ g/mL) were added dropwise into a dilute DNA solution ($C = 14.5$ µg/mL), resulting in different molar ratios of nitrogen from PEI to phosphate from DNA (N:P). Each resultant DNA/PEI dispersion was incubated for 5 min at room temperature before its administration to the cell culture medium.

2.2.3 DNA Binding Assay

The binding of DNA to PEI to form polyplexes in the solution mixture was evaluated by the gel-shift assay. Each DNA/PEI dispersion with a desired N:P ratio was mixed with $6 \times$ loading buffer and then loaded on a 0.8 % (w/v) agarose gel containing ethidium bromide in tris–borate EDTA buffer. The amount of DNA loaded into each well was 0.4 μg in a total volume of 10 μL . The electrophoresis was performed under 100 V for 45 min. DNA bands were visualized under UV. The DNA binding was also characterized using laser light scattering (LLS).

2.2.4 POPO-3 Fluorescence Quenching Assay

Plasmid was labeled with POPO-3 dye at 1 dye to 100 base pair ratio. DNA/PEI polyplexes with different N:P ratios were prepared as described above. The final DNA concentration is 1.45 $\mu\text{g}/100 \mu\text{L}$ solution. After 5-min incubation, the relative fluorescence intensity (F) of each solution mixture was collected using a Hitachi F4500 fluorescence spectrophotometer (excitation 530 nm, emission 570 nm). The fraction of the uncomplexed DNA chains (x) was determined by $x = (F_{\text{polyplex}} - F_{\text{blank}})/(F_{\text{DNA}} - F_{\text{blank}})$.

2.2.5 Laser Light Scattering

A commercial LLS instrument (ALV5000) with a vertically polarized 22-mV He–Ne laser (632.8 nm, Uniphase) was used. The measurable angular range is $15\text{--}154^\circ$. In dynamic LLS, the intensity–intensity time correlation function ($G^{(2)}(\tau)$) of each DNA/PEI polyplex solution mixture was measured at different scattering angles. The Laplace inversion of each $G^{(2)}(\tau)$ can lead to a line-width distribution of $G(\Gamma)$ by the CONTIN program or by the double-exponential fitting, if there are only two relaxation modes, as

$$\{[G^{(2)}(q, \tau) - B]/B\}^{1/2} = A_1(q)e^{-\langle\Gamma\rangle_1\tau} + A_2(q)e^{-\langle\Gamma\rangle_2\tau} \quad (2.1)$$

where B is the measured baseline; $\langle\Gamma\rangle$ and $A(q)$ are the average line-width and the normalized intensity contribution of each relaxation mode, respectively, and $A_1(q) + A_2(q) \equiv 1$; q is the scattering vector defined as $q \equiv (4\pi n/\lambda_0)\sin(\theta/2)$ with θ , λ_0 and n , the scattering angle, the incident wavelength in vacuum and the refractive index of solvent, respectively. $\langle\Gamma\rangle$ can be related to the average translational diffusion coefficient $\langle D \rangle$ as $\langle\Gamma\rangle = \langle D \rangle q^2$. Using the Stokes–Einstein equation [48], $\langle D \rangle$ is further related to the average hydrodynamic radius ($\langle R_h \rangle$) by $\langle R_h \rangle = k_B T / (6\pi\eta \langle D \rangle)$, where η is the solvent viscosity. In this way, each $G(\Gamma)$ can also be converted into a hydrodynamic radius distribution $f(R_h)$.

On the other hand, the time-averaged scattering intensity from each relaxation mode can be calculated from the total time-averaged scattering intensity ($\langle I(q) \rangle$) and $A(q)$, respectively, measured from static and dynamic LLS, i.e., $\langle I(q) \rangle_1 = \langle I(q) \rangle A_1(q)$ and $\langle I(q) \rangle_2 = \langle I(q) \rangle A_2(q)$. The plot of $1/\langle I(q) \rangle_1$ or $1/\langle I(q) \rangle_2$ versus q^2 leads to an average radius of gyration ($\langle R_g \rangle_1$ or $\langle R_g \rangle_2$) and the time-averaged scattering intensity at $q \rightarrow 0$ ($\langle I(0) \rangle_1$ or $\langle I(0) \rangle_2$) using [48, 49]

$$\frac{1}{\langle I(q) \rangle} \approx \frac{1}{\langle I(0) \rangle} \left(1 + \frac{1}{3} \langle R_g^2 \rangle q^2 \right) \quad (2.2)$$

Where $\langle I(0) \rangle$ is proportional to the weight-averaged molar mass. Therefore, individual short PEI chains with a much lower molar mass are invisible in LLS when long DNA chains exist in the solution mixture. The formation of DNA/PEI polyplexes with some DNA chains collapsed inside leads to a relaxation mode faster than that of individual swollen DNA chains. This is why we mainly focused on how N:P ratio affects the fast mode in the current study.

2.2.6 Zeta-Potential Measurement

The average mobility (μ_E) of the polyplexes under an electric field in an aqueous solution was determined from the frequency shift in a laser Doppler spectrum using a commercial zeta-potential spectrometer (ZetaPlus, Brookhaven) with two platinum-coated electrodes and one He-Ne laser as the light source. Each data point in the mobility measurement was averaged over 20 times at 20 °C. The zeta-potential ($\zeta_{\text{potential}}$) can be calculated from μ_E using $\mu_E = 2\epsilon\zeta_{\text{potential}}f(\kappa R_b)/(3\eta)$, where ϵ is the permittivity of water and $1/\kappa$ is the Debye screening length [50]. When $\kappa R_b \ll 1$ (the Hückel limit), $f(\kappa R_b) \approx 1$; while $\kappa R_b \gg 1$ (the Smoluchowski limit), $f(\kappa R_b) \approx 1.5$. In the current study, the Hückel and Smoluchowski limits are respectively used to calculate $\zeta_{\text{potential}}$ in water and PBS.

2.2.7 Determination of the Content of Free PEI in DNA/PEI Dispersion

The content of free PEI in the DNA/PEI dispersion was determined using a combination of filtration and copper complex assay. The DNA/PEI dispersion at N:P = 10 was prepared as described above, with a final concentration of 219 μg pGL3/1 mL PBS. After 4-h incubation, the DNA/PEI dispersion was centrifuged and the supernatant was carefully filtered using a 20-nm filter. The content of uncomplexed PEI chains in the filtrate was determined using a copper complex assay by adding 100 μL of the filtrate into 900 μL of 0.02 M cupric acetate solution containing 5 % potassium acetate (pH 5.5). The absorption of the filtrate

was recorded at 630 nm (A_{630}) using a UV–Vis spectrometer (Hitachi, Japan), and converted to the corresponding PEI concentration according to the calibration curve.

2.2.8 *In Vitro Gene Transfection*

The in vitro gene transfection efficiency was quantified by using the luciferase transfection assays, in which plasmid pGL3 was used as an exogenous reporter gene. 293T cells were plated in 48-well plate at an initial density of 1.5×10^5 per well, 24 h prior to transfection. The DNA/PEI dispersion with a desired N:P ratio was further diluted in serum-free medium and then added at a final concentration of 0.4 μg DNA per well. The complete DMEM medium (800 μL per well) was added 6 h after the gene transfection. Using a GloMax 96 microplate luminometer (Promega, USA) and the Bio-Rad protein assay reagent, we respectively determined the transgene expression level and corresponding protein concentration in each well 48 h after the polyplex administration. The gene transfection efficiency is expressed as a relative luminescence unit (RLU) per cellular protein (mean $\pm$ SD of triplicates). Alternatively, the transfection efficiency can be directly visualized under a fluorescence microscopy when the plasmid pEGFP-N1 is used.

2.2.9 *Cytotoxicity Assay*

The cytotoxicity of free PEI chains and DNA/PEI dispersion was evaluated on 293T cells by using the MTT assay. 293T cells were seeded in a 96-well plate at an initial density of 5000 cells/well. After 24 h, free PEI chains alone or with DNA to form DNA/PEI dispersion were respectively added to cells at different chosen concentrations and N:P ratios. For the DNA/PEI dispersion, the final DNA concentration is 0.2 μg /well in a total volume of 100 μL . The treated cells were incubated in a humidified environment with 5 % CO_2 at 37 $^\circ\text{C}$ for 48 h. The MTT reagent (in 20 μL PBS, 5 mg/mL) was then added to each well. The cells were further incubated for 4 h at 37 $^\circ\text{C}$. The medium in each well was then removed and replaced by 100 μL DMSO. The plate was gently agitated for 15 min before the absorbance (A) at 490 nm was recorded by a microplate reader (Bio-rad, USA). The cell viability (y) was calculated by $y = (A_{\text{treated}}/A_{\text{control}}) \times 100 \%$, where A_{treated} and A_{control} are the absorbance of the cells cultured with polymer/polyplex and fresh culture medium, respectively. Each experiment condition was done in quadruple. The data was shown as the mean value plus a standard deviation ($\pm$ SD).

2.2.10 DNA, PEI Labeling and Flow Cytometry

Plasmid pGL3 was covalently labeled with the fluorophore Cy5 using the Label IT kit (Mirus, Madison, WI) according to the manufacturer's instructions. For labeling of PEI, 11.2 mg of FITC (28.8 μmol) was mixed with 360 mg of *b*PEI-25 K (14.4 μmol) in EtOH, and the reaction mixture was stirred overnight at room temperature in the dark place. In order to separate labeled PEI from unreacted FITC, size exclusion chromatography (SEC) was performed using a gel-filtration column (Sephadex G-25 superfine) preequilibrated with HBS (20 mM HEPES, pH 7.4, 150 mM NaCl). The separated sample was freeze-dried and resolved in water. The gel electrophoresis results confirmed that the labeling on DNA and PEI chains has no obvious impact on the formation of DNA/PEI polyplexes (data not shown).

To explore the effect of free PEI chains on the cellular uptake of polyplexes, FITC-PEI, Cy5-pGL3/PEI polyplexes alone (N:P = 3) or polyplexes plus 7 portions of free FITC-PEI (N:P = 10) were added to the cells in serum-free DMEM and, after incubation at 37 °C, cells were harvested at the indicated time points. Briefly, cells were first rinsed twice with PBS containing 0.001 % SDS and then PBS to remove the extracellularly attached polyplexes [51]. Further, cells were detached by 0.05 % trypsin/EDTA supplemented with 20 mM sodium azide to prevent further endocytosis [52]. Finally, cells were washed twice by pelleting and then resuspended in ice-cold PBS containing 2 % FBS. Confocal microscopy images of cells were taken to ensure the effective removal of polyplexes from the cell surfaces (data not shown). Cellular uptake of polyplexes and free PEI chains was assayed by flow cytometry using a FC 500 flow cytometry system (Beckman Coulter, USA). The fluorophores FITC and Cy5 were excited at 488 and 635 nm, respectively, and the corresponding emissions were detected at 525/10 and 675/15 nm, respectively. To discriminate viable cells from dead cells and to exclude doublets, the cells were appropriately gated by forward/side scattering and pulse width. 1×10^4 gated events per sample were collected. Experiments were performed at least in triplicates.

2.2.11 Confocal Laser Scanning Microscopy

6×10^5 cells were seeded in a $\mu\text{-Dish}^{35\text{mm, high}}$ (ibidi GmbH, Germany). After 24 h, the cellular membrane was stained with di-4-ANEPPDHQ for 15 min and then washed 2–3 times. Afterwards, the cell culture medium was carefully aspirated and FITC-labeled *b*PEI-25 K in serum-free DMEM was applied at a final concentration of 1×10^{-5} g/mL. Live cell imaging was performed for 1.5 h using a Nikon C1si confocal laser scanning microscope equipped with a spectral imaging detector (Nikon, Japan) and a INU stage-top incubator (Tokai Hit, Japan). Image sequences were captured at approximately 30 s intervals. FITC and di-4-ANEPPDHQ were visualized by the 488-nm excitation and the corresponding emission

spectral was acquired in the 495–650 nm range at 5-nm wavelength resolution. The mean FITC-PEI fluorescence intensities at the cellular membrane and inside the cell are recorded at 520 nm and normalized by that outside the cell. All the spectral data was analyzed using the Nikon EZ-C1 software.

2.3 Results and Discussion

To investigate the complexation profiles of PEI-mediated vectors, we first monitored the formation of DNA/PEI polyplexes in PBS using a combination of static and dynamic LLS. Figure 2.1 shows the N:P ratio dependence of hydrodynamic radius distribution ($f(R_h)$) of pGL3 without and with the addition of different amounts of PEI. In the pure pGL3 solution (N:P = 0.00), the peak located at $\sim 1.4 \mu\text{m}$ represents the swollen and extended DNA chains. The addition of a small amount of branched PEI (N:P = 0.25) leads to a new peak located at $\sim 120 \text{ nm}$, presumably corresponding to the newly formed contracted DNA/PEI polyplexes. The shifting of the DNA peak from ~ 1.4 to $\sim 1 \mu\text{m}$ indicates the partial contraction of DNA chains due to the incomplete complexation. Further addition of PEI leads to polyplexes of a larger size with more DNA and PEI chains incorporated into the individual complexes. When N:P ~ 3 , the DNA peak completely disappears, suggesting that nearly all the DNA chains are complexed with PEI, i.e., no DNA chains are free in the solution mixture.

Fig. 2.1 N:P ratio dependence of the normalized hydrodynamic radius distribution ($f(R_h)$) of DNA/PEI polyplexes formed in PBS, where different colors represent variation of the time-averaged scattering intensity of the solution mixture ($\langle I(q) \rangle$)





Fig. 2.2 Complexation profile between PEI and DNA in PBS at different N:P ratios, evaluated by **a** gel-shift assay and **b** POPO-3 fluorescence quenching assay



The complexation between PEI and DNA was also evaluated by the gel-shift assay (Fig. 2.2a). Progressive condensation can be revealed from the retarded mobility of the DNA bands and their reduced fluorescence intensity. Clearly, the two DNA bands (supercoiled and relaxed forms) disappear when $N:P \sim 3$. Furthermore, we quantitatively estimated the extent of DNA complexation at each given N:P ratio using a double-strand DNA intercalating dye, POPO-3, whose fluorescence is dramatically increased upon its binding to DNA. The complexation of DNA with PEI excludes POPO-3 molecules from the DNA double helix, resulting in a reduced fluorescence emission intensity. Figure 2.2b shows that over 90 % of the DNA chains have been condensed into the polyplexes when $N:P \sim 3$. In static LLS, we also found that the average scattering intensity ($\langle I(0) \rangle$) nearly remains a constant after $N:P \geq 3$.

It is known that $\langle I(0) \rangle \sim KCM_w \sim (dn/dc)^2 CM_w$. Therefore, we can calculate the apparent weight-averaged molar mass ($M_{w, \text{polyplex}}$) of the polyplexes at each given N:P ratio using

$$M_{w, \text{polyplex}} = M_{w, \text{DNA}} \times \frac{\langle I(0) \rangle_{\text{polyplex}} C_{\text{DNA}} \left(\frac{dn}{dc} \right)_{\text{DNA}}^2}{\langle I(0) \rangle_{\text{DNA}} C_{\text{polyplex}} \left(\frac{dn}{dc} \right)_{\text{polyplex}}^2} \quad (2.3)$$

where C is concentration and (dn/dc) is the differential refractive index increment. Note that $(dn/dc)_{\text{polyplex}} = w_{\text{DNA}}(dn/dc)_{\text{DNA}} + w_{\text{PEI}}(dn/dc)_{\text{PEI}}$, where w_{DNA} and w_{PEI} are two normalized weight fractions of DNA and PEI inside the polyplexes, i.e., $w_{\text{DNA}} + w_{\text{PEI}} = 1$. It is necessary to estimate C_{polyplex} and w_{PEI} in order to calculate $M_{w, \text{polyplex}}$. When $N:P \leq 3$, we can assume that nearly all the PEI chains are complexed with DNA, while the fraction of DNA associated with PEI can be estimated using the POPO-3 fluorescence assay mentioned above. Therefore,

$$C_{\text{polyplex}} = C_{\text{PEI}} + (1 - x)C_{\text{DNA}} \text{ and } W_{\text{PEI}} = C_{\text{PEI}}/C_{\text{polyplex}} \quad (2.4)$$

where C_{PEI} and C_{DNA} are the initial PEI and DNA concentrations, respectively; and x is the fraction of those uncomplexed DNA chains in the solution mixture. On the other hand, when $3 < \text{N:P} \leq 10$, nearly all the DNA chains are complexed with PEI so that C_{polyplex} is a constant in this range and $C_{\text{polyplex}} \approx C_{\text{polyplex}} (\text{N:P} = 3)$.

Figure 2.3 summaries the N:P ratio dependence of the average hydrodynamic radius ($\langle R_h \rangle$), the apparent weight-averaged molar mass ($M_{w, \text{polyplex}}$), the zeta-potential ($\zeta_{\text{potential}}$) and the gene transfection efficiency (RLU/mg protein) of the DNA/PEI polyplexes formed in PBS. The initial decrease of $\langle R_h \rangle$ and increases of both $M_{w, \text{polyplex}}$ and $\zeta_{\text{potential}}$ upon the addition of PEI ($\text{N:P} = 0.125\text{--}1.50$) indicate that the positively charged PEI chains are gradually complexed with the negatively charged DNA chains so that both PEI and DNA contract in the solution mixture. The sharp increases of both $\langle R_h \rangle$ and $M_{w, \text{polyplex}}$ at $\text{N:P} \sim 2$ reflect the merge of initially formed small polyplexes. Further addition of PEI in the range of $\text{N:P} > 3$ has nearly no effect on $\langle R_h \rangle$, $M_{w, \text{polyplex}}$ and $\zeta_{\text{potential}}$, clearing revealing that those PEI chains added afterwards are free in the solution mixture.

Only few previous studies have noted the existence of free PEI, and estimated the amount of those free polycationic chains in the DNA/PEI dispersion used for gene delivery. Clamme et al. found that, by using a two photon fluorescence correlation spectroscopy, $\sim 86\%$ of the PEI chains are in the free form at $\text{N:P} = 10$ (i.e., polyplexes is completely formed at $\text{N:P} \sim 1.4$) [43], whereas Wagner and his coworkers showed that the purified polyplexes by SEC have an ultimate N:P ratio of ~ 2.5 [44]. Our LLS results here clearly shows that the anionic DNA chains are not fully condensed by the cationic PEI chains when $\text{N:P} \sim 1.4$, also reflected in its corresponding negative zeta-potential of the resultant polyplexes. Besides LLS, herein we used a simple filtration method combined with copper complex assay to determine the content of uncomplexed PEI chains in the DNA/PEI dispersion. Figure 2.4 shows that the PEI concentration after the removal of the polyplexes by a 20-nm filter is $\sim 68.6\%$ of the initial PEI concentration at $\text{N:P} = 10$, further confirming that DNA is fully condensed by PEI only when $\text{N:P} \geq 3$. For $\text{N:P} = 10$, $\sim 70\%$ of the PEI chains are free in the solution mixture of PEI and DNA. On the other hand, our results in Fig. 2.3d consolidate that the polyplexes with 7 portions of free PEI chains ($\text{N:P} = 10$) are $\sim 10^3$ times more efficient in the gene transfection than those formed at $\text{N:P} = 3$ without free PEI chains.

Previous literature also repeatedly showed that when $\text{N:P} \geq 3$, nearly all the DNA chains are completely complexed with PEI, but a higher N:P ratio leads to much better gene transfection. Putting these two experimental facts together, one should ask an obvious, but certainly overlooked, question; namely, is it the free PEI chains, rather than those inside the DNA/PEI polyplexes, that play a vital role in the gene transfection? To generalize such a question, we have tested PEI-mediated vectors with different chain lengths in both salt-free water and salt-rich PBS. All of our results (not shown here) confirm that the solution mixture with polyplexes alone have a much lower gene transfection efficiency.

Fig. 2.3 N:P ratio dependence of **a** average hydrodynamic radius ($\langle R_h \rangle$); **b** apparent weight-averaged molar mass ($M_{w, \text{ polyplex}}$); **c** average zeta-potential ($\zeta_{\text{potential}}$); and **d** gene transfection efficiency (RLU/mg protein) of polyplexes formed in PBS. *Inset* Fluorescent microscopic image of 293T cells transfected with pEGFP-N1 after 48 h. Scale bar: 200 μm





Fig. 2.4 Concentration calibration curve of *b*PEI-25 K chains in PBS, where symbols ▲ and ★ refer to the concentration of PEI in DNA/PEI dispersion (N:P = 10) before and after the removal of the polyplexes, respectively



On the other hand, we evaluated the cytotoxicity of free PEI chains and the corresponding DNA/PEI dispersion on 293T cells using the MTT assay (Fig. 2.5). The initial cell viability of DNA/PEI at N:P = 3 is ~85 %, and is stably reduced as the increasing amount of free PEI chains, especially when $C_{PEI} \geq 2.7 \mu\text{g/mL}$, corresponding to N:P ≥ 10 , clearing revealing that those free PEI chains are indeed the major cause of toxicity of DNA/PEI at high N:P ratios. It is also worth noting that in the general N:P range for transfection, i.e., N:P ≤ 10 , both the DNA/PEI dispersion and free PEI chains exhibit relatively low cytotoxicity, with the cell viability well above 70 %.

Furthermore, we decided to add the 7 portions of free PEI chains at different times; namely, hours before or after the administration of the DNA/PEI polyplexes (N:P = 3), so that the final and total N:P ratio remains 10. We define $t = 0$ for the simultaneous addition of the DNA/PEI polyplexes (N:P = 3) and 7 portions of free PEI chains. Therefore, the negative time means that free PEI chains are added before the polyplex administration. Figure 2.6 shows that in the presence of free PEI chains, the transfection efficiency is typically 10^2 – 10^3 times higher than that without free chains no matter whether these free polycations are introduced before or after the polyplex administration. To our knowledge, this is the first demonstration that the addition of free PEI chains even prior to the polyplexes can facilitate the gene transfection. It is clear that the simultaneous addition of the polyplexes (N:P = 3) and free polycationic chains (i.e., $t = 0$) leads to the highest

Fig. 2.5 PEI concentration dependence of 293T cell viability in terms of free PEI chains alone and DNA/PEI dispersions, where $C_{PEI} = 2.7 \mu\text{g/mL}$ is corresponding to N:P = 10





Fig. 2.6 Effect of 7 portions of free PEI chains added at different times (prior to or post the administration of polyplexes formed at $N:P = 3$) on the gene transfection efficiency in 293T cells, where the total and final $N:P$ ratio is kept to be 10 and the cell culture medium is not replaced before the addition of free PEI chains



transfection level, indicating a possibly cooperative role of the bound and free PEI chains in gene delivery.

Figure 2.6 also reveals that in the delayed addition of free PEI chains, i.e., $t > 0$, the gene transfection efficiency gradually decreases with the time interval between the administration of polyplexes and the addition of free PEI chains. To explain such a decrease, we have to find whether this decline is due to the insufficient incubation time since the incubation time was fixed to be 48 h, starting from the polyplex administration. The incubation-time dependence of the transgene expression, as shown in Fig. 2.7, clearly excludes such a possibility because the luciferase expression reaches its maximum at ~ 36 h after the polyplex administration. Therefore, the total 48-h incubation time is sufficient. The next question is whether these free PEI chains promote the cellular uptake in the extracellular space or other processes inside the cell. In the extracellular pathway, free PEI chains could facilitate the cellular internalization by increasing either the uptake rate or the endocytosis amount. To estimate the internalization timescale of the DNA/PEI dispersion, the cell culture medium with the polyplexes, if any, was replaced at different times after the polyplex administration. Figure 2.8 shows that the cellular uptake nearly ceases after ~ 6 h regardless of whether there are free PEI chains, but these free polycations enhance the final transfection levels by ~ 500 times even though they do not shorten the time required for internalization.

Fig. 2.7 Incubation-time dependence of luciferase expression of DNA/PEI polyplexes formed at $N:P = 10$. *Inset* Fluorescent microscopic image of 293T cells transfected with pEGFP-N1 after 48 h. Scale bar: 200 μ m





Fig. 2.8 Effect of removal of cell culture medium on the gene transfection efficiency of DNA/PEI polyplexes formed in PBS with and without free PEI chains in the solution mixture, where “ $t = 0$ ” means that no cell culture medium was replaced




Furthermore, flow cytometry was used to explore the internalization kinetics of DNA/PEI polyplexes and free PEI chains in the gene transfection. Generally, three characteristics can be extracted from the flow cytometry data; namely, (1) the fraction of cells containing Cy5 fluorescence (Fig. 2.9a); (2) the median of Cy5-DNA fluorescence intensity of Cy5-positive cell population ($F_{\text{Cy5-DNA, Cy5+}}$, Fig. 2.9b); and (3) the median of Cy5-DNA fluorescence intensity of total cell population ($F_{\text{Cy5-DNA, tot}}$, Fig. 2.9c). Figure 2.9a shows that in the presence of 7 portions of free PEI chains, more than 80 % of the cells have internalized polyplexes after a 30-min incubation, whereas the internalized polyplexes without free PEI chains are detected in only ~10 % of the cells. At 4 h post-incubation, ~95 % of the cells are Cy5-positive regardless of the addition of free PEI chains. In addition, $F_{\text{Cy5-DNA, Cy5+}}$, which is indicative of the polyplex amount per Cy5-positive cell, is stably 3–6 times higher during the first 6-h incubation when the extra 7 portions of PEI chains are applied (Fig. 2.9b). In combination, the addition of free PEI leads to a significant faster and remarkably increased internalization (Fig. 2.9c).

Moreover, different cellular uptake kinetics is revealed for polyplexes in the absence/presence of the free PEI chains (Fig. 2.10). When the polyplexes are applied alone, the number of cells they entered and their content per cell elevate simultaneously in the first 6 h of incubation. With the aid of 7 portions of free PEI chains, in contrast, the fraction of Cy5-positive cells quickly rises to ~80 % within just 0.5 h, and the enhanced internalization afterwards (1.5–6 h) is mainly attributed to the increasing amount of polyplexes inside one cell.

On the other hand, the 7 portions of free PEI chains, whether applied alone or with DNA/PEI polyplexes at N:P = 3, exhibit similar internalization kinetics (Fig. 2.11). Within just 30 min, these free PEI chains enter ~80 % of the cells, with their intracellular fluorescence intensity almost reaching the maximum plateau. Moreover, our CLSM result also confirms that the concentrations of free PEI chains at the cellular membrane and inside the cell quickly approach to the maximum ~30 min after they are added outside the cell (Fig. 2.12). This indicates that the free PEI chains have a significantly faster uptake rate, and perhaps a different internalization pathway, compared to the DNA/PEI polyplexes (~6 h). Notably, the major uptake timescale of free PEI (~0.5–1.5 h) is in good

Fig. 2.9 Effect of free PEI chains on the internalization of DNA/PEI polyplexes by flow cytometry. 293T cells were transfected with PEI/Cy5-DNA polyplexes alone ($N:P = 3$), or polyplexes plus 7 portions of free FITC-PEI ($N:P = 10$), harvested, and analyzed at the indicated time points. Internalization extent is expressed as **a** percentage of Cy5-positive cells; **b** median of Cy5-DNA fluorescence intensity of Cy5-positive cell population ($F_{\text{Cy5-DNA, Cy5+}}$); and **c** median of Cy5-DNA fluorescence intensity of total cell population ($F_{\text{Cy5-DNA, tot}}$)



accordance with the sharp increase in the number of polyplex-containing cells at $N:P = 10$ (Fig. 2.9a), suggesting that these excessive PEI chains might mainly play their roles in the first 1.5 h. It is still to be elucidated that how the free PEI chains promote the cellular internalization. They might either destabilize the cellular membrane to allow direct make the endocytosis of polyplexes more easily, or help the polyplexes to utilize an internalization pathway more favorable for intracellular trafficking [27, 28, 53].

To compare the contribution of free PEI chains in the extracellular and intracellular pathways, we added free PEI chains with or without removing the cell culture medium that contains the polyplexes (Fig. 2.13). Since the cellular internalization stops after ~ 6 h, those polyplexes in the extracellular space, if any,




Fig. 2.10 Schematic illustration of cellular internalization of DNA/PEI polyplexes (N:P = 3) in the absence and presence of 7 portions of free PEI chains, where different colors represent the median of DNA fluorescence intensity (MFI) inside one cell. Note that the depictions of different components do not reflect their actual sizes

should still remain there. For N:P = 10, when the polyplexes are removed, at 6 h post-incubation, before the addition of 7 portions of free PEI chains, the gene transfection efficiency is only attenuated by 2-fold in comparison with the non-removal group, indicating that those free PEI chains promote the cellular uptake a little. However, it is worth noting that this reduced transfection efficiency is still ~ 50 -fold higher than that without the addition of free PEI chains (N:P = 3), clearly indicating that the free PEI chains mainly play their role inside the cells.

Finally, we explored the effect of FBS on the gene transfection efficiency of polyplexes alone (N:P = 3), or polyplexes with 7 portions of free PEI chains added at different times (N:P = 10). When the cell culture medium is supplemented with 10 % FBS, the initial transfection efficiency of polyplexes at N:P = 3 is reduced by ~ 5 times (Fig. 2.14a). Moreover, the 7 portions of free PEI chains added afterwards only increase the transfection level by 1–150 folds, whereas, in the serum-free medium, they can enhance the transfection efficiency by 10–300 times (Fig. 2.14b). This decreased efficacy is mainly attributed to the association of the cationic polyplexes and PEI chains with the negatively charged proteins in FBS; namely, it reduces the effective amount of free PEI chains, which indirectly supports our finding that free PEI chains play a significant role in promoting the transfection. Therefore, for the future in vivo study, we have to consider how to incorporate free PEI chains together with polyplexes and keep their integrity in the blood stream so that they can be safely delivered to the targeted cells or organs.

Fig. 2.11 Comparison of cell internalization kinetics of free FITC-PEI chains applied alone or with DNA/PEI polyplexes of N:P = 3 in the gene transfection. Internalization extent is expressed as **a** percentage of FITC-positive cells; **b** median of FITC-PEI fluorescence intensity of FITC-positive cells ($F_{\text{FITC-PEI, FITC+}}$); and **c** median of FITC-PEI fluorescence intensity of total cell population ($F_{\text{FITC-PEI, tot}}$)




Fig. 2.12 Cellular membrane translocation kinetics of FITC-labeled *b*PEI-25 K chains, where the mean FITC-PEI fluorescence intensities at the membrane (on the circle 2) and inside the cell (inside circle 1, scale bar: 10 μm) are normalized by that outside the cell




Fig. 2.13 Effect of free PEI chains on the intracellular trafficking of DNA/PEI polyplexes formed in PBS, where 7 portions of free PEI chains were added, respectively, at 6 h and 8 h post-administration of the polyplexes (N:P = 3), with or without the removal of cell culture medium





Fig. 2.14 a Effect of FBS (10 % in the cell culture medium) on the transfection efficiency of polyplexes alone (N:P = 3), or polyplexes with 7 portions of free PEI added at different times (N:P = 10). **b** Relative transfection efficiency of polyplexes plus free PEI ($E_{N:P=10}$) to polyplex alone ($E_{N:P=3}$) in the serum-free (–) and serum-containing (+) medium, respectively. The cell culture medium is not replaced before the addition of free PEI chains



In sum, our results reveal the existence of some cooperative actions between the polyplexes and free polycationic chains. The cationic PEI chains inside each polyplex condense anionic DNA chains into a small and compact particle so that the endocytosis is facilitated. These bound PEI chains also shield DNA from the intracellular degradation; while those free PEI chains in the solution mixture mainly navigate some unknown intracellular trafficking barriers, including endolysosomal formation and/or escape and the subsequent nuclear localization. It is a remaining challenge to map these intracellular trafficking pathways and understand how free PEI chains exactly promote the gene transfection inside the cell by using various existing tools of molecular biology.

2.4 Conclusion

A combination of LLS and gel electrophoresis results confirms that most of anionic DNA chains are complexed and condensed by cationic PEI chains when the molar ratio of nitrogen from PEI to phosphate from DNA (N:P) reaches ~ 3 , independent of the PEI chain length and solvent (pure water or PBS), revealing that the charge neutrality (electrostatic interaction) is the main driving force for the polyplex formation. In the solution mixture with $N:P > 3$, there are two kinds of PEI chains: bound to DNA and free in the solution mixture. The bound PEI chains inside the polyplexes provide charge compensation so that DNA is condensed and protected from degradation. Our current study convincingly reveals that it is those free PEI chains, rather than other physical properties of the DNA/PEI polyplexes, that play a vital role in promoting the gene transfection. The addition of free PEI leads to a significant faster and more efficient cellular uptake of polyplexes, but these free PEI chains mainly contribute to the subsequent intracellular trafficking. Our finding leads to a different thinking in the development of non-viral vectors; namely, we might not need to invest much of our effort on the synthesis of different cationic polymers, but focus on how the free polycationic chains promote the intracellular trafficking of the polyplexes. As for in vivo experiments, we have to consider how to incorporate free chains together with polyplexes and keep their integrity during circulation so that they can be safely delivered to the targeted cells or organs.

References

1. Check, E. (2002). *Nature*, 420, 116–118.
2. Raper, S. E., Chirmule, N., Lee, F. S., Wivel, N. A., Bagg, A., Gao, G. P., et al. (2003). *Molecular Genetics and Metabolism*, 80, 148–158.
3. Hacia-Bey-Abina, S. (2003). *Science*, 302, 568–568.
4. Pack, D. W., Hoffman, A. S., Pun, S., & Stayton, P. S. (2005). *Nature Reviews Drug Discovery*, 4, 581–593.
5. Li, S. D., & Huang, L. J. (2007). *Journal of Controlled Release*, 123, 181–183.
6. Mintzer, M. A., & Simanek, E. E. (2009). *Chemical Reviews*, 109, 259–302.
7. Boussif, O., Lezoualch, F., Zanta, M. A., Mergny, M. D., Scherman, D., Demeneix, B., et al. (1995). *Proceedings of the National Academy of Sciences of the United States of America*, 92, 7297–7301.
8. Lungwitz, U., Breunig, M., Blunk, T., & Gopferich, A. (2005). *European Journal of Pharmaceutics and Biopharmaceutics*, 60, 247–266.
9. Neu, M., Fischer, D., & Kissel, T. J. (2005). *Gene Medicine*, 7, 992–1009.
10. Breunig, M., Lungwitz, U., Liebl, R., & Gopferich, A. (2007). *Proceedings of the National Academy of Sciences of the United States of America*, 104, 14454–14459.
11. Deng, R., Yue, Y., Jin, F., Chen, Y. C., Kung, H. F., Lin, M. C. M., et al. (2009). *Controlled Release*, 140, 40–46.
12. Ogris, M., Brunner, S., Schuller, S., Kircheis, R., & Wagner, E. (1999). *Gene Therapy*, 6, 595–605.

13. Cheng, H., Zhu, J. L., Zeng, X., Jing, Y., Zhang, X. Z., & Zhuo, R. X. (2009). *Bioconjugate Chemistry*, 20, 481–487.
14. Zanta, M. A., Boussif, O., Adib, A., & Behr, J. P. (1997). *Bioconjugate Chemistry*, 8, 839–844.
15. Diebold, S. S., Kursa, P., Wagner, E., Cotten, M., & Zenke, M. J. (1999). *Biological Chemistry*, 274, 19087–19094.
16. Pollard, H., Remy, J. S., Loussouarn, G., Demolombe, S., Behr, J. P., & Escande, D. J. (1998). *Biological Chemistry*, 273, 7507–7511.
17. Godbey, W. T., Wu, K. K., & Mikos, A. G. (1999). *Proceedings of the National Academy of Sciences of the United States of America*, 96, 5177–5181.
18. Bieber, T., Meissner, W., Kostin, S., Niemann, A., & Elsasser, H. P. J. (2002). *Controlled Release*, 82, 441–454.
19. Sonawane, N. D., Szoka, F. C., & Verkman, A. S. J. (2003). *Biological Chemistry*, 278, 44826–44831.
20. Suh, J., Wirtz, D., & Hanes, J. (2003). *Proceedings of the National Academy of Sciences of the United States of America*, 100, 3878–388.
21. Akinc, A., Thomas, M., Klibanov, A. M., & Langer, R. J. (2005). *Gene Medicine*, 7, 657–663.
22. Kulkarni, R. P., Wu, D. D., Davis, M. E., & Fraser, S. E. (2005). *Proceedings of the National Academy of Sciences of the United States of America*, 102, 7523–7528.
23. de Bruin, K., Ruthardt, N., von Gersdorff, K., Bausinger, R., Wagner, E., Ogris, M., et al. (2007). *Molecular Therapy*, 15, 1297–1305.
24. de Bruin, K. G., Fella, C., Ogris, M., Wagner, E., Ruthardt, N., & Brauchle, C. J. (2008). *Controlled Release*, 130, 175–182.
25. Gabrielson, N. P., & Pack, D. W. J. (2009). *Controlled Release*, 136, 54–61.
26. Won, Y., Sharma, R., & Konieczny, S. F. J. (2009). *Controlled Release*, 139, 88–93.
27. Lee, H., Kim, I. K., & Park, T. G. (2010). *Bioconjugate Chemistry*, 21, 289–295.
28. Wagner, E., Cotten, M., Foisner, R., & Birnstiel, M. L. (1991). *Proceedings of the National Academy of Sciences of the United States of America*, 88, 4255–4259.
29. Godbey, W. T., Wu, K. K., & Mikos, A. G. (1999). *Journal of Biomedical Materials Research*, 45, 268–275.
30. Wightman, L., Kircheis, R., Rossler, V., Carotta, S., Ruzicka, R., Kursa, M., et al. (2001). *Gene Medicine*, 3, 362–372.
31. Kunath, K., von Harpe, A., Fischer, D., Peterson, H., Bickel, U., Voigt, K., et al. (2003). *Controlled Release*, 89, 113–125.
32. Honore, I., Grosse, S., Frison, N., Favatier, F., Monsigny, M., & Fajac, I. J. (2005). *Controlled Release*, 107, 537–546.
33. Pun, S. H., Bellocq, N. C., Liu, A. J., Jensen, G., Machemer, T., Quijano, E., et al. (2004). *Bioconjugate Chemistry*, 15, 831–840.
34. Ooya, T., Choi, H. S., Yamashita, A., Yui, N., Sugaya, Y., Kano, A., et al. (2006). *Journal of the American Chemical Society*, 128, 3852–3853.
35. Tian, H. Y., Xiong, W., Wei, J. Z., Wang, Y., Chen, X. S., Jing, X. B., et al. (2007). *Biomaterials*, 28, 2899–2907.
36. Sun, Y. X., Xiao, W., Cheng, S. X., Zhang, X. Z., & Zhuo, R. X. J. (2008). *Controlled Release*, 128, 171–178.
37. Liang, B., He, M. L., Chan, C. Y., Chen, Y. C., Li, X. P., Li, Y., et al. (2009). *Biomaterials*, 30, 4014–4020.
38. Bloomfield, V. A. (1997). *Biopolymers*, 44, 269–282.
39. Lai, E., & van Zanten, J. H. (2001). *Biophysical Journal*, 80, 864–873.
40. Vuorimaa, E., Urtti, A., Seppanen, R., Lemmetyinen, H., & Yliperttula, M. (2008). *Journal of the American Chemical Society*, 130, 11695–11700.
41. Ziebarth, J., & Wang, Y. M. (2009). *Biophysical Journal*, 97, 1971–1983.
42. Behr, J. P. (1997). *Chimia*, 51, 34–36.
43. Clamme, J. P., Azoulay, J., & Mely, Y. (2003). *Biophysical Journal*, 84, 1960–1968.

44. Boeckle, S., von Gersdorff, K., van der Piepen, S., Culmsee, C., Wagner, E., & Ogris, M. J. (2004). *Gene Medicine*, 6, 1102–1111.
45. Fahrmeir, J., Gunther, M., Tietze, N., Wagner, E., & Ogris, M. J. (2007). *Controlled Release*, 122, 236–245.
46. Clamme, J. P., Krishnamoorthy, G., & Mely, Y. (2003). *Biochimica et Biophysica Acta, Biomembranes*, 1617, 52–61.
47. Saul, J. M., Wang, C. H. K., Ng, C. P., & Pun, S. H. (2008). *Advanced Materials*, 20, 19–25.
48. Berne, B., & Pecora, R. (1976). *Dynamic light scattering*. New York: Plenum Press.
49. Chu, B. (1991). *Laser light scattering* (2nd ed.). New York: Academic Press.
50. Hunter, R. J. (2000). *Foundations of colloid science* (2nd ed.). Oxford: Oxford Press.
51. Forrest, M. L., & Pack, D. W. (2002). *Molecular Therapy*, 6, 57–66.
52. Breunig, M., Lungwitz, U., Liebl, R., Fontanari, C., Klar, J., Kurtz, A., et al. (2005). *Journal of Gene Medicine*, 7, 1287–1298.
53. Rejman, J., Bragonzi, A., & Conese, M. (2005). *Molecular Therapy*, 12, 468–474.

How Free Cationic Polymer Chains Promote Gene
Transfection

Yanan, Y.

2013, XVIII, 94 p. 45 illus., 38 illus. in color., Hardcover

ISBN: 978-3-319-00335-1