

Chapter 8

Minimizing Movements Along a Sequence of Functionals

Gradient flows, and hence minimizing movements, and the convergence of functionals trivially do not commute even when the convergence is uniform. As a simple example, take $X = \mathbb{R}$ and

$$F_\varepsilon(x) = x^2 - \rho \sin\left(\frac{x}{\varepsilon}\right),$$

with $\rho = \rho_\varepsilon \rightarrow 0$ as $\varepsilon \rightarrow 0$, uniformly converging to $F(x) = x^2$. If also

$$\varepsilon \ll \rho,$$

then for fixed x_0 the solutions u_ε to the equation

$$\begin{cases} u'_\varepsilon = -2u_\varepsilon + \frac{\rho}{\varepsilon} \cos\left(\frac{u_\varepsilon}{\varepsilon}\right) \\ u_\varepsilon(0) = x_0 \end{cases}$$

converge to the constant function $u_0(t) = x_0$ as $\varepsilon \rightarrow 0$. This is easily seen by studying the stationary solutions of

$$-2x + \frac{\rho}{\varepsilon} \cos\left(\frac{x}{\varepsilon}\right) = 0.$$

Conversely, the gradient flow of the limit is

$$\begin{cases} u' = -2u \\ u(0) = x_0, \end{cases}$$

for which the constant functions are not solutions if $x_0 \neq 0$.

With the remark above in mind, in order to give a meaningful limit for the energy-driven motion along a sequence of functionals it may be useful to vary the definition

of minimizing movement. This will be done in the following section. As in the previous chapter, we will limit our analysis to a Hilbert setting for simplicity.

8.1 Minimizing Movements Along a Sequence

In this section we will give a notion of minimizing movement along a sequence F_ε , which will depend in general on the interaction between the time scale τ and the parameter ε in the energies.

Definition 8.1 (minimizing movements along a sequence). Let X be a separable Hilbert space, let $F_\varepsilon : X \rightarrow [0, +\infty]$ be equicoercive and lower semicontinuous, $x_0^\varepsilon \rightarrow x_0$ with

$$F_\varepsilon(x_0^\varepsilon) \leq C < +\infty, \quad (8.1)$$

and let $\tau_\varepsilon > 0$ converge to 0 as $\varepsilon \rightarrow 0$. With fixed $\varepsilon > 0$ we define x_k^ε recursively as a minimizer for the problem

$$\min \left\{ F_\varepsilon(x) + \frac{1}{2\tau} \|x - x_{k-1}^\varepsilon\|^2 \right\}, \quad (8.2)$$

and the piecewise-constant trajectory $u^\varepsilon : [0, +\infty) \rightarrow X$ given by

$$u^\varepsilon(t) = x_{[t/\tau_\varepsilon]}. \quad (8.3)$$

A *minimizing movement* for F_ε from x_0^ε is any limit of a subsequence u^{ε_j} uniform on compact sets of $[0, +\infty)$.

After remarking that the Hölder continuity estimates in Proposition 7.1 only depend on the bound on $F_\varepsilon(x_0^\varepsilon)$, with the same proof we can show the following result.

Proposition 8.1. *For every F_ε and x_0^ε as above there exist minimizing movements for F_ε from x_0^ε in $C^{1/2}([0, +\infty); X)$.*

Remark 8.1 (Growth conditions). As for the case of a single functional, the positiveness of F_ε can be substituted by the requirement that for all \bar{x} the functionals

$$x \mapsto F_\varepsilon(x) + \frac{1}{2\tau} \|x - \bar{x}\|^2$$

be bounded from below; i.e., that there exists $C > 0$ such that

$$x \mapsto F_\varepsilon(x) + C \|x - \bar{x}\|^2$$

be bounded from below.

Example 8.1. We give a simple example that shows how the limit minimizing movement may depend on the choice of the mutual behavior of ε and τ . We consider the functions

$$F_\varepsilon(x) = \begin{cases} -x & \text{if } x \leq 0 \\ 0 & \text{if } 0 \leq x \leq \varepsilon \\ \varepsilon - x & \text{if } x \geq \varepsilon, \end{cases}$$

which converge uniformly to $F(x) = -x$. Note that the energies are not bounded from below, but their analysis falls within the framework in the previous remark. For this example a direct computation is immediately carried on. We consider a fixed initial datum x_0 .

If $x_0 > 0$, then for $\varepsilon < x_0$ we have $x_k^\varepsilon = x_{k-1}^\varepsilon + \tau$ for all $k \geq 0$.

If $x_0 \leq 0$ then we have $x_k^\varepsilon = x_{k-1}^\varepsilon + \tau$ if $x_{k-1}^\varepsilon \leq -\tau$. If $0 \geq x_{k-1}^\varepsilon > -\tau$ then $x_k^\varepsilon - x_{k-1}^\varepsilon$ is obtained by minimizing the function

$$f(y) = \begin{cases} -y + \frac{1}{2\tau}y^2 & \text{if } 0 \leq y \leq -x_{k-1}^\varepsilon \\ x_{k-1}^\varepsilon + \frac{1}{2\tau}y^2 & \text{if } -x_{k-1}^\varepsilon \leq y \leq -x_{k-1}^\varepsilon + \varepsilon \\ \varepsilon - y + \frac{1}{2\tau}y^2 & \text{if } y \geq -x_{k-1}^\varepsilon + \varepsilon, \end{cases}$$

whose minimizer is always $y = \tau + x_{k-1}^\varepsilon$ if $\varepsilon - x_{k-1}^\varepsilon > \tau$. In this case $x_k^\varepsilon = 0$. If otherwise $\varepsilon - x_{k-1}^\varepsilon \leq \tau$ the other possible minimizer is $y = \tau$. We then have to compare the values

$$f(-x_{k-1}^\varepsilon) = x_{k-1}^\varepsilon + \frac{1}{2\tau}(x_{k-1}^\varepsilon)^2, \quad f(\tau) = \varepsilon - \frac{1}{2}\tau.$$

We have three cases:

- (a) $\varepsilon - \frac{1}{2}\tau > 0$. In this case we have $x_k^\varepsilon = 0$ (and this holds for all subsequent steps).
- (b) $\varepsilon - \frac{1}{2}\tau < 0$. In this case we either have $f(\tau) < f(-x_{k-1}^\varepsilon)$, in which case $x_k^\varepsilon = x_{k-1}^\varepsilon + \tau$ (and this then holds for all subsequent steps); otherwise $x_k^\varepsilon = 0$ and $x_{k+1}^\varepsilon = x_k^\varepsilon + \tau$ (and this holds for all subsequent steps).
- (c) $\varepsilon - \frac{1}{2}\tau = 0$. If $x_{k-1}^\varepsilon < 0$ then $x_k^\varepsilon = 0$ (otherwise we already have $x_{k-1}^\varepsilon = 0$). Then, since we have the two solutions $y = 0$ and $y = \tau$, we have $x_j^\varepsilon = 0$ for $k \leq j \leq k_0$ for some $k_0 \in \mathbb{N} \cup +\infty$ and $x_j^\varepsilon = x_{j-1}^\varepsilon + \tau$ for $j > k_0$.

We can summarize the possible minimizing movements with initial datum $x_0 \leq 0$ as follows:

- (i) If $\tau < 2\varepsilon$ then the unique minimizing movement is $x(t) = \min\{x_0 + t, 0\}$.
- (ii) If $\tau > 2\varepsilon$ then the unique minimizing movement is $x(t) = x_0 + t$.
- (iii) If $\tau = 2\varepsilon$ then we have the family of minimizing movements (parameterized by $x_1 \leq x_0$) $x(t) = \max\{\min\{x_0 + t, 0\}, x_1 + t\}$.

For $x_0 > 0$ we always have the only minimizing movement $x(t) = x_0 + t$.

8.2 Commutability Along ‘Fast-Converging’ Sequences

We now show that, by suitably choosing the ε - τ regimes, the minimizing movement along the sequence F_ε from x_ε converges to a minimizing movement for the limit F from x_0 (‘fast-converging ε ’), while for other choices (‘fast-converging τ ’) the minimizing movement converges to a limit of minimizing movements for F_ε as $\varepsilon \rightarrow 0$. Heuristically, minimizing movements for all other regimes are ‘trapped’ between these two extrema.

Theorem 8.1. *Let F_ε be a equi-coercive sequence of (non-negative) lower-semicontinuous functionals on a Hilbert space X Γ -converging to F , let $x_\varepsilon \rightarrow x_0$. Then:*

- (i) *There exists a choice of $\varepsilon = \varepsilon(\tau)$ such that every minimizing movement along F_ε (and with time-step τ) with initial data x_ε is a minimizing movement for F from x_0 on $[0, T]$ for all T .*
- (ii) *There exists a choice of $\tau = \tau(\varepsilon)$ such that every minimizing movement along F_ε (and with time-step τ) with initial data x_ε is a limit of a sequence of minimizing movements for F_ε (for ε fixed) from x_ε on $[0, T]$ for all T .*

Proof. (i) Note that if $y_\varepsilon \rightarrow y_0$ then the solutions of

$$\min\left\{F_\varepsilon(x) + \frac{1}{2\tau}\|x - y_\varepsilon\|^2\right\} \quad (8.4)$$

converge to solutions of

$$\min\left\{F(x) + \frac{1}{2\tau}\|x - y_0\|^2\right\} \quad (8.5)$$

since we have a continuously converging perturbation of a Γ -converging sequence.

Let now $x_\varepsilon \rightarrow x_0$. Let τ be fixed. We consider the sequence $\{x_k^{\tau, \varepsilon}\}$ defined by iterated minimization of F_ε with initial point x_ε . Since $x_\varepsilon \rightarrow x_0$, up to subsequences we have $x_1^{\tau, \varepsilon} \rightarrow x_1^{\tau, 0}$, which minimizes

$$\min\left\{F(x) + \frac{1}{2\tau}\|x - x_0\|^2\right\}. \quad (8.6)$$

The point $x_2^{\tau,\varepsilon}$ converge to $x_2^{\tau,0}$. Since they minimize

$$\min \left\{ F_\varepsilon(x) + \frac{1}{2\tau} \|x - x_1^{\tau,\varepsilon}\|^2 \right\} \quad (8.7)$$

and $x_1^{\tau,\varepsilon} \rightarrow x_1^{\tau,0}$, their limit is a minimizer of

$$\min \left\{ F(x) + \frac{1}{2\tau} \|x - x_1^{\tau,0}\|^2 \right\}. \quad (8.8)$$

This operation can be repeated iteratively, obtaining (upon subsequences) $x_k^{\tau,\varepsilon} \rightarrow x_k^{\tau,0}$, and $\{x_k^{\tau,0}\}$ iteratively minimizes F with initial point x_0 . Since up to subsequences the trajectories $\{x_k^{\tau,0}\}$ converge to a minimizing movement for F with initial datum x_0 , the thesis follows by a diagonal argument.

- (ii) For fixed ε , the piecewise-constant functions $u^{\varepsilon,\tau}(t) = x_{\lfloor t/\tau \rfloor}^{\varepsilon,\tau}$ converge uniformly to a minimizing movement u^ε for F_ε with initial datum x_ε . By compactness, these u^ε converge uniformly to some function u as $\varepsilon \rightarrow 0$. Again, a diagonal argument gives the thesis. \square

Remark 8.2. Note that, given x_ε and F_ε , if F has more than one minimizing movement from x_0 then the approximation gives a choice criterion. As an example, take $F(x) = -|x|$, $F_\varepsilon(x) = -|x + \varepsilon|$ and $x_0 = x_\varepsilon = 0$.

Remark 8.3 (The convex case). If all F_ε are convex then it can be shown that, actually, the minimizing movement along the sequence F_ε always coincides with the minimizing movement for their Γ -limit. This (exceptional) case will be dealt with in detail separately in Chap. 11.

Example 8.2. In dimension one, we can take

$$F_\varepsilon(x) = \frac{1}{2}x^2 + \varepsilon W\left(\frac{x}{\varepsilon}\right),$$

where W is a one-periodic odd Lipschitz function with $\|W'\|_\infty = 1$. Up to addition of a constant is not restrictive to suppose that the average of W is 0. We check that the critical regime for the minimizing movements along F_ε is $\varepsilon \sim \tau$. Indeed, if $\varepsilon \ll \tau$ then from the estimate

$$\left| F_\varepsilon(x) - \frac{1}{2}x^2 \right| \leq \frac{\varepsilon}{2}$$

we deduce that

$$\frac{x_k - x_{k-1}}{\tau} = -x_k + O\left(\frac{\varepsilon}{\tau}\right),$$

and hence that the limit minimizing movement satisfies $u' = -u$, so that it corresponds to the minimizing movement of the limit $F_0(x) = \frac{1}{2}x^2$.

Conversely, if $\tau \ll \varepsilon$ then it may be seen that for $|x_0| \leq 1$ the motion is *pinned*; i.e., the resulting minimizing movement is the trivial solution $u(t) = x_0$ for all t . If $W \in C^2$ this is easily checked, since in this case the stationary solutions, corresponding to x satisfying

$$x + W'\left(\frac{x}{\varepsilon}\right) = 0$$

tend to be dense in the interval $[-1, 1]$ as $\varepsilon \rightarrow 0$. Moreover, in this regime the minimizing movement corresponds to the limit as $\varepsilon \rightarrow 0$ of the minimizing movements of F_ε for ε fixed; i.e., solutions u_ε of the gradient flow

$$u'_\varepsilon = -u_\varepsilon - W'\left(\frac{u_\varepsilon}{\varepsilon}\right).$$

Integrating between t_1 and t_2 we have

$$\int_{u_\varepsilon(t_1)}^{u_\varepsilon(t_2)} \frac{1}{s + W'(s/\varepsilon)} ds = t_1 - t_2.$$

By the uniform convergence $u_\varepsilon \rightarrow u$ we can pass to the limit, recalling that the integrand weakly converges to the function $1/g$ defined by

$$\frac{1}{g(s)} = \int_0^1 \frac{1}{s + W'(\sigma)} d\sigma,$$

and obtain the equation

$$u' = -g(u).$$

This equation corresponds to the minimizing movement for the even energy \tilde{F}_0 given for $x \geq 0$


$$\tilde{F}_0(x) = \begin{cases} 0 & \text{if } x \leq 1 \\ \int_1^x g(w) dw & \text{if } x \geq 1. \end{cases}$$

The plot of the derivatives of F_ε , F_0 and \tilde{F}_0 is reproduced in Fig. 8.1

We can explicitly compute the minimizing movement for $\tau \ll \varepsilon$; e.g., in the case

$$W(x) = \frac{1}{2\pi} \sin(2\pi x),$$

Fig. 8.1 The derivatives of F_ε , F_0 and \tilde{F}_0



which gives the equation

$$u' = \sqrt{u^2 - 1},$$

for $|x_0| \geq 1$, and

$$\tilde{F}_0(x) = \frac{1}{2} \left(|x| \sqrt{x^2 - 1} - \log(|x| + \sqrt{x^2 - 1}) \right)$$

for $|x| > 1$, and in the case

$$W(x) = \left| x - \frac{1}{2} \right| - \frac{1}{4} \quad \text{for } 0 \leq x \leq 1. \quad (8.9)$$

In the latter, the solutions with initial datum $x_0 > 1$ satisfy the equation

$$u' = \frac{1}{u} - u.$$

Integrating this limit equation we conclude that the minimizing movement along F_ε corresponds to that of the effective energy

$$\tilde{F}_0(x) = \left(\frac{1}{2} x^2 - \log |x| - \frac{1}{2} \right)^+.$$

Example 8.3 (Pinning threshold). In the previous example we have computed the critical regime $\varepsilon \sim \tau$, but we have not computed the minimizing movement for a fixed ratio ε/τ . In this case, a simpler interesting problem is the computation of the pinning threshold; i.e., the maximal value T such that $|x_0| \leq T$ gives in the limit a stationary minimizing movement. We have seen that for $\varepsilon \ll \tau$ we have $T = 0$, while for $\tau \ll \varepsilon$ we have $T = 1$. After considering the linearization of the problem above, the pinning threshold can be characterized as the greatest value T such that we have only stationary minimizing movements for the energies

$$F_\varepsilon^T(x) = Tx + \varepsilon W\left(\frac{x}{\varepsilon}\right).$$

In order to have an explicit description of $T = T(\gamma)$ in terms of $\gamma := \varepsilon/\tau$, we only treat the case of

$$W(x) = |x| \quad \text{for } |x| \leq \frac{1}{2}, \quad (8.10)$$

which gives the same limit as the one in (8.9). By comparison with the case $\tau \ll \varepsilon$, we have $T(\gamma) \leq 1$ for all γ .

By a comparison argument, it is not restrictive to suppose that $x_0 \in \varepsilon\mathbb{Z}$, and then by translation that $x_0 = 0$. The problem is then translated in the existence of negative minimizers for the problem

$$\min \left\{ Tx + \varepsilon W\left(\frac{x}{\varepsilon}\right) + \frac{1}{2\tau}x^2 \right\}.$$

Since $T \leq 1$ and $W' = -1$ in $[-\varepsilon/2, 0]$, this holds only if we have a negative value in $[-\varepsilon, -\varepsilon/2]$, or equivalently if

$$\begin{aligned} 0 &> \min \left\{ Tx + \varepsilon W\left(\frac{x}{\varepsilon}\right) + \frac{1}{2\tau}x^2 : -\varepsilon \leq x \leq -\varepsilon/2 \right\} \\ &= \min \left\{ (T+1)x + \varepsilon + \frac{1}{2\tau}x^2 : -\varepsilon \leq x \leq -\varepsilon/2 \right\}. \end{aligned}$$

Taking again into account that $T \leq 1$, it is easily seen that this minimum must be taken for $x = -\varepsilon$, so that the condition is equivalent to

$$0 > -T\varepsilon + \frac{1}{2\tau}\varepsilon^2; \text{ i.e., } T > \frac{\varepsilon}{2\tau}.$$


This proves that we have pinning for $T \leq \gamma/2$. In conclusion, the pinning threshold is

$$T(\gamma) = \min \left\{ \frac{\gamma}{2}, 1 \right\}$$

(see Fig. 8.2). As $\gamma \rightarrow 0$ and $\gamma \rightarrow +\infty$ we recover the thresholds in the limit cases.

8.2.1 Relaxed Evolution

In Theorem 8.1 we have considered, as usual for simplicity, the Γ -convergence with respect to the topology in X . In this way we characterize the convergence of solutions to problems (8.4) to solutions of problems (8.5) in terms of the Γ -limit. This is the only argument where we have used the definition of F in the proof of

Fig. 8.2 Pinning threshold in dependence of the ratio ε/τ 

Theorem 8.1(i). We may consider the Γ -limits with respect to weaker topologies, for which we have coerciveness but the distance term is not a continuous perturbation. In analogy with what already observed for quasistatic motions in Chap. 3 (see, e.g., Sect. 3.1.5), the proof of Theorem 8.1(i) can be repeated, upon defining a *relaxed limit motion*, where the minimizing movement for F is replaced by the limit of u^τ defined by successive minimizing

$$\min_X \mathcal{F}_\tau^{x_{k-1}}(x),$$

where

$$\mathcal{F}_\tau^y(x) = \Gamma\text{-}\lim_{\varepsilon \rightarrow 0} \left(F_\varepsilon(x) + \frac{1}{2\tau} \|x - y\|^2 \right). \quad (8.11)$$

The study of this more general minimizing movements is beyond the scope of these notes. We only give a simple example.

Example 8.4. Consider $X = L^2(0, 1)$ and

$$F_\varepsilon(u) = \int_0^1 a\left(\frac{x}{\varepsilon}\right) u^2 dx,$$

where a is 1-periodic and $0 < \alpha \leq a(y) \leq \beta < +\infty$ for some constants α and β . Then F_ε is equicoercive with respect to the *weak- L^2* topology, and its limit is $\underline{a} \int_0^1 u^2 dx$ (\underline{a} the harmonic mean of a). However, the perturbations with the L^2 -distance are not continuous, and the limits in (8.11) with respect to the weak topology are easily computed as

$$\begin{aligned} \mathcal{F}_\tau^v(u) &= \Gamma\text{-}\lim_{\varepsilon \rightarrow 0} \left(F_\varepsilon(u) + \frac{1}{2\tau} \|u - v\|^2 \right) \\ &= \Gamma\text{-}\lim_{\varepsilon \rightarrow 0} \int_0^1 \left(a\left(\frac{x}{\varepsilon}\right) + \frac{1}{2\tau} \right) u^2 + \frac{(v^2 - 2uv)}{2\tau} dx \end{aligned}$$

$$\begin{aligned}
&= \int_0^1 \left(\underline{a}_\tau u^2 + \frac{(v^2 - 2uv)}{2\tau} \right) dx \\
&= \int_0^1 \left(\underline{a}_\tau - \frac{1}{2\tau} \right) u^2 dx + \frac{1}{2\tau} \|u - v\|^2,
\end{aligned}$$

where

$$\underline{a}_\tau = \left(\int_0^1 \frac{1}{\left(a(y) + \frac{1}{2\tau}\right)} dy \right)^{-1}.$$

A series expansion argument easily yields that

$$\begin{aligned}
\underline{a}_\tau &= \frac{1}{2\tau} \left(\int_0^1 \frac{1}{2\tau a(y) + 1} dy \right)^{-1} \\
&= \frac{1}{2\tau} \left(\int_0^1 \left(1 - 2\tau a(y) + O(\tau^2) \right) dy \right)^{-1} \\
&= \frac{1}{2\tau} \left(1 + 2\tau \int_0^1 a(y) dy + O(\tau^2) \right) = \frac{1}{2\tau} + \bar{a} + O(\tau),
\end{aligned}$$

where \bar{a} is the arithmetic mean of a . We then obtain that the limit of u^τ coincides with the minimizing motion for \tilde{F} given by

$$\tilde{F}(u) = \bar{a} \int_0^1 u^2 dx.$$

The same argument leading to an effective motion can be applied to varying distances as in the following example.

Example 8.5. We consider $X_\varepsilon = X = L^2(0, 1)$ equipped with the distance d_ε given by

$$d_\varepsilon^2(u, v) = \int_0^1 a\left(\frac{x}{\varepsilon}\right) |u - v|^2 dx,$$

and $F_\varepsilon(u) = F(u) = \int_0^1 |u'|^2 dx$. For fixed v the square distances can be seen as functionals depending on v , weakly equicoercive in L^2 and Γ -converging to $\frac{a}{2} \|u - v\|^2$ ($\|u\|$ the L^2 -norm). Nevertheless, in this case the functionals $F_\varepsilon(u) + \frac{1}{2\varepsilon} d_\varepsilon^2(u, v)$ are coercive with respect to the strong L^2 -norm and Γ -converge to $F(u) + \frac{1}{2\tau} \bar{a} \|u - v\|^2$. As a conclusion, the minimizing movement coincide with the minimizing movement for F with respect to the norm $\sqrt{\bar{a}} \|u\|$ or, equivalently, with the minimizing movement for $\frac{1}{\bar{a}} F$ with respect to the L^2 -norm.

8.3 An Example: ‘Overdamped Dynamics’ of Lennard-Jones Interactions

We now give an example of a sequence of non-convex energies which commute with the minimizing movement procedure.

Let J be as in Sect. 4.4 and $\frac{1}{\varepsilon} = N \in \mathbb{N}$. We consider the energies

$$F_\varepsilon(u) = \sum_{i=1}^N J\left(\frac{u_i - u_{i-1}}{\sqrt{\varepsilon}}\right)$$

with the periodic boundary condition $u_N = u_0$. As proved in Sect. 4.4, after identification of u with a piecewise-constant function on $[0, 1]$, these energies Γ -converge to the energy

$$F(u) = \int_0^1 |u'|^2 dt + \#(S(u) \cap [0, 1)), \quad u^+ > u^-,$$

defined on piecewise- H^1 functions, in this case extended 1-periodically on the whole real line.

In this section we apply the minimizing movements scheme to F_ε as a sequence of functionals in $L^2(0, 1)$. In order to have initial data u_0^ε with equibounded energy, we may suppose that these are the discretization of a single piecewise- H^1 function u_0 (with a slight abuse of notation we will continue to denote all those discrete functions by u_0).

With fixed ε and τ , the time-discretization scheme consists in defining recursively u^k as a minimizer of

$$u \mapsto \sum_{i=1}^N J\left(\frac{u_i - u_{i-1}}{\sqrt{\varepsilon}}\right) + \frac{1}{2\tau} \sum_{i=1}^N \varepsilon |u_i - u_i^{k-1}|^2. \quad (8.12)$$

By Proposition 8.1, upon extraction of a subsequence, the functions $u^\tau(t) = u_{\lfloor t/\tau \rfloor}$ converge uniformly in L^2 to a function $u \in C^{1/2}([0, +\infty); L^2(0, 1))$. Moreover, since we have $F(u(t)) \leq F(u_0) < +\infty$, $u(t)$ is a piecewise- H^1 function for all t .

We now describe the motion of the limit u . For the sake of simplicity we suppose that u_0 is a piecewise-Lipschitz function and that $S(u_0) \cap \{\varepsilon i : i \in \{1, \dots, N\}\} = \emptyset$ (so that we do not have any ambiguity in the definition of the interpolations of u_0).

We first write down the Euler–Lagrange equations for u^k , which simply amount to a N -dimensional system of equations obtained by deriving (8.12) with respect to u_i

$$\frac{1}{\sqrt{\varepsilon}} \left(J'\left(\frac{u_i^k - u_{i-1}^k}{\sqrt{\varepsilon}}\right) - J'\left(\frac{u_{i+1}^k - u_i^k}{\sqrt{\varepsilon}}\right) \right) + \frac{\varepsilon}{\tau} (u_i^k - u_i^{k-1}) = 0. \quad (8.13)$$

- With fixed $i \in \{1, \dots, N\}$ let v_k be defined by

$$v_k = \frac{u_i^k - u_{i-1}^k}{\varepsilon}.$$

For simplicity of notation, we set

$$J_\varepsilon(w) = \frac{1}{\varepsilon} J(\sqrt{\varepsilon} w).$$

By (8.13) and the corresponding equation for $i - 1$, which can be rewritten as

$$J'_\varepsilon\left(\frac{u_{i-1}^k - u_{i-2}^k}{\varepsilon}\right) - J'_\varepsilon\left(\frac{u_i^k - u_{i-1}^k}{\varepsilon}\right) + \frac{\varepsilon}{\tau}(u_{i-1}^k - u_{i-1}^{k-1}) = 0,$$

we have

$$\begin{aligned} \frac{v_k - v_{k-1}}{\tau} &= \frac{1}{\tau} \left(\frac{u_i^k - u_{i-1}^k}{\varepsilon} - \frac{u_i^{k-1} - u_{i-1}^{k-1}}{\varepsilon} \right) \\ &= \frac{1}{\varepsilon} \left(\frac{u_i^k - u_i^{k-1}}{\tau} - \frac{u_{i-1}^k - u_{i-1}^{k-1}}{\tau} \right) \\ &= \frac{1}{\varepsilon^2} \left(\left(J'_\varepsilon\left(\frac{u_{i-1}^k - u_{i-2}^k}{\varepsilon}\right) - J'_\varepsilon\left(\frac{u_i^k - u_{i-1}^k}{\varepsilon}\right) \right) \right. \\ &\quad \left. - \left(J'_\varepsilon\left(\frac{u_i^k - u_{i-1}^k}{\varepsilon}\right) - J'_\varepsilon\left(\frac{u_{i+1}^k - u_i^k}{\varepsilon}\right) \right) \right), \end{aligned}$$

so that

$$\begin{aligned} \frac{v_k - v_{k-1}}{\tau} - \frac{2}{\varepsilon^2} J'_\varepsilon(v_k) &= -\frac{1}{\varepsilon^2} \left(J'_\varepsilon\left(\frac{u_{i-1}^k - u_{i-2}^k}{\varepsilon}\right) + J'_\varepsilon\left(\frac{u_{i+1}^k - u_i^k}{\varepsilon}\right) \right) \\ &\geq -\frac{2}{\varepsilon^2} J'_\varepsilon\left(\frac{w_0}{\sqrt{\varepsilon}}\right). \end{aligned} \tag{8.14}$$

We recall that we denote by w_0 the maximum point of J' .

We can read (8.14) as an inequality for the difference system

$$\frac{v_k - v_{k-1}}{\eta} - 2J'_\varepsilon(v_k) \geq -2J'_\varepsilon\left(\frac{w_0}{\sqrt{\varepsilon}}\right),$$

where $\eta = \tau/\varepsilon^2$ is interpreted as a discretization step. Note that $v_k = w_0/\sqrt{\varepsilon}$ for all k is a stationary solution of the equation

$$\frac{v_k - v_{k-1}}{\eta} - 2J'_\varepsilon(v_k) = -2J'_\varepsilon\left(\frac{w_0}{\sqrt{\varepsilon}}\right)$$

and that J'_ε are equi-Lipschitz functions on $[0, +\infty)$. If $\eta \ll 1$ this implies that if $v_{k_0} \leq w_0/\sqrt{\varepsilon}$ for some k_0 then

$$v_k \leq \frac{w_0}{\sqrt{\varepsilon}} \quad \text{for } k \geq k_0,$$

or, equivalently, that if $\tau \ll \varepsilon^2$ the set

$$S_\varepsilon^k = \left\{ i \in \{1, \dots, N\} : \frac{u_i^k - u_{i-1}^k}{\varepsilon} \geq \frac{w_0}{\sqrt{\varepsilon}} \right\}$$

is decreasing with k . By our assumption on u_0 , for ε small enough we then have

$$S_\varepsilon^0 = \left\{ i \in \{1, \dots, N\} : [\varepsilon(i-1), \varepsilon i] \cap S(u_0) \neq \emptyset \right\},$$

so that, passing to the limit

$$S(u(t)) \subseteq S(u_0) \quad \text{for all } t \geq 0. \quad (8.15)$$

- Taking into account that we may define

$$u^\tau(t, x) = u_{\lfloor \frac{t}{\tau} \rfloor}^{\lfloor \frac{x}{\varepsilon} \rfloor},$$

we may choose functions $\phi \in C_0^\infty(0, T)$ and $\psi \in C_0^\infty(x_1, x_2)$, with $(x_1, x_2) \cap S(u_0) = \emptyset$, and obtain from (8.13)

$$\begin{aligned} & \int_0^T \int_{x_1}^{x_2} u^\tau(t, x) \left(\frac{\phi(t) - \phi(t + \tau)}{\tau} \right) \psi(x) dx dt \\ &= - \int_0^T \int_{x_1}^{x_2} \left(\frac{1}{\sqrt{\varepsilon}} J' \left(\sqrt{\varepsilon} \frac{u^\tau(t, x) - u^\tau(t, x - \varepsilon)}{\varepsilon} \right) \right) \\ & \quad \times \phi(t) \left(\frac{\psi(x) - \psi(x + \varepsilon)}{\varepsilon} \right) dx dt. \end{aligned}$$

Taking into account that

$$\lim_{\varepsilon \rightarrow 0} \frac{1}{\sqrt{\varepsilon}} J'(\sqrt{\varepsilon} w) = 2w,$$

we can pass to the limit and obtain that

$$- \int_0^T \int_{x_1}^{x_2} u(t, x) \phi'(t) \psi(x) dx dt = \int_0^T \int_{x_1}^{x_2} 2 \frac{\partial u}{\partial x} \phi(t) \psi'(x) dx dt;$$

i.e., that

$$\frac{\partial u}{\partial t} = -2 \frac{\partial^2 u}{\partial x^2} \quad (8.16)$$

in the sense of distributions (and hence also classically) in $(0, T) \times (x_1, x_2)$. By the arbitrariness of the interval (x_1, x_2) we have that equation (8.16) is satisfied for x in $(0, 1) \setminus S(u_0)$.

- We now derive boundary conditions on $S(u(t))$. Let $i_0 + 1$ belong to S_ε^0 , and suppose that $u^+(t, x) - u^-(t, x) \geq c > 0$. Then we have

$$\lim_{\tau \rightarrow 0} \frac{1}{\sqrt{\varepsilon}} J' \left(\frac{u_{i_0}^{\lfloor t/\tau \rfloor} - u_{i_0-1}^{\lfloor t/\tau \rfloor}}{\sqrt{\varepsilon}} \right) = 0.$$

If $i < i_0$, from (8.13) it follows, after summing up the indices from i to i_0 , that

$$\sum_{j=i}^{i_0} \frac{\varepsilon}{\tau} (u_j^k - u_j^{k-1}) = -\frac{1}{\sqrt{\varepsilon}} J' \left(\frac{u_i^k - u_{i-1}^k}{\sqrt{\varepsilon}} \right). \quad (8.17)$$

We may choose $i = i_\varepsilon$ such that $\varepsilon i_\varepsilon \rightarrow \bar{x}$ and we may deduce from (8.17) that

$$\int_{\bar{x}}^{x_0} \frac{\partial u}{\partial t} dx = -2 \frac{\partial u}{\partial x}(\bar{x}),$$


where $x_0 \in S(u(t))$ is the limit of εi_0 . Letting $\bar{x} \rightarrow x_0^-$ we obtain

$$\frac{\partial u}{\partial x}(x_0^-) = 0.$$

Similarly we obtain the homogeneous Neumann condition at x_0^+ .

Summarizing, the minimizing movement along the scaled Lennard-Jones energies F_ε from a piecewise- H^1 function consists in a piecewise- H^1 motion, following the heat equation on $(0, 1) \setminus S(u_0)$, with homogeneous Neumann boundary conditions on $S(u_0)$ (as long as $u(t)$ has a discontinuity at the corresponding point of $S(u_0)$).

Note that for $\varepsilon \rightarrow 0$ sufficiently fast Theorem 8.1 directly ensures that the minimizing movement along F_ε coincides with the minimizing movement for the functional F . The computation above shows that this holds also for $\tau \ll \varepsilon^2$ (i.e., $\varepsilon \rightarrow 0$ ‘sufficiently slow’), which then must be regarded as a technical condition.

Fig. 8.3 The function F_ε 

8.4 Homogenization of Minimizing Movements

We now examine minimizing movements along oscillating sequences (with many local minima), treating two model cases in the real line.

8.4.1 Minimizing Movements for Piecewise-Constant Energies

We apply the minimizing-movement scheme to the functions

$$F_\varepsilon(x) = -\left\lfloor \frac{x}{\varepsilon} \right\rfloor \varepsilon$$

converging to $F(x) = -x$ (see Fig. 8.3). This is a prototype of a function with many local minimizers (actually, in this case all points are local minimizers) converging to a function with few local minimizers (actually, none).

Note that, with fixed ε , for any initial datum x_0 the minimizing movement for F_ε is trivial: $u(t) = x_0$, since all points are local minimizers. Conversely, the corresponding minimizing movement for the limit is $u(t) = x_0 + t$.


We now fix an initial datum x_0 , the space scale ε and the time scale τ , and examine the successive-minimization scheme from x_0 . Note that it is not restrictive to suppose that $0 \leq x_0 < 1$ up to a translation in $\varepsilon\mathbb{Z}$.

The first minimization, giving x_1 , is

$$\min \left\{ F_\varepsilon(x) + \frac{1}{2\tau}(x - x_0)^2 \right\}. \quad (8.18)$$

The function to minimize is pictured in Fig. 8.4 in normalized coordinates ($\varepsilon = 1$); note that it equals $-x + \frac{1}{2\tau}(x - x_0)^2$ if $x \in \varepsilon\mathbb{Z}$.

Fig. 8.4 The function in the minimization problem (8.18)



Apart from some exceptional cases that we deal separately below, we have two possibilities:

- (i) If $\frac{\tau}{\varepsilon} < \frac{1}{2}$ then the motion is trivial. The value $1/2$ is the **pinning threshold**.

Indeed, after setting $x_0 = s\varepsilon$ with $0 \leq s < 1$, we have two sub-cases:

- (a) The minimizer x_1 belongs to $[0, \varepsilon)$. This occurs exactly if $F_\varepsilon(\varepsilon) + \frac{1}{2\varepsilon}(\varepsilon - x_0)^2 > 0$; i.e.,

$$\tau < \frac{(s-1)^2\varepsilon}{2}. \quad (8.19)$$

In this case the only minimizer is the initial datum x_0 . This implies that we have $x_k = x_0$ for all k .

- (b) We have that $x_1 = \varepsilon$. This implies that, up to a translation we are in the case $x_0 = 0$ with $s = 0$, and (8.19) holds since $\tau < \frac{\varepsilon}{2}$. Hence, $x_k = x_1$ for all $k \geq 1$.

- (ii) If $\frac{\tau}{\varepsilon} > \frac{1}{2}$ then for ε small the minimum is taken on $\varepsilon\mathbb{Z}$. So that again we may suppose that $x_0 = 0$.

Note that we are leaving out for the time being the case when $x_0 = 0$ and $\frac{\tau}{\varepsilon} = \frac{1}{2}$. In that case we have a double choice for the minimizer; such situations will be examined separately.

If $x_0 = 0$ then x_1 is computed by solving

$$\min \left\{ F_\varepsilon(x) + \frac{1}{2\varepsilon}x^2 : x \in \varepsilon\mathbb{Z} \right\}, \quad (8.20)$$

and is characterized by

$$x_1 - \frac{1}{2}\varepsilon \leq \tau \leq x_1 + \frac{1}{2}\varepsilon.$$

We then have

$$x_1 = \left\lfloor \frac{\tau}{\varepsilon} + \frac{1}{2} \right\rfloor \varepsilon \quad \text{if } \frac{\tau}{\varepsilon} + \frac{1}{2} \notin \mathbb{Z}$$

(note again that we have two solutions for $\frac{\tau}{\varepsilon} + \frac{1}{2} \in \mathbb{Z}$, which also includes the case $\frac{\tau}{\varepsilon} = \frac{1}{2}$ already set aside, and we examine those cases separately). The same computation is repeated at each k giving

$$\frac{x_k - x_{k-1}}{\tau} = \left\lfloor \frac{\tau}{\varepsilon} + \frac{1}{2} \right\rfloor \frac{\varepsilon}{\tau}.$$

We can now choose τ and ε tending to 0 simultaneously and pass to the limit. The behaviour of the limit minimizing movements is governed by the quantity

$$w = \lim_{\varepsilon \rightarrow 0} \frac{\tau}{\varepsilon}, \quad (8.21)$$

which we may suppose exists up to subsequences. If $w + \frac{1}{2} \notin \mathbb{Z}$ then the minimizing movement along F_ε from x_0 is uniquely defined by

$$u(t) = x_0 + vt, \text{ with } v = \left\lfloor w + \frac{1}{2} \right\rfloor \frac{1}{w}, \quad (8.22)$$


so that the whole sequence converges if the limit in (8.21) exists. Note that

- **(pinning)** we have $v = 0$ exactly when $\frac{\tau}{\varepsilon} < \frac{1}{2}$ for ε small. In particular this holds for $\tau \ll \varepsilon$ (i.e., for $w = 0$).
- **(limit motion for slow times)** if $\varepsilon \ll \tau$ then the motion coincides with the gradient flow of the limit, with velocity 1.
- **(discontinuous dependence of the velocity)** the velocity is a discontinuous function of w at points of $\frac{1}{2} + \mathbb{Z}$. Note moreover that it may be actually greater than the limit velocity 1. The graph of v is pictured in Fig. 8.5.
- **(non-uniqueness at $w \in \frac{1}{2} + \mathbb{Z}$)** in these exceptional cases we may have either of the two velocities $1 + \frac{1}{2w}$ or $1 - \frac{1}{2w}$ in the cases $\frac{\varepsilon}{\tau} + \frac{1}{2} > w$ or $\frac{\varepsilon}{\tau} + \frac{1}{2} < w$ for all ε small respectively, but we may also have any $u(t)$ with

$$1 - \frac{1}{2w} \leq u'(t) \leq 1 + \frac{1}{2w}$$

if we have precisely $\frac{\varepsilon}{\tau} + \frac{1}{2} = w$ for all ε small, since in this case at every time step we may choose any of the two minimizers giving the extremal velocities, and then obtain any such u' as a weak limit of piecewise constant functions taking only those two values. Note therefore that in this case the limit is not determined only by w , and in particular it may depend on the subsequence even if the limit (8.21) exists.

Fig. 8.5 The velocity v in terms of w



We remark that the functions F_ε above can be substituted by functions with isolated local minimizers; e.g. by taking ($\alpha > 0$)

$$F_\varepsilon(x) = -\left\lfloor \frac{x}{\varepsilon} \right\rfloor \varepsilon + \alpha \left(x - \left\lfloor \frac{x}{\varepsilon} \right\rfloor \varepsilon \right),$$

with isolated local minimizers at $\varepsilon\mathbb{Z}$ (for which the computations run exactly as above), or

$$F_\varepsilon(x) = -x + (1 + \alpha)\varepsilon \sin\left(\frac{x}{\varepsilon}\right).$$

Note that the presence of an energy barrier between local minimizers does not influence the velocity of the final minimizing movement, that can always be larger than 1 (the velocity as $\varepsilon \ll \tau$).


We also remark that the same result can be obtained by a ‘discretization’ of F ; i.e., taking

$$F_\varepsilon(x) = \begin{cases} -x & \text{if } x \in \varepsilon\mathbb{Z} \\ +\infty & \text{otherwise.} \end{cases} \quad (8.23)$$

8.4.2 A Heterogeneous Case

We briefly examine a variation of the previous example obtained by introducing a heterogeneity parameter $1 \leq \lambda \leq 2$ and defining

$$F^\lambda(x) = \begin{cases} -2\left\lfloor \frac{x}{2} \right\rfloor & \text{if } 2\left\lfloor \frac{x}{2} \right\rfloor \leq x < 2\left\lfloor \frac{x}{2} \right\rfloor + \lambda \\ -2\left\lfloor \frac{x}{2} \right\rfloor - \lambda & \text{if } 2\left\lfloor \frac{x}{2} \right\rfloor + \lambda \leq x < 2\left\lfloor \frac{x}{2} \right\rfloor + 1. \end{cases} \quad (8.24)$$

Fig. 8.6 The function F^λ 

If $\lambda = 1$ we are in the previous situation; for general λ the function F^λ is pictured in Fig. 8.6.

We apply the minimizing-movement scheme to the functions

$$F_\varepsilon(x) = F_\varepsilon^\lambda(x) = \varepsilon F^\lambda\left(\frac{x}{\varepsilon}\right).$$

Arguing as above, we can reduce to the two cases

$$(a) x_k \in 2\varepsilon\mathbb{Z}, \quad \text{or} \quad (b) x_k \in 2\varepsilon\mathbb{Z} + \varepsilon\lambda.$$

Taking into account that x_{k+1} is determined as the point in $2\varepsilon\mathbb{Z} \cup (2\varepsilon\mathbb{Z} + \varepsilon\lambda)$ closer to τ (as above, we only consider the cases when we have a unique solution to the minimum problems in the iterated procedure), we can characterize it as follows.

In case (a) we have the two sub cases:

(a₁) If we have

$$2n < \frac{\tau}{\varepsilon} - \frac{\lambda}{2} < 2n + 1$$

for some $n \in \mathbb{N}$ then

$$x_{k+1} = x_k + (2n + \lambda)\varepsilon.$$

In particular $x_{k+1} \in 2\varepsilon\mathbb{Z} + \varepsilon\lambda$.

(a₂) If we have

$$2n - 1 < \frac{\tau}{\varepsilon} - \frac{\lambda}{2} < 2n$$

for some $n \in \mathbb{N}$ then

$$x_{k+1} = x_k + 2n\varepsilon.$$

In particular $x_{k+1} \in 2\varepsilon\mathbb{Z}$. Note that $x_{k+1} = x_k$ (pinning) if $\frac{\tau}{\varepsilon} < \frac{\lambda}{2}$.

In case (b) we have the two sub cases:

(b₁) If we have

$$2n < \frac{\tau}{\varepsilon} + \frac{\lambda}{2} < 2n + 1$$

for some $n \in \mathbb{N}$ then

$$x_{k+1} = x_k + 2n\varepsilon.$$

In particular $x_{k+1} \in 2\varepsilon\mathbb{Z} + \varepsilon\lambda$. Note that $x_{k+1} = x_k$ (pinning) if $\frac{\tau}{\varepsilon} < 1 - \frac{\lambda}{2}$, which is implied by the pinning condition in (a₂).

(b₂) If we have

$$2n - 1 < \frac{\tau}{\varepsilon} + \frac{\lambda}{2} < 2n$$

for some $n \in \mathbb{N}$ then

$$x_{k+1} = x_k + 2n\varepsilon - \varepsilon\lambda.$$

In particular $x_{k+1} \in 2\varepsilon\mathbb{Z}$.

Eventually, we have the two cases:

(1) When

$$\left| \frac{\tau}{\varepsilon} - 2n \right| < \frac{\lambda}{2}$$


for some $n \in \mathbb{N}$ then, after possibly one iteration, we are either in the case (a₂) or (b₁). Hence, either $x_k \in 2\varepsilon\mathbb{Z}$ or $x_k \in 2\varepsilon\mathbb{Z} + \varepsilon\lambda$ for all k . The velocity in this case is then

$$\frac{x_{k+1} - x_k}{\tau} = 2n \frac{\varepsilon}{\tau}.$$

(2) When

$$\left| \frac{\tau}{\varepsilon} - (2n + 1) \right| < 1 - \frac{\lambda}{2}$$

Fig. 8.7 The function f describing the effective velocity



for some $n \in \mathbb{N}$ then we are alternately in case (a₁) or (b₂). In this case we have an

- **averaged velocity:** the speed of the orbit $\{x_k\}$ oscillates between two values with an average speed given by

$$\frac{x_{k+2} - x_k}{2\tau} = \frac{2n\varepsilon + \lambda\varepsilon}{2\tau} + \frac{2(n+1)\varepsilon - \lambda\varepsilon}{2\tau} = (2n+1)\frac{\varepsilon}{\tau}.$$


This is an additional feature with respect to the previous example.

Summarizing, if we define w as in (8.21) then (taking into account only the cases with a unique limit) the minimizing movement along the sequence F_ε with initial datum x_0 is given by $x(t) = x_0 + vt$ with $v = \frac{1}{w}f(w)$, and f is given by

$$f(w) = \begin{cases} 2n & \text{if } |w - 2n| \leq \frac{\lambda}{2}, n \in \mathbb{N} \\ 2n + 1 & \text{if } |w - (2n + 1)| < 1 - \frac{\lambda}{2}, n \in \mathbb{N} \end{cases}$$

(see Fig. 8.7). Note that the pinning threshold is now $\lambda/2$. We can compare this minimizing movement with the one given in (8.22) by examining the graph of $w \mapsto \lfloor w + 1/2 \rfloor - f(w)$ in Fig. 8.8. For $2n + 1/2 < w < 2n + \lambda/2$ the new minimizing movement is slower, while for $2n + 2 - \lambda/2 < w < 2n + 2 - 1/2$ it is faster.

Fig. 8.8 Comparison with the homogeneous case



8.4.3 A Proposal for Some Random Models

From the heterogeneous example above we may derive two possible random models, of which we may then study the corresponding minimizing movement. We only give a heuristic proposal, which can then be correctly formalized by introducing suitable random variables.

1. Random environment. Let $\lambda \in (1/2, 1)$ and $p \in [0, 1]$. We consider a random array of points $\{x_i^\omega\}$ in \mathbb{R} such that, e.g.,

$$x_i^\omega - x_{i-1}^\omega = \begin{cases} \lambda & \text{with probability } p \\ 2 - \lambda & \text{with probability } 1 - p. \end{cases} \quad (8.25)$$

With fixed ω we may consider the minimizing movement related to

$$F_\varepsilon^\omega(x) = \begin{cases} -x & \text{if } x \in \{\varepsilon x_i^\omega : i \in \mathbb{Z}\} \\ +\infty & \text{otherwise,} \end{cases}$$

or equivalently (as in the definition (8.23))

$$F_\varepsilon^\omega(x) = -\varepsilon x_i^\omega \text{ if } x \in [\varepsilon x_i^\omega, \varepsilon x_{i+1}^\omega), i \in \mathbb{Z}.$$

In the case $p = 0$ or $p = 1$ we almost surely have a homogeneous environment as in Sect. 8.4.1. For $p = 1/2$ we have a random version of the heterogeneous model of Sect. 8.4.2. Note that in this case for all $p \in (0, 1)$ the pinning threshold for the ratio τ/ε is almost surely $\lambda/2$, since below that value, the motion will be pinned at the first index i with $x_i^\omega - x_{i-1}^\omega = \lambda$; i.e., almost surely after a finite number of steps. For $\tau/\varepsilon = \lambda/2$ and $\lambda < 2/3$ (with this condition we always move of one index) then the (maximal) velocity after pinning is $v = \lambda p + (1-p)$ (for $\lambda > 2/3$ the computation of the velocity involves the probability of m -consecutive points x_i^ω at distance $2 - \lambda$).

2. Random movements. Let $\lambda \in (1/2, 1)$ and $p \in [0, 1]$. Contrary to the model above, we suppose that at every time step k we may make a random choice of points $\{x_i^{\omega_k}\}$ satisfying (8.25) such that $x_k^\varepsilon \in \{x_i^{\omega_k}\}$; i.e., this choice now represents the random possibility of motion of the point itself (and not a characteristic of the medium). Note that in this case for $p \in (0, 1)$ the pinning threshold for the ratio τ/ε is almost surely the lower value $1 - \frac{\lambda}{2}$, and the (maximal) velocity after pinning is $v = (2 - \lambda)(1 - p)$.

8.5 Time-Dependent Minimizing Movements

Following the arguments of Sect. 7.2 we can define a minimizing movement along a time-dependent sequence of energies $F_\varepsilon(x, t)$, upon some technical assumptions as in (7.10). In this case we fix a sequence of initial data x_0^ε and $\tau = \tau_\varepsilon \rightarrow 0$, and define recursively x_k^ε as minimizing

$$\min \left\{ F_\varepsilon(x, k\tau) + \frac{1}{2\tau} \|x - x_{k-1}^\varepsilon\|^2 \right\}. \quad (8.26)$$

A minimizing movement is then any limit u of u^ε defined by $u^\varepsilon(t) = x_{\lfloor t/\tau \rfloor}^\varepsilon$.

We only give a simple one-dimensional example with a time-dependent forcing term.

Example 8.6. We consider

$$F_\varepsilon(x, t) = \varepsilon W\left(\frac{x}{\varepsilon}\right) - tx$$

with W as in Example 8.2. Similarly to that example we can check that $\varepsilon \sim \tau$ is the critical case, and we can explicitly describe the minimizing movement in the extreme cases:

- ($\varepsilon \ll \tau$) the minimizing movement is that corresponding to $F_0(x, t) = -tu$; i.e., to the equation $u' = t$.
- ($\tau \ll \varepsilon$) the minimizing movement is that corresponding to the function

$$\tilde{F}_0(t, x) = \begin{cases} 0 & \text{if } t \leq 1 \\ g(t)u & \text{if } t \geq 1, \end{cases}$$

where g is now defined by

$$\frac{1}{g(t)} = \int_0^1 \frac{1}{W'(\sigma) - t} d\sigma.$$

8.6 Varying Dissipations: BV-Solutions of Evolution Equations

In the previous sections of this chapter, we have limited ourselves to a Hilbert setting. This often rules out interesting applications, in particular a viscosity approach to quasistatic motion as a limit of gradient flows, which is obtained by perturbing a positively one-homogeneous dissipation \mathcal{D} by a sequence $\mathcal{D} + \frac{1}{\tau}\mathcal{D}_\varepsilon$ for which a gradient flow-type motion can be defined using the minimizing-movement approach. In general, the limit of these gradient flows gives a motion, called *BV-solution*, which is different from the energetic solution as defined in Sect. 3.2, and can be characterized in a variational way different from the energy balance. A treatment of this subject is beyond the scope of these notes, since it would need a too refined introduction to the theory of gradient flows in metric spaces, even though it would fit the spirit of the book since it may be stated in terms of Γ -limits. Many of the arguments followed above for varying energies also hold for varying dissipations.

We only deal with a simple example, in order to highlight the differences with energetic solutions.

Example 8.7 (Nonconvex mechanical play). We can consider the double-well potential in Example 3.3 and the perturbed dissipations

$$\mathcal{F}(t, x) = \frac{1}{2} \min\{(x-1)^2, (x+1)^2\} - tx, \quad \mathcal{D}_{\varepsilon, \tau}(x) = |x| + \frac{\varepsilon}{2\tau} x^2,$$

with $x_0 \in [-2, -1]$. Then the sequence x_k^τ is increasing and minimizes

$$\min \left\{ \frac{1}{2} \min\{(x-1)^2, (x+1)^2\} - (k\tau - 1)x - x_{k-1}^\tau + \frac{\varepsilon}{2\tau} (x - x_{k-1}^\tau)^2 : x \geq x_{k-1}^\tau \right\}.$$

We fix the ratio


$$\gamma = \frac{\varepsilon}{\tau}. \tag{8.27}$$

With a computation similar to the one in Example 3.3, we obtain as limit the solution

$$x(t) = \begin{cases} x_0 & \text{if } t \leq x_0 + 2 \\ t - 2 & \text{if } x_0 \leq t \leq 2 - \frac{1}{\gamma+1} \\ t & \text{if } t > 2 - \frac{1}{\gamma+1} \end{cases}$$

or the one equal to this except for $t = 2 - \frac{1}{\gamma+1}$ where $x = t$.

Fig. 8.9 Interpolation between energetic and BV solutions



Remark 8.4 (Interpolations of energetic and BV solutions). In the previous example, the case $\varepsilon \ll \tau$ (formally, $\gamma = 0$) gives the energetic solution obtained in Example 3.3. The case $\tau \ll \varepsilon$ (formally, $\gamma = +\infty$) corresponds to the BV-solution hinted at above. The case in which (8.27) holds can be interpreted as an interpolation between these two extreme case, and is pictured in Fig. 8.9.

Appendix

The definition of minimizing movement along a sequence of functionals formalizes a natural extension to the notion of minimizing movement, and follows the definition given in the paper by Braides et al. [2].

The energies in Examples 8.2 and 8.6 have been taken as a prototype to model plastic phenomena by Puglisi and Truskinovsky [7]. More recently, that example has been recast in the framework of quasistatic motion in the papers by Mielke and Truskinovsky [4, 6].

The example of the minimizing movement for Lennard-Jones interactions is part of results of Braides et al. [1]. It is close in spirit to a semi-discrete approach (i.e., the study of the limit of the gradient flows for the discrete energies) by Gobbino [3].

For the notion of BV-solution we refer to Mielke et al. [5].

References

1. Braides, A., Defranceschi, A., Vitali, E.: Variational evolution of one-dimensional Lennard-Jones systems. Preprint Scuola Normale Superiore, Pisa (2013). <http://cvgmt.sns.it/paper/2210/>
2. Braides, A., M.S. Gelli, M.S., Novaga, M.: Motion and pinning of discrete interfaces. Arch. Ration. Mech. Anal. **95**, 469–498 (2010)

3. Gobbino, M.: Gradient flow for the one-dimensional Mumford-Shah functional. *Ann. Scuola Norm. Sup. Pisa (IV)* **27**, 145–193 (1998)
4. Mielke, A.: Emergence of rate-independent dissipation from viscous systems with wiggly energies. *Continuum Mech. Thermodyn.* **24**, 591–606 (2012)
5. Mielke, A., Rossi, R., Savaré, G.: BV solutions and viscosity approximations of rate-independent systems. *ESAIM Control Optim. Calc. Var.* **18**, 36–80 (2012)
6. Mielke, A., Truskinovsky, L.: From discrete visco-elasticity to continuum rate-independent plasticity: rigorous results. *Arch. Ration. Mech. Anal.* **203**, 577–619 (2012)
7. Puglisi, G., Truskinovsky, L.: Thermodynamics of rate-independent plasticity. *J. Mech. Phys. Solids* **53**, 655–679 (2005)



<http://www.springer.com/978-3-319-01981-9>

Local Minimization, Variational Evolution and
 Γ -Convergence

Braides, A.

2014, XI, 174 p. 42 illus., Softcover

ISBN: 978-3-319-01981-9