

Chapter 2

Potassium Channels Implicated in the Short QT Syndrome

The work presented in this thesis centres on the *in silico* investigation of arrhythmia substrates in an inherited cardiac condition: the short QT syndrome (SQTS). Chapter 3 provides detailed background information on this syndrome (which consequently will be discussed only briefly here). The three variants of the SQTS examined in this thesis—SQT1, SQT2 and SQT3—involve gene mutations that affect proteins forming different potassium channels [1–6]. SQT1 affects the *hERG* channel, which is responsible for the rapid-delayed outward rectifier potassium current (I_{Kr}) [7–9], SQT2 affects the *KCNQ1* gene, which encodes the α -subunit of channels mediating slow-delayed outward rectifier potassium channel (I_{Ks}) [10]. SQT3 affects *KCNJ2*, which encodes the Kir2.1 protein that contributes to inwardly-rectifying potassium channel current (I_{K1}) [11]. Building upon the brief descriptions of these channels in Chap. 1, this chapter gives a detailed consideration of their structure and characteristics.

2.1 Rectification

In Sect. 1.3, it was established that the lipid bilayer of the cell membrane can be thought of as a capacitor that separates charges in the intracellular and extracellular regions. The charges separated are the ions in intracellular/extracellular fluid that can flow across the membrane through ion channels, when these are in a conducting state. The ion channels are thus resistors. Simple resistors follow Ohm's law:

$$V = IR \quad (2.1)$$

where V is the voltage (membrane potential), I is the current and R is the resistance. To describe the biophysical properties of ion channels during voltage clamp experiments, the conductance of the channels (inverse of resistance) is normally employed. If an ion channel has a linear current-voltage (I – V) relationship, i.e., the slope (conductance) is linear, the channel is said to show 'ohmic' behaviour (Fig. 2.1).




Fig. 2.1 Schematic representation of the I-V relationship for an ohmic channel and channels showing rectification. The black dashed line represents an ohmic channel with a linear I-V relationship. The green line shows the I-V relationship for I_{Kr} , which shows outward rectification. The blue line represents the I-V relationship for I_{Ks} , which also shows outward rectification and the red line represents I_{K1} , which is an inwardly rectifying current because it passes current preferentially in the inward direction. Note that currents are not drawn to scale

When the I-V relationship of the channel is nonlinear, i.e., channel conductance changes with voltage, the channel is said to show voltage-dependent rectification (Fig. 2.1). In this event, the channel passes current preferentially in one direction over another. For example, Fig. 2.1 shows a schematic representation of the I-V relationships for four channels: an ohmic channel, hERG/ I_{Kr} (outward rectifier), KCNQ1-KCNE1/ I_{Ks} (outward rectifier) and KCNJ2/ I_{K1} (inward rectifier). hERG/ I_{Kr} and KCNQ1-KCNE1/ I_{Ks} pass current preferentially in the outward direction while KCNJ2/ I_{K1} passes current preferentially in the inward direction over the outward direction.

2.2 The hERG/ I_{Kr} Potassium Channel

The significance of the hERG/ I_{Kr} channel for normal human cardiac electrical activity was discovered when inherited gene mutations to hERG resulted in long QT syndrome (LQTS) [12–14]; a cardiac repolarization disorder. In the LQTS, patients have a lengthened QT interval on the ECG and become susceptible to the potentially fatal arrhythmia *torsades de pointes* [15–17]. hERG is also now known to be responsible for pharmacologically induced (“acquired”) Long QT syndrome

[16, 18–20]. This is due to structural features of the channel that have been established to render it particularly susceptible to pharmacological blockade [15, 16, 21, 22]. The consideration of hERG here focuses on its electrophysiological properties and for detailed consideration of the basis of its pharmacological promiscuity the reader is referred to [15, 16, 21, 22]. Some antihistamines and antibiotics have also been known to cause arrhythmia and sudden death through blockade of the hERG/I_{Kr} channel [23].

hERG/I_{Kr} is expressed in several tissue and cell types including cardiac [13], neurons, neuroendocrine glands [24, 25], smooth muscle [26] and tumour cells [27, 28]. Its expression is greatest in cardiac cells and this is the region that has received the greatest research focus and from which its properties and function are best understood. Heterologous expression studies have shown that hERG encodes the α -subunit of the I_{Kr} channel [13, 14, 29–31].

2.2.1 hERG/I_{Kr} Potassium Channel Gating

hERG/I_{Kr} plays a significant role in cardiac action potential repolarization; it is the channel that is largely responsible for the early and middle stages of ventricular action potential repolarization (phase 3; Fig. 1.1) after which I_{K1} (being responsible for terminal repolarization) takes over, bringing the membrane potential back to its normal resting value (phase 4; Fig. 1.1) [13, 14]. It is characterised by slow activation but fast and profound voltage-dependent inactivation [14, 29, 32–35]. Its inactivation (a non-conducting state) is considerably faster than its activation.

Figure 2.2 shows characteristic features of I_{hERG}. With depolarisation to more positive membrane potentials, outward I_{hERG} is elicited, which increases rapidly and then plateaus for the duration of the pulse. A step repolarization of the membrane potential (to a negative potential) results in a resurgent increase in I_{hERG} (tail current); this increase in I_{hERG} is despite the decrease in the driving force for K⁺ out of the cell membrane. This is due to the rapid recovery of I_{hERG} from inactivation [13, 14, 33]. Eventually, via channel deactivation, I_{hERG} decreases bi-exponentially [36].

Further insight can be gleaned from the current-voltage (I–V) relation of I_{hERG}/I_{Kr} (Fig. 2.3). At the membrane resting potential (~ -80 mV), hERG/I_{Kr} channels are closed and in that state conduct no current. As the membrane is depolarised to more positive membrane potentials greater than ~ -60 mV, the channels get activated (open) and conduct current (Fig. 2.3); K⁺ ions flow out of the cell across the cell membrane according to their electrochemical gradient. At the same time, the channel begins to inactivate. As hERG/I_{Kr} inactivation with progressive depolarisation is faster than its activation, channel inactivation with increasing depolarisation eventually overtakes channel activation resulting in the channel entering the non-conducting, inactivated state. This gives the current-voltage relationship a region of negative slope (Fig. 2.3). This helps to prolong phase 2 of the action potential. Eventually, the channel begins to deactivate (close) and also

Fig. 2.2 Representative current trace for hERG elicited by a standard voltage protocol from a hERG-expressing Chinese Hamster Ovary (CHO) cell (recording made at 37 °C). Outward I_{hERG}/I_{Kr} current is elicited when the membrane potential is stepped to 20 mV from a holding potential of -80 mV. On stepping the membrane potential to -40 mV, I_{hERG}/I_{Kr} (tail current) of greater amplitude than that seen at the 20 mV are produced. Modified from [35]





Fig. 2.3 Current-voltage (I-V) relationship for I_{hERG}/I_{Kr} (recorded in *Xenopus* oocytes). Depolarisation activates I_{hERG}/I_{Kr} causing an increase in outward current. At around 0 mV, I_{hERG}/I_{Kr} reaches a maximum then declines in amplitude with further depolarisation (negative slope region). Modified from [13]



begins to recover from inactivation. However, recovery from inactivation for I_{hERG}/I_{Kr} is faster than deactivation, thus allowing the channel to again conduct current, which repolarises the cell membrane (phase 3 of the action potential), returning it to its resting state (phase 4).

2.2.2 I_{hERG}/I_{Kr} Channel Structure

The I_{hERG}/I_{Kr} K^+ channel is comprised of a protein tetramer consisting of four identical α -subunits [29, 37–39]. Each subunit contains six α -helical transmembrane domains (S1–S6). Functionally, each α -subunit can be divided into two parts: a voltage-sensing region and a pore-forming K^+ -selective filter (Fig. 2.4). Segments S1–S4 form the transmembrane potential sensor region with S4 in particular having




Fig. 2.4 Schematic diagram showing a representative hERG/I_{Kr} α -subunit. hERG channels consist of four identical α -subunits. Each subunit contains six transmembrane segments S1–S6. S4 has positively charged amino acids and acts as the main voltage sensor for transmembrane potential changes. The blue cylinder indicates the pore of the P-loop that acts as a K⁺ selectivity filter by blocking or unblocking the pore. Intracellularly, the α -subunit has N and C termini




Fig. 2.5 Membrane depolarisation to voltages more positive than ~ -60 mV activates (opens) the channel slowly. With greater depolarisation, the channel rapidly inactivates (it undergoes C-type inactivation—a slight constriction of the K⁺ selectivity filter). Repolarization reverses the whole process

positive charges, which allow it to react to a change in membrane potential by moving its position [40–43]. Voltage-sensor movement leads to conformational changes that open, inactivate or close the channel (Fig. 2.5). Segments S5–S6 form the ion permeation pathway and consist of the P-loop (S5) that penetrates the membrane and creates a pore through which ions enter or leave. Together, these traits confer voltage dependence and ion selectivity on the channel [40–43].

Below the P-loop is a water-filled cavity that is lined by S6 α -helices. In a conducting (open) state, i.e., in response to membrane depolarisation, all four S6 α -helices are spread out allowing the passage of K⁺ ions. They crisscross in the closed state, forming a narrow cleft that blocks entry of K⁺ ions [41]. hERG also

has a pair of intracellular termini (an N terminus and a C terminus), which contribute to its function. The N-terminus is responsible for channel deactivation following membrane depolarisation while mutations to the C-terminus affect trafficking and disrupt the processing of hERG channels [44, 45].

Voltage-gated K^+ channels can inactivate via two different mechanisms: rapid ‘N-type’ inactivation (also known as “ball and chain” type) and slow ‘C-type’ inactivation. N-type inactivation occurs when a ball-like structure (an intracellular protein segment) anchored to the channel’s N-terminus blocks the channel pore [33] whereas ‘C-type’ inactivation has been suggested to occur via a slight narrowing of the K^+ -selectivity filter [46] at the extracellular mouth. Deletion of the N-terminus removes ‘N-type’ inactivation but has no significant effect on hERG inactivation [33, 47] suggesting that hERG inactivates via ‘C-type’ inactivation. This was confirmed by removing ‘C-type’ inactivation via the Ser 631 Val mutation [48], which resulted in the elimination hERG inactivation. In addition, application of extracellular tetraethylammonium (TEA) blockade but not intracellular TEA blockade has been shown to slow inactivation suggesting that the process involves changes towards the channel exterior [49].

In addition to the four α -subunits, hERG also interacts with a β -subunit, MiRP1 (minK-related protein 1) in heterologous expression systems. The expression of MiRP1 in atrial or ventricular tissue is quite low but it is highly expressed in Purkinje fibres and the atrial pacemaking cells [50, 51]. The effect of this co-expression of the hERG α -subunit and MiRP1 is to reduce the trafficking of the channel to the cell membrane surface, reduce channel conductance and to increase channel deactivation [52]. As MiRP1 expression is low outside of the conduction system [53], it has been suggested that it may not function as hERG β -subunit in other regions and comparison of hERG with and without MiRP1 has shown that when hERG is heterologously expressed in mammalian instead of amphibian cells (*Xenopus* oocytes), it adequately recapitulates I_{Kr} [54]. Also, MiRP1 can co-assemble with a variety of other cardiac ion channels [53, 55], and so may exhibit broad interactions rather than being specific for hERG.

hERG is heteromeric and consists of two isoforms: hERG 1a (the major isoform) and hERG 1b, which differ by a truncated N-terminus in the latter. This heteromeric form of hERG (hERG 1a/1b) has recently been proposed to recapitulate native I_{Kr} more accurately than hERG 1a expressed alone [56, 57].

2.3 The KCNQ1-KCNE1/ I_{Ks} Potassium Channel

Similar to the hERG/ I_{Kr} potassium channel, the KCNQ1 potassium channel also plays a significant role in cardiac repolarization. The KCNQ1 α -subunit co-expresses with the KCNE1 β -subunit (minK) [40, 58, 59], the result of which is a protein complex that is responsible for the slow-delayed rectifier potassium channel I_{Ks} , which is partially responsible for action potential repolarization (phase 3) along with hERG/ I_{Kr} [40, 58, 59]. In the event of an impairment to I_{Kr} ,





Fig. 2.6 Representative superimposed current traces for KCNQ1 (purple) and KCNQ1 + KCNE1 (black) elicited by a standard voltage protocol from KCNQ1-expressing oocytes (recording made at 36 °C). The *left arrow* shows the slower activation of KCNQ1 when co-expressed with KCNE1 when the membrane potential is stepped to 60 mV from a holding potential of -80 mV. On stepping the membrane potential to -60 mV, KCNQ1-KCNE1/ I_{Ks} (tail current) deactivates more slowly than KCNQ1 (*right arrow*). Modified from [78]

e.g., channel block or gene mutation, I_{Ks} is a key component of repolarization reserve [40, 60, 61]. A *loss-of-function* mutation to the channel results in the first variant of the long QT syndrome LQTS1 [62–64].

In the ventricular wall, KCNQ1-KCNE1/ I_{Ks} is heterogeneously distributed with its expression being very low in the mid-myocardial cells [60, 65–67]. Consequently, mid-myocardial cells have the longest APD and repolarise later than epicardial and endocardial cells. This leads to a transmural dispersion of repolarization across the ventricular wall [68–70] (meaning repolarization occurs at different rates in the different cell types that make up the wall of the myocardium), which is a substrate for re-entry. Re-entry is the continuous (usually several cycles) re-excitation of a cardiac tissue region by a single electrical signal and it usually results in arrhythmias.

In addition to being expressed in cardiac tissue, the *KCNQ1* protein is also expressed in epithelial tissue of different organs: stomach, cochlea, lungs, intestine and kidney. Its function in these tissue types is the transport of salt and water [65]. In humans, impairment to *KCNQ1* expression in epithelial tissue has been known to cause deafness [65, 71, 72], while in knockout mice, it causes deafness, balance problems and morphological abnormalities in the gastrointestinal tract and inner ear [65, 73, 74].

Fig. 2.7 I–V relationship of I_{Ks} tail current (recording in HEK 293 cells at 36 °C) measured from peak tail current to complete deactivation. Depolarisation activates KCNQ1-KCNE1/ I_{Ks} causing an increase in outward current. Modified from [79]



2.3.1 KCNQ1-KCNE1/ I_{Ks} Potassium Channel Gating

Similar to hERG, KCNQ1 is a voltage-gated channel, though its kinetic properties differ considerably from those of hERG (Fig. 2.8). Depolarisation activates (opens) KCNQ1 channels but very slowly. Upon further depolarisation, a fraction of the open channels inactivates [40, 60, 75–77] (Fig. 2.6). The channel is also characterised by slow deactivation kinetics [40, 60, 75–77]. Recovery from inactivation, however, is faster than deactivation and is seen as a hook on the current trace during the repolarization (Fig. 2.6). On co-expression with KCNE1, thus reproducing I_{Ks} , the current is enhanced (Fig. 2.6) because KCNE1 increases the single channel conductance causing a positive shift in voltage activation threshold, i.e., KCNE1 slows activation (Fig. 2.6) considerably and inactivation is completely eliminated [28, 60, 76]. Figure 2.7 shows the tail current I–V relationship for KCNQ1-KCNE1/ I_{Ks} measured from peak tail current to complete deactivation.

2.3.2 KCNQ1-KCNE1/ I_{Ks} Channel Structure

Structurally, KCNQ1 is a voltage-gated potassium channel like hERG described in Sect. 2.2.2 above. It consists of four identical subunits, each of which is comprised of six α -helical transmembrane domains (S1–S6). Segments S1–S4 form the voltage sensor while segments S5 and S6 form the ion permeation pathway with a pore loop and K^+ selectivity filter [58, 59, 65, 80] (Fig. 2.8). Each α -subunit also has intracellular ‘N’ and ‘C’ terminals. A major difference from hERG is KCNQ1’s co-assembly with KCNE1 β -subunit, which together encode the human I_{Ks} . A representation of the KCNQ1-KCNE1 structure is shown in Fig. 2.8. The effect of the KCNE1 subunit is to stabilise the open state of the channel by altering the interaction between the pore loop, the K^+ selectivity filter and the S5/S6




Fig. 2.8 A structural representation of the α and β -subunits KCNQ1-KCNE1/ I_{Ks} . KCNQ1 channels consist of four identical α -subunits. Each subunit contains six transmembrane segments S1–S6. S4 has positively charged amino acids and acts as the main voltage sensor for transmembrane potential changes. The blue cylinder indicates the pore of the P-loop that acts as a K^+ selectivity filter by blocking or unblocking the pore. KCNQ1 has intracellular N and C termini


segment [81]. However, the stoichiometry of KCNQ1:KCNE1 is still a matter of debate because there is no extant crystal structure for KCNQ1 with or without KCNE1 [82, 83]. Stoichiometries of 4:4 [84], 4:2 [85] and other forms [77, 83] have been proposed. A recent review can be found in [83].

2.4 The KCNJ2/ I_{K1} Potassium Channel

The KCNJ2-encoded Kir2.1 protein belongs to the family of inwardly rectifying potassium (Kir) channels [86, 87]. Specifically, it belongs to the Kir2.x channel family. It is expressed in skeletal muscle, blood vessels, neurons and richly in cardiac tissue, where it is expressed in Purkinje fibres, atrial and ventricular tissues [87–92]. This family of channels allows K^+ to flow more easily into the cell than out of it. They therefore, preferentially pass current in the inward direction over the outward direction. A schematic representation of the I–V relation for I_{K1} is shown in Fig. 2.1 (red line) and an experimentally observed I–V relation is shown in Fig. 2.9.

Unlike voltage-gated K^+ channels such as I_{Kr} and I_{Ks} , the behaviour of Kir channels depends not only on the membrane potential but also predominantly on the electrochemical gradient of K^+ in the cell, i.e., the difference between the membrane potential and the K^+ reversal potential (E_K). They therefore have greater K^+ conductance at potentials negative to E_K and pass comparatively little current at depolarised membrane potentials, i.e., potentials positive to E_K (see red

Fig. 2.9 Current-voltage (I - V) relations of the KCNJ2-encoded Kir2.1 channel expressed in HEK cells elicited by the ascending voltage ramp command shown inset. Modified from [93]



line in Figs. 2.1 and 2.9) [86–88, 94–97]. As membrane depolarisation increases further, the outward I_{K1} current decreases due to its inward rectification.

These characteristics make myocardial cells expressing I_{K1} have resting potentials close to E_K whereas cells without I_{K1} or those with an insignificant expression of I_{K1} have depolarised resting potentials and tend to show spontaneous activity, e.g., the SAN [98]. This is because with little I_{K1} present, the cells have high membrane resistances at negative voltages and so small changes in current can produce substantial changes in voltage (i.e. membrane potential is more ‘labile’ in the absence of I_{K1}). Consequently, in cardiac cells, I_{K1} plays a role in stabilizing the resting potential of the cell and in the duration of the action potential [86–88, 94–97]. As I_{K1} channels pass little outward current at depolarised potentials, there is little K^+ efflux through I_{K1} channels during phase 2 (the plateau phase) of the ventricular action potential [99] (Fig. 2.10b); therefore, the rectification property of I_{K1} channels also serves to maintain membrane depolarisation, thereby facilitating prolonged action potential duration. When phase 3 repolarization commences (via I_{Kr} and I_{Ks}) and the membrane becomes more and more hyperpolarised, relatively large outward I_{K1} current is generated. This serves to accelerate terminal repolarization [100–103]. Figure 2.10b shows the I - V relationship of I_{K1} and relates it to the action potential, which is drawn sideways to give a better indication of the current amplitude at different phases of the action potential.

2.4.1 Kir2.1/ I_{K1} Potassium Channel Gating

The inward rectification of I_{K1} and Kir2.x channels is due to blockade of outward K^+ movement through the channel pore by intracellular Mg^{2+} ions and polyamines





Fig. 2.10 A: structure of the Kir2.1/KCNJ2/ I_{K1} channel. Each of its four α -subunits consists of two transmembrane segments (TM1 and TM2), a pore-forming loop (H5) and intracellular N and C terminals. B: I_{K1} current-voltage relationship related to the membrane action potential. The action potential is drawn sideways to give a better understanding of the contribution of I_{K1} during the action potential. The reader should focus on the repolarization phase and resting potential (the *non-dashed* parts of the action potential)

[87, 97, 104–107]. At depolarised membrane potentials, the Mg^{2+} ions and polyamines such as spermine and spermidine, which are present in sub-micromolar quantities within the cell, reduce outward K^+ current. On hyperpolarisation, the Mg^{2+} ions and polyamines unblock the pore, thereby leading to increased current. This inward current first increases time-independently due to the unblocking of Mg^{2+} ions and then increases time-dependently due to unblocking of the polyamines. The unblocking of the Mg^{2+} ions is fast while the polyamine unblocking is slow [87, 108].

2.4.2 KCNJ2/ I_{K1} Channel Structure

Functional Kir2.x channels (similarly to hERG/ I_{Kr} and KCNQ1/ I_{Ks}) also have four α -subunits but each subunit consists of only two transmembrane segments, TM1 and TM2 (Fig. 2.10a), which are highly homologous to the S5 and S6 domains of the K^+ channels discussed previously [87, 109, 110]. These two segments are

Fig. 2.11 A schematic diagram of the current profiles of I_{Kr} , I_{Ks} and I_{K1} ionic currents during a ventricular action potential. Also shown are the major gene candidates responsible for these currents. Modified from [111]



linked by the extracellular pore-forming loop or P-loop (H5), which acts as the K^+ selectivity filter. The intracellular ‘N’ and ‘C’ termini are also present. Unlike hERG/ I_{Kr} and KCNQ1/ I_{Ks} , there is no voltage sensor region or S4 segment; hence the channel is not truly voltage-gated in a traditional sense, with voltage-dependence instead occurring through voltage-dependent channel block/unblock by Mg^{2+} and polyamines (as described above).

2.5 Current Profiles During an Action Potential

Figure 2.11 shows a schematic diagram of the current profiles of each potassium channel currents discussed in this chapter during a ventricular action potential. I_{Kr} contributes during most of phase 3 repolarization, I_{K1} contributes later in phase 3 than I_{Kr} while I_{Ks} contributes earlier during phase 3 and some part of phase 2.

References

1. Hancox JC, McPate MJ, Harchi A, Duncan RS, Dempsey CE, Witchel HJ, et al (2011) The short QT syndrome. In: Tripathi ON, Ravens U, Sanguinetti MC (eds). Heart rate and rhythm [Internet]. Springer, Berlin pp 431–49. <http://www.springerlink.com/content/m8l86l8n3h81w43m/>
2. Cross B, Homoud M, Link M, Foote C, Garlitski A, Weinstock J et al (2011) The short QT syndrome. J Intervent Card Electrophysiol 31(1):25–31
3. Schimpf R, Wolpert C, Gaita F, Giustetto C, Borggrefe M (2005) Short QT syndrome. Cardiovasc Res 67(3):357–366
4. McPate MJ, Witchel HJ, Hancox JC (2006) Short QT syndrome. Future Cardiol 2(3):293–301
5. Maury P, Extramiana F, Sbragia P, Giustetto C, Schimpf R, Duparc A et al (2008) Short QT syndrome. Update on a recent entity. Arch Cardiovasc Dis 101(11–12):779–786
6. Lu LX, Zhou W, Zhang X, Cao Q, Yu K, Zhu C (2006) Short QT syndrome: a case report and review of literature. Resuscitation 71(1):115–121

7. Brugada R, Hong K, Dumaine R, Cordeiro J, Gaita F, Borggrefe M et al (2004) Sudden death associated with short-QT syndrome linked to mutations in HERG. *Circulation* 109(1):30–35
8. Hong K, Bjerregaard P, Gussak I, Brugada R (2005) Short QT syndrome and atrial fibrillation caused by mutation in KCNH2. *J Cardiovasc Electrophysiol* 16(4):394–396
9. Sun Y, Quan X-Q, Fromme S, Cox RH, Zhang P, Zhang L et al (2011) A novel mutation in the KCNH2 gene associated with short QT syndrome. *J Mol Cell Cardiol* 50(3):433–441
10. Belloccq C, van Ginneken ACG, Bezzina CR, Alders M, Escande D, Mannens MMAM et al (2004) Mutation in the KCNQ1 gene leading to the short QT-interval syndrome. *Circulation* 109(20):2394–2397
11. Priori SG, Pandit SV, Rivolta I, Berenfeld O, Ronchetti E, Dhamoon A et al (2005) A novel form of short QT syndrome (SQT3) is caused by a mutation in the KCNJ2 gene. *Circ Res* 96(7):800–807
12. Curran ME, Splawski I, Timothy KW, Vincent GM, Green ED, Keating MT (1995) A molecular basis for cardiac arrhythmia: HERG mutations cause long QT syndrome. *Cell* 80(5):795–803
13. Sanguinetti MC, Jiang C, Curran ME, Keating MT (1995) A mechanistic link between an inherited and an acquired cardiac arrhythmia: HERG encodes the IKr potassium channel. *Cell* 81(2):299–307
14. Trudeau MC, Warmke JW, Ganetzky B, Robertson GA (1995) HERG, a human inward rectifier in the voltage-gated potassium channel family. *Science* 269(5220):92–95
15. Kramer DB, Zimetbaum PJ (2011) Long-QT syndrome. *Cardiol Rev* 19(5):217–225
16. Kallergis EM, Goudis CA, Simantirakis EN, Kochiadakis GE, Vardas PE (2012) Mechanisms, risk factors, and management of acquired long QT syndrome: a comprehensive review. *Sci World J* 2012:212178
17. Roden DM, Lazzara R, Rosen M, Schwartz PJ, Towbin J, Vincent GM (1996) Multiple mechanisms in the long-QT syndrome. Current knowledge, gaps, and future directions. The SADS foundation task force on LQTS. *Circulation* 94(8):1996–2012
18. Dennis A, Wang L, Wan X, Ficker E (2007) hERG channel trafficking: novel targets in drug-induced long QT syndrome. *Biochem Soc Trans* 35(Pt 5):1060–1063
19. Wehrens XHT, Vos MA, Doevendans PA, Wellens HJJ (2002) Novel insights in the congenital long QT syndrome. *Ann Intern Med* 137(12):981–992
20. Camm AJ, Janse MJ, Roden DM, Rosen MR, Cinca J, Cobbe SM (2000) Congenital and acquired long QT syndrome. *Eur Heart J* 21(15):1232–1237
21. Sanguinetti MC, Mitcheson JS (2005) Predicting drug-hERG channel interactions that cause acquired long QT syndrome. *Trends Pharmacol Sci* 26(3):119–124
22. Mitcheson JS, Chen J, Lin M, Culberson C, Sanguinetti MC (2000) A structural basis for drug-induced long QT syndrome. *Proc Natl Acad Sci USA* 97(22):12329–12333
23. Haverkamp W, Breithardt G, Camm AJ, Janse MJ, Rosen MR, Antzelevitch C et al (2000) The potential for QT prolongation and proarrhythmia by non-antiarrhythmic drugs: clinical and regulatory implications. Report on a policy conference of the European Society of Cardiology. *Eur Heart J* 21(15):1216–1231
24. Warmke JW, Ganetzky B (1994) A family of potassium channel genes related to eag in *Drosophila* and mammals. *Proc Natl Acad Sci USA* 91(8):3438–3442
25. Shen XZ, Wu J, Lin JJ (2008) HERG and arrhythmia. *Adv Cardiovasc Dis* 29:436–440
26. Farrelly AM, Ro S, Callaghan BP, Khoyi MA, Fleming N, Horowitz B et al (2003) Expression and function of KCNH2 (HERG) in the human jejunum. *Am J Physiol Gastrointest Liver Physiol* 284(6):G883–G895
27. Chiesa N, Rosati B, Arcangeli A, Olivetto M, Wanke E (1997) A novel role for HERG K⁺ channels: spike-frequency adaptation. *J Physiol (Lond)* 501(Pt 2):313–318
28. Smith GAM, Tsui H-W, Newell EW, Jiang X, Zhu X-P, Tsui FWL et al (2002) Functional up-regulation of HERG K⁺ channels in neoplastic hematopoietic cells. *J Biol Chem* 277(21):18528–18534

29. Tristani-Firouzi M, Sanguinetti MC (2003) Structural determinants and biophysical properties of HERG and KCNQ1 channel gating. *J Mol Cell Cardiol* 35(1):27–35
30. Mitcheson JS, Sanguinetti MC (1999) Biophysical properties and molecular basis of cardiac rapid and slow delayed rectifier potassium channels. *Cell Physiol Biochem* 9(4–5):201–216
31. Sanguinetti MC, Tristani-Firouzi M (2006) hERG potassium channels and cardiac arrhythmia. *Nature* 440(7083):463–469
32. McPate MJ, Duncan RS, Milnes JT, Witchel HJ, Hancox JC (2005) The N588 K-HERG K⁺ channel mutation in the short QT syndrome: mechanism of gain-in-function determined at 37 °C. *Biochem Biophys Res Commun* 334(2):441–449
33. Spector PS, Curran ME, Zou A, Keating MT, Sanguinetti MC (1996) Fast inactivation causes rectification of the IKr channel. *J Gen Physiol* 107(5):611–619
34. Hancox JC, McPate MJ, El Harchi A, Zhang YH (2008) The hERG potassium channel and hERG screening for drug-induced torsades de pointes. *Pharmacol Ther* 119(2):118–132
35. McPate MJ, Duncan RS, Hancox JC, Witchel HJ (2008) Pharmacology of the short QT syndrome N588 K-HERG K⁺ channel mutation: differential impact on selected class I and class III antiarrhythmic drugs. *Br J Pharmacol* 155(6):957–966
36. Zhou Z, Gong Q, Ye B, Fan Z, Makielski JC, Robertson GA et al (1998) Properties of HERG channels stably expressed in HEK 293 cells studied at physiological temperature. *Biophys J* 74(1):230–241
37. Stuhmer W, Conti F, Stocker M, Pongs O, Heinemann SH (1991) Gating currents of inactivating and non-inactivating potassium channels expressed in *Xenopus* oocytes. *Pflügers Arch* 418(4):423–429
38. Stefani E, Toro L, Perozo E, Bezanilla F (1994) Gating of Shaker K⁺ channels: I Ionic and gating currents. *Biophys J* 66(4):996–1010
39. Bezanilla F, Perozo E, Stefani E (1994) Gating of Shaker K⁺ channels: II. The components of gating currents and a model of channel activation. *Biophys J* 66(4):1011–1021
40. Rudy Y, Silva JR (2006) Computational biology in the study of cardiac ion channels and cell electrophysiology. *Q Rev Biophys* 39(1):57–116
41. Doyle DA, Morais Cabral J, Pfuetzner RA, Kuo A, Gulbis JM, Cohen SL, et al (1998) The structure of the potassium channel: molecular basis of K⁺ conduction and selectivity. *Science* 280(5360):69–77
42. Long SB, Campbell EB, Mackinnon R (2005) Voltage sensor of Kv1.2: structural basis of electromechanical coupling. *Science* 309(5736):903–908
43. Jiang Y, Lee A, Chen J, Cadene M, Chait BT, MacKinnon R (2002) The open pore conformation of potassium channels. *Nature* 417(6888):523–526
44. Akhavan A, Atanasiu R, Noguchi T, Han W, Holder N, Shrier A (2005) Identification of the cyclic-nucleotide-binding domain as a conserved determinant of ion-channel cell-surface localization. *J Cell Sci* 118(Pt 13):2803–2812
45. Zhou P, Babcock J, Liu L, Li M, Gao Z (2011) Activation of human ether-a-go-go related gene (hERG) potassium channels by small molecules. *Acta Pharmacol Sin* 32(6):781–788
46. Baukrowitz T, Yellen G (1995) Modulation of K⁺ current by frequency and external [K⁺]: a tale of two inactivation mechanisms. *Neuron* 15(4):951–960
47. Smith PL, Baukrowitz T, Yellen G (1996) The inward rectification mechanism of the HERG cardiac potassium channel. *Nature* 379(6568):833–836
48. Fan JS, Jiang M, Dun W, McDonald TV, Tseng GN (1999) Effects of outer mouth mutations on hERG channel function: a comparison with similar mutations in the Shaker channel. *Biophys J* 76(6):3128–3140
49. Smith PL, Yellen G (2002) Fast and slow voltage sensor movements in HERG potassium channels. *J Gen Physiol* 119(3):275–293
50. Pourrier M, Zicha S, Ehrlich J, Han W, Nattel S (2003) Canine ventricular KCNE2 expression resides predominantly in Purkinje fibers. *Circ Res* 93(3):189–191
51. Lundquist AL, Manderfield LJ, Vanoye CG, Rogers CS, Donahue BS, Chang PA et al (2005) Expression of multiple KCNE genes in human heart may enable variable modulation of I(Ks). *J Mol Cell Cardiol* 38(2):277–287

52. Abbott GW, Sesti F, Splawski I, Buck ME, Lehmann MH, Timothy KW et al (1999) MiRP1 forms IKr potassium channels with HERG and is associated with cardiac arrhythmia. *Cell* 97(2):175–187
53. Yu H, Wu J, Potapova I, Wymore RT, Holmes B, Zuckerman J et al (2001) MinK-related peptide 1: a beta subunit for the HCN ion channel subunit family enhances expression and speeds activation. *Circ Res* 88(12):E84–E87
54. Weerapura M, Nattel S, Chartier D, Caballero R, Hébert TE (2002) A comparison of currents carried by HERG, with and without coexpression of MiRP1, and the native rapid delayed rectifier current. Is MiRP1 the missing link? *J Physiol (Lond)* 540(Pt 1):15–27
55. Zhang M, Jiang M, Tseng GN (2001) MinK-related peptide 1 associates with Kv4.2 and modulates its gating function: potential role as beta subunit of cardiac transient outward channel? *Circ Res* 88(10):1012–1019
56. Jones EMC, Roti Roti EC, Wang J, Delfosse SA, Robertson GA (2004) Cardiac IKr channels minimally comprise hERG 1a and 1b subunits. *J Biol Chem* 279(43):44690–44694
57. Sale H, Wang J, O'Hara TJ, Tester DJ, Phartiyal P, He J-Q et al (2008) Physiological properties of hERG 1a/1b heteromeric currents and a hERG 1b-specific mutation associated with Long-QT syndrome. *Circ Res* 103(7):e81–e95
58. Barhanin J, Lesage F, Guillemare E, Fink M, Lazdunski M, Romey G (1996) K(V)LQT1 and IsK (minK) proteins associate to form the I(Ks) cardiac potassium current. *Nature* 384(6604):78–80
59. Sanguinetti MC, Curran ME, Zou A, Shen J, Spector PS, Atkinson DL et al (1996) Coassembly of K(V)LQT1 and minK (IsK) proteins to form cardiac I(Ks) potassium channel. *Nature* 384(6604):80–83
60. Silva J, Rudy Y (2005) Subunit interaction determines IKs participation in cardiac repolarization and repolarization reserve. *Circulation* 112(10):1384–1391
61. Jost N, Virág L, Bitay M, Takács J, Lengyel C, Biliczki P et al (2005) Restricting excessive cardiac action potential and QT prolongation: a vital role for IKs in human ventricular muscle. *Circulation* 112(10):1392–1399
62. Mikuni I, Torres CG, Bienengraeber MW, Kwok W-M (2011) Partial restoration of the long QT syndrome associated KCNQ1 A341 V mutant by the KCNE1 β -subunit. *Biochim Biophys Acta* 1810(12):1285–1293
63. Moric-Janiszewska E, Głogowska-Ligus J, Paul-Samojedny M, Smolik S, Woźniak M, Markiewicz-Łoskot G et al (2011) Expression of genes KCNQ1 and HERG encoding potassium ion channels Ikr, Iks in long QT syndrome. *Kardiol Pol.* 69(5):423–429
64. Eldstrom J, Xu H, Werry D, Kang C, Loewen ME, Degenhardt A et al (2010) Mechanistic basis for LQT1 caused by S3 mutations in the KCNQ1 subunit of IKs. *J Gen Physiol* 135(5):433–448
65. Jespersen T, Grønnet M, Olesen S-P (2005) The KCNQ1 potassium channel: from gene to physiological function. *Physiol (Bethesda)* 20:408–416
66. Akar FG, Yan G-X, Antzelevitch C, Rosenbaum DS (2002) Unique topographical distribution of M cells underlies reentrant mechanism of torsade de pointes in the long-QT syndrome. *Circulation* 105(10):1247–1253
67. Antzelevitch C, Dumaine R (2002) Electrical heterogeneity in the heart: physiological, pharmacological and clinical implications. In: Page E, Fozzard H, Solaro J (eds). *Handbook of physiology, section 2: the cardiovascular system, Volume I: The Heart*. Oxford University Press, New York, pp 654–92
68. Viswanathan PC, Shaw RM, Rudy Y (1999) Effects of IKr and IKs heterogeneity on action potential duration and its rate dependence: a simulation study. *Circulation* 99(18):2466–2474
69. Di Diego JM, Belardinelli L, Antzelevitch C (2003) Cisapride-induced transmural dispersion of repolarization and torsade de pointes in the canine left ventricular wedge preparation during epicardial stimulation. *Circulation* 108(8):1027–1033
70. Xue J, Chen Y, Han X, Gao W (2010) Electrocardiographic morphology changes with different type of repolarization dispersions. *J Electrocardiol* 43(6):553–559

71. Jervell A, Lange-Nielsen F (1957) Congenital deaf-mutism, functional heart disease with prolongation of the Q-T interval and sudden death. *Am Heart J* 54(1):59–68
72. Neyroud N, Tesson F, Denjoy I, Leibovici M, Donger C, Barhanin J et al (1997) A novel mutation in the potassium channel gene KVLQT1 causes the Jervell and Lange-Nielsen cardioauditory syndrome. *Nat Genet* 15(2):186–189
73. Casimiro MC, Knollmann BC, Ebert SN, Vary JC Jr, Greene AE, Franz MR et al (2001) Targeted disruption of the *Kcnq1* gene produces a mouse model of Jervell and Lange-Nielsen Syndrome. *Proc Natl Acad Sci USA* 98(5):2526–2531
74. Lee MP, Ravenel JD, Hu RJ, Lustig LR, Tomaselli G, Berger RD et al (2000) Targeted disruption of the *Kvlqt1* gene causes deafness and gastric hyperplasia in mice. *J Clin Invest* 106(12):1447–1455
75. Pusch M (1998) Increase of the single-channel conductance of KvLQT1 potassium channels induced by the association with minK. *Pflugers Arch* 437(1):172–174
76. Splawski I, Tristani-Firouzi M, Lehmann MH, Sanguinetti MC, Keating MT (1997) Mutations in the *hminK* gene cause long QT syndrome and suppress IKs function. *Nat Genet* 17(3):338–340
77. Wang W, Xia J, Kass RS (1998) MinK-KvLQT1 fusion proteins, evidence for multiple stoichiometries of the assembled IsK channel. *J Biol Chem* 273(51):34069–34074
78. Jiang M, Xu X, Wang Y, Toyoda F, Liu X-S, Zhang M et al (2009) Dynamic partnership between KCNQ1 and KCNE1 and influence on cardiac IKs current amplitude by KCNE2. *J Biol Chem* 284(24):16452–16462
79. Dong M-Q, Lau C-P, Gao Z, Tseng G-N, Li G-R (2006) Characterization of recombinant human cardiac KCNQ1/KCNE1 channels (I (Ks)) stably expressed in HEK 293 cells. *J Memb Biol* 210(3):183–192
80. Wang Q, Curran ME, Splawski I, Burn TC, Millholland JM, VanRaay TJ et al (1996) Positional cloning of a novel potassium channel gene: KVLQT1 mutations cause cardiac arrhythmias. *Nat Genet* 12(1):17–23
81. Seeböhm G, Sanguinetti MC, Pusch M (2003) Tight coupling of rubidium conductance and inactivation in human KCNQ1 potassium channels. *J Physiol (Lond)* 552(Pt 2):369–378
82. Osteen JD, Sampson KJ, Kass RS (2010) The cardiac IKs channel, complex indeed. *Proc Natl Acad Sci USA* 107(44):18751–18752
83. Wang M, Kass R (2012) Stoichiometry of the slow I(Ks) potassium channel in human embryonic stem cell-derived myocytes. *Pediatr Cardiol* 33(6):938–942
84. Nakajo K, Ulbrich MH, Kubo Y, Isacoff EY (2010) Stoichiometry of the KCNQ1—KCNE1 ion channel complex. *Proc Natl Acad Sci USA* 107(44):18862–18867
85. Morin TJ, Kobertz WR (2008) Counting membrane-embedded KCNE beta-subunits in functioning K⁺ channel complexes. *Proc Natl Acad Sci USA* 105(5):1478–1482
86. Sakmann B, Trube G (1984) Conductance properties of single inwardly rectifying potassium channels in ventricular cells from guinea pig heart. *J Physiol* 347:641–657
87. Hibino H, Inanobe A, Furutani K, Murakami S, Findlay I, Kurachi Y (2010) Inwardly rectifying potassium channels: their structure, function, and physiological roles. *Physiol Rev* 90(1):291–366
88. Noble D (1965) Electrical properties of cardiac muscle attributable to inward going (anomalous) rectification. *J Cell Comp Physiol* 66(S2):127–135
89. Beeler GW Jr, Reuter H (1970) Voltage clamp experiments on ventricular myocardial fibres. *J Physiol (Lond)* 207(1):165–190
90. Kurachi Y (1985) Voltage-dependent activation of the inward-rectifier potassium channel in the ventricular cell membrane of guinea-pig heart. *J Physiol (Lond)* 366:365–85
91. McAllister RE, Noble D (1966) The time and voltage dependence of the slow outward current in cardiac Purkinje fibres. *J Physiol (Lond)* 186(3):632–662
92. Rougier O, Vassort G, Stämpfli R (1968) Voltage clamp experiments on frog atrial heart muscle fibres with the sucrose gap technique. *Pflugers Arch Gesamte Physiol Menschen Tiere* 301(2):91–108

93. Dhamoon AS, Pandit SV, Sarmast F, Parisian KR, Guha P, Li Y et al (2004) Unique Kir2.x properties determine regional and species differences in the cardiac inward rectifier K⁺ current. *Circ Res* 94(10):1332–1339
94. Hagiwara S, Takahashi K (1974) The anomalous rectification and cation selectivity of the membrane of a starfish egg cell. *J Membr Biol* 18(1):61–80
95. Miyazaki SI, Takahashi K, Tsuda K, Yoshii M (1974) Analysis of non-linearity observed in the current-voltage relation of the tunicate embryo. *J Physiol (Lond)* 238(1):55–77
96. Sakmann B, Trube G (1984) Voltage-dependent inactivation of inward-rectifying single-channel currents in the guinea-pig heart cell membrane. *J Physiol (Lond)* 347:659–83
97. Dhamoon AS, Jalife J (2005) The inward rectifier current (IK1) controls cardiac excitability and is involved in arrhythmogenesis. *Heart Rhythm* 2(3):316–324
98. Schram G, Pourrier M, Melnyk P, Nattel S (2002) Differential distribution of cardiac ion channel expression as a basis for regional specialization in electrical function. *Circ Res* 90(9):939–950
99. Nichols CG, Makhina EN, Pearson WL, Sha Q, Lopatin AN (1996) Inward rectification and implications for cardiac excitability. *Circ Res* 78(1):1–7
100. Miake J, Marbán E, Nuss HB (2003) Functional role of inward rectifier current in heart probed by Kir2.1 overexpression and dominant-negative suppression. *J Clin Invest* 111(10):1529–1536
101. Nerbonne JM, Kass RS (2005) Molecular physiology of cardiac repolarization. *Physiol Rev* 85(4):1205–1253
102. Lopatin AN, Nichols CG (2001) Inward rectifiers in the heart: an update on I(K1). *J Mol Cell Cardiol* 33(4):625–638
103. Nichols CG, Lopatin AN (1997) Inward rectifier potassium channels. *Annu Rev Physiol* 59:171–191
104. Lopatin AN, Makhina EN, Nichols CG (1994) Potassium channel block by cytoplasmic polyamines as the mechanism of intrinsic rectification. *Nature* 372(6504):366–369
105. Matsuda H, Saigusa A, Irisawa H (1987) Ohmic conductance through the inwardly rectifying K channel and blocking by internal Mg²⁺. *Nature* 325(7000):156–159
106. Vandenberg CA (1987) Inward rectification of a potassium channel in cardiac ventricular cells depends on internal magnesium ions. *Proc Natl Acad Sci USA* 84(8):2560–2564
107. Yamada M, Kurachi Y (1995) Spermine gates inward-rectifying muscarinic but not ATP-sensitive K⁺ channels in rabbit atrial myocytes. Intracellular substance-mediated mechanism of inward rectification. *J Biol Chem* 270(16):9289–9294
108. Lopatin AN, Makhina EN, Nichols CG (1995) The mechanism of inward rectification of potassium channels: long-pore plugging by cytoplasmic polyamines. *J Gen Physiol* 106(5):923–955
109. Glowatzki E, Fakler G, Brändle U, Rexhausen U, Zenner HP, Ruppersberg JP et al (1995) Subunit-dependent assembly of inward-rectifier K⁺ channels. *Proc Biol Sci* 261(1361):251–261
110. Yang J, Jan YN, Jan LY (1995) Determination of the subunit stoichiometry of an inwardly rectifying potassium channel. *Neuron* 15(6):1441–1447
111. James AF, Choisy SCM, Hancox JC (2007) Recent advances in understanding sex differences in cardiac repolarization. *Prog Biophys Mol Biol* 94(3):265–319

Modelling the Short QT Syndrome Gene Mutations
And Their Role in Cardiac Arrhythmogenesis

Adeniran, I.

2014, XVII, 217 p. 102 illus., 38 illus. in color.,

Hardcover

ISBN: 978-3-319-07199-2