

Carbon Footprint Estimation in the Agriculture Sector

Divya Pandey and Madhoolika Agrawal

Abstract The term “carbon footprint” has evolved as an important expression of greenhouse gas (GHG) intensity for diverse activities and products. Widespread public acceptance and the ease of conveying information about GHG intensity with this term has also attracted scientists and policy makers to review and refine its calculations. Standard methods for carbon footprinting have been prepared, and sector-specific standards are under development. These standards direct the procedures to carry out carbon footprinting through life cycle assessment in conjunction with GHG accounting, classifies activities into three tiers based on the order of emissions. Agriculture is the largest contributor to anthropogenic emissions of greenhouse gases, so the quantification of different agricultural practices is essential for identification of more sustainable practices. Carbon footprinting has potential as a tool for assessing and comparing GHG performances of different agricultural products along with identification of points to improve environmental efficiencies. Case studies on the application of carbon footprinting to cultivation practices are increasing in the scientific literature, but the majority of studies do not comply with the standard three-tier methodology. This leads to nonuniformity among different studies and their comparisons. Hence, a standard guideline addressing carbon footprinting specifically for agriculture is essential for the effective application of this tool in the quantification of GHG intensity, mitigation of global warming, and adaptation against future climate change scenarios.

Keywords Cultivation practices • Mitigation • Agricultural management • Three-tier methodology

D. Pandey · M. Agrawal (✉)

Laboratory of Air Pollution and Global Climate Change, Department of Botany,
Banaras Hindu University, Varanasi 221005, India
e-mail: madhoo.agrawal@gmail.com

1 Introduction

Climate change has emerged as the biggest environmental and developmental challenge of the present time; it also influences the focal possibilities for sustainable development. The effects of climate change have already been felt all over the world, in diverse forms ranging from shifting weather patterns, receding ice caps, crop losses, altered distribution of precipitation, increased frequency and intensities of floods and droughts, and serious ecological imbalances. All of these effects also have resulted in significant economic losses (Stern 2006). To prevent projected and unforeseen disasters, global temperatures must not exceed 2 °C more than 1990 levels. For this, the atmospheric stock of greenhouse gases (GHGs) should be controlled to remain below 550 ppm in terms of CO₂ equivalents (CO₂-e). Among different GHGs, carbon dioxide (CO₂), methane (CH₄), nitrous oxide (N₂O), sulfur hexafluoride (SF₆), perfluorocarbons (PFCs), and hydrofluorocarbons (HFCs) are the six important anthropogenic GHGs. GHG inventories can identify, quantify, and manage all sources and sinks of GHGs. Among different quantitative indicators, the carbon footprint has gained popularity and widespread application. Moreover, because of its ease of conveying information about the GHG intensity of variety of products and activities among the general public, carbon footprint also offers a simple mode of communication about climate responsibility of different entities between people, scientists, and policy makers. Scientific analyses of carbon footprinting are being conducted, mainly for consumer products and industrial processes; its application to agricultural systems is less, despite the fact that agriculture alone is responsible for GHG emissions to the largest degree. Here, we review the available scientific literature on the concept and calculations of carbon footprint, and its application to the agriculture sector. We begin with an overview of agriculture's role in regulating GHG fluxes, followed by the concept and general principle of carbon footprinting. Applications and challenges in using carbon footprinting in agriculture are also discussed.

2 Agriculture as a Source of Greenhouse Gases

Covering about 35 % of the land area, agriculture accounts for nearly 13.5 % of the total global anthropogenic GHG emissions, contributing about 25, 50, and 70 % of CO₂, CH₄, and N₂O, respectively (Montzka et al. 2011). As it is recognized that cereal production must increase at a rate not less than 1.3 % annually (Cassman et al. 2003), related emissions are also expected to increase. GHG emissions from agriculture originate mainly in the form of CH₄ from rice cultivating systems and cattle rearing and N₂O from fertilizer management practices.

Rice fields alone emit 32 to 44 Tg CH₄ yr⁻¹ (Le Mer and Roger 2001). Del Grosso et al. (2008) estimated that agricultural activities add into the atmosphere about 4.2 to 7 Tg N annually in the form of N₂O. Due to their high global warming

potential of 298, emissions of N_2O , even in a small quantity, cause significant radiative forcing. Increased soil temperatures coupled with high moisture conditions during cooler months will increase N_2O production in soil. Elevation in CO_2 concentrations is also projected to increase N_2O emissions from upland agricultural soils (Van Groenigen et al. 2011). Regarding CO_2 , soil respiration is an important source, but the majority of the farm operations and inputs, such as fertilizers, pesticides, and energy, also have embodied CO_2 content.

The majority of GHG estimates cover only soilborne emissions, generally of CH_4 and N_2O only, whereas numerous studies have been carried out targeting only CH_4 measurements (Le Mer and Roger 2001) and its mitigation from rice fields, mainly through water (Pathak et al. 2003), fertilizer, and manure managements (Linguist et al. 2012). Among different management techniques, mulching and organic manure applications are found to increase the emissions of CH_4 (Ma et al. 2007), whereas midseason drainage can cut CH_4 emissions significantly (Zou et al. 2005). Aerobic soils, on the contrary, may act as CH_4 sinks (Le Mer and Roger 2001; Smith et al. 2008) or sources (Ma et al. 2013), but they too are poorly quantified (Robertson 2000).

As a widely recognized effect, application of mineral nitrogen increased the emissions of N_2O . However, the effects of different management practices on emissions of all the GHGs are highly inconsistent, depending on the cultivation system and environmental conditions. Some inhibitors to methanogenesis and nitrification have also been tested in agricultural soils (Liu et al. 2010). It is found that frequency and timing of tillage also influence fluxes of soilborne GHGs. In the long term, the elimination of tillage reduced the emissions of CH_4 and N_2O , but increased CO_2 from rice cultivation (Pandey et al. 2012) as compared to conventional practice of regular tillage.

3 Agricultural Management as a Carbon Offsetting Option

Although, agriculture is an emissions source, there are opportunities for reducing the emissions and even using cultivated soils as a GHG offsetting tool if better management practices are identified and adopted (Hutchinson et al. 2007). Soils are the largest terrestrial carbon store; they hold carbon in the form of organic and inorganic molecules. Due to erosion and oxidation, a significant part of soil organic carbon has been lost. Scientific evidence suggests that 50–66 % of the cumulative historic carbon loss from soil can be recovered if managed intelligently (Lal 2004b). Increasing the organic carbon content in soil may lock the carbon out of the atmosphere for centuries a phenomenon is termed as carbon sequestration. The two fundamental keys to support carbon sequestration in soils is minimization of soil disturbance and increasing inputs of organic matter. Therefore, cover crops, mulching, no tillage, organic manure, and decreasing the fallow period are among the recommended management practices (Lal 2004a). Improvement in nutrient status, particularly of nitrogen and phosphorous, also strengthens carbon

sequestration; hence, intercropping with legumes and certain permutations of crop rotations are found to be greatly effective (Nishimura et al. 2008). It is estimated that under recommended management, soils of the European Union and UK, respectively, can offset 0.09–0.12 and 0.010 Pg C annually (Smith et al. 2005); at the global level, soils offer an annual sequestration potential of 0.6–2.0 Pg C (Lal 2000). Long-term studies have shown that organic manure application increases the carbon sequestration capacity of soil in the range of 70–551 kg C ha⁻¹ as compared to mineral fertilizer use (Mandal et al. 2007, 2008).

4 Understanding Product Carbon Footprints: Concept, Scope, and Calculation

Carbon footprints originated as a subset of the “ecological footprint” proposed by Wackernagel and Rees (1996). *Ecological footprint* referred to the biologically productive land and sea area required to sustain a given human population, expressed as global hectares. According to this concept, carbon footprint was the land area that will assimilate the CO₂ produced during the lifetime of a person or total global population. The calculation of carbon footprint as a part of the ecological footprint was very tedious and complex. But as the issue of global warming gradually gained prominence on the global environmental forefront, carbon footprinting emerged independently, in a modified form (East 2008). The present form of carbon footprints can be regarded as a hybrid that derives its name from “ecological footprint” but conceptually is a global warming potential indicator. However, few studies still report carbon footprints in terms of global hectares with regard to its origin (Browne et al. 2009).

In spite of its widespread popularity among the public as an indicator of contribution of an entity to the global warming, until few years back, there was confusion over what carbon footprints exactly meant (Wiedmann and Minx 2007; Pandey et al. 2011). This was particularly due to the lack of a standard methodology for carbon footprint calculation and its scientific analyses. Most studies have been carried out by private organizations and companies for business purposes rather than environmental responsibility (Kleiner 2007; East 2008). However, recognizing the public response to carbon footprinting studies and increasing financial transactions in the carbon market, standards are now under construction by the International Organization for Standardization (ISO); British Standards Institution (BSI) is also developing and upgrading their guidelines. Scientific literature is also growing, with more and more case studies of carbon footprinting, thus adding to the development of standard methods.

Based on a survey, Wiedmann and Minx (2007) recognized that definitions of carbon footprints were also different among different studies. They suggested that the term carbon footprint should reflect measure of the exclusive total amount of CO₂ emissions that is directly and indirectly caused by an activity or is accumulated

over the life stages of a product. A similar indicator, “climate footprint,” was proposed to be used, if all the GHGs were included in the calculation instead of only CO₂. But keeping in mind the motive of carbon footprinting, (i.e., assessing the impact of the activity on global climate), new studies and guidelines suggested inclusion of all the GHGs that are covered under the Kyoto Protocol (Kelly et al. 2009; Eshel and Martin 2006). However, still there are terms that are used interchangeably with carbon footprints, such as embodied carbon, carbon content, embedded carbon, carbon flows, virtual carbon, GHG footprint, and climate footprint (Courchene and Allan 2008; Peters 2010). Selection of direct and embodied emissions is also inconsistent among different studies. Direct emissions take place onsite. For example, in an industrial unit, CO₂ released during the combustion of gasoline fired in boiler is a direct emission. On the other hand, if the boiler was electrically powered, no direct emissions will be observed on the site. But during production of that electricity in a thermal power plant, a certain amount of CO₂ should have been released. Such an emission is referred as the embodied or indirect emission. In most cases, it becomes too complicated to include all possible indirect emissions; hence, many carbon footprinting case studies report only direct or first-order indirect emissions (Carbon Trust 2007; Wiedmann and Minx 2007; Matthews et al. 2008). But indirect emissions may constitute the major share of carbon footprints for many activities (Matthews et al. 2008). In spite of prevailing differences among the calculations, the CO₂ equivalent (CO₂-e) mass based on 100 years global warming potential of GHGs is used as the reporting unit of carbon footprints (WRI/WBCSD 2004; Carbon Trust 2007; BSI 2008), although there had been certain critical comments over it. Hammond (2007) and Global Footprint Network (2007) hold the opinion that “footprints are spatial indicators”; therefore, the carbon footprint should precisely be called a “carbon weight” or “carbon mass” (Jarvis 2007). However, convenient calculations and widespread acceptance makes CO₂-e mass the practical unit of carbon footprints.

The definition of carbon footprints is therefore proposed as follows: “The quantity of GHGs expressed in terms of CO₂-e, emitted into the atmosphere by an individual, organization, process, product or event from within a specified boundary” (Pandey et al. 2011).

4.1 Scope of Product Carbon Footprinting

The main drivers of carbon footprint calculations are legislative requirements, carbon trading, corporate social responsibility, and scientific analyses for devising effective policies to combat global warming (Carbon Trust 2007). The scope of carbon footprinting is wide and includes virtually all kinds of products, services, activities, and processes. Carbon footprinting of products and services has proven useful in not only managing the emissions more effectively across the supply chain, but also as a business tool (Kleiner 2007). It is proven by the continually

increasing number of companies participating in the Carbon Disclosure Project (CDP 2009). This rush gained momentum from changing marketing strategies as more consumers began to prefer products with low carbon footprints (LEK Consulting 2007). Therefore, regulated carbon labeling of products has also been introduced. Suh (2006) calculated carbon footprints for different products in the USA and concluded that lime was the most GHG intensive product ($22.1 \text{ kg CO}_2\text{-e } \$^{-1}$), followed by chemicals, fertilizers, and meat production. Services such as health care, water supply, computing and data processing, and amusement left smaller carbon footprints, ranging from 42.1 to 46.1 Tg $\text{CO}_2\text{-e}$ for an average household. Hoefnagels et al. (2010) used product carbon footprinting for comparing the overall performances of different energy options. Under optimum conditions, biofuels production emitted between 17 and 140 g $\text{CO}_2\text{-e MJ}^{-1}$. Carbon footprints of different fuels are calculated to decide about the import of nonconventional vehicular fuels in California (Courchene and Allan 2008). Gemechu et al. (2012) also advocated the application of carbon tax based on product carbon footprinting for kraft pulp production, in which energy usage was the most polluting sector with nearly $0.32 \text{ kg CO}_2\text{-e kg}^{-1}$ of pulp produced.

Among services, aviation has been identified among the highest GHG emitters; hence, the carbon footprinting of airlines is ongoing, covering different aspects such as aircraft types, load factors, and seat configurations (Miyoshi and Mason 2009). The European Union has taken the lead in formulating legal bindings for reduction in emissions embodied in aviation. Schools and universities are also participating in such calculations. GAP et al. (2006) in the UK Schools Carbon Footprint Scoping Study estimated that, in 2001, all schools in the United Kingdom left a carbon footprint of $9.2 \times 10^9 \text{ kg CO}_2\text{-e}$. Elsewhere, the University of British Columbia and University of Pennsylvania left carbon footprints of 8.2750×10^7 and $3.0 \times 10^8 \text{ kg CO}_2\text{-e}$, respectively (Ferris et al. 2007; TC Chan Centre for Building simulation and Energy Studies/Penn Praxis 2007). Carbon footprints have also been included in the management of cities and organizations to improve environmental policies (Courchene and Allan 2008; Good Company 2008). The UNDP (2007) and Edgar and Peters (2009) used per capita CF of different countries to compare the contributions of countries, cities, and sectors to global warming. These reports clearly indicated that high-income countries leave the biggest footprint, while it was substantially lower for developing countries. Carbon footprint is now used as an indicator in event management as well (London-2012 Sustainability Plan 2007).

In addition to the above, voluntary carbon footprinting by organizations as well as individuals is growing at a fast rate. Consultancies and online calculators have further promoted individual carbon footprinting, particularly in developed countries (Padgett et al. 2008; Kenny and Gray 2008). Such calculators also offer carbon offsetting options, mainly through supporting forestation and renewable energy resources (Murray and Day 2009). A dramatic growth in the voluntary carbon market has been reported since 1989 (Hamilton et al. 2007). Carbon footprinting is also extended to the natural and semi-natural systems, which may

help compare natural versus anthropogenic impacts on the environment (Chambers et al. 2007). Hence, we see that there is hardly any entity that cannot be a candidate for carbon footprinting.

4.2 Calculation of Product Carbon Footprints

Being a quantitative expression of GHG emissions, carbon footprinting helps in emission management and evaluation of mitigation measures (Carbon Trust 2007). Through carbon footprint analyses, important sources of emissions can be identified and areas of emission reductions can be prioritized. For carbon footprint calculation, estimates of GHGs emitted/embodyed at each identified step of the product's/activity's/individual's life cycle are conducted, which is technically known as GHG accounting. Standards and guidance are available for GHG accounting. Common resources are:

- (a) GHG protocol of World Resource Institute (WRI)/World Business Council on Sustainable Development (WBCSD): Nearly all GHG accounting guidelines, including ISO 14064 and PAS 2050 of BSI (2008), are based on this protocol. The GHG protocol provides separate guidelines for GHG accounting and reporting during the life cycles of products and corporate organizations. ISO 14064 (parts 1 and 2): International Organization for Standardization has developed this standard for determination of boundaries, quantification of GHG emissions, and removal (ISO 2006a, b). Part 1 deals with carbon footprinting of organizations, addressing guidance for the quantification, monitoring, and reporting of GHG emissions. Part 2 deals specifically with well-defined activities and projects.
- (b) Publicly Available Specifications-2050 (PAS 2050) of British Standard Institution (BSI): It specifies the requirements for assessing the life cycle GHG emissions of goods and services (BSI 2008). PAS 2050 is preparing a standard method to calculate carbon footprints of agricultural systems as well.
- (c) Intergovernmental Panel on ClimateChange (IPCC) guidelines for National Greenhouse Gas inventories: IPCC categorizes all anthropogenic sources of GHG emissions into four sectors—energy, industrial process and product use, agriculture, forestry, and other land use and waste.

All of these guidelines and standards proceed through life cycle assessment (LCA) or 'cradle-to-grave analyses' for the activity for which the carbon footprint is to be calculated. ISO formulated standard methods for conducting LCA as a part of the ISO 14000 series. ISO 14040 provides the principles and framework for carrying out LCA (ISO 2006c), whereas ISO 14044 provides guidelines on detailed methodology (ISO 2006d). It also directed the Life Cycle Impact Assessment (LCIA) as the last and compulsory stage of LCA. For effective application of ISO 14044, two technical revisions have been made: ISO 14047

(ISO 2012a) and 14049 (ISO 2012b). They focus on the key points of LCIA that are important for carbon footprinting, with specific examples and sample practices.

To provide guidelines and principles of product carbon footprinting, ISO 14067 is under development (ISO 2013). This technical specification is based on the ISO's standards of LCA and environmental labeling of products. Although there are provisions of different modes of communicating product carbon footprints and performance tracking, it is under critical review and evaluation so that an international standard can be developed.

According to the available standards, following structured framework is suggested for carbon footprinting (WRI/WBCSD 2004; Carbon Trust 2007; BSI 2008):

- a. Selection of GHGs
- b. Setting boundaries
- c. Collection of GHG emission data
- d. Footprint calculation

4.2.1 Selection of GHGs

Selection of the set of GHGs covered in the calculation depends on the guideline followed, the need for carbon footprinting, and the type of activity. For example, in a thermal power plant, where CO₂ is a predominant emission and other gases are almost negligibly emitted, only CO₂ emission measurement will be feasible, whereas for a cattle farm, CH₄, CO₂, and N₂O emissions may be significant. Although some studies include only CO₂ emissions in carbon footprinting (Patel 2006; Wiedmann and Minx 2007; Craeynest and Streatfeild 2008), the guidelines recommend all six Kyoto gases (Bokowski et al. 2007; Garg and Dornfeld 2008; Good company 2008; Matthews et al. 2008). All guidance and standards also direct to include all Kyoto gases.

4.2.2 Setting Boundaries

A boundary refers to an imaginary line drawn around the activities that will be used for calculating carbon footprints. It depends on the objective of footprinting and characteristics of the entity for which footprinting will be done. Defining the boundary is crucial as it determines the activities, which will be included in the study. To facilitate convenient accounting, the following tiers have been suggested (WRI/WBCSD 2004; Carbon Trust 2007; BSI 2008):

- Tier₁: direct, i.e., onsite emissions
- Tier₂: emissions embodied in purchased energy
- Tier₃: all indirect emissions not covered under tier₂, such as those associated with the transport of purchased and sold goods, business travels, waste disposal, etc. Carbon footprint has also been divided into two parts: basic/primary and full carbon footprint. Primary carbon footprint is calculated from tier₁ and tier₂ only, whereas full carbon footprint covers emissions up to tier₃ (Carbon Trust 2007; Lynas 2007).

Most carbon footprinting studies limit up to tier₂ because going beyond tier₂ increases the complexity and uncertainty in estimates (Matthews et al., 2008). Even during trading of carbon offsets, only tier₁ and tier₂ emissions are important. It is also advocated that embodied emissions are beyond the control of the organization of process for which carbon footprints are to be calculated and hence tier₃ should be left out during carbon footprinting (Lenzen 2001). For this reason, PAS-2050, GHG protocol, and other registries and consultancies based on these have kept tier₃ optional. Critical analyses of carbon footprinting case studies, however, reveal that indirect emissions compose a significant part of total carbon footprint (Matthews et al. 2008). Hence, attempts must be taken to count tier₃ emissions. To make the definition of tier₃ more clear, Mathews et al. (2008) proposed that emissions exclusively related to delivery, use, and disposal of products also should be kept out of tier₃. An additional tier₄ can be used for the same.

Advancement in the tracking and management of emissions in the supply chain is expected to promote tier₃ accounting (Matthews et al. 2008; CDP 2009). In the Carbon Disclosure Project (CDP), 72 % of respondents among 500 companies reported their basic carbon footprints, but the number of companies reporting up to tier₃ is increasing (CDP 2009). As more and more organizations carry out their complete LCA, a database can be developed through which average sector-specific emission factors can be calculated (Matthews et al. 2008; Weidema et al. 2008). International trade of raw materials and finished products poses further challenges in tier₃ estimation (Courchene and Allan 2008). Appropriate assumptions over sharing of responsibilities of countries and organization related with emissions associated with international trade of goods and services need to be developed (Peters 2010).

Regarding natural systems and land uses, almost all the carbon footprinting studies focus on emissions; the amount of GHG removal and sequestration appears to be neglected (Peters 2010).

4.2.3 Collection of GHG Data

Estimation of GHG emissions and removals associated with all the activities identified within the boundary can be carried out by direct measurements or estimated using emission factors or models. Direct measurements yield near

accurate estimates and are clearly prescribed in globally accepted protocols, but their cost and application may be prohibitive in certain cases (WRI/WBCSD 2004). Under such conditions, indirect estimations through models and emission factors are applicable. If developed or modified specifically for a particular region or sector, they yield fairly accurate results. Usually, customized tools relying on combinations of direct measurements, emission factors, and models are popular and practicable (USCCTP 2009). For large-scale GHG estimation, observation networks such as FLUXNET have been initiated (Sundareshwar et al. 2007), but due to high costs and nonuniform global distribution of sites, they are still far away from global representativeness. To overcome the patchy coverage of ground-based monitoring networks, satellites have been launched to monitor sources and sinks of CO₂ and other GHGs (Haag 2007). A Japanese satellite, (the “greenhouse gas observing satellite”) and Vulcan, a joint project of NASA and the US Department of Energy, are two such examples (Gurney et al. 2009; Kelly et al. 2009). Remote sensing and geographic information systems are also used extensively for large and relatively less accessible areas. Such an example is the case of carbon footprinting of Hurricane Katrina on the US coast, carried out by Chambers et al. (2007) using LANDSAT imageries.

4.2.4 Footprint Calculation

The collected GHG data is translated into CO₂-e using global warming potentials of different GHGs as provided by IPCC (2007). The final unit of the carbon footprint depends on the nature of the entity. For individuals and dynamic processes, carbon footprints need to be calculated periodically, but events such as conferences, sports events, etc. have one-time emissions. Some entities have a combination of both; for example, for building, a one-time emission take place during construction phase, while periodic calculations are needed during the operation phase. For such activities, there is a provision of sharing one-time emissions over the operation phase. Natural processes are highly complex; hence, they have a temporally as well as spatially dynamic carbon footprint.

5 Carbon Footprinting as a Tool to Estimate Agriculture’s Contribution to Atmospheric Stock of Greenhouse Gases

As discussed, the emissions as well as sink capacity of the agriculture sector are still highly uncertain, and available estimates must be refined through an extensive monitoring network covering different geographical regions, environmental conditions, and management practices (Seip 2011). In addition to soilborne GHGs and carbon sequestration, keeping in mind the increasing energy and chemical inputs in farming, the boundaries of agriculture must be expanded to include all relevant

emissions and/or removals of GHGs. Carbon footprinting therefore can be utilized for cultivation systems by producing a detailed map of different sources and sinks of GHGs. This will identify the points where environmental efficiencies can be improved. This also facilitates a comparison of different management options and their environmental cost-benefit analyses. Although scientific literature is still sparse in carbon footprinting studies targeting cultivation practices, such estimates are essential for upgrading national GHG budgets and to improve environmental efficiency of the agriculture sector.

6 Calculating Carbon Footprints for Agricultural Products

The GHG protocol acts as a common resource for carbon footprinting, but it is important to keep in mind the role of the agriculture sector in anthropogenic GHG emissions and sensitivity of this sector to a number of environmental and social factors. Therefore, the development of an agriculture-specific carbon footprinting method is proposed. BSI is developing PAS 2050 specifically for agriculture.

6.1 *Selection of Boundary and Tiers*

For carbon footprinting of agricultural practices, all activities associated with farming must be identified. A generalized illustration of different activities involved in cultivation practices that are relevant to carbon footprinting is presented in Fig. 1. Because there is no agriculture-specific standard for carbon footprint calculation, the generalized standard three-tier approach of the GHG protocol must be followed in order to maintain uniformity among different studies. The selection of the boundary will depend on the level up to which carbon footprints are to be calculated, as presented in Table 1. For carbon footprinting in the production of cereals, vegetables, fruits, etc., the activities related to cultivation of the concerned crop up to the final harvest and readiness for use as raw material will be covered. To cover the activities up to the shelf of the store, activities related to processing, packaging, and transportation of farm produce must also be included. To calculate carbon footprints of food, the boundary is set to cover home preparations also. Among the three proposed boundaries, carbon footprints of cultivation (i.e., up to the farm gate) is more helpful for comparing different agricultural practices and efficacies of different management systems on GHG performances. Extending the boundary beyond this introduces activities such as the transportation of products to the market, their distribution, and food preparation techniques and preferences, which are more sensitive to local and personal conditions.

Direct emissions to be covered under tier₁ for agricultural systems include CH₄, N₂O, and CO₂ emissions from soil and onsite CO₂ emissions from fossil fuel-powered farm machines such as tractors, harvesters, threshers, grain cleaning




Fig. 1 A generalized illustration of activities and inputs associated with cultivation of a crop to be considered in the boundary (sowing to farm gate) for carbon footprinting

Table 1 Choice of boundaries for carbon footprinting of agricultural products

Objective	Boundary
Carbon footprinting of cultivation	Up to the farm gate
Carbon footprinting of finished farm products	Up to the shelf
Carbon footprinting of food	Up to the table

systems, etc. Electricity use in activities such as irrigation constitutes tier₂. Table 2 lists common farm activities and their classification into different tiers. Most of these activities are also performed manually, but human labor is not considered under carbon footprint calculations (WRI/WBCSD 2004). Because agricultural soils can sequester atmospheric CO₂ (Lal 2004b), it is proposed to be a part of tier₁.

In addition to these, agricultural inputs such as fertilizers, pesticides, herbicides, and soil conditioners carry embodied emissions, and hence they constitute tier₃.

6.2 Estimation of GHG Emissions/Removals

Because agricultural practices depend significantly on region, traditional practices, economic conditions of farmers, and the crop under cultivation, emission factors and models developed to express emissions of GHGs need to be validated and refined before their application in a particular agricultural system. Particularly for soilborne GHG emissions, which are also sensitive to environmental conditions, direct measurements are the most reliable (Pandey et al. 2012). For this purpose,

Table 2 Farm activities and their classification into tiers

Activity	Cultivation practices	Energy source	Tier
Land preparation	Plow	Diesel	Tier ₁
	Harrow	Diesel	Tier ₁
	Spader	Diesel	Tier ₁
	Subsoiler	Diesel	Tier ₁
	Spreader	Diesel	Tier ₁
Sowing	Seed drill	Diesel	Tier ₁
	Broadcast	Diesel	Tier ₁
	Seeders/spreaders	Diesel	Tier ₁
	Transplanter	Diesel	Tier ₁
Irrigation	Channel	Electricity	Tier ₂
	Sprinkler drip	Electricity	Tier ₂
		Electricity	Tier ₂
Fertilizer application	Spreader	Diesel	Tier ₁
	Self-propelled sprayers	Diesel	Tier ₁
	Agricultural aircrafts	Petroleum spirit	Tier ₁
Pesticide application	Self-propelled sprayers	Diesel	Tier ₁
	Agricultural aircrafts	Diesel	
Irrigation	Channel	Diesel/electricity	Tier ₁ /Tier ₂
	Drip	Electricity	Tier ₂
	Sprinkler/central pivot	Electricity	Tier ₂
Harvesting	Harvester (reaper, thresher)	Diesel/electricity	Tier ₂
Threshing	Thresher	Diesel/electricity	Tier ₂
Seed processing	Seed processing systems	Diesel/electricity	Tier ₂

closed chamber systems are a simple, low-cost, and most applied technique (Parkin et al. 2012). There have been different shapes, sizes, and sampling procedures for estimating GHG fluxes using chamber systems depending upon the crop, rates of GHG fluxes, and analyses procedures. The periodically sampled gases are usually analyzed through gas chromatography and infrared gas analyzers; however, with advancements in chamber technique, GHG flux rates can now be measured in situ (Arnold et al. 2001). Automated chambers are capable of carrying out continuous monitoring of GHGs covering diurnal and daily variations under different environmental conditions. However, their cost, maintenance, and electricity requirement prohibits their application in the farmer's field and remote rural areas. The sensitivity of chambers also poses a challenge; in many cases, emission rates are very low and remain below the detection limits of chambers (Parkin et al. 2012). This is particularly important when soil acts as a net sink of GHGs. Also, due to poor chamber sensitivities, low or negative fluxes are often discarded as experimental errors (Chapuis-Lardy et al. 2007). Such low positive or negative fluxes become significant for large cultivated areas. With more refined chamber designs and flux calculation methods, sensitivity has been improved significantly. Flux towers are meant for large farming areas under similar cultivation practices and cropping systems because they provide the cumulative flux of the entire coverage area (Sundareshwar et al. 2007).

Regarding changes in soil carbon, which also constitutes tier₁, actual measurement of stock difference in soil over a long period of time is needed. This is because the changes in soil carbon are too slow to be measured reliably over time scales of years (Post et al. 2001). It is observed that changes in soil carbon are a function of management practices, environmental conditions, crop cultivation, and depth of measurement. Another issue related to carbon sequestration measurement is the question of permanence, i.e., for how long the carbon accrued in the soil will stay out of the atmosphere. Therefore, this part is usually left out during GHG accounting, but some studies have shown that carbon sequestration in soils can offset a part of carbon footprint of cultivation systems significantly.

For emissions taking place from farm machines and electricity consumption, emission factors are available for most countries. According to the GHG protocol, an activity data sheet should be maintained, keeping records of different farm activities, fuel consumption, hours of operations, etc. From this activity data sheet, emissions associated with different activities can be calculated using emission factors or models. As a requirement of carrying out GHG emission inventories for countries signatory to the United Nations Framework Convention on Climate Change (UNFCCC), emissions associated with combustion of fossil fuels and electricity generation have to be calculated under their national communications to UNFCCC. These emission factors have been modified according to the types of engines and machines used on the field. However, for tier₃ emissions, production technique and country-specific emission factors are not available in most of the countries. In such cases, IPCC (2006) National GHG inventory guidelines provide average and default emission factors for production of fertilizers, GHG emissions from soil under different irrigation, and manure applications etc. Based on these factors, GHG emissions embodied in Urea production have been calculated (Tirado et al. 2010; Lal 2004a) as the most commonly used resource, in which emission factors for CO₂ emissions associated with different activities on farm and inputs of fertilizers and pesticides were derived on the basis of extensive literature survey. Such estimates nevertheless need to be refined and updated. Nelson et al. (2009) also calculated the onsite and offsite CO₂ emissions from different farm activities in USA during 1990–2004. Results indicated that onsite and total CO₂ emissions ranged from 23 to 176 kg C ha⁻¹ yr⁻¹ and from 91 to 365 kg C ha⁻¹ yr⁻¹, respectively. Such region-specific emission factors are necessary for reducing uncertainties in the calculations.

6.3 Footprint Calculation

Global warming potential (GWP) of all the tiers is calculated individually using the conversion factors of IPCC (2007) corresponding to a 100-year time horizon. The formula for the calculation of GWP of tier_{*i*} (*i* = 1, 2 or 3) is given by:

$$\text{GWP}(\text{tier}_i) = \text{emission/removal of CH}_4 \times 25 + \text{emission/removal of N}_2\text{O} \times 298 + \text{emission/removal of CO}_2$$

where GWP is in kg CO₂-e ha⁻¹.

Emissions are taken as positive while removal as negative. Values are given in kg ha⁻¹.

Carbon footprint is calculated by adding the GWP of all tiers. The final representation of the carbon footprint of agricultural systems can be made as spatial or yield scaled carbon footprints according to the formulae given below:

$$\text{CF}_s = \sum_{i=1}^3 [\text{GWP}(\text{tier}_i)]$$

$$\text{CF}_y = \frac{\text{CF}_s}{\text{Grain yield}}$$

where CF_s is the spatial carbon footprint. Units are (kg CO₂-e ha⁻¹); CF_y is yield scaled carbon footprint. Units are (kg CO₂-e Kg⁻¹ yield).

These that two units differ by the factor of yield, which is the prime motive of cultivation. Spatial carbon footprints are helpful in comparing agricultural practices that are already under high yielding conditions. Under such cases, the better practices emit less per unit area under cultivation without declining the yield. Yield scaled carbon footprints are considered a better indicator for intercomparison of different cropping systems (Linguist et al. 2011, 2012).

7 Case Studies

In the last few years, there has been an increase in number of case studies of carbon footprinting of cultivation systems and food. Some of them considered all three tiers, but none of the studies defined them. Similarly, there was no mention of boundary selection. Table 3 presents different carbon footprinting studies for crop cultivation. Even though it is remarked that GHG emissions from soil are highly sensitive to environmental conditions and management practices, none of the carbon footprinting studies was based on actual measurements. Furthermore, CH₄ emissions were considered only for rice cultivation; for the rest, only N₂O emissions were covered (Gan et al. 2011a, b, c, 2012). In light of many studies demonstrating that crops other than rice act as significant CH₄ sinks or sources, it becomes essential to monitor CH₄ fluxes under such systems. In an attempt to calculate the yield scaled carbon footprint of barley, Gan et al. (2012), calculated that nearly 26 % of the GHG emissions were contributed by farm operations. Gan et al. (2012) however, did not measure CH₄ emissions. In most of the studies, tier₃ emissions, particularly of fertilizer application alone, contributed from 45 to 85 % of the total yield scaled carbon footprints (Gan et al. 2011a, b, 2012), whereas

Table 3 Carbon footprints of some agricultural systems and cultivation practices

Agricultural system (crop)	Region	Activity and tiers	Estimation protocol	Carbon footprint (kg CO ₂ -e kg ⁻¹) ^a	Reference
1 Canola-mustard (sowing to farm gate)	Canada	Tier ₁ and Tier ₂ : soilborne N ₂ O, land preparation, pesticide and fertilizer spray, harvester Tier ₃ : fertilizers and pesticides (factory to farm)	Rochette et al. (2008); Lal (2004a)	0.548–0.966	Gan et al. (2011a)
2 Durum wheat (sowing to farm gate)	Canada	Tier ₁ and Tier ₂ : soil borne N ₂ O, land preparation, pesticide and fertilizer spray, harvester Tier ₃ : fertilizers and pesticides (factory to farm)	Rochette et al. (2008); Lal (2004a)	0.383–0.533	Gan et al. (2011c)
3 Barley (sowing to farm gate)	Canada	Tier ₁ and Tier ₂ : soilborne N ₂ O, land preparation, pesticide and fertilizer spray, harvester Tier ₃ : fertilizers and pesticides (factory to farm)	Rochette et al. (2008); Lal (2004)	0.252–0.456	Gan et al. (2011b)
4 Spring wheat (sowing to farm gate)	Canada	Tier ₁ and Tier ₂ : soilborne N ₂ O, land preparation, pesticide and fertilizer spray, harvester, changes in soil C Tier ₃ : fertilizers and pesticides (factory to farm)	Gregorich et al. (2005); Rochette et al. (2008); EF from previous studies	0.357–0.140	Gan et al. (2012)
5 Potato (farm to table)	Sweden	Tier ₁ and Tier ₂ : soilborne N ₂ O, soil borne CO ₂ changes in soil C Electricity	SEPA (2009); IPCC (2006); EC (2007); Swedenergy, (2009); Andrén et al. (2004)	0.12	Roos et al. (2010)

(continued)

Table 3 (continued)

Agricultural system (crop)	Region	Activity and tiers	Estimation protocol	Carbon footprint (kg CO ₂ -e kg ⁻¹) ^a	Reference
6 Rice (sowing to farm gate)	Italy	Soilborne CH ₄ and N ₂ O	EF from previous studies	2.90	Blengini and Busto (2009)
7 Biofuels (ethanol, methyl ester, FT diesel) (Cultivation to power generation)	Europe, Canada, S.E. Asia, Brazil, USA	Fertilizer, farm machines (land preparation, harvester, grain processing), irrigation, pesticide and fertilizer spray, harvester Tier ₁ and Tier ₂ : no demarcation between tiers	IPCC (2007); Croezen et al. (2008)	In kg CO ₂ -e 70-140 MJ ⁻¹	Hoefnagels et al. (2010)
8 57 farms with different organic, conventional and integrated farming (sowing to farm gate)	Scotland	Soil borne CH ₄ , N ₂ O, CO ₂ Change in soil carbon Power generation Tier ₃ : fertilizers and pesticides (factory to farm) Tier ₁ and Tier ₂ : soilborne N ₂ O; Land preparation; Harvesting; Fertilizer, pesticide spray	Lal (2004a)	In ×10 ² kg CO ₂ -e ha ⁻¹ ; Organic farming: 7.49; Conventional farming: 16.06; Integrated farming: 12.38; Leguminous crops: 4.50; Potato: 19.44; Cereals: 11.16–15.69	Hillier et al. (2012)

EF Emission factor

^a Units of carbon footprints are kg CO₂-e kg⁻¹ unless stated otherwise

changes in soil carbon could turn the carbon footprints of wheat cultivation negative, which was otherwise $0.34 \text{ kg CO}_2\text{-e kg}^{-1}$ (Gan et al. 2012). On basis of this result, PAS 2050 is recommended to include changes in soil carbon in their upcoming guidelines for carbon footprint calculation for agricultural systems.

Practicing different permutations of conventional tillage and no tillage in rice–wheat systems showed that although no tillage led to significant reductions in cumulative GWP of rice cultivation under continuously no-tilled systems, it also reduced the yield; hence the yield scaled GWP was increased, resulting in a higher carbon footprint compared to the conventional practice. On the contrary, during wheat cultivation, the conventional practice acted as a net sink of CH_4 , thereby leaving a negative carbon footprint of -8.11 to $125.2 \text{ kg CO}_2\text{-e kg}^{-1}$. Under no-tillage practice, emission of GHGs increased along with yield; hence, the carbon footprint became positive (Pandey et al. 2013).

Food as a commodity has independently become an important candidate of carbon footprinting. Kim and Neff (2009) showed that carbon footprint calculators for food items had different scopes and calculations were based on different emission factors. Hence, they could not address effectively the diet-related preferences. Pathak et al. (2010) calculated the carbon footprints of Indian food items, taking into account cultivation of crops, processing, transportation, and kitchen preparations. The average emission factors they used did not address CH_4 emissions from non-rice crops, different management conditions, and changes in soil carbon. They calculated the average daily carbon footprint of $723.7 \text{ g CO}_2\text{-e}$ for an Indian adult male.

8 Sources of Uncertainty

According to the GHG protocol, sources of uncertainties should be mentioned when reporting carbon footprints. For agricultural practices, nonavailability of activity-specific emission factors is an important source of error. In addition, the associated land use changes and alternative scenarios under different agricultural practices are not easy to predict confidently. For example, in the case of no-tillage cultivation, the stubble left over the soil could have been used as cattle fodder. Loss of fodder under no-tillage practice might put pressure on the cattle rearing; hence, there may be requirements to arrange extra land to compensate for the fodder demand. Land use change to compensate for fodder demand and shifts in yield should be considered. Because agriculture is largely affected by the climate, long-term monitoring and calculation are required to generate better footprint estimates, and how it modulates with changes in different components. Although, soil carbon sequestration is regarded as a ‘win–win strategy,’ there are certain controversies over quantification and assessment of sink capacity reliability (Lehmann 2009). It is argued that sequestration must be able to keep the carbon out of the atmosphere for a relevant time period, conventionally assumed to be at least 100 years. Determination of the degree of permanence of sequestered carbon

has not yet been established convincingly; however, scientists have adopted fractionation of the carbon pool physically and chemically (Post et al. 2001; Rovira and Vellejo 2007).

9 Conclusions

Carbon footprinting has appeared as a strong and popular indicator of the GHG intensity of any activity or organization. Due to its important role in raising awareness regarding responsibility toward global warming, scientists and policy makers are trying to use it as a management tool. However, its application over the agricultural sector is still limited. A standard methodology is required to address the emissions associated with soil, carbon sequestration in soil, emissions associated with farm equipments, and other relevant activities. Due to widespread differences in agricultural activities over the world, it is essential to have guidelines on the selection of boundaries. In addition, there is also an immediate need for uniformities in GHG estimation techniques. The lack of sector- and region-specific emission factors for important agricultural inputs add to the uncertainty. The standard method must address how to deal with alternative scenarios and land use changes. The number of carbon footprinting studies of agricultural systems is increasing, but due to widespread differences, their comparison remains difficult. Nevertheless, such studies represent the contribution of cultivation practices in a better way than merely focusing on soilborne GHG emissions, carbon sequestration, or energy intensity individually.

Acknowledgments The authors thank the University Grants Commission (UGC), New Delhi for providing financial help in the form of a research project. Divya Pandey is grateful to Council for Scientific and Industrial Research, New Delhi for senior research fellowship and UGC for Junior Research Fellowship.

References

- Arnold S, Parkin TB, Doran JW, Mosier AR (2001) Automated gas sampling system for laboratory analysis of CH₄ and N₂O. *Commun Soil Sci Plant Anal* 32:2795–2807. doi:[10.1081/CSS-120000962](https://doi.org/10.1081/CSS-120000962)
- Browne D, O'Regan B, Moles R (2009) Use of carbon footprinting to explore alternative household waste policy scenarios in an Irish city-region. *Resour Conserv Recycling* 54:113–122
- Blengini GA, Busto M (2009) The life cycle of rice: LCA of alternative agri-food chain management systems in Vercelli (Italy). *J Environ Manag* 90:1512–1522
- BSI. (2008) Publicly Available Specification 2050. Specification for the assessment of the life cycle greenhouse gas emissions of goods and services. British Standards Institute
- Carbon Trust (2007) Carbon Footprint Measurement Methodology, Version 1.1. The Carbon Trust, London, UK

- Cassman KG, Dobermann A, Walters DT, Yang H (2003) Meeting cereal demand while protecting natural resources and improving environmental quality. *Ann Rev Environ Resour* 28:315–358
- CDP (2009) Carbon disclosure project report 2009: Global 500, On behalf of 385 investors with assets of \$57 trillion. Carbon Disclosure Project
- Chambers JQ, Fisher JJ, Zeng H, Chapman EL, Baker DB, Hurtt GC (2007) Hurricane Katrina's carbon footprint on U.S. gulf coast forests. *Science* 318:1107
- Courchene TJ, Allan J (2008) Climate change: The case of carbon tariff/tax. Policy options. March issue:59–64
- Craeynest L, Streatfeild D (2008) The World Bank and its carbon footprint: why the World Bank is still far from being an environment bank. *World Wildlife Fund*, pp. 35
- Del Grosso SJ, Wirt T, Ogle SM, Parton WJ (2008) Estimating agricultural nitrous oxide emissions. *Eos, Trans Am Geophys Union* 89:529–530
- East AJ (2008) What is a carbon footprint? An overview of definitions and methodologies. Vegetable industry carbon footprint scoping study—discussion papers and workshop, 26 September 2008. Horticulture Australia Limited, Sydney
- GAP, SEI, Eco-Logica (2006) UK Schools Carbon Footprint Scoping Study. Report by Global Action Plan, Stockholm Environment Institute and Eco-Logica Ltd. for the Sustainable Development Commission, London
- Edgar GH, Peters GP (2009) Carbon footprint of nations: a global, trade linked analysis. *Environ Sci Technol* 43:6414–6420
- Eshel G, Martin P (2006) Diet, energy and global warming, University of Chicago study. University of Chicago, Chicago
- Ferris L, Best J, Scholefield H, Marques J, Sawada B, Nemetz P (2007) Carbon neutrality and University of British Colombia: a first glance. University of British Colombia, Colombia
- Gan Y, Liang BC, Hamel C, Cutforth H, Wang H (2011a) Strategies for reducing the carbon footprint of field crops for semiarid areas. A review. *Agron Sust Dev* 31:643–656
- Gan Y, Liang L, Huang G, Malhi SS, Brandt SA, Katepa-Mupondwa F (2011b) Carbon footprint of canola and mustard is a function of the rate of N fertilizer. *Int J Life Cycle Assess* 17:58–68
- Gan YT, Liang BC, Wang XY, McConkey BG (2011c) Lowering carbon footprint of durum wheat through diversifying cropping systems. *Field Crops Res* 122:199–206
- Gan YT, Liang BC, May W, Malhi SS, Niu J, Wang X (2012) Carbon footprint of spring barley in relation to preceding oilseeds and N fertilization. *Int J Life Cycle Assess* 17:635–645
- Garg S, Dornfeld D (2008) An indigenous application for estimating carbon footprint of academia library based on life cycle assessment. University of California, Berkeley: Laboratory for manufacturing and sustainability
- Gemechu ED, Butnar I, Llop M, Castells F (2012) Environmental tax on products and services based on their carbon footprint: a case study of the pulp and paper sector. *Energy Policy* 50:336–344
- Global Footprint Network (2007) Ecological Footprint Glossary. Global Footprint Network, Oakland. <http://www.footprintnetwork.org>, Assessed November 02, 2008
- Chapuis-Lardy L, Wrange N, Metay, A, Chotte JL, Bernoux M (2007) Soils, a sink for N₂O? A review. *Glob Change Biol* 13:1–17
- Bokowski G, White D, Pacifico A, Talbot S, DuBelko A, Phipps A, Utley C, Mui, T, Lewis G. (2007) Towards campus climate neutrality. Simon Fraser University's Carbon Footprint. Simon Fraser University
- Good Company. (2008) The City of Vancouver, Washington: Corporate Greenhouse Gas Emissions Inventory For Calendar Years 2006 and 2007-Report and Analysis. Good Company, Oregon, USA
- Gurney KR, Mendoza DL, Zhou Y, Fischler ML, Miller CC, Geethkumar S (2009) Can S.D. L.R.D. High Resolution Fossil Fuel Combustion CO₂ Emission Fluxes for the United States. *Environ Sci technol* 43:5535–5541
- Haag A (2007) The crucial measurement. *Nature* 450:6
- Hammond G (2007) Time to give due weight to the 'carbon footprint' issue. *Nature* 445:256

- Hamilton K, Bayon R, Turner G, Higgins D (2007) State of the voluntary carbon markets 2007: picking up steam. *Ecosystem Marketplace and New Carbon Finance*
- Hoefnagels R, Smeets E, Faaij A (2010) Greenhouse gas footprints of different biofuel production systems. *Renew Sustain Energy Rev* 14:1661–1694
- Hutchinson JJ, Campbell CA, Desjardins RL (2007) Some perspectives on carbon sequestration in agriculture. *Agric Forest Meteorol* 142:288–302
- IPCC (2006) National Greenhouse gas inventories: Land use, land use change and forestry. Institute of global environmental strategies. Hayama, Japan
- IPCC (2007) Climate change: the physical science basis. In: (Eds) Solomon S, Qin D, Manning M, Chen Z, Marquis M, Averyt KB, Tignor M, Miller HL. Contribution of working group I to the fourth assessment report of the intergovernmental panel on climate change. Cambridge University Press, Cambridge, U.K
- ISO (2006a) ISO 14064-1:2006. Greenhouse gases Part 1: Specification with guidance at the organization level for quantification and reporting of greenhouse gas emissions and removals. <http://www.iso.org> (Assessed 09 December, 2008)
- ISO (2006b) ISO 14064-2:2006. Greenhouse gases Part 2: Specification with guidance at the project level for quantification, monitoring and reporting of greenhouse gas emission reductions or removal enhancements. <http://www.iso.org> (Assessed 05 December, 2009)
- ISO (2006c) ISO 14040: Environmental management-Life cycle assessment-Principles and framework. <http://www.iso.org> (Assessed 07 January, 2009)
- ISO (2006d) ISO 14044: Environmental management-Life cycle assessment-Requirements and guidelines. <http://www.iso.org> (Assessed 07 January, 2009)
- ISO (2012a) ISO/TR 14047: Environmental management-Life cycle assessment-Illustrative examples on how to apply ISO 14044 to impact assessment situations. <http://www.iso.org> (Assessed 10 June, 2013)
- ISO (2012b) ISO/TR 14049: Environmental management-Life cycle assessment-Illustrative examples on how to apply ISO 14044 to goal and scope definition and inventory analysis. (Assessed 10 June, 2013)
- ISO (2013) ISO/TS 14067: Greenhouse gases-Carbon footprint of products-Requirements and guidelines for quantification and communication. (Assessed 15 June, 2013)
- Jarvis P (2007) Never mind the footprints, get the mass right. *Nature* 446:24
- Kelly LM, Shepson PB, Strim BP, Karion A, Sweeney C, Gurney KR (2009) Aircraft-based measurements of the carbon footprint of Indianapolis. *Environ Sci Technol* 43:7816–7823
- Kenny T, Gray NF (2008) Comparative performance of six carbon footprint models for use in Ireland. *Environ Impact Assess Rev* 29:1–6
- Kim B, Neff R (2009) Measurement and communication of greenhouse gas emissions from U.S. food consumption via carbon calculators. *Ecol Econ* 69:186–196
- Kleiner K (2007) The corporate race to cut carbon. *Nature* 3:40–43
- Lal R (2000) World cropland soils as a source or sink for atmospheric carbon. *Adv Agron* 71:145–191
- Lal R (2004a) Carbon emission from farm operations. *Environ Int* 30:981–990
- Lal R (2004b) Soil carbon sequestration impacts on global climate change and food security. *Science* 304:1623–1627
- Le Mer JL, Roger P (2001) Production, oxidation, emission and consumption of methane by soils: a review. *Eur J Soil Biol* 37:25–50
- Lehmann J (2009) Biological carbon sequestration must and can be a win-win approach. *Clim Change* 97:459–463
- Lenzen M (2001) Errors in Conventional and Input-Output Based Life-Cycle Inventories. *J Indus Ecol* 4:127–148
- Linquist BA, Van Groenigen KJ, Adviento-Borbe MA, Pittelkow C, Van Kessel C (2011) An agronomic assessment of greenhouse gas emissions from major cereal crops. *Glob Change Biol* 18:194–209

- Linquist BA, Adviento-Borbe MA, Pittelkow CM, Kessel C, Groenigen KJ (2012) Fertilizer management practices and greenhouse gas emissions from rice systems: a quantitative review and analysis. *Field Crop Res* 135:10–21
- Liu C, Zheng X, Zhou Z, Han S, Wang Y, Wang K, Liang W, Li M, Chen D, Yang Z (2010) Nitrous oxide and nitric oxide emissions from an irrigated cotton field in Northern China. *Plant Soil* 332:123–134
- London-2012 Sustainability Plan (2007) Towards a one planet. London 2012 organizing committee, London, pp 63
- Lynas M (2007) Carbon counter. Harper Collins, Glasgow
- Matthews SC, Hendrickson C, Weber CL (2008) The importance of carbon footprint estimation boundaries. *Environ Sci Technol* 42:5839–5842
- Ma J, Li XL, Xu H, Han Y, Cai ZC, Yagi K (2007) Effects of nitrogen fertilizer and wheat straw application on CH₄ and N₂O emissions from a paddy rice field. *Aust J Soil Res* 45:359–367
- Ma YC, Kong XW, Yang B, Zhang XL, Yan XY, Yang JC, Xiong ZQ (2013) Net global warming potential and greenhouse gas intensity of annual rice–wheat rotations with integrated soil–crop system management. *Agric Ecosys Environ* 164:209–219
- Majumder B, Mandal B, Bandyopadhyay PK, Gangopadhyay A, Mani PK, Kundu AL, Majumdar D (2008) Organic amendments influence soil organic carbon pools and crop productivity in a 19 years old rice–wheat agroecosystem. *Soil Sci Soc Am J* 72:775–785
- Mandal B, Majumder B, Bandyopadhyay PK, Hazra GC, Gangopadhyay A, Samantaray RN, Mishra A, Chaudhury J, Saha MN, Kundu S (2007) The potential of cropping systems and soil amendments for carbon sequestration in soils under long-term experiments in subtropical India. *Glob Change Biol* 13:357–369
- Mandal B, Majumder B, Adhya TK, Bandyopadhyay PK, Gangopadhyay A, Sarkar D, Kundu MC, Gupta Choudhury S, Hazra GC, Kundu S, Samantaray RN, Mishra AK (2008) Potential of double-cropped rice ecology to conserve organic carbon under subtropical climate. *Glob Change Biol* 14:2139–2151
- Montzka SA, Dlugokencky EJ, Butler JH (2011) Non-CO₂ greenhouse gases and climate change. *Nature* 476:46–50
- Miyoshi C, Mason KJ (2009) *J Air Transp Manage* 15:138–147
- Sundareshwar PV, Murtugudde R, Srinivasan G, Singh J, Rajmeh JK, Ramesh R, Verma SB, Agrawal D, Baldocchi D, Baru CK, Baruah KK, Chowdhury GR, Dadhwal VK, Dutt CBS, Fuentes J, Gupta PK, Hargrove WW, Howard M, Jha CS, Lal S, Michener WK, Mitra AP, Morris JT, Myneni RR, Naja M, Nemani R, Purvaja, R, Raha S, Santhana Vanan SK, Sharma M, Subramaniam A, Sukumar R, Twilley RR, Zimmerman PR (2007) Environmental monitoring network for India. *Science*, 316:204–205
- Murray J, Day C (2009) The carbon neutral free for all. *Int J GHG Control* 3:237–248
- Nelson R, Hellwinckel CC, Brandt TO, West DG, De La Torre U, Marland G (2009) Energy use and carbon dioxide emissions from cropland production in the United States, 1990–2004. *J Environ Qual* 38:418–425
- Nishimura S, Yonemura S, Sawamoto S, Shirato Y, Akiyama H, Sudo S, Yagi K (2008) Effect of land use change from paddy rice cultivation to upland crop cultivation on soil carbon budget of a cropland in Japan. *Agric Ecosys Environ* 125:9–20
- Padgett JP, Steinemann AC, Clarke JH, Vandenberg MP (2008) A comparison of carbon calculators. *EIA Rev* 28:106–115
- Pandey D, Agrawal M, Pandey JS (2011) Carbon footprints: current methods of estimation. *Environ Monit Assess* 178:135–160
- Pandey D, Agrawal M, Bohra JS (2012) Greenhouse gas emissions from rice crop with different tillage permutations in rice-wheat system. *Agric Ecosyst Environ* 159:133–144
- Pandey D, Agrawal M, Bohra JS (2013) Impact of four tillage permutations in rice-wheat system on GHG performance of wheat cultivation through carbon footprinting. *Ecol Eng* (accepted). doi:10.1016/j.ecoleng.2013.07.020
- Parkin TB, Venterea RT, Hargreaves SK (2012) Calculating the detection limits of chamber-based soil greenhouse gas flux measurements. *J Environ Qual* 41:705–715

- Patel J (2006) Green sky thinking. *Environ Bus* 122:32
- Pathak H, Prasad S, Bhatia A, Singh S, Kumar S, Singh J, Jain MC (2003) Methane emission from rice-wheat cropping system of India in relation to irrigation, farmyard manure and dicyandiamide application. *Agric Ecosyst Environ* 97:309–316
- Pathak H, Jain N, Bhatia A, Patel J, Aggarwal PK (2010) Carbon footprints of Indian food items. *Agric Ecosyst Environ* 139:66–73
- Peters GP (2010) Carbon footprints and embodied carbon at multiple scales. *Curr Opin Environ Sustain* 2:245–250
- Post WM, Izaurralde RC, Mann LK, Bliss N (2001) Monitoring and verifying changes of organic carbon in soil. *Clim Change* 51:73–99
- Rochette R (2008) No-till only increases N₂O emissions in poorly-aerated soils. *Soil Till Res* 101:97–100
- Roos E, Sundberg C, Hansson PA (2010) Uncertainties in the carbon footprint of food products: a case study on table potatoes. *Int J Life Cycle Assess* 15:478–488
- Rovira P, Vallejo VR (2007) Labile, recalcitrant, and inert organic matter in Mediterranean forest soils. *Soil Biol Biochem* 39:202–215
- Smith P, Andren O, Karlsson T, Peñala P, Regina K, Rounsevell M, Wesemael BV (2005) Carbon sequestration potential in European croplands has been overestimated. *Glob Change Biol* 11:2153–2163
- Smith P, Martino D, Cai Z, Gwary D, Janzen H, Kumar P, McCarl B, Ogle S, O'Mara F, Rice C, Scholes B, Sirotenko O, Howden M, McAllister T, Pan G, Romanenkov, V, Schneider U, Towprayoon S, Wattenbach M, Smith P (2008) J. Greenhouse gas mitigation in agriculture. *Phil Trans R Soc B* 363:789–813
- Seip HM (2011) Carbon footprints: relative drop for farming emissions. *Nature* 480:321
- Stern N (2006) *Stern review: the economics of climate change*. Cambridge University Press, Cambridge
- Suh S (2006) Are services better for climate change? *Environ Sci Technol* 40:6555–6560
- T C Chan Center for Building Simulation and Energy Studies/Penn Praxis (2007). University of Pennsylvania carbon footprint. Pennsylvania: University of Pennsylvania
- Tirado R, Gopikrishna SR, Krishnan R, Smith P (2010) Greenhouse gas emissions and mitigation potential from fertilizer manufacture and application in India. *Int J Agric Sustain* 8:176–185
- UNDP (2007) Human development report 2007/2008: fighting climate change: human solidarity in a divided world. Palgrave, Macmillan, New York
- USCCTP (2009) Technology options for near and long term future. <http://www.climatechange.gov/library/2005/tech-options/index.htm>. (Assessed on 02 January, 2009)
- Van Groenigen KJ, Osenberg CW, Hungate BA (2011) Increased soil emissions of potent greenhouse gases under increased atmospheric CO₂. *Nature* 475:214–216
- Wackernagel M, Rees WE. (1996) *Our Ecological footprint: reducing human impact on the earth*. New Society Publishers, Gabriola Island, p 160
- Weidema BP, Thrane M, Christensen P, Schmidt J, Løkke S (2008) Carbon footprint: a catalyst for life cycle assessment. *J Ind Ecol* 12:3–6
- Wiedmann T, Minx JA (2007) Definition of Carbon Footprint. ISAUK Research Report 07-01, ISAUK Research & Consulting, Durham
- WRI/WBCSD (2004) *The Greenhouse Gas Protocol: A Corporate Accounting and Reporting Standard Revised Edition*. World Business Council for Sustainable Development and World Resource Institute, Geneva
- Zou JW, Huang Y, Jiang JY, Zheng XH, Sass RL (2005) A 3-year field measurement of methane and nitrous oxide emissions from rice paddies in China: effects of water regime, crop residue, and fertilizer application. *Glob Biogeochem Cycles* 19:GB2021

Assessment of Carbon Footprint in Different Industrial
Sectors, Volume 1

Muthu, S.S. (Ed.)

2014, VIII, 292 p. 89 illus., 74 illus. in color., Hardcover

ISBN: 978-981-4560-40-5