

Undergraduate Texts in Mathematics

Undergraduate Texts in Mathematics

Series Editors:

Sheldon Axler

San Francisco State University, San Francisco, CA, USA

Kenneth Ribet

University of California, Berkeley, CA, USA

Advisory Board:

Colin Adams, *Williams College*

David A. Cox, *Amherst College*

Pamela Gorkin, *Bucknell University*

Roger E. Howe, *Yale University*

Michael Orrison, *Harvey Mudd College*

Lisette G. de Pillis, *Harvey Mudd College*

Jill Pipher, *Brown University*

Fadil Santosa, *University of Minnesota*

Undergraduate Texts in Mathematics are generally aimed at third- and fourth-year undergraduate mathematics students at North American universities. These texts strive to provide students and teachers with new perspectives and novel approaches. The books include motivation that guides the reader to an appreciation of interrelations among different aspects of the subject. They feature examples that illustrate key concepts as well as exercises that strengthen understanding.


More information about this series at <http://www.springer.com/series/666>

Matthias Beck • Sinai Robins

Computing the Continuous Discretely

Integer-Point Enumeration in Polyhedra

Second Edition



With illustrations by David Austin



Springer

Matthias Beck
Department of Mathematics
San Francisco State University
San Francisco, CA, USA

Sinai Robins
Department of Mathematics
Brown University
Providence, RI, USA

ISSN 0172-6056 ISSN 2197-5604 (electronic)
Undergraduate Texts in Mathematics
ISBN 978-1-4939-2968-9 ISBN 978-1-4939-2969-6 (eBook)
DOI 10.1007/978-1-4939-2969-6

Library of Congress Control Number: 2015942640

Springer New York Heidelberg Dordrecht London
© Matthias Beck and Sinai Robins 2007, 2015

This work is subject to copyright. All rights are reserved by the Publisher, whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, reuse of illustrations, recitation, broadcasting, reproduction on microfilms or in any other physical way, and transmission or information storage and retrieval, electronic adaptation, computer software, or by similar or dissimilar methodology now known or hereafter developed.

The use of general descriptive names, registered names, trademarks, service marks, etc. in this publication does not imply, even in the absence of a specific statement, that such names are exempt from the relevant protective laws and regulations and therefore free for general use.

The publisher, the authors and the editors are safe to assume that the advice and information in this book are believed to be true and accurate at the date of publication. Neither the publisher nor the authors or the editors give a warranty, express or implied, with respect to the material contained herein or for any errors or omissions that may have been made.

Printed on acid-free paper

Springer Science+Business Media LLC New York is part of Springer Science+Business Media (www.springer.com)

To Tendai

To my mom, Michal Robins

with all our love

Preface

The world is continuous, but the mind is discrete.

David Mumford

The mathematical interplay between polytopes and lattices comes to life when we study the relationships between the continuous volume of a polytope and its discrete volume. Since the humbling and positive reception of the first edition of this book, published in 2007, the field of integer-point enumeration in polyhedra has gained considerable momentum. Many fields of mathematics have begun to interact in even more surprising ways, and a beautifully simple unification among a multitude of classical problems continues to emerge when we use this combinatorial-geometric lens.

In this second edition, and with encouragement from many wonderful readers, we have added two new chapters: Chapter 9 introduces zonotopes, an extremely useful class of polytopes, and Chapter 10 explores some deep yet elegant relationships that are satisfied by the h -polynomials and h^* -polynomials. We have also added many new exercises, new and updated open problems, and graphics.

Examples of polytopes in three dimensions include crystals, boxes, tetrahedra, and any convex object whose faces are all flat. It is amusing to see how many problems in combinatorics, number theory, and many other mathematical areas can be recast in the language of polytopes that exist in some Euclidean space. Conversely, the versatile structure of polytopes gives us number-theoretic and combinatorial information that flows naturally from their geometry.

The *discrete volume* of a body \mathcal{P} can be described intuitively as the number of grid points that lie inside \mathcal{P} , given a fixed grid in Euclidean space. The *continuous volume* of \mathcal{P} has the usual intuitive meaning of volume that we attach to everyday objects we see in the real world.




Fig. 0.1 Continuous and discrete volume.

Indeed, the difference between the two realizations of volume can be thought of in physical terms as follows. On the one hand, the quantum-level grid imposed by the molecular structure of reality gives us a discrete notion of space and hence discrete volume. On the other hand, the Newtonian notion of continuous space gives us the continuous volume. We see things continuously at the Newtonian level, but in practice, we often compute things discretely at the quantum level. Mathematically, the grid we impose in space—corresponding to the grid formed by the atoms that make up an object—helps us compute the usual continuous volume in very surprising and charming ways, as we shall discover.

In order to see the continuous/discrete interplay come to life among the three fields of combinatorics, number theory, and geometry, we begin our focus with the simple-to-state *coin-exchange problem* of Frobenius. The beauty of this concrete problem is that it is easy to grasp, it provides a useful computational tool, and yet it has most of the ingredients of the deeper theories that are developed here.

In the first chapter, we give detailed formulas that arise naturally from the Frobenius coin-exchange problem in order to demonstrate the interconnections between the three fields mentioned above. The coin-exchange problem provides a scaffold for identifying the connections between these fields. In the ensuing chapters, we shed this scaffolding and focus on the interconnections themselves:

- (1) Enumeration of integer points in polyhedra—combinatorics,
- (2) Dedekind sums and finite Fourier series—number theory,
- (3) Polygons and polytopes—geometry.

We place a strong emphasis on computational techniques and on computing volumes by counting integer points using various old and new ideas. Thus, the formulas we get should not only be pretty (which they are!) but also allow us to compute volumes efficiently using some nice functions. In the very rare instances of mathematical exposition when we have a formulation that is both “easy to write” and “quickly computable,” we have found a mathematical nugget. We have endeavored to fill this book with such mathematical nuggets.

Much of the material in this book is developed by the reader in the more than three hundred exercises. Most chapters contain warmup exercises that

do not depend on the material in the chapter and can be assigned before the chapter is read. Some exercises are central, in the sense that current or later material depends on them. Those exercises are marked with ♣, and we give detailed hints for them at the end of the book. Most chapters also contain lists of open research problems.

It turns out that even a fifth grader can write an interesting paper on integer-point enumeration [192], while the subject lends itself to deep investigations that attract the current efforts of leading researchers. Thus, it is an area of mathematics that attracts our innocent childhood questions as well as our refined insight and deeper curiosity. The level of study is highly appropriate for a junior/senior undergraduate course in mathematics. In fact, this book is ideally suited to be used for a *capstone course*. Because the three topics outlined above lend themselves to more sophisticated exploration, our book has also been used effectively for an introductory graduate course.

To help the reader fully appreciate the scope of the connections between continuous volume and discrete volume, we begin the discourse in two dimensions, where we can easily draw pictures and quickly experiment. We gently introduce the functions we need in higher dimensions (Dedekind sums) by looking at the coin-exchange problem geometrically as the discrete volume of a generalized triangle, called a simplex.

The initial techniques are quite simple, essentially nothing more than expanding rational functions into partial fractions. Thus, the book is easily accessible to a student who has completed a standard college calculus and linear algebra curriculum. It would be useful to have a basic understanding of partial fraction expansions, infinite series, open and closed sets in \mathbb{R}^d , complex numbers (in particular, roots of unity), and modular arithmetic.

An important computational tool that is harnessed throughout the text is the *generating function* $f(x) = \sum_{m=0}^{\infty} a(m) x^m$, where the $a(m)$'s form a sequence of numbers that we are interested in analyzing. When the infinite sequence of numbers $a(m)$, $m = 0, 1, 2, \dots$, is embedded into a single generating function $f(x)$, it is often true that for hitherto unforeseen reasons, we can rewrite the whole sum $f(x)$ in a surprisingly compact form. It is the rewriting of these generating functions that allows us to understand the combinatorics of the relevant sequence $a(m)$. For us, the sequence of numbers might be the number of ways to partition an integer into given coin denominations, or the number of points in an increasingly large body, and so on. Here we find yet another example of the interplay between the discrete and the continuous: we are given a *discrete* set of numbers $a(m)$, and we then carry out analysis on the generating function $f(x)$ in the *continuous* variable x .

What Is the Discrete Volume?

The physically intuitive description of the discrete volume given above rests on a sound mathematical footing as soon as we introduce the notion of a

lattice. The grid is captured mathematically as the collection of all integer points in Euclidean space, namely $\mathbb{Z}^d = \{(x_1, \dots, x_d) : \text{all } x_k \in \mathbb{Z}\}$. This discrete collection of equally spaced points is called a *lattice*. If we are given a geometric body $\mathcal{P} \subset \mathbb{R}^d$, its discrete volume is simply defined as the number of lattice points inside \mathcal{P} , that is, the number of elements in the set $\mathbb{Z}^d \cap \mathcal{P}$.

Intuitively, if we shrink the lattice by a factor k and count the number of newly shrunk lattice points inside \mathcal{P} , we obtain a better approximation for the volume of \mathcal{P} , relative to the volume of a single cell of the shrunk lattice. It turns out that after the lattice is shrunk by an integer factor k , the number $\#(\mathcal{P} \cap \frac{1}{k}\mathbb{Z}^d)$ of shrunk lattice points inside an *integral polytope* \mathcal{P} is magically a polynomial in k . This counting function $\#(\mathcal{P} \cap \frac{1}{k}\mathbb{Z}^d)$ is known as the *Ehrhart polynomial* of \mathcal{P} . If we kept shrinking the lattice by taking a limit, we would of course end up with the continuous volume that is given by the usual Riemannian integral definition of calculus:

$$\text{vol } \mathcal{P} = \lim_{k \rightarrow \infty} \# \left(\mathcal{P} \cap \frac{1}{k}\mathbb{Z}^d \right) \frac{1}{k^d}.$$

However, pausing at fixed dilations of the lattice gives surprising flexibility for the computation of the volume of \mathcal{P} and for the number of lattice points that are contained in \mathcal{P} .

Thus, when the body \mathcal{P} is an integral polytope, the error terms that measure the discrepancy between the discrete volume and the usual continuous volume are quite nice; they are given by Ehrhart polynomials, and these enumeration polynomials are the content of Chapter 3.

The Fourier–Dedekind Sums Are the Building Blocks: Number Theory

Every polytope has a discrete volume that is expressible in terms of certain finite sums that are known as *Dedekind sums*. Before giving their definition, we first motivate these sums with some examples that illustrate their building-block behavior for lattice-point enumeration. To be concrete, consider, for example, a 1-dimensional polytope given by an interval $\mathcal{P} = [0, a]$, where a is any positive real number. It is clear that we need the greatest integer function $\lfloor x \rfloor$ to help us enumerate the lattice points in \mathcal{P} , and indeed, the answer is $\lfloor a \rfloor + 1$.

Next, consider a 1-dimensional line segment that is sitting in the 2-dimensional plane. Let's choose our segment \mathcal{P} so that it begins at the origin and ends at the lattice point (c, d) . As becomes apparent after a moment's thought, the number of lattice points on this finite line segment involves an old friend, namely the greatest common divisor of c and d . The exact number of lattice points on the line segment is $\gcd(c, d) + 1$.

To unify both of these examples, consider a triangle \mathcal{P} in the plane whose vertices have rational coordinates. It turns out that a certain finite sum is completely natural because it simultaneously extends both the greatest integer function and the greatest common divisor, although the latter is less obvious. An example of a Dedekind sum in two dimensions that arises naturally in the formula for the discrete volume of the rational triangle \mathcal{P} is the following:

$$s(a, b) = \sum_{m=1}^{b-1} \left(\frac{m}{b} - \frac{1}{2} \right) \left(\frac{ma}{b} - \left\lfloor \frac{ma}{b} \right\rfloor - \frac{1}{2} \right).$$

The definition makes use of the greatest integer function. Why do these sums also resemble the greatest common divisor? Luckily, the Dedekind sums satisfy a remarkable reciprocity law, quite similar to the Euclidean algorithm that computes the greatest common divisor. This reciprocity law allows the Dedekind sums to be computed in roughly $\log(b)$ steps rather than the b steps that are implied by the definition above. The reciprocity law for $s(a, b)$ lies at the heart of some amazing number theory that we treat in an elementary fashion, but that also comes from the deeper subject of modular forms and other modern tools.

We find ourselves in the fortunate position of viewing an important summit of an enormous mountain of ideas, submerged by the waters of geometry. As we delve more deeply into these waters, more and more hidden beauty unfolds for us, and the Dedekind sums are an indispensable tool that allow us to see farther as the waters get deeper.

The Relevant Solids Are Polytopes: Geometry

The examples we have used, namely line segments and polygons in the plane, are special cases of polytopes in all dimensions. One way to define a polytope is to consider the *convex hull* of a finite collection of points in Euclidean space \mathbb{R}^d . That is, suppose someone gives us a set of points $\mathbf{v}_1, \dots, \mathbf{v}_n$ in \mathbb{R}^d . The polytope determined by the given points \mathbf{v}_j is defined by all linear combinations $c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \dots + c_n\mathbf{v}_n$, where the coefficients c_j are nonnegative real numbers that satisfy the relation $c_1 + c_2 + \dots + c_n = 1$. This construction is called the *vertex description* of the polytope.

There is another equivalent definition, called the *hyperplane description* of the polytope. Namely, if someone hands us the linear inequalities that define a finite collection of half-spaces in \mathbb{R}^d , we can define the associated polytope as the simultaneous intersection of the half-spaces defined by the given inequalities.

There are some “obvious” facts about polytopes that are intuitively clear to most students but are, in fact, subtle and often nontrivial to prove from first principles. One of these facts, namely that every polytope has both a vertex and a hyperplane description, forms a crucial basis to the material we

will develop in this book. We carefully prove this fact in the appendix. The statement is intuitively clear, so that novices can skip over its proof without any detriment to their ability to compute continuous and discrete volumes of polytopes. All theorems in the text (including those in the appendix) are proved from first principles, with the exception of Chapter 14, where we assume basic notions from complex analysis.




Fig. 0.2 The partially ordered set of chapter dependencies

The text naturally flows into two parts, which we now explicate.

Part I

We have taken great care in making the content of the chapters flow seamlessly from one to the next, over the span of the first six chapters.

- Chapters 1 and 2 introduce some basic notions of generating functions, in the visually compelling context of discrete geometry, with an abundance of detailed motivating examples.
- Chapters 3, 4, and 5 develop the full Ehrhart theory of discrete volumes of rational polytopes.
- Chapter 6 is a “dessert” chapter, in that it enables us to use the theory developed to treat the enumeration of *magic squares*, an ancient topic that enjoys active current research.

Part II

We now begin anew.

- Having attained experience with numerous examples and results about integral polytopes, we are ready to learn about the *Dedekind sums* of Chapter 8, which form the atomic pieces of the discrete-volume polynomials. On the other hand, to fully understand Dedekind sums, we need to understand *finite Fourier analysis*, which we therefore develop from first principles in Chapter 7, using only partial fractions.
- In Chapter 9, we study a concrete class of polytopes—projections of cubes, which go by the name *zonotopes*—whose discrete volume is tractable and has neat connections to number theory and graph theory.
- Chapter 10 develops inequalities among the coefficients of an Ehrhart polynomial, based on a *polynomial decomposition* formula that arises naturally from the arithmetic and combinatorial data of triangulations.
- Chapter 11 answers a simple yet tricky question: how does a finite geometric series in one dimension extend to higher-dimensional polytopes? *Brion’s theorem* gives an elegant and decisive answer to this question.
- In Chapter 12, we extend the interplay between the continuous volume and the discrete volume of a polytope (already studied in detail in Part I) by introducing *Euler–Maclaurin summation* formulas in all dimensions. These formulas compare the continuous Fourier transform of a polytope to its discrete Fourier transform, yet the material is completely self-contained.
- Chapter 13 develops an exciting extension of Ehrhart theory that defines and studies the *solid angles* of a polytope; these are the natural extensions of 2-dimensional angles to higher dimensions.
- Finally, we end with another “dessert” chapter that uses complex-analytic methods to find an integral formula for the discrepancy between the discrete and continuous areas enclosed by a closed curve in the plane.

Because polytopes are both theoretically useful (in triangulated manifolds, for example) and practically essential (in computer graphics, for example) we use them to link results in number theory and combinatorics. Many research papers have been written on these interconnections, and it is impossible to capture them all here, especially since some are being written even as we are writing this sentence! However, we hope that these modest beginnings will give the reader who is unfamiliar with these fields a good sense of their beauty, inexorable connectedness, and utility. We have written a gentle invitation to what we consider a gorgeous world of counting and of links between the fields of combinatorics, number theory, and geometry for the general mathematical reader.

There are a number of excellent books that have a nontrivial intersection with ours and contain material that complements the topics discussed here. We heartily recommend the monographs of Barvinok [20, 21] (on general convexity topics and Ehrhart theory), Bruns–Gubeladze [73] (on commutative algebra and K -theory connected with polytopes), De Loera–Hemmecke–Köppe [95] (on connections to optimization), De Loera–Rambau–Santos [98] (on triangulations of point configurations), Ehrhart [111] (the historic introduction to Ehrhart theory), Ewald [112] (on connections to algebraic geometry), Hibi [134] (on the interplay of algebraic combinatorics with polytopes), Miller–Sturmfels [177] (on combinatorial commutative algebra), and Stanley [230] (on general enumerative problems in combinatorics).

Acknowledgments

We have had the good fortune of receiving help from many gracious people in the process of writing this book. First and foremost, we thank the students of the classes in which we could try out this material, at Binghamton University (SUNY), Nanyang Technological University, San Francisco State University, and Temple University. We are indebted to our MSRI/Banff 2005 graduate summer school students. We give special thanks to Kristin Camenga and Kevin Woods, who ran the problem sessions for this summer school, detected numerous typos, and provided us with many interesting suggestions for this book. We are grateful for the generous support for the summer school from the Mathematical Sciences Research Institute, the Pacific Institute of Mathematics, and the Banff International Research Station.

Many colleagues have supported this endeavor, and we are particularly grateful to everyone notifying us of mistakes and typos and to those who have made useful suggestions: Daniel Antonetti, Alexander Barvinok, Nathanael Berglund, Andrew Beyer, Tristram Bogart, Garry Bowlin, Benjamin Braun, Swee Hong Chan, Robin Chapman, Yitwah Cheung, Jessica Cuomo, Dimitros Dais, Jesus De Loera, David Desario, Mike Develin, Ricardo Diaz, Michael Dobbins, Jeff Doker, Han Minh Duong, Richard Ehrenborg, David Einstein, Joseph Gubeladze, Christian Haase, Mary Halloran, Friedrich Hirzebruch,

Brian Hopkins, Serkan Hosten, Benjamin Howard, Piotr Maciak, Evgeny Materov (Chapter 5 rocks!), Asia Matthews, Peter McMullen, Martín Mereb, Ezra Miller, Mel Nathanson, Julian Pfeifle, Peter Pleasants, Jorge Ramírez Alfonsín, Bruce Reznick, Adrian Riskin, Junro Sato, Melissa Simmons, Richard Stanley, Bernd Sturmfels, Thorsten Theobald, Read Vanderbilt, Andrew Van Herick, Sven Verdoolaege, Michèle Vergne, Julie Von Bergen, Neil Weickel, Carl Woll, Zhiqiang Xu, Jon Yaggie, Ruriko Yoshida, Thomas Zaslavsky, Günter Ziegler, and two anonymous referees.

We are indebted to the Springer editorial staff, first and foremost to Mark Spencer, for facilitating the publication process in an always friendly and supportive spirit. We thank David Kramer for the impeccable copyediting, Frank Ganz for sharing his L^AT_EX expertise, and Felix Portnoy for the seamless production process. The Springer editors in Germany and Japan kindly suggested translations of our book, and we are particularly grateful to Kord Eickmeyer and Yoshio Okamoto for taking on the monumental task of translating this work.

Many colleagues have sent us suggestions and updates for this second edition; in addition to the ones listed above, we thank Ana Berrizbeitia, Steven Collazos, Aaron Dall, Maryam Farahmand, Katharina Jochemko, Michael Joswig, Victor Katsnelson, Jim Lawrence, Nhat Le Quang, Matthias Lenz, Pieter Moree, Bruce Sagan, Steven Sam, Raman Sanyal, Kim Seashore, and Ed Swartz. Special thanks go to Philippe Clauss for permission to include Eugène Ehrhart's self-portrait.

Perhaps the most visible change can be seen in the figures, whose new and much-improved looks owe their beauty to David Austin. We are huge fans of his work and thank him for his time and energy on this project.

We thank Eugene Ha for facilitating the publication process of this second edition. We will collect further corrections, updates, etc., at the Internet website math.sfsu.edu/beck/ccd.html.

Matthias Beck's work has been partially supported by the United States National Science Foundation (DMS-0810105 & DMS-1162638). Sinai Robins's work has been partially supported by the Singapore Ministry of Education (Tier 2 Research Grant MOE2011-T2-1-090), by the Institute for Constructive and Experimental Mathematics (ICERM), and by Brown University.

Matthias Beck would like to express his deepest gratitude to Tendai Chitewere for her patience, support, and unconditional love. He thanks his family for always being there for him. Sinai Robins would like to thank his mother, Michal Robins, and his brothers, Shani Robins and Gabriel Robins, for their relentless support and understanding during the completion of this project. We both thank all the cafés we have inhabited over the past 14 years for enabling us to turn their coffee into theorems.

San Francisco, CA
 Providence, RI
 September 2015

Matthias Beck
 Sinai Robins

Contents

Part I The Essentials of Discrete Volume Computations

1	The Coin-Exchange Problem of Frobenius	3
1.1	Why Use Generating Functions?	3
1.2	Two Coins	5
1.3	Partial Fractions and a Surprising Formula	8
1.4	Sylvester's Result	12
1.5	Three and More Coins	14
	Notes	16
	Exercises	19
	Open Problems	26
2	A Gallery of Discrete Volumes	27
2.1	The Language of Polytopes	27
2.2	The Unit Cube	29
2.3	The Standard Simplex	31
2.4	The Bernoulli Polynomials as Lattice-Point Enumerators of Pyramids	34
2.5	The Lattice-Point Enumerators of the Cross-Polytopes	38
2.6	Pick's Theorem	40
2.7	Polygons with Rational Vertices	43
2.8	Euler's Generating Function for General Rational Polytopes	48
	Notes	51
	Exercises	53
	Open Problems	58
3	Counting Lattice Points in Polytopes: The Ehrhart Theory	59
3.1	Triangulations	59
3.2	Cones	62
3.3	Integer-Point Transforms for Rational Cones	64
3.4	Expanding and Counting Using Ehrhart's Original Approach	68

3.5	The Ehrhart Series of an Integral Polytope	71
3.6	From the Discrete to the Continuous Volume of a Polytope ..	76
3.7	Interpolation	78
3.8	Rational Polytopes and Ehrhart Quasipolynomials	80
3.9	Reflections on the Coin-Exchange Problem and the Gallery of Chapter 2	81
	Notes	81
	Exercises	82
	Open Problems	88
4	Reciprocity	89
4.1	Generating Functions for Somewhat Irrational Cones	90
4.2	Stanley's Reciprocity Theorem for Rational Cones	92
4.3	Ehrhart–Macdonald Reciprocity for Rational Polytopes	93
4.4	The Ehrhart Series of Reflexive Polytopes	94
4.5	More “Reflections” on the Coin-Exchange Problem and the Gallery of Chapter 2	96
	Notes	96
	Exercises	98
	Open Problems	100
5	Face Numbers and the Dehn–Sommerville Relations in Ehrhartian Terms	101
5.1	Face It!	101
5.2	Dehn–Sommerville Extended	103
5.3	Applications to the Coefficients of an Ehrhart Polynomial ...	105
5.4	Relative Volume	106
	Notes	108
	Exercises	109
6	Magic Squares	113
6.1	It's a Kind of Magic	114
6.2	Semimagic Squares: Points in the Birkhoff–von Neumann Polytope	115
6.3	Magic Generating Functions and Constant-Term Identities ...	119
6.4	The Enumeration of Magic Squares	124
	Notes	125
	Exercises	127
	Open Problems	128

Part II Beyond the Basics

7	Finite Fourier Analysis	133
7.1	A Motivating Example	133
7.2	Finite Fourier Series for Periodic Functions on \mathbb{Z}	135
7.3	The Finite Fourier Transform and Its Properties	139
7.4	The Parseval Identity	141
7.5	The Convolution of Finite Fourier Series	143
	Notes	145
	Exercises	146
8	Dedekind Sums, the Building Blocks of Lattice-Point Enumeration	149
8.1	Fourier–Dedekind Sums and the Coin-Exchange Problem Revisited	149
8.2	The Dedekind Sum and Its Reciprocity and Computational Complexity	153
8.3	Rademacher Reciprocity for the Fourier–Dedekind Sum	154
8.4	The Mordell–Pommersheim Tetrahedron	158
	Notes	160
	Exercises	161
	Open Problems	164
9	Zonotopes	167
9.1	Definitions and Examples	167
9.2	Paving a Zonotope	170
9.3	The Permutohedron	172
9.4	The Ehrhart Polynomial of a Zonotope	176
	Notes	178
	Exercises	179
	Open Problems	181
10	h-Polynomials and h^*-Polynomials	183
10.1	Simplicial Polytopes and (Unimodular) Triangulations	183
10.2	Fundamental Parallelepipeds Open Up, with an h -Twist	187
10.3	Palindromic Decompositions of h^* -Polynomials	189
10.4	Inequalities for h^* -Polynomials	191
	Notes	192
	Exercises	193
	Open Problems	196
11	The Decomposition of a Polytope into Its Cones	199
11.1	The Identity “ $\sum_{m \in \mathbb{Z}} z^m = 0$ ” ... Or “Much Ado About Nothing”	199
11.2	Technically Speaking...	201
11.3	Tangent Cones and Their Rational Generating Functions	204
11.4	Brion’s Theorem	205

11.5 Brion Implies Ehrhart	207
Notes	209
Exercises	210
12 Euler–Maclaurin Summation in \mathbb{R}^d	213
12.1 Todd Operators and Bernoulli Numbers	214
12.2 A Continuous Version of Brion’s Theorem	216
12.3 Polytopes Have Their Moments	218
12.4 Computing the Discrete Continuously	221
Notes	223
Exercises	224
Open Problems	225
13 Solid Angles	227
13.1 A New Discrete Volume Using Solid Angles	227
13.2 Solid-Angle Generating Functions and a Brion-Type Theorem	231
13.3 Solid-Angle Reciprocity and the Brianchon–Gram Relations	232
13.4 The Generating Function of Macdonald’s Solid-Angle Polynomials	236
Notes	237
Exercises	238
Open Problems	239
14 A Discrete Version of Green’s Theorem Using Elliptic Functions	241
14.1 The Residue Theorem	241
14.2 The Weierstraß \wp - and ζ -Functions	243
14.3 A Contour-Integral Extension of Pick’s Theorem	245
Notes	246
Exercises	247
Open Problems	248
Appendix: Vertex and Hyperplane Descriptions of Polytopes	249
A.1 Every h-Cone Is a v-Cone	250
A.2 Every v-Cone Is an h-Cone	252
Hints for ♣ Exercises	255
References	267
List of Symbols	279
Index	281

Computing the Continuous Discretely
Integer-Point Enumeration in Polyhedra

Beck, M.; Robins, S.

2015, XX, 285 p., Hardcover

ISBN: 978-1-4939-2968-9