

# The Neonatal Neuromechanical Unit: Generalities of Operation

## 2

Jacopo P. Mortola

### Educational Aims

- General principles of operation of the neonatal respiratory system examined as a neuromechanical unit, that is, the translation of neural output into mechanical events, and the neurochemical feedback
- Implications of the highly compliant chest wall on the neonatal breathing pattern
- Dynamic elevation of the functional residual capacity: mechanisms and implications
- Importance of hypometabolism in neonatal hypoxia and its implications on the level of pulmonary ventilation

events accompanying birth. With respect to body size, the newborn's small body mass reduces the heat capacitance of the body and the high body surface-to-volume ratio favors heat loss. Hence, a large weight ( $W$ )-specific metabolic rate is required to offset the propensity for hypothermia. This translates into high values of oxygen uptake ( $\dot{V}_{O_2}/kg$ ), which imply a proportionally large pulmonary ventilation ( $\dot{V}_E$ ), achieved mostly by breathing at high rates. Indeed, fast breathing is a common characteristic of all newborn mammals. With respect to maturation, the development of the lung and respiratory apparatus is never completed at birth, even in the most precocial species. The lungs continue to grow postnatally, in a centripetal direction, with formation of peripheral airways and alveoli, implying that in newborns the central airways comprise a larger proportion of the total air space and contribute to a relatively large anatomical dead space. The incomplete myelination and low conduction velocity of neural fibers, including those of the laryngeal, vagi, and carotid sinus nerves, limit the afferent sensory information involved in the regulation of breathing. Finally, with respect to the adaptation to air breathing at birth, a multitude of events among the most dramatic of the whole life occur quite rapidly, such as the transition from filtration to absorption of the pulmonary fluid, the drastic changes in pulmonary circulation, and the rise in oxygenation with its implications on the function of the chemoreceptors.

This chapter touches on the general principles of operation of the neonatal respiratory system

## 2.1 Generalities

Many examples could be offered to illustrate the concept that the respiratory physiology of the newborn is markedly distinct from that of the adult. Some of the differences are determined by body size, others by the stage of development and maturation, and others yet by the tumultuous

---

J.P. Mortola  
Department of Physiology, McGill University,  
3655 Sir William Osler promenade, Montreal,  
QC H3G 1Y6, Canada  
e-mail: [jacopo.mortola@mcgill.ca](mailto:jacopo.mortola@mcgill.ca)

**Table. 2.1** Terminology, abbreviations adopted in text, common units

Cw	Chest wall compliance	$\Delta\text{Volume}/\Delta\text{Pressure}$
CL	Lung compliance	$\Delta\text{Volume}/\Delta\text{Pressure}$
f	Breathing frequency	Breaths/min
FRC	Functional residual capacity	ml
P	Pressure	cm H <sub>2</sub> O or mmHg
Pab	Abdominal pressure	cm H <sub>2</sub> O
PaO <sub>2</sub>	Arterial pressure of oxygen	mmHg
PACO <sub>2</sub>	Alveolar pressure of carbon dioxide	mmHg
Pdi	Transdiaphragmatic pressure	cm H <sub>2</sub> O
PEEP	Positive end-expiratory pressure	cm H <sub>2</sub> O
Ppl	Pleural pressure	cm H <sub>2</sub> O
T	Tension	dyne/cm
T <sub>E</sub>	Expiratory time	s
T <sub>I</sub>	Inspiratory time	s
V	Volume	ml
$\dot{V}$	Flow	ml/s
$\dot{V}_A$	Alveolar ventilation	ml/min
$\dot{V}_E$	Pulmonary ventilation	ml/min
$\dot{V}_{O_2}$	Oxygen consumption	ml/min
$\dot{V}_{CO_2}$	Carbon dioxide production	ml/min
V <sub>r</sub>	Resting volume of the respiratory system	ml
V <sub>T</sub>	Tidal volume	ml
W	Weight	kg

examined as a neuromechanical unit that operates to generate the optimal  $\dot{V}_E$  responses by integrating peripheral information with metabolic requirements. The general principles of operation of the neuromechanical unit (that is, translation of neural output into mechanical events, evaluation of the adequacy of  $\dot{V}_E$  from neurochemical feedback) are the same throughout the whole postnatal life. However, some functional aspects are peculiar to the neonatal period. The goal of this chapter is to highlight the functional properties characteristics of the newborn. References are not extensive, with preference given to review articles where the interested reader can trace the background information and technical details. A short list of terminology and abbreviations adopted in the text is given in Table 2.1.

## 2.2 Neural Output

The mechanisms involved in the generation of the neural output responsible for the breathing rhythm are still largely unexplained. Several approaches have attempted to unveil the origin of respiratory rhythmogenesis with reduced preparations, such as brainstem with or without the spinal cord or brainstem slices, commonly from neonatal rats. In vitro, it appears that a respiratory rhythm can be generated in very discrete brainstem regions (Haddad 2003), and the maturation of the putative respiratory centers has been studied extensively (Hilaire and Duron 1999). Nevertheless, discrepancies between the in vitro neural pattern and the natural breathing pattern in vivo are common and difficult to explain (Achard et al. 2005). Partly, these differences could be due to technical issues, adequate oxygenation of the preparation, and modification or abolition of the normal afferent inputs. Partly, they could reflect a fundamental flaw in the assumption that, normally, rhythmogenesis is independent of peripheral metabolic, neural, or chemical inputs. The major obstacle for a unifying interpretation of the results regarding the development of the neonatal network is the lack of an accepted model of respiratory rhythmogenesis.

What are the mechanisms that make breathing a continuous process at birth after the intermittent pattern of the fetal period is an unsolved mystery in perinatal physiology. The number and intensity of stimuli at birth – tactile, pressure, thermal, visual, and acoustic – are large. In addition, internal afferent inputs, like those of the chemoreceptors and of the airway mechanoreceptors, change markedly because of the rapid changes in oxygenation and in transpulmonary pressure. Under controlled experimental conditions, each of the putative stimuli in isolation can be shown to be important in initiating breathing. In fact, it is reasonable to assume that any stimulus can take a dominant role in specific situations (Jansen and Chernick 1991). Arousal, oxygenation, and a general state of stress (Lagercrantz and Slotkin 1986) increase metabolic rate, which by itself (through the increase in CO<sub>2</sub> production) stimulates breathing. Experiments on animal fetuses have indicated that exteriorization and clamping of the chord are not sufficient to trigger

and maintain continuous breathing unless also  $\text{CO}_2$  is allowed to rise (reviewed in Mortola 2001). In birds, the trigger, increase, and maintenance of  $\dot{V}_E$  at hatching are believed to be caused by the rise of metabolically produced  $\text{CO}_2$  (Mortola 2009). Hence, it is quite possible that the risen level in metabolic rate at birth represents the main mechanism sustaining continuous  $\dot{V}_E$ .

In newborns, the phrenic output is typically an abrupt and short burst of activity with few spikes, quite different from the activity-reach ramp-shaped firing of the adult phrenic nerve. The rapid and short-lasting neural burst leaves little room for fine modulation. Presumably, this is one reason for the fact that in newborns peripheral inputs often play only a coarse control on the breathing pattern, which is typically irregular and variable. In fact, in the neonatal period peripheral neural afferents, despite being less in number and with lower firing frequency than in adults, can generate quite drastic reflex effects on breathing (2.5.1).


## 2.3 Translation of Muscle Contraction into Pulmonary Ventilation

The  $\dot{V}_E$  generated by the contraction of the respiratory muscle results from three key steps (Fig. 2.1). First, the intrinsic mechanical characteristics of the muscle fibers determine the force

that the muscle is able to generate. Then, the force translates into pressure, the magnitude of which depends on the geometrical characteristics of the muscle and the structure over which the force is applied. Finally, the pressure generates airflow and causes changes in lung volume according to the flow-resistive and elastic properties of the respiratory system.

### 2.3.1 Step 1. Force Generation

For a long time it was suspected that the respiratory muscles of the newborn operate close to the fatigue threshold, as if they were barely capable to fulfill the ventilatory requirements. This view probably stemmed from the observation that breathing in neonates, especially premature infants, often is irregular and interspersed with periods of short apneas. The finding that the so-called “fatigue-resistant” muscle fibers were underrepresented fueled the idea that muscle failure was a contributor to periodic breathing and apneas (Keens et al. 1978; Le Souëf et al. 1988; Watchko et al. 1992; Vazquez et al. 1993). In reality, diaphragm fiber typing does not necessarily correlate with muscle fatigability (Sieck and Fournier 1991), and changes in diaphragmatic activity do not correlate with the infant’s periodic breathing or apneas (Nugent and Finley 1985). Studies on newborns of various species have shown the neonatal diaphragm to be at



**Fig. 2.1** Schematic representation of the neuromechanical unit for the operation of the respiratory system, with a generator, an output, and a feedback control. At right are

indicated the main steps involved in the translation of muscle activation into pulmonary ventilation

least as fatigue-resistant as the adult's diaphragm, capable of performing well even in the face of major workloads (Lieberman et al. 1972; Maxwell et al. 1983; Powers et al. 1991). Blood perfusion of the neonatal diaphragm is at least as adequate as it is in adult (Soust et al. 1989; Berger et al. 1994), even at high contraction rates or against resistive loads.

### 2.3.2 Step 2. Pressure Generation

In mammals, the total mass of the respiratory muscles, in relation to body mass, is almost a fixed proportion irrespective of species and animal age. The force produced by a muscle is proportional to its cross-sectional area, and the resulting pressure is the ratio between the force produced and the surface which the force is applied on. Because of their small size, the neonatal respiratory muscles produce little force by comparison to adults (Sieck et al. 2002). However, the surface over which such force is applied is proportionally small; hence, mammals of all sizes and ages do not differ much in the pleural pressure that their respiratory muscles can generate (Mortola 2001).

The shape of the diaphragmatic dome is an additional factor in the translation of force into pressure. In fact, by application of the Young-Laplace relationship, for a given diaphragmatic tension  $T$ , the resulting transdiaphragmatic pressure  $P_{di}$ <sup>1</sup> depends on the radius of curvature  $r$  of the dome ( $P_{di} = T/r$ ). In newborns, the fact that  $r$  is much smaller than in adults favors the generation of a greater  $P_{di}$  for a similar  $T$ . In conclusion, despite the fact that muscle mass and muscle force are undoubtedly less in newborns than in adults, the tidal swings in pleural pressure ( $P_{pl}$ ) are similar at all ages, about 5–7 cm H<sub>2</sub>O.<sup>2</sup> If needed, the healthy newborn can generate much higher  $P_{pl}$  values, up to 100 cm H<sub>2</sub>O, as is the case during the


first inspiration (Mortola 2001). The infant's maximal inspiratory pressures during crying (Shardonofsky et al. 1989) are not much lower than the maximal static inspiratory pressures developed by adults.

Contraction of the diaphragm raises abdominal pressure ( $P_{ab}$ ). This increase causes the outward motion of the frontal abdominal wall during inspiration and the expansion of the lower portion of the rib cage. The latter occurs because of the mechanical interdependence between the abdomen and rib cage and because  $P_{ab}$  gets transmitted to the thoracic wall through the apposition area. The apposition area is the lowermost region of the rib cage that the diaphragmatic dome faces without interposed lungs. In infants, differently from the adult, in inspiration the expansion of the lower rib cage is small because of two reasons. First, the high abdominal compliance limits the rise in  $P_{ab}$  during diaphragmatic contraction. Second, the rather rounded shape of the ribs and their almost perpendicular attachment to the vertebral column limit the size of the area of apposition (Allen and Gripp 2002). With growth, the area of apposition increases because of the gradual downward orientation of the ribs, presumably caused by the gravitational pull. The result of this anatomical arrangement is that in newborns the increase in  $P_{ab}$  during diaphragmatic contraction does not contribute to the expansion of the lower ribs as much as it does in adults. Switching from the supine to the prone position stiffens the abdomen; in fact, this postural change is functionally equivalent to binding the infant's abdomen and improves diaphragmatic efficiency (Fleming et al. 1979; Guslits et al. 1987; Laing et al. 1988; Wolfson et al. 1992) with positive effects on blood oxygenation (Numa et al. 1997). As an aside, it is interesting to note that the sudden infant death syndrome (SIDS) has a greater prevalence in the prone position (Silvestri and Weese-Mayer 2003). This implies that in the prone position the posture-related factors involved in the physiopathology of SIDS, whatever they may be, more than offset the advantage in the mechanical operation of the respiratory system.

<sup>1</sup>  $P_{di}$  is the pressure difference between the abdomen and the pleural space, or  $P_{ab} - P_{pl}$ .

<sup>2</sup> For analogous reasons, the tidal pleural pressure swings have similar values in species of very different body size.

**Fig. 2.2** An example of dynamic lung compliance progressively lower than the static value the higher the frequency of ventilation. The phenomenon, often labeled “frequency dependence of compliance,” is contributed by the stress relaxation of the pulmonary structures and by peripheral resistance. These experimental data refer to newborn and adult cats (Redrawn from the data of Sullivan and Mortola (1986))



### 2.3.3 Step 3. Lung and Chest Wall Mechanics


As it is the case for many other internal organs, the W-specific mass of the lung (lung weight-body weight ratio) is large in newborns and decreases during postnatal growth. Despite the large mass, air spaces are still incompletely developed, the pulmonary elastin and collagen contents are low, and during the first postnatal hours or days, some fluid is trapped in the peripheral airways and lung interstitium. All these factors contribute to values of lung compliance (CL,  $\Delta\text{volume}/\Delta\text{pressure}$ ) lower in newborns than in adults, when the comparison is made on the basis of lung volume or lung weight. The peripheral inequalities in airway resistance and the viscous properties of the pulmonary tissue prolong the response time of the lungs both in inspiration and in expiration. For these reasons, the differences between dynamic and static CL (and, possibly, the phenomenon of frequency dependence of CL<sup>3</sup>) are more pronounced in newborns than in adults (Fig. 2.2).

The chest wall anatomically and functionally comprises two main compartments, the rib cage (or thorax) and the abdomen-diaphragm. Chest wall compliance (Cw) has been measured in

newborns of many species and in infants; the results have been uniform in revealing high values (after normalization by body W) in comparison to the adult. The high Cw in newborns is a structural necessity for the passage through the birth canal. Cw decreases gradually during postnatal growth (Papastamelos et al. 1995), probably because of the stiffening of the cartilaginous structures and changes in abdominal and thoracic configuration.

Because the Cw-CL ratio is a dimensionless parameter, it can be compared directly among individuals of different age or body size, with no needs for normalization. In adult humans, because Cw and CL have similar values, their ratio is approximately equal to one. In infants, Cw is about five times higher than CL (Polgar and Weng 1979). In fact, the high Cw-CL ratio is a characteristic feature of the respiratory system of all neonatal mammals investigated (Mortola 2001). The fact that Cw is so high relative to CL implies that in infants CL is the major determinant of the compliance of the respiratory system (Crs) and that changes in Crs are an excellent indicator of changes in CL. From a practical viewpoint, this is quite convenient because measurements of CL in infancy can be difficult (due to the uncertainties in the measurements of Ppl), while measurements of Crs can be performed easily (England 1988; Wohl 1991; Mortola 2004).

<sup>3</sup> Whereby CL decreases, the faster the ventilation rate.



**Fig. 2.3** Schematic representation of the two main compartments of the chest wall (dashed line), the rib cage and abdomen, separated by the diaphragm. During passive inflation (panel at left), the pressure across the diaphragm (transdiaphragmatic pressure,  $P_{di}$ ) is zero because the diaphragm is relaxed. In this condition, pleural pressure

( $P_{pl}$ ) and abdominal pressure ( $P_{ab}$ ) increase by similar amounts, and both compartments expand. When the diaphragm contracts ( $P_{di} > 0$ , panel at right),  $P_{ab}$  increases and  $P_{pl}$  decreases. The result is that the abdomen expands while the rib cage moves little in its lower portion and caves inward in its upper portion

## 2.4 Mechanical Constraints and Breathing Pattern

The high Cw-CL ratio of the newborn has two major implications. First, in inspiration, the tendency of the chest wall to distort is greater than in adults. Second, during expiration, the low ratio reduces the outward pull of the chest on the lungs and facilitates lung emptying and a low resting volume of the respiratory system ( $V_r$ ). These mechanical characteristics have an impact on the neonatal pattern of breathing.


### 2.4.1 Chest Wall Distortion

In any solid structure, distortion can be defined as a configuration that differs from that requiring minimal energy. In the case of the chest wall, distortion occurs in active conditions (i.e., during respiratory muscle contraction) when the shape deviates from the configuration assumed in the passive situation.<sup>4</sup> This latter is the configuration

attained at any lung volume when the respiratory muscles are relaxed, as is the case, for example, during mechanical ventilation. By definition, the occurrence of distortion implies that the respiratory muscles need to spend extra energy to generate a given tidal volume ( $V_T$ ). In inspiration, distortion is the unavoidable consequence of the location of the diaphragm between the two main chest wall compartments, the abdomen and thorax. In fact, during contraction (i.e., with  $P_{di} > 0$ ) the diaphragm operates simultaneously as a positive pressure pump for the abdomen, raising  $P_{ab}$ , and as a negative pressure pump for the thorax, lowering  $P_{pl}$ . Therefore, differently from the passive inflation in which  $P_{ab}$  and  $P_{pl}$  rise homogeneously and expand both abdomen and thorax, during active inspiration the abdomen expands while the rib cage caves inward (Fig. 2.3). The suction on the rib cage can be appreciated visually in tetraplegic patients, who have no control of the extradiaphragmatic muscles (Mortola and Sant'Ambrogio 1978; Thach et al. 1980). In normal conditions, the tendency to distortion is less in adults than in newborns because adults have a more rigid chest wall and greater compensatory action of the extradiaphragmatic muscles. In newborns, the highly compliant chest wall, the small area of apposition (2.3.2), and the limited mechanical linkage between lower and upper rib

<sup>4</sup>The terms *active* and *passive*, *static* and *dynamic*, refer to the modes of operation of the respiratory system. *Active* and *passive* indicate, respectively, the presence or absence of respiratory muscle contraction. *Static* and *dynamic* indicate, respectively, the absence or presence of airflow.





**Fig. 2.4** Schematic pressure-volume (P-V) relations of the lungs, chest wall, and respiratory system (*heavy line*) in adult (*left*) and newborn (*right*). For simplicity, the P-V relations are indicated by *straight lines*, the slope of which represents the compliance. At left, the compliance of the chest wall (Cw) is similar to that of the lungs (CL). At right, the Cw-CL ratio is about five times higher. The

result is that in the newborn the resting volume of the respiratory system ( $V_r$ , *dashed line*) is lower than in adults. Tidal volume ( $V_T$ , with inspiratory-expiratory loops indicated by *arrows*) is approximately at the same absolute lung volume in both newborns and adults, because newborns keep their end-expiratory level dynamically elevated


cage are the main factors responsible for chest distortion, of which the most obvious aspect is the paradoxical inward motion of the rib cage during inspiration and expansion in expiration. In addition, the poor activity and mechanical coordination of the intercostals muscles, especially during some phases of sleep (Muller et al. 1979), further contribute to the limited stability of the neonatal thorax.

It is difficult to estimate the energetic price of distortion. Taking the abdominal expansion as an index of diaphragmatic shortening during inspiration, in newborns during resting breathing, chest wall distortion reduces by half the inspiratory efficiency of the diaphragm (Mortola 1995). This means that, in first approximation, to achieve a given  $V_T$ , the diaphragm must contract twice as much than it would need had the system expanded along its passive configuration. An additional burden caused by chest distortion is the deformation of the lungs, which reduces CL (Sullivan and Mortola 1985) and probably worsens the ventilation-perfusion matching. As mentioned above (3.2.), the prone position, by stiffening the chest wall, reduces its distortion

during inspiration and improves the mechanical efficiency of breathing.

#### 2.4.2 Low Resting Volume: Problems and Solutions

A highly compliant chest wall exerts a lesser outward pull on the lungs than a stiffer chest does, with the result that the passive resting volume of the respiratory system ( $V_r$ , the volume at which lungs and chest recoil pressures offset each other) is low (Fig. 2.4). Indeed,  $V_r$  (after normalization by lung weight) is lower in newborns than in adults of several species and in infants by comparison to the adult man (Cook et al. 1958; Fisher and Mortola 1980). The fact that  $V_r$  is low carries consequences on the functional residual capacity (FRC), which has a prominent role in the efficient operation of the mammalian respiratory system. FRC buffers the oscillations and maintains the stability of the alveolar and arterial gases and is a reserve of oxygen during occasional periods of hypoventilation or apneas. Most important is the fact that inflation of the lungs



**Fig. 2.5** Schematic representation of the dynamic elevation of the end-expiratory level (FRC). (a) represents the breathing pattern with expiration ending at resting volume ( $V_r$ ); this is the common resting pattern of adult

humans. (b, c) Represent patterns commonly seen in infants, in whom FRC is above  $V_r$  because of fast breathing (b) or because of laryngeal expiratory breaking (c)

from FRC requires substantially less pressure than inflation from a collapsed state. In adults during resting breathing, FRC is essentially equal to  $V_r$ . Infants, on the contrary, compensate the potentially disadvantageous mechanical situation of the low  $V_r$  by keeping FRC dynamically elevated; hence, differently from adults, in infants FRC exceeds  $V_r$  (Fig. 2.4).

The core mechanism that permits a dynamic elevation of FRC above  $V_r$  is a mismatch between the time needed for expiratory flow (mechanical expiratory time) and the neural expiratory time ( $T_E$ ), with the former being longer than the latter. In this way, inspiration begins before the air is fully exhaled, causing lung hyperinflation. At least three mechanisms operate to achieve this goal, the post-inspiratory activity of the expiratory muscles, laryngeal braking in expiration, and high breathing frequency (Fig. 2.5). The former two prolong the mechanical  $T_E$  and the latter shortens the neural  $T_E$ . All of them are operative, either together or individually, in infants (Eichenwald and Stark 2003) and in newborns of many other species (Mortola 2001). In infants, the  $FRC - V_r$  difference is 10–15 ml, or about 3 ml/kg. In cases of apnea (long neural  $T_E$ ) FRC invariably decreases toward  $V_r$ . During mechanical ventilation, the presence of an endotracheal tube eliminates the newborn's laryngeal control of expiratory flow; in this case, the application of an end-expiratory load (or positive end-expiratory pressure, PEEP) of a few cm  $H_2O$  is necessary to counteract the otherwise unavoidable drop in FRC. This becomes even more necessary in conditions of low CL because of lung disease, which further increases the Cw-CL ratio (Gregory et al.


1971; Berman et al. 1976). In fact, the  $FRC - V_r$  difference can be seen as a mechanism that generates an internal PEEP of a few cm  $H_2O$ . Although small, this positive airway pressure probably contributes to the absorption of the pulmonary fluid from the alveolar spaces into the lung interstitium during the first hours after birth (Strang 1991).

## 2.5 Feedback Regulation

Because gas exchange is the primary purpose of breathing,  $O_2$  and  $CO_2$  chemoreceptors could be considered sufficient to provide the pertinent information about the function of the neuromechanical unit (Fig. 2.1). However, although essential, the information on blood gases is not adequate for immediate responses to perturbations on breathing, because of the long delay built in the chemoreceptor feedback system. In fact, chemical feedback operates on a time scale of minutes and is most effective in managing the proper matching between alveolar ventilation ( $\dot{V}_A$ ) and metabolic requirements. A separate and purely neural feedback provides instantaneous information on the status of lung expansion, with reflex responses operational within the breath itself (Fig. 2.6). Some of this feedback originates from airway mechanoreceptors (pulmonary stretch receptors), specifically concerned with the status of lung expansion.<sup>5</sup> Proprioceptors within the respiratory muscles provide additional

<sup>5</sup>The tension in the airway wall is the stimulus for these receptors.





**Fig. 2.6** Schema of the main feedback loops regulating breathing. The neural control is a rapid loop primarily concerned with the breathing pattern, tidal volume ( $V_T$ ), inspiratory ( $T_I$ ) and expiratory time ( $T_E$ ), and breathing frequency ( $f$ ). The chemical control is a slower loop aiming to stabilize oxygen, carbon dioxide, and acid–base in blood and

cerebrospinal fluid (CSF), by matching alveolar ventilation ( $\dot{V}_A$ ) to gaseous metabolism (oxygen consumption  $\dot{V}_{O_2}$  and carbon dioxide production  $\dot{V}_{CO_2}$ ). Metabolism can provide a feedforward stimulus on ventilation, although this is a controversial issue because no plausible mechanisms for metabolism detection are known

neural information used predominantly to optimize the integration of breathing with non-respiratory functions, posture, and locomotion.

### 2.5.1 Vagal Feedback

The vagal feedback from the slowly adapting airway receptors is of primary importance in the regulation of the breathing pattern and of lung volume. This rapid neural information is essential for the stability of breathing when the respiratory system is confronted by mechanical loads, such as a reduction in compliance or increase in resistance, or changes in chest configuration, as with changes in body posture or muscle exercise (Milic-Emili and Zin 1986; Daubenspeck 1995).

Many experiments in several species have indicated that in newborns the response to a respiratory load that limits lung expansion happens within one breath, with a compensatory ability at least as efficient as that of adults (Mortola 2001). Premature infants have good reflex compensatory mechanisms (Kosch et al. 1986; Fox et al. 1988). Following lung expansion

or during PEEP, breathing becomes slow, a response common to all newborn species investigated, including infants; a similar response has been observed in fetuses (Ponte and Purves 1973). These compensatory responses to respiratory loads or lung inflation are mediated by the vagi nerves; indeed, many animal experiments have shown that bilateral section of the vagi not only causes deeper and slower breathing pattern but also diminishes or abolishes the compensatory responses to mechanical loads. In addition to its role in the regulation of the breathing pattern, the vagal afferent innervation seems to be of crucial importance in the transition from fetal to postnatal life (Lalani et al. 2001). According to some reports, cervical bilateral vagotomy in newborns can cause severe or fatal respiratory insufficiency (Coombs and Pike 1930; Schwieler 1968). One important caveat in the interpretation of these latter experiments, though, is that cervical section of the vagi not only abolishes the afferent information from the airways but also interrupts efferent neural traffic that could have an impact on breathing. For example, vagotomy eliminates the motor control of the larynx via the

recurrent laryngeal nerve, which is important in the control of FRC (2.4.2), and the efferent control of airway smooth muscle, with effects on the activity of the airway receptors (Fisher et al. 2003). In addition, the vagal pulmonary innervation influences the secretion of surfactant and lung compliance.

In newborns, the activity of the pulmonary airway receptors at end expiration and during lung expansion is less than in adults (Fisher et al. 1991). Many neural fibers are of small diameter and are not myelinated, with low conduction velocity (Schwieler 1968; Marlot and Duron 1979; Hasan et al. 1993). As mentioned above (2.2), during the neonatal period the combination of a limited peripheral afferent input with a strong reflex response is not infrequent, and the pulmonary vagal reflexes are one example of this combination. Hypoxia and hyperthermia, either singly or combined, strengthen the vagal reflex inhibition of breathing, a phenomenon that could have some importance in the genesis of breathing irregularities (Mortola 1999).

### 2.5.2 Extrapulmonary Ventilatory Reflexes

From studies in adult humans and animals, it is clear that stimuli from the most disparate sources – internal (e.g., carotid sinus, bladder and abdominal viscera, nasal, pharyngeal, and laryngeal regions) and external (e.g., visual, tactile, thermal, auditory) – can have reflex effects on breathing. Out of this multitude of extrapulmonary ventilatory reflexes, only a few have been studied with sufficient depth in the neonatal period.

The upper airways are involved in multiple functions; in newborns, reflexes from this area exert mostly inhibitory influences on breathing. Experiments on neonatal mammals have shown that liquids instilled in the pharyngolaryngeal area trigger apneas and bradycardia, the severity of which decreases with postnatal growth. Small  $[Cl^-]$ , high  $[K^+]$ , and extreme deviations of the pH of the liquid cause the strongest episodes of apnea (Boggs and Bartlett 1982). Observations in infants, although very limited (Plaxico and


Loughlin 1981; Davies et al. 1989; Wennergren et al. 1989), suggest that the results in animals are applicable to the human case and may be part of the reflexes lowering  $\dot{V}_E$  during oral feeding (Mathew 2003). Cooling of the laryngeal mucosa or sustained negative pressures in the upper airways are other powerful inhibitory stimuli on breathing (Fisher et al. 1991), responsible for reflex apneas that can persist beyond the removal of the stimulus. The evolutionary significance of the ventilatory inhibition triggered by upper airway reflexes is unclear. The inhibition of breathing may be considered the most obvious response against foreign bodies in the upper airways, irritants, or gastroesophageal reflux. The associated reflexes designed to clear the upper airway passages, like sneezing, coughing, and other responses common in adults, in newborns have modest efficacy (Mortola and Fisher 1988; Chernick 2002).

Gentle touching of the skin can facilitate breathing; presumably, this is one purpose for the maternal grooming and licking of the pups at birth practiced by many mammals. In humans, the skin-to-skin contact between infant and mother (“kangaroo care”) seems to reduce apneas and increase the regularity of the breathing pattern (Anderson 1991). However, the reflex response to skin stimulation is not necessarily facilitating breathing, since inhibitory effects, including apnea, have been described (reviewed in Mortola 2001). The large range of responses probably reflects the interplay of numerous factors of difficult control in an experimental setting, the intensity and location of the stimulus, association with other inputs, metabolic condition, state of alertness or sleep stage, etcetera. It seems probable that under normal circumstances, inputs from the chest wall have low priority on the regulation of breathing by comparison to inputs from the lungs and the chemoreceptors.

### 2.5.3 Chemical Feedback

The  $O_2$  and  $CO_2$  chemoreception important for respiratory control is organized at two sites, the arterial blood (carotid body) and the ventral

**Fig. 2.7** Metabolism-ventilation diagram. The oblique *short-dashed line* joins various combinations of metabolism and ventilation during normo-ventilation, of which the open circle represents one case. Hyperventilation is the increase in the ventilation-metabolism ratio, as represented by the steeper *long-dashed line*. In newborns, hypercapnic hyperventilation (*filled circle*) is obtained by an increase in  $\dot{V}_E$  (hyperpnea) with no hypometabolism. Differently, hypoxic hyperventilation is obtained by various combinations of hypometabolism and hyper- or hypopnea (*squares*)



surface of the medulla in close contact with the respiratory neurons. Only the former senses changes in oxygenation by monitoring the arterial partial pressure of  $O_2$  ( $PaO_2$ ). With some variations, this arrangement is common at all mammals and ages; however, the degree of operation of the receptors and their ventilatory reflexes differ between newborns and adults.

### 2.5.3.1 Hypoxia

The ventilatory response to hypoxia during the early postnatal period has been the object of extensive investigation after the earliest observations that in newborn infants the hypoxic increase in  $\dot{V}_E$  is minimal by comparison to adult humans. In fact, during hypoxia  $\dot{V}_E$  can even drop below the normoxic value (e.g., Brady and Ceruti 1966; Cohen et al. 1997; Horne et al. 2005). These results have been confirmed on a large number of neonatal species. The interpretation of the  $\dot{V}_E$  response to hypoxia is complex (Mortola 1996; Powell et al. 1998). In newborns, a major factor is the metabolic response to hypoxia, usually quantified as a change in oxygen consumption ( $\dot{V}_{O_2}$ ). Measurements in infants (Brady and Ceruti 1966; Rigatto and Brady 1972) and in neonatal animals have shown that arterial  $PCO_2$  does not increase in hypoxia even when  $\dot{V}_E$  is below the normoxic level. The only explanation is that the drop in  $\dot{V}_E$  is accompanied by a parallel decrease

in metabolism.<sup>6</sup> In fact, many studies have indicated that newborn mammals drop  $\dot{V}_{O_2}$  during hypoxia, as reported first in human infants (Cross et al. 1958). Hyperventilation, defined as an increase in the ventilation-metabolism ratio and expressed by the drop in alveolar or arterial  $PCO_2$ , equally results from hyperpnea (increase in  $\dot{V}_E$ ) or hypometabolism (drop in  $\dot{V}_{O_2}$ ), in any combination (Fig. 2.7). In adult humans the hyperpnea represents the common way to hyperventilate; in newborns, hypometabolism is the main mechanism to achieve hypoxic hyperventilation, usually with minimal hyperpnea (Mortola 1999). The hypometabolism results from the downregulation of various energy-consuming functions, especially thermogenesis (with a drop in body temperature) and body growth. The benefit of hypoxic hypometabolism is a hypoxic resistance far superior to adults. The physician's failure to recognize hypometabolism as the key strategy adopted by the newborn to cope with hypoxia can lead to unnecessary or mistaken interventions on the infant's ventilation and body temperature.

<sup>6</sup>In fact,  $PaCO_2$  is in equilibrium with the alveolar pressure of  $CO_2$  ( $PACO_2$ ), which, for a given barometric pressure  $P_b$ , is solely determined by the ratio between  $CO_2$  production ( $\dot{V}_{CO_2}$ ) and alveolar ventilation ( $\dot{V}_A$ ), according to the alveolar gas equation  $PACO_2 = (\dot{V}_{CO_2} / \dot{V}_A) \cdot P_b$ .

Recordings from neural afferent fibers of the carotid sinus nerve or from its cell bodies in the sensory petrosal ganglion have indicated that the carotid body is tonically active in newborns and probably in the fetus (Mulligan 1991). However, neither the tonic activity at resting  $\text{PaO}_2$  nor the carotid body neural afferent response to hypoxia or hypercapnia is as marked as at older ages (Carroll et al. 1993). During maintained hypoxia, the excitation of the carotid sinus nerve is sustained over time, even when the ventilatory output is declining; this indicates that the modest or absent hyperpnea of the hypoxic newborn cannot be attributed to adaptation of the chemoreceptors. Hyperoxia lowers or completely silences the tonic activity of the carotid body. The rapid oxygenation at birth in the transition from the fetal to postnatal life probably saturates the chemoreceptors and silences them temporarily, until they gradually reset to the postnatal value of  $\text{PaO}_2$ .

Chemodenervation, either surgically or functionally induced by hyperoxia, reduces  $\dot{V}_E$  and causes irregularities in breathing. Whether or not the absence of this peripheral chemo-feedback in the newborn results in severe respiratory insufficiency and eventually death is unclear; in fact, experimental studies have produced conflicting results (reviewed in Mortola 2001).

### 2.5.3.2 Hypercapnia

All newborn species investigated respond to hypercapnia with hyperpneic hyperventilation (absolute increase in  $\dot{V}_E$ , Fig. 2.7), although the postnatal temporal evolution of the response may vary from species to species (Putnam et al. 2005). The neonatal hypercapnic response has two qualitative differences from the hypoxic response. First, the hyperventilation is entirely contributed by the hyperpnea even in those newborns that do not increase  $\dot{V}_E$  in hypoxia, with minimal changes in  $\dot{V}_{\text{O}_2}$ . Second, the hyperpnea is characterized predominantly by an increase in  $V_T$  in both animals and human infants (Brady and Dunn 1970; Haddad et al. 1980; Martin et al. 1985; Mortola and Lanthier 1996), contrary to the shallow and rapid breathing of the response to hypoxia.

After peripheral chemodenervation, the  $\dot{V}_E$  response to  $\text{CO}_2$  is still present, meaning that the chemoreceptors located centrally close to the medulla are functional (Purves 1966). Other approaches, on the whole animal or in reduced preparations, have confirmed that the central chemosensory areas are active in the newborn and that are responsive to changes in  $\text{CO}_2$  and pH (Wolsink et al. 1991; Issa and Remmers 1992; Wennergren and Wennergren 1980, 1983; Whittaker et al. 1990). With  $\text{CO}_2$  exposure, neurons expressing Fos-protein, taken as a marker of the immediate early gene c-Fos, were found in the ventrolateral medulla of neonatal rats; the incidence of these neurons was not lower than in older rats (Belegu et al. 1999). All together, these and other observations indicate that the central chemosensory area is active since birth and probably in the late fetal phases (Nattie 1991) and that the  $\dot{V}_E$  response to its stimulation is, at least qualitatively, similar to the adult.

## Conclusions

As mentioned at the onset of this chapter, respiratory physiology in the neonate differs from that of the adult in many aspects intertwined in a complex web. Any arbitrary attempt to point at just a few of them is bound to be restrictive and superficial. Nevertheless, after having focused our attention to the generation of mechanical output and the feedback operation, in the opinion of this author, the most fundamental peculiarities of the neuro-mechanical unit in the neonatal period can be summarized as follows:

- (a) The high  $C_w$  (and high  $C_w$ -CL ratio) is a functional characteristic responsible for the low  $V_b$ , with consequences on inspiration (chest distortion and extra inspiratory work) and expiration (dynamic elevation of FRC).
- (b) The low number of myelinated fibers (small conduction velocity) in the vagus, laryngeal, and carotid sinus nerves limits the effectiveness of peripheral feedback control.
- (c) The abrupt and brief burst of neural output activity contributes to the fact that changes

in feedback information produce rather course changes in amplitude and duration of inspiration. This could contribute to the variable breathing pattern and to the apparent paradox that in newborns, despite (b), peripheral inputs can provoke dramatic effects on  $\dot{V}_E$ .

- (d) During hypoxia, the hypometabolism is a neonatal strategy of fundamental importance and great effectiveness. It has implications on the control of the set point of body temperature and permits adequate hyperventilation with minimal hyperpnea.

### Essentials to Remember

- In infancy, chest wall compliance (Cw) relative to lung compliance (CL) is higher than in adults. A useful practical consequence is that measurements of the compliance of the respiratory system (which can be performed easily both during spontaneous breathing and mechanical ventilation) are excellent indicators of CL.
- During mechanical ventilation, the presence of an endotracheal tube eliminates the newborn's laryngeal control of expiratory flow; in this case, the application of an end-expiratory load (or positive end-expiratory pressure) of a few cm H<sub>2</sub>O is necessary to counteract the otherwise unavoidable drop in end-expiratory volume. This becomes even more necessary with lung diseases that lower CL and further increase the Cw-CL ratio.
- In the neonatal period, peripheral neural afferents, despite being less in number and with lower firing frequency than in adults, can generate drastic reflex effects on breathing. During mechanical ventilation, reflex inhibition on inspiration originates from the distention of the lungs and central airways, especially at elevated PEEP. The reflex inhibition on

breathing may be desirable to eliminate infant-ventilator interaction, but may be misleading in assessing the infant's capacity for spontaneous breathing.

- Liquids instilled in the pharyngolaryngeal area trigger apneas and bradycardia, especially at the youngest ages.
- In newborn infants, hypoxia decreases metabolic rate much more readily than it does in adults. This metabolic drop lowers the inspiratory output and the level of pulmonary ventilation and is often accompanied by a decrease in thermogenesis and body temperature. Failure to recognize hypometabolism as the key strategy adopted by the newborn to cope with hypoxia can lead to unnecessary or mistaken interventions on the infant's ventilation and body temperature.

### References

- Achard P, Zanella S, Rodriguez R, Hilaire G (2005) Perinatal maturation of the respiratory rhythm generator in mammals, from experimental results to computational simulation. *Respir Physiol Neurobiol* 149:17–27
- Allen J, Gripp KW (2002) Chapter 9. Development of the thoracic cage. In: Haddad GG, Abman SH, Chernick V (eds) *Basic mechanisms of pediatric respiratory disease*, 2nd edn. BC Decker Inc, Hamilton, pp 124–138
- Anderson GC (1991) Current knowledge about skin-to-skin (kangaroo) care for preterm infants. *J Perinatol* 11:216–226
- Belegu R, Hadžiefendić S, Dreshaj IA et al (1999) CO<sub>2</sub>-induced c-fos expression in medullary neurons during early development. *Respir Physiol* 117:13–28
- Berger PJ, Soust M, Smolich JJ et al (1994) Respiratory muscle blood flow in the fetal lamb during apnoea and breathing. *Respir Physiol* 97:111–121
- Berman LS, Fox WW, Raphaely RC et al (1976) Optimum levels of CPAP for tracheal extubation of newborn infants. *J Pediatr* 89:109–112
- Boggs DF, Bartlett D Jr (1982) Chemical specificity of a laryngeal apneic reflex in puppies. *J Appl Physiol* 53:455–462
- Brady JP, Ceruti E (1966) Chemoreceptor reflexes in the new-born infant: effects of varying degrees of hypoxia on heart rate and ventilation in a warm environment. *J Physiol Lond* 184:631–645

- Brady JP, Dunn PM (1970) Chemoreceptor reflexes in the newborn infant: effect of CO<sub>2</sub> on the ventilatory response to hypoxia. *Pediatrics* 45:206–214
- Carroll JL, Bamford OS, Fitzgerald RS (1993) Postnatal maturation of carotid chemoreceptor responses to O<sub>2</sub> and CO<sub>2</sub> in the cat. *J Appl Physiol* 75:2383–2391
- Chernick V (2002) Chapter 13. Physiology of cough. In: Haddad GG, Abman SH, Chernick V (eds) *Basic mechanisms of pediatric respiratory disease*, 2nd edn. BC Decker Inc, Hamilton, pp 179–183
- Cohen G, Malcolm G, Henderson-Smart D (1997) Ventilatory response of the newborn infant to mild hypoxia. *Pediatr Pulmonol* 24:163–172
- Cook CD, Helliesen PJ, Agathon S (1958) Relation between mechanics of respiration, lung size and body size from birth to young adulthood. *J Appl Physiol* 13:349–352
- Coombs HC, Pike FH (1930) The nervous control of respiration in kittens. *Am J Physiol* 95:681–693
- Cross KW, Tizard JPM, Trythall DAH (1958) The gaseous metabolism of the new-born infant breathing 15% oxygen. *Acta Paediatr* 46:265–285
- Daubenspeck JA (1995) Part A. Physiology. Chapter 32. Mechanical aspects of loaded breathing. In: Roussos C (ed) *The thorax*. Second revision, revised and expanded, vol 85, Lung biology in health and disease series. Marcel Dekker, New York, pp 953–985
- Davies AM, Koenig JS, Thach BT (1989) Characteristics of upper airway chemoreflex prolonged apnea in human infants. *Am Rev Respir Dis* 139:668–673
- Eichenwald EC, Stark AR (2003) Chapter 15. Maturation of respiratory control. In: Mathew OP (ed) *Respiratory control and disorders in the newborn*, vol 173, Lung biology in health and disease series. Marcel Dekker, New York, pp 355–372
- England SJ (1988) Current techniques for assessing pulmonary function in the newborn infant: advantages and limitations. *Pediatr Pulmonol* 5:48–53
- Fisher JT, Mortola JP (1980) Statics of the respiratory system in newborn mammals. *Respir Physiol* 41:155–172
- Fisher JT, Mathew OP, Sant'Ambrogio G (1991) Chapter 7. Morphological and neurophysiological aspects of airway and pulmonary receptors. In: Haddad GG, Farber JP (eds) *Developmental neurobiology of breathing*, vol 53, Lung biology in health and disease series. Marcel Dekker, New York, pp 219–244
- Fisher JT, Schweitzer CE, Weichselbaum M et al (2003) Chapter 3. Ontogeny of upper- and lower-airway innervation. In: Mathew OP (ed) *Respiratory control and disorders in the newborn*, vol 173, Lung biology in health and disease series. Marcel Dekker, New York, pp 39–81
- Fleming PJ, Muller NL, Bryan MH et al (1979) The effects of abdominal loading on rib cage distortion in premature infants. *Pediatrics* 64:425–428
- Fox RE, Kosch PC, Feldman HA et al (1988) Control of inspiratory duration in premature infants. *J Appl Physiol* 64:2597–2604
- Gregory GA, Kitterman JA, Phibbs RH et al (1971) Treatment of the idiopathic respiratory-distress syndrome with continuous positive airway pressure. *N Engl J Med* 284:1333–1340
- Guslits BG, Gaston SE, Bryan MH et al (1987) Diaphragmatic work of breathing in premature human infants. *J Appl Physiol* 62:1410–1415
- Haddad GG (2003) Chapter 1. Respiratory control in the newborn. Comparative physiology and clinical disorders. In: Mathew OP (ed) *Respiratory control and disorders in the newborn*, vol 173, Lung biology in health and disease series. Marcel Dekker, New York, pp 1–15
- Haddad GG, Leistner HL, Epstein RA et al (1980) CO<sub>2</sub>-induced changes in ventilation and ventilatory pattern in normal sleeping infants. *J Appl Physiol* 48:684–688
- Hasan SU, Sarnat HB, Auer RN (1993) Vagal nerve maturation in the fetal lamb: an ultrastructural and morphometric study. *Anat Rec* 237:527–537
- Hilaire G, Duron B (1999) Maturation of the mammalian respiratory system. *Physiol Rev* 79:325–360
- Horne RSC, Parslow PM, Harding R (2005) Postnatal development of the ventilator and arousal responses to hypoxia in human infants. *Respir Physiol Neurobiol* 149:257–271
- Issa FG, Remmers JE (1992) Identification of a subsurface area in the ventral medulla sensitive to local changes in PCO<sub>2</sub>. *J Appl Physiol* 72:439–446
- Jansen AH, Chernick V (1991) Fetal breathing and development of control of breathing. *J Appl Physiol* 70:1431–1446
- Keens TG, Bryan AC, Levison H et al (1978) Developmental pattern of muscle fiber types in human ventilatory muscles. *J Appl Physiol* 44:909–913
- Kosch PC, Davenport PW, Wozniak JA et al (1986) Reflex control of inspiratory duration in newborn infants. *J Appl Physiol* 60:2007–2014
- Lagercrantz H, Slotkin TA (1986) The “stress” of being born. *Sci Am* 254:100–107
- Laing IA, Teele RL, Stark AR (1988) Diaphragmatic movement in newborn infants. *J Pediatr* 112:638–643
- Lalani S, Remmers JE, Hasan SU (2001) Breathing patterns, pulmonary mechanics and gas exchange: role of vagal innervation in neonatal lamb. *Exp Physiol* 86:803–810
- LeSouëf PN, England SJ, Stogryn HAF et al (1988) Comparison of diaphragmatic fatigue in newborn and older rabbits. *J Appl Physiol* 65:1040–1044
- Lieberman DA, Maxwell LC, Faulkner JA (1972) Adaptation of guinea pig diaphragm muscle to aging and endurance training. *Am J Physiol* 222:556–560
- Marlot D, Duron B (1979) Postnatal maturation of phrenic, vagus, and intercostal nerves in the kitten. *Biol Neonate* 36:264–272
- Martin RJ, Carlo WA, Robertson SS et al (1985) Biphasic response of respiratory frequency to hypercapnea in preterm infants. *Pediatr Res* 19:791–796
- Mathew OP (2003) Chapter 16. Respiratory control during oral feeding. In: Mathew OP (ed) *Respiratory control and disorders in the newborn*, vol 173, Lung biology in health and disease series. Marcel Dekker, New York, pp 373–393



- Maxwell LC, McCarter RJM, Kuehl TJ et al (1983) Development of histochemical and functional properties of baboon respiratory muscles. *J Appl Physiol* 54:551–561
- Milic-Emili J, Zin WA (1986) Part 2. Chapter 23. Breathing responses to imposed mechanical loads. In: Cherniack NS, Widdicombe JG (eds) *Handbook of physiology. The respiratory system, vol II, Control of breathing*. American Physiological Society, Bethesda, pp 751–769
- Mortola JP (1995) Part A. Physiology. Chapter 22. Chest wall mechanics in newborns. In: Roussos C (ed) *The thorax. Second edition, revised and expanded, vol 85, Lung biology in health and disease series*. Marcel Dekker, New York, pp 617–631
- Mortola JP (1996) Chapter 15. Ventilatory responses to hypoxia in mammals. In: Haddad GG, Lister G (eds) *Tissue oxygen deprivation: developmental, molecular and integrated function, vol 95, Lung biology in health and disease series*. Marcel Dekker, New York, pp 433–477
- Mortola JP (1999) How newborn mammals cope with hypoxia. *Respir Physiol* 116:95–103
- Mortola JP (2001) *Respiratory physiology of newborn mammals. A comparative perspective*. The Johns Hopkins University Press, Baltimore
- Mortola JP (2004) Chapter 84. Mechanics of breathing. In: Polin RA, Fox WW, Abman SH (eds) *Fetal and neonatal physiology, vol 1, 3rd edn*. Saunders, Philadelphia, pp 865–870
- Mortola JP (2009) Gas exchange in avian embryos and hatchlings. *Comp Biochem Physiol A* 153:359–377
- Mortola JP, Fisher JT (1988) Chapter 9. Upper airway reflexes in newborns. In: Mathew OP, Sant'Ambrogio G (eds) *Respiratory function of the upper airways, vol 35, Lung biology in health and disease series*. Marcel Dekker, New York, pp 303–357
- Mortola JP, Lanthier C (1996) The ventilatory and metabolic response to hypercapnia in newborn mammalian species. *Respir Physiol* 103:263–270
- Mortola JP, Sant'Ambrogio G (1978) Motion of the rib cage and the abdomen in tetraplegic patients. *Clin Sci Mol Med* 54:25–32
- Muller N, Gulston G, Cade D et al (1979) Diaphragmatic muscle fatigue in the newborn. *J Appl Physiol* 46: 688–695
- Mulligan EM (1991) Chapter 11. Discharge properties of the carotid bodies. Developmental aspects. In: Haddad GG, Farber JP (eds) *Developmental neurobiology of breathing, vol 53, Lung biology in health and disease series*. Marcel Dekker, New York, pp 321–340
- Nattie EE (1991) Chapter 12. Central respiratory chemoreceptors. Cellular mechanisms and developmental aspects. In: Haddad GG, Farber JP (eds) *Developmental neurobiology of breathing, vol 53, Lung biology in health and disease series*. Marcel Dekker, New York, pp 341–371
- Nugent ST, Finley JP (1985) Spectral analysis of the EMG and diaphragmatic muscle fatigue during periodic breathing in infants. *J Appl Physiol* 58:830–833
- Numa AH, Hammer J, Newth CJL (1997) Effect of prone and supine positions on functional residual capacity, oxygenation, and respiratory mechanics in ventilated infants and children. *Am J Respir Crit Care Med* 156:1185–1189
- Papastamelos C, Panitch HB, England SE et al (1995) Developmental changes in chest wall compliance in infancy and early childhood. *J Appl Physiol* 78: 179–184
- Plaxico DT, Loughlin GM (1981) Nasopharyngeal reflux and neonatal apnea. *Am J Dis Child* 135:793–794
- Polgar G, Weng TR (1979) The functional development of the respiratory system. From the period of gestation to adulthood. *Am Rev Respir Dis* 120:625–695
- Ponte J, Purves MJ (1973) Types of afferent nervous activity which may be measured in the vagus nerve of the sheep foetus. *J Physiol Lond* 229:51–76
- Powell FL, Milsom WK, Mitchell GS (1998) Time domains of the hypoxic ventilatory response. *Respir Physiol* 112:123–134
- Powers SK, Lawler J, Criswell D et al (1991) Age-related changes in enzyme activity in the rat diaphragm. *Respir Physiol* 83:1–10
- Purves MJ (1966) The respiratory response of the newborn lamb to inhaled CO<sub>2</sub> with and without accompanying hypoxia. *J Physiol Lond* 185:78–94
- Putnam RW, Conrad SC, Gdovin MJ et al (2005) Neonatal maturation of the hypercapnic ventilatory response and central neural CO<sub>2</sub> chemosensitivity. *Respir Physiol Neurobiol* 149:165–179
- Rigatto H, Brady JP (1972) Periodic breathing and apnea in preterm infants. II. Hypoxia as a primary event. *Pediatrics* 50:219–227
- Schwieler GH (1968) Respiratory regulation during postnatal development in cats and rabbits and some of its morphological substrate. *Acta Physiol Scand Suppl* 304:3–123
- Shardonofsky FR, Perez-Chada D, Carmuega E et al (1989) Airway pressures during crying in healthy infants. *Pediatr Pulmonol* 6:14–18
- Sieck GC, Fournier M (1991) Chapter 13. Developmental aspects of diaphragm muscle cells. Structural and functional organization. In: Haddad GG, Farber JP (eds) *Developmental neurobiology of breathing, vol 53, Lung biology in health and disease series*. Marcel Dekker, New York, pp 375–428
- Sieck GC, Vandenboom R, Mantilla CB et al (2002) Chapter 10. Structural and functional development of the respiratory muscles. In: Haddad GG, Abman SH, Chernick V (eds) *Basic mechanisms of pediatric respiratory disease, 2nd edn*. BC Decker Inc, Hamilton, pp 139–153
- Silvestri JM, Weese-Mayer DE (2003) Chapter 23. SIDS and the newborn infant. In: Mathew OP (ed) *Respiratory control and disorders in the newborn, vol 173, Lung biology in health and disease series*. Marcel Dekker, New York, pp 525–539
- Soust M, Walker AM, Berger PJ (1989) Blood flow to the respiratory muscles during hyperpnoea in the newborn lamb. *Respir Physiol* 76:93–105

- Strang LB (1991) Fetal lung liquid: secretion and reabsorption. *Physiol Rev* 71:991–1016
- Sullivan KJ, Mortola JP (1985) Effect of distortion on the mechanical properties of the newborn piglet lung. *J Appl Physiol* 59:434–442
- Sullivan KJ, Mortola JP (1986) Dynamic lung compliance in newborn and adult cats. *J Appl Physiol* 60:743–750
- Thach BT, Abroms IF, Frantz ID et al (1980) Intercostal muscle reflexes and sleep breathing patterns in the human infant. *J Appl Physiol* 48:139–146
- Vazquez R, Daood M, Watchko JF (1993) Regional distribution of myosin heavy chain isoforms in rib cage muscles as a function of postnatal development. *Pediatr Pulmonol* 16:289–296
- Watchko JF, Daood MJ, Vazquez RL et al (1992) Postnatal expression of myosin isoforms in an expiratory muscle – external abdominal oblique. *J Appl Physiol* 73:1860–1866
- Wennergren G, Wennergren M (1980) Respiratory effects elicited in newborn animals via the central chemoreceptors. *Acta Physiol Scand* 108:309–311
- Wennergren G, Wennergren M (1983) Neonatal breathing control mediated via the central chemoreceptors. *Acta Physiol Scand* 119:139–146
- Wennergren G, Hertzberg T, Milerad J et al (1989) Hypoxia reinforces laryngeal reflex bradycardia in infants. *Acta Paediatr Scand* 78:11–17
- Whittaker JAC, Trouth CO, Pan Y et al (1990) Age differences in responsiveness of brainstem chemosensitive neurons to extracellular pH changes. *Life Sci* 46:1699–1705
- Wohl MEB (1991) Lung mechanics in the developing human infant. In: Chernick V, Mellins RB (eds) *Basic mechanisms of pediatric respiratory disease, cellular and integrative*. B.C. Decker Inc, Philadelphia, pp 89–99
- Wolfson MR, Greenspan JS, Deoras KS et al (1992) Effect of position on the mechanical interaction between the rib cage and abdomen in preterm infants. *J Appl Physiol* 72:1032–1038
- Wolsink JG, Berkenbosch A, DeGoede J et al (1991) Ventilatory sensitivities of peripheral and central chemoreceptors of young piglets to inhalation of CO<sub>2</sub> in air. *Pediatr Res* 30:491–495

<http://www.springer.com/978-3-642-01218-1>

**Pediatric and Neonatal Mechanical Ventilation  
From Basics to Clinical Practice**

Rimensberger, P. (Ed.)

2015, X, 1642 p. 405 illus., 109 illus. in color.,  
Hardcover

ISBN: 978-3-642-01218-1