

## Chapter 2

# The Taxonomy, Phylogeny, and Distribution of Lampreys

Ian C. Potter, Howard S. Gill, Claude B. Renaud and Dalal Haoucher

**Abstract** The lampreys (Petromyzontiformes), one of the two surviving groups of agnathan (jawless) vertebrates, currently consist of 41 recognized species. This group has an antitropical distribution, with the 37 species of Northern Hemisphere lampreys assigned to the Petromyzontidae, whereas the four species of Southern Hemisphere lampreys are separated into either the Geotriidae (one species) or Mor-daciidae (three species). All lamprey species have a blind and microphagous, burrowing larva (ammocoete), which spends a number of years in the soft sediment of creeks and rivers, after which it undergoes a radical metamorphosis. Eighteen lamprey species then embark on an adult parasitic phase (nine at sea and nine in fresh water) during which they increase markedly in size, whereas the other 23 species do not feed as adults and remain in fresh water. On the basis of morphology, 17 of the 23 non-parasitic species each evolved from a particular parasitic species whose descendants are still represented in the contemporary fauna. The remaining six non-parasitic species, the so-called “southern relict” species, have no obvious potential ancestral parasitic species, implying they have diverged markedly from their parasitic ancestor or that the parasitic ancestor is now extinct. Many of the main taxonomic characteristics reside in features that are associated with parasitic feeding, for example, the type and arrangement of the teeth on the suctorial disc and tongue-like piston. The phylogenetic relationships, derived by maximum parsimony analyses of morphological and anatomical data for the 18 parasitic species, were similar in most respects to those obtained by subjecting molecular data (cytochrome *b* mitochondrial DNA sequence data) for those species to Bayesian analyses. However, in contrast to the results of morphological analyses, the genera

---

I. C. Potter (✉) · H. S. Gill · D. Haoucher  
Centre for Fish and Fisheries Research, School of Veterinary and Life Sciences,  
Murdoch University, Murdoch, WA, 6150 Australia  
e-mail: I.Potter@murdoch.edu.au

H. S. Gill  
e-mail: H.Gill@murdoch.edu.au

C. B. Renaud  
Research and Collections Division, Canadian Museum of Nature,  
PO Box 3443, Station D, Ottawa, ON, K1P 6P4 Canada  
e-mail: crenaud@mus-nature.ca


*Eudontomyzon* and *Lampetra* were not monophyletic when using molecular analyses. When non-parasitic species were included in the molecular analyses, some of the six relict non-parasitic species formed clades with parasitic species which, from their morphology, had been allocated by taxonomists to different genera. More genes, and particularly nuclear genes, should be used to help resolve the basis for these differences between the morphological and molecular phylogenies.

**Keywords** Evolution · Geotriidae · Mordaciidae · Morphological and molecular analyses · Paired species · Petromyzontidae

## 2.1 Introduction

The lampreys, together with the hagfishes, are the two sole surviving groups of agnathan (jawless) vertebrates (Janvier 1981; Hardisty 2006; see Chap. 1). The possession by these two groups of “round mouths” led to them being termed by Duméril (1806), collectively, as the Cyclostomata, a term retained as a class by Holly (1933) in his important taxonomic treatise on these animals. The implication that lampreys and hagfishes formed a monophyletic group was accepted for many years. However, detailed comparisons of their anatomy, morphology, and physiology, in conjunction with comparisons to the morphology of extinct agnathans, led to an alternative viewpoint (Hardisty 1979, 1982; Janvier 1981). The latter authors came independently to the conclusion that lampreys were more closely related to the gnathostomatous (jawed) vertebrates than to the hagfishes. Since that time, however, the majority of the numerous molecular studies undertaken on the two surviving groups of agnathans have supported the monophyly of lampreys and hagfishes (e.g., Stock and Whitt 1992; Mallatt and Sullivan 1998; Kuraku et al. 1999; Delarbre et al. 2002; Takezaki et al. 2003; Blair and Hedges 2005; Kuraku and Kuratani 2006). The question of whether or not cyclostomes are considered to constitute a monophyletic group was subsequently shown by Near (2009) to be influenced by the characters used and the types of analyses employed. A subsequent study, however, by Heimberg et al. (2010), employing microRNAs and a reanalysis of morphological characters, provided such overwhelming evidence for cyclostome monophyly that it convinced Janvier (2010) that this was indeed the case.

The first fossil lamprey to be described was the beautifully-preserved *Mayomyzon pieckoensis* from the upper Carboniferous (c. 280 million years ago, mya) deposits of Mazon Creek in Illinois (Bardack and Zangerl 1968, 1971). This fossil clearly possessed many of the morphological and anatomical characters of the adults of extant lampreys, such as an annular cartilage, which maintains the structural integrity of the suctorial disc, a piston cartilage, dorsolateral eyes, and seven gill apertures on either side of the body. Since the landmark discovery of *M. pieckoensis*, a further three definitive fossil lampreys have been found. The youngest of these is *Mesomyzon mengae* from the lower Cretaceous of China c. 125 mya (Chang et al. 2006), followed in age by *Hardistiella montanensis* from lower Carboniferous deposits in Montana c. 320 mya (Janvier and Lund 1983), and then *Priscomyzon riniensis*



**Fig. 2.1** Lateral views of **a** a larval lamprey (ammocoete), **b** an adult lamprey, and **c** a hagfish. This figure was originally published in Hardisty et al. (1989). (Reproduced by permission of The Royal Society of Edinburgh from Transactions of the Royal Society of Edinburgh: Earth Sciences volume 80 (1989), pp. 241–254)

from upper Devonian deposits in South Africa c. 360 mya (Gess et al. 2006). The first indisputable fossil hagfish to be discovered was *Myxinikela siroka*, which was found in the same geological horizon and general locality as the lamprey *M. pieckoensis*, and thus likewise dates back c. 300 mya (Bardack 1991, 1998). More recently, another hagfish fossil, *Myxineidus gononorum*, was discovered in upper Carboniferous deposits in France and is therefore also of approximately the same age as the above two fossils (Poplin et al. 2001). Germain et al. (2014) have cast doubt, however, on whether *M. gononorum* is a hagfish and provide evidence that it could be a lamprey.

Both groups of extant cyclostomes possess a similar body shape (Fig. 2.1) and typically have an antitropical distribution (Hubbs and Potter 1971; Hardisty 1979). Although lampreys are thereby essentially confined to temperate regions of the world, two species (genus *Tetrapleurodon*) are found in elevated cooler waters in a restricted sub-tropical area (Álvarez del Villar 1966). The living lampreys are represented by three families (Mordaciidae, Geotriidae, and Petromyzontidae) and 41 species (Table 2.1; Potter et al. 2014) and the hagfishes by two subfamilies (Eptatretinae and Myxiniinae) and approximately 60 species (Fernholm 1998). However, whereas the Mordaciidae and Geotriidae are confined to the Southern Hemisphere and the Petromyzontidae to the Northern Hemisphere, the two subfamilies of hagfishes are represented in both hemispheres.

The aim of this chapter is to provide a comprehensive list of the species, genera, subfamilies, and families of extant lampreys, providing details of the types of morphological characters used in taxonomic studies and the distributions of each species. Emphasis is also placed on outlining the schemes that have been proposed for the interrelationships of the various species, based on morphological and molecular criteria, and discussing the implications of any differences between those schemes.

**Table 2.1** Classification, common names, life cycle types, and distributions of the 41 extant species of lampreys, following Potter et al. (2014) in all respects except that *hubbsi* is assigned to *Lampetra* rather than *Entosphenus*. Authorities for each taxon are given in Appendix 2.1. In the life cycle column, attention is drawn to those non-parasitic species that can unambiguously be paired with a particular parasitic species. Drainage refers to a river and its tributaries. Other frequently used common names, for example, those adopted by the American Fisheries Society (AFS; Page et al. 2013a) or Food and Agriculture Organization (FAO; see FishBase: Froese and Pauly 2013) or explanations regarding alternative or previous classifications are also provided

Classification	Common name	Life cycle type	Distribution	Comments
<b>Family Mordacidae</b> (southern top-eyed lampreys)				
Genus <i>Mordacia</i> (3 species)				
<i>Mordacia mordax</i>	Short-headed lamprey	Anadromous; parasitic	Drainages and coastal waters of south-eastern Australia, including Tasmania	Also known as Australian lamprey, but not recommended as it is imprecise
<i>Mordacia praecox</i>	Precocious lamprey	Freshwater; non-parasitic derivative of <i>M. mordax</i>	Drainages of southeastern Australia	Also known as Australian brook lamprey, but not recommended as it is imprecise
<i>Mordacia lapicida</i>	Chilean lamprey	Anadromous; parasitic	Drainages and coastal waters of Chile	
<b>Family Geotriidae</b> (southern striped lamprey)				
Genus <i>Geotria</i> (1 species)				
<i>Geotria australis</i>	Pouched lamprey	Anadromous; parasitic	Drainages of southern Australia, New Zealand, Chile, Argentina and widespread in intervening oceans	
<b>Family Petromyzontidae</b> (Northern Hemisphere lampreys)				
<b>Subfamily Petromyzontinae</b>				
Genus <i>Caspiomyzon</i> (1 species)				
<i>Caspiomyzon wagneri</i>	Caspian lamprey	Anadromous; parasitic	Caspian Sea and its drainages	Nelson (2006) excluded <i>Caspiomyzon</i> from this subfamily (see Sect. 2.4)
Genus <i>Petromyzon</i> (1 species)				
<i>Petromyzon marinus</i>	Sea lamprey	Anadromous and freshwater; parasitic	Atlantic drainages in North America from Newfoundland southwards to Florida and into the Gulf of Mexico and in Europe from Varanger Fjord southwards into the Mediterranean Sea. Widespread in the North Atlantic Ocean	

Table 2.1 (continued)

Classification	Common name	Life cycle type	Distribution	Comments
Genus <i>Ichthyomyzon</i> (6 species)				
<i>Ichthyomyzon unicuspis</i>	Silver lamprey	Freshwater; parasitic	Hudson Bay, Great Lakes, St. Lawrence River, and Mississippi River drainages	Nelson (2006) included <i>Caspiomyzon</i> in this subfamily; Vladikov (1972) separated these five genera into subfamilies Entospheninae ( <i>Entosphenus</i> and <i>Tetrapleurodon</i> ), and Lampetrinae ( <i>Lampetra</i> , <i>Lethenteron</i> , and <i>Eudontomyzon</i> ) Formerly synonymized with <i>Lampetra</i> (see Sect. 2.4)
<i>Ichthyomyzon fossor</i>	Northern brook lamprey	Freshwater; non-parasitic derivative of <i>I. unicuspis</i>	As for <i>I. unicuspis</i>	
<i>Ichthyomyzon castaneus</i>	Chestnut lamprey	Freshwater; parasitic	Hudson Bay, Great Lakes, St. Lawrence River, and Gulf of Mexico drainages	
<i>Ichthyomyzon gagei</i>	Southern brook lamprey	Freshwater; non-parasitic derivative of <i>I. castaneus</i>	Gulf of Mexico drainages	
<i>Ichthyomyzon bdellium</i>	Ohio lamprey	Freshwater; parasitic	Ohio River drainage	
<i>Ichthyomyzon greeleyi</i>	Mountain brook lamprey	Freshwater; non-parasitic derivative of <i>I. bdellium</i>	As for <i>I. bdellium</i>	
Subfamily Lampetrinae				
Genus <i>Tetrapleurodon</i> (2 species)				
<i>Tetrapleurodon spadiceus</i>	Mexican lamprey	Freshwater; parasitic	Celio, Duero, Zula, and Lerma rivers, and Lake Chapala, Mexico	Also known as Chapala lamprey
<i>Tetrapleurodon geminis</i>	Mexican brook lamprey	Freshwater; non-parasitic derivative of <i>T. spadiceus</i>	Celio and Duero rivers, and Rio Grande de Morelia drainage, Mexico	Also known as Jacona lamprey

Table 2.1 (continued)

Classification	Common name	Life cycle type	Distribution	Comments
Genus <i>Entosphenus</i> (6 species)				
<i>Entosphenus tridentatus</i>	Pacific lamprey	Anadromous and freshwater; parasitic	Drainages of western Canada, USA and Mexico, and Japan. Widespread in the North Pacific Ocean	Formerly synonymized with <i>Lampetra</i> (see Sect. 2.4)
<i>Entosphenus minimus</i>	Miller Lake lamprey	Freshwater; parasitic	Upper Klamath River drainage, Oregon	
<i>Entosphenus similis</i>	Klamath lamprey	Freshwater; parasitic	Klamath River drainage, Oregon and California	
<i>Entosphenus macrostomus</i>	Vancouver lamprey	Freshwater; parasitic	Lake Cowichan drainage, Vancouver Island, British Columbia	Cowichan lamprey more precise than Vancouver lamprey (see Beamish and Wade 2008)
<i>Entosphenus folletti</i>	Northern California brook lamprey	Freshwater; non-parasitic	Klamath River drainage, California	Formerly considered by AFS to be synonymous with <i>En. lethophagus</i> (Robins et al. 1980), but recognized as distinct on recent list (Page et al. 2013a)
<i>Entosphenus lethophagus</i>	Pit-Klamath brook lamprey	Freshwater; non-parasitic	Klamath River drainage, Oregon, and Pit River, California	Appears to be recent non-parasitic derivative but not clear whether <i>En. tridentatus</i> or <i>En. similis</i> is the ancestor (see Sect. 2.2)
Genus <i>Lethenteron</i> (6 species)				
<i>Lethenteron camtschaticum</i>	Arctic lamprey	Anadromous and freshwater; parasitic	Drainages of Arctic and North Pacific oceans	Formerly synonymized with <i>Lampetra</i> (see Sect. 2.4)
<i>Lethenteron alaskense</i>	Alaskan brook lamprey	Freshwater; non-parasitic derivative of <i>Le. camtschaticum</i>	Drainages of Brooks and Chatanika rivers, Alaska, and Mackenzie River, Canada	Formerly known as <i>Lethenteron japonicum</i> (see Renaud et al. 2009b)

Table 2.1 (continued)

Classification	Common name	Life cycle type	Distribution	Comments
<i>Lethenteron appendix</i>	American brook lamprey	Freshwater; non-parasitic derivative of <i>Le. camtschaticum</i>	Great Lakes drainages and eastern USA, St. Lawrence, and Mississippi river drainages	
<i>Lethenteron reissneri</i>	Far Eastern brook lamprey	Freshwater; non-parasitic derivative of <i>Le. camtschaticum</i>	Drainages of Amur River, Sakhalin Island and Kamchatka Peninsula, Russia and in South Korea and Japan	
<i>Lethenteron kessleri</i>	Siberian brook lamprey	Freshwater; non-parasitic derivative of <i>Le. camtschaticum</i>	Drainages between Ob and Anadyr rivers, and of Sakhalin Island, Russia and Hokkaido Island, Japan	Also known as Siberian lamprey, but “brook” lamprey makes clear that this species is non-parasitic
<i>Lethenteron ninae</i>	Western Transcaucasian brook lamprey	Freshwater; non-parasitic	Drainages of the Black Sea	
Genus <i>Eudontomyzon</i> (6 species)				
<i>Eudontomyzon danfordi</i>	Carpathian lamprey	Freshwater; parasitic	Danube River drainage	Formerly synonymized with <i>Lampeira</i> (see Sect. 2.4)
<i>Eudontomyzon mariae</i>	Ukrainian brook lamprey	Freshwater; non-parasitic derivative of <i>Eu. danfordi</i>	Drainages of Baltic, Azov, Black, Adriatic, and Aegean seas	
<i>Eudontomyzon stankokaramani</i>	Drin brook lamprey	Freshwater; non-parasitic derivative of <i>Eu. danfordi</i>	Drainages of Adriatic Sea	Synonymized with <i>Eu. mariae</i> in Renaud (2011)
<i>Eudontomyzon morii</i>	Korean lamprey	Freshwater; parasitic	Yalu River drainage, China and North Korea	
<i>Eudontomyzon hellenicus</i>	Macedonia brook lamprey	Freshwater; non-parasitic	Strymon River drainage, Greece	Also known as Greek brook lamprey, but Macedonia brook lamprey distinguishes this species from the more recently-described <i>Eu. graecus</i>
<i>Eudontomyzon graecus</i>	Epirus brook lamprey	Freshwater; non-parasitic	Louros River drainage, Greece	

Table 2.1 (continued)

Classification	Common name	Life cycle type	Distribution	Comments
Genus <i>Lampetra</i> (9 species)				
<i>Lampetra ayresii</i>	Western river lamprey	Anadromous and possibly freshwater; parasitic	Drainages of North American Pacific Coast	Western river lamprey proposed as common name by Renaud et al. (2009b) and adopted by AFS, but also commonly referred to as river lamprey
<i>Lampetra pacifica</i>	Pacific brook lamprey	Freshwater; non-parasitic derivative of <i>La. ayresii</i>	Drainages of Columbia River, Oregon and Sacramento-San Joaquin rivers, California (but see Sect. 2.5)	Formerly considered by AFS to be synonymous with <i>La. richardsoni</i> (Robins et al. 1991), but recognized as distinct on recent list (Page et al. 2013a)
<i>Lampetra richardsoni</i>	Western brook lamprey	Freshwater; non-parasitic derivative of <i>La. ayresii</i>	Drainages of Pacific Ocean, British Columbia, Washington, Oregon and Alaska	Formerly referred to <i>Entosphenus</i> (see Sect. 2.4.2)
<i>Lampetra hubbsi</i>	Kern brook lamprey	Freshwater; non-parasitic	Friant-Kern Canal and Merced River, California	
<i>Lampetra aepyptera</i>	Least brook lamprey	Freshwater; non-parasitic	Drainages of northwestern Atlantic Ocean and Gulf of Mexico, USA	
<i>Lampetra fluviatilis</i>	European river lamprey	Anadromous and freshwater; parasitic	Drainages of northeastern Atlantic Ocean	Also known as Lombardy brook lamprey Sometimes referred to <i>Leihenteron</i> (see Sect. 2.4.2)
<i>Lampetra planeri</i>	European brook lamprey	Freshwater; non-parasitic derivative of <i>La. fluviatilis</i>	As for <i>La. fluviatilis</i> , plus Danube and Volga river drainages	
<i>Lampetra lanceolata</i>	Turkish brook lamprey	Freshwater; non-parasitic derivative of <i>La. fluviatilis</i>	Iyidere River, Turkey	
<i>Lampetra zanandreae</i>	Po brook lamprey	Freshwater; non-parasitic	Drainages of the Adriatic Sea	




## 2.2 Life Cycles and “Paired Species”

The ability to describe accurately a species of lamprey and thereby facilitate its allocation to the appropriate genus and family requires both a thorough understanding of the features that characterize the divergent larval and adult stages and recognition that, in some species, the morphology changes markedly during adult life. It should also be recognized that the types of life cycle vary amongst lampreys, with some containing a parasitic adult phase whereas others do not feed after the completion of larval life (see later).

The life cycle of all lamprey species contains a protracted larval phase that is spent in fresh water (Hardisty and Potter 1971a; Potter 1980a; see Chap. 3). The larva, termed an ammocoete, has a worm-like body shape and is blind and toothless (Fig. 2.1a). The ammocoete spends most of its time burrowed in the soft substrata in the slower-flowing regions of streams and rivers, feeding on the detritus and microorganisms (e.g., diatoms) that it extracts from the water overlying its burrow (Moore and Mallatt 1980; Yap and Bowen 2003). After typically between 3 and 7 years, the ammocoete undergoes a radical metamorphosis, which leads to the development of functional eyes, a suctorial disc and protrusible tongue-like piston (both of which are armed with teeth), and enlargement of the dorsal fins (Figs. 2.1b, 2.2, and 2.3; Hardisty and Potter 1971b; Potter 1980a; Youson 1980; see Chap. 4), with metamorphosis typically occurring at body lengths of 80–200 mm (Hardisty and Potter 1971a).

Following the completion of the larval phase, the life cycle of the lamprey diverges in one of two main directions. One course leads to the development, during metamorphosis, of a sexually immature young adult (Fig. 2.1b) that embarks on a parasitic feeding phase (Renaud et al. 2009a; Renaud and Cochran *in press*). The young adults of nine of these eighteen parasitic species feed at sea following a downstream migration. When fully grown they cease feeding and return to rivers, but not necessarily their natal systems, where they become sexually mature, spawn and die (Table 2.1; Hardisty and Potter 1971b; Potter et al. 2014; see Chap. 5). Five of these nine anadromous species have given rise to freshwater-resident or landlocked forms, whose immature adults feed in lakes or in the wider regions of large rivers (Table 2.1; Applegate 1950; Nursall and Buchwald 1972; see Docker and Potter *in press*). The remaining nine parasitic species are confined to fresh water and have essentially the same life cycle as the landlocked forms of anadromous species (Table 2.1; Hubbs and Trautman 1937; Chappuis 1939; Álvarez del Villar 1966; Renaud and Cochran *in press*). The maximum total length attained by parasitic species varies markedly, ranging from 145 mm in the freshwater Miller Lake lamprey *Entosphenus minimus* to 310–490 mm in small anadromous species, such as the western and European river lampreys (*Lampetra ayresii* and *La. fluviatilis*, respectively) to between 780 and 1,200 mm in the large anadromous pouched lamprey *Geotria australis*, Pacific lamprey *Entosphenus tridentatus*, and sea lamprey *Petromyzon marinus* (Oliva 1953; Vladykov and Follett 1958; Hardisty and Potter 1971b; Potter et al. 1983; Hardisty 1986; Lorion et al. 2000).



**Fig. 2.2** The suctorial disc and dentition of **a** a fully-metamorphosed *Mordacia mordax*, **b** an early upstream migrant of *Geotria australis*, **c** a young feeding adult of anadromous *Petromyzon marinus*, and **d** a recently-metamorphosed *Lampetra fluviatilis*. (Photos b–d: David Bird)

The second main direction exhibited by the lamprey life cycle involves a shifting in the timing of sexual maturation relative to metamorphosis, such that it commences during the transition from the ammocoete to the adult rather than after the completion of a parasitic phase as with the species above. The parasitic phase thus becomes eliminated and spawning takes place soon after the completion of metamorphosis (Hardisty 2006; Docker 2009). Consequently, these non-parasitic species breed at a length no greater than that of their longest ammocoetes. As most of these non-parasitic species are morphologically similar to a particular parasitic species in all aspects other than body size, it has been assumed that each evolved from that parasitic species (Potter 1980b; Docker 2009). On this basis, 15 of the 23 non-parasitic species listed in Table 2.1 can be “paired” with a congeneric parasitic species (in some cases, with a single parasitic “stem” species giving rise

Lampreys: Biology, Conservation and Control

Volume 1

Docker, M.F. (Ed.)

2015, XX, 438 p. 51 illus., 14 illus. in color., Hardcover

ISBN: 978-94-017-9305-6