

Chapter 2

Thermal Processing in Food Industries and Chemical Transformation

Maria Micali and Marco Fiorino

Abstract This chapter reviews thermal processes in the food industry—pasteurisation, sterilisation, UHT processes and others. It evaluates the effects on a chemical level and possible failures from a safety viewpoint and discusses in how far the effects can be predicted. In addition, historical preservation techniques—smoking, addition of natural additives, irradiation, etc.—are compared with current industrial systems, like fermentation, irradiation and addition of food grade chemicals. This chapter critically discusses storage protocols—cooling, freezing, etc.—and packing systems (modified atmosphere technology, active and intelligent packaging). Can undesired chemical effects on the food products be reliably predicted? This chapter elucidates on this important question. On that basis, new challenges, which currently arise in the food sector, can be approached.

Keywords Freezing · Modified atmosphere · Pasteurisation · Refrigeration · Semi-preserved food · Sterilisation

Abbreviations

CAS Controlled atmosphere storage
HTST High-temperature short-time
HHST Higher heat shorter time
HPP High-pressure processing
IQF Individual quick freezing
MTA Minimum temperature accretion
MAP Modified atmosphere packaging

M. Micali (✉)
Industrial Consultant, Messina, Italy
e-mail: marinamicali@virgilio.it

M. Fiorino
Technical Consultant, Siracusa, Italy
e-mail: marco.fiorino.ingegnere@gmail.com

RH	Relative humidity
TTI	Time temperature indicator
UHT	Ultra-High Temperature
a_w	Water activity

2.1 Introduction to Food Preservation Methods. High-Temperature Systems

Food preservation is crucial when speaking of food hygiene and health: each single preservation treatment is mainly designed with the aim of delaying the inevitable alteration of quickly perishable foods between production and consumption. Generally, food alterations are considered in relation with microbial spoilage, with the additional presence of potential pathogen agents. In addition, enzymatic reactions have an important ‘weight’ when speaking of food alterations, on condition that a certain amount of bioavailable water is present. As a result, food preservation should be designed and performed with two basic aims at least:

- The destruction of pathogen agents and the reduction of spoilage microorganisms
- The inactivation of microbial enzymes, if already synthesised.

The chosen preservation strategy (or strategies) should satisfy the following conditions:

- Treated food items have to be optimal, when speaking of microbial counts (low numbers) and freshness
- The respect of times is crucial. Treatments have to be carried on as early as possible to avoid the onset of microbial or enzymatic alterations
- Manipulation and general processing steps should be minimised because each step or sup-step without a clear preservation effect may be cause of microbial contamination, recontamination or spoilage
- Preliminary washing operations are performed when needed, if high-level superficial contamination is demonstrated and/or forecasted.

Food preservation methods imply the creation of unfavourable conditions to the development, activity and life of various pathogens, spoilage microorganisms and other life forms. On the other side, treated foods have to be and remain constantly safe and legally compliant until the end of labelled durability periods.

This chapter is dedicated to the description of many preservation systems: Sect. 2.1 discusses high-temperature methods, including pasteurisation, sterilisation and similar techniques. Subsequently, approaches based on low-temperature storages (refrigeration and freezing) are discussed. Finally, this chapter considers traditional preservation treatments and systems based on water removal. Different

techniques may be also used at present, including pulsed electric fields technology, ultrasounds and high-pressure treatments, but these technologies are not described in this book.

2.1.1 Pasteurisation: General Features

Basically, two different food preservation categories are performed with heat application: pasteurisation and sterilisation (Rahman 2007). The first method of preservation is defined which is able to destroy pathogenic life forms, and the most part of those vegetative microorganisms in foods (Alais 1984); moreover, enzymes are reported to be inactivated with pasteurisation treatments. Historically, the process is ascribed to Louis Pasteur in relation to his experiences in 1863 with 60 °C—treated wines and ameliorated shelf life periods (Alais 1984; Fry 2001; Rogers 2013).

In detail, pasteurisation has to be described as a microbiostatic and partially microbicidal treatment; for these reasons, it has to be synergically accompanied with other preservation methods such as refrigeration, addition of chemical preservatives, or vacuum packaging (Andersen et al. 1990; Degraeve et al. 2003; Mangalassary et al. 2008; Murphy et al. 2006). Anyway, heat treatments should be followed by the rapid cooling of treated products with the aim of avoiding storage temperatures that could favour enzyme-catalysed or simple chemical reactions. Generally, processing times depend on the nature and typology of foods: at present, low-temperature pasteurisation may be used when speaking of beers and wines, while milk for cheesemaking purposes must be heated at 75 °C for 15 s. On the other hand, high-temperature short-time (HTST) and high-temperature pasteurisation techniques are used for milk and fresh milk because of higher thermal values and lower processing times (Table 2.1).

Table 2.1 A brief overview of pasteurisation and sterilisation processes

Pasteurisation and sterilisation processes	Minimum temperature, °C	Maximum temperature, °C	Processing times
Low-temperature pasteurisation	60	65	30 min
High-temperature pasteurisation	75	85	≤ 3 min
High-temperature short-time (HTST) pasteurisation	75	85	≤ 20 s
Classical sterilisation (appertisation)	100	120	>20 min
Indirect UHT system	140	150	<10 s
Direct UHT system	140	150	<10 s

Low-temperature pasteurisation may be used for beers and wines, while milk for cheesemaking purposes is expected to be heated at 75 °C at least for 15 s. HTST and high-temperature pasteurisation techniques are used for milk and fresh milk. Appertisation can be used when speaking of canned vegetables and animal products. Indirect and direct UHT systems are recommended for aseptic packaged foods (UHT milk, soups, baby foods, etc.); direct UHT is preferred for low-viscosity products because of the use of injection or infusion techniques

Four different pasteurisation treatments are currently used in the food and beverage industry (Alais 1984; Murphy et al. 2003; Rahman 2007):

- (a) In-package pasteurisation. Temperature gradients can be used in certain situations. Fluids can be treated directly into bottles or cans. The process may be explained in the following way: the fluid is initially placed in the ‘cold zone’ of the container at about 2.2 °C; subsequently, it continues its diffusion towards hotter areas of the containers with a final 60 °C value. After a certain period at 60 °C, the fluid is progressively cooled until the final temperature is between 21.1 and 26.7 °C (Engelman and Sani 1983). In-package systems can be used with reliable effects on solid products such as ready-to-eat meat products with the aim of destroying pathogens and reducing post-process contamination caused by handling (Mangalassary et al. 2004)
- (b) Pasteurisation before the final packaging. High-temperature gradients are recommended when speaking of thermally sensible foods. For this reason, preheating stages are a useful option. Products such as processed juices and spreadable butter-like compositions may be treated with this system (Ahmed and Luksas 1988; Samani et al. 2016)
- (c) ‘Batch’ pasteurisation, also named low-temperature short-time pasteurisation. This system is specifically designed for fluids (milk, orange and apple juices, liquid whole eggs, etc.): the intermediate food is pasteurised at 62.8 °C for 30 min (Brant et al. 1968; Cinquanta et al. 2010; Cunningham 1986; Potter and Hotchkiss 1995; Wheeler et al. 1987). Interestingly, this treatment has been also used for waste raw materials into farms with good results when speaking of peculiar microbial contamination (Stabel 2001)
- (d) Continuous or HTST pasteurisation (Sect. 2.1.1.1). Fluids are treated at 71.7 °C at least for 15 s or more (heat exchange is required). This method is discussed in Sect. 2.1.1.1.

2.1.1.1 Pasteurisation: The HTST Method

The HTST method, also known as ‘flash’ pasteurisation, is performed (Rigo 2010) on liquid foods passing through a two-heated walls system (heat exchanger plates) with an empty space of 1 mm thickness only (temperatures between 75 and 85 °C, times: 15–20 s). Heat exchange occurs immediately in a uniform manner, while germs tend to adhere to surfaces (wall effect). Another similar system, ascribed to Stassano, implies the following time and temperature conditions: 14 s, 75 °C (Rigo 2010). Anyway, rapid cooling is required after heating (International Dairy Foods Association 2006). An interesting variation is the Higher Heat Shorter Time (HHST) system; it requires different machines and slightly different conditions (temperature >89 °C, holding time: one second or less) with consequent higher durabilities and enhanced enzymatic inhibition (Chandan 2015; International Dairy Foods Association 2006). Should HHST system be used, treated products would be sometimes defined ‘extended shelf life’ foods. The extreme solution is represented

by ultra-pasteurisation systems with temperature equal to 138 °C (minimum value) for at least 2 s (Chandan 2015). Shelf life values are enhanced: however, refrigeration is always required.

2.1.2 Sterilisation: General Features

Sterilisation is a more drastic treatment of food and non-food products, if compared with pasteurisation. The main difference is represented by the desired objective: the destruction of all microbial forms, including spores. Sterilisation may be defined as a commercial heat treatment, which can be able to destroy all living microorganisms in food during storage and distribution. On the other side, treated foods cannot be defined completely ‘aseptic’; additionally, organoleptic and nutritional characteristics are expected to suffer some alteration. It should be also noted that treatments can differ according to the pH of preserved foods (Sect. 2.3): in detail, intermediate foods with pH values < 4.5 should require temperatures around 100 °C, while products with pH > 4.5 would need at least 120 °C for 20 min (minimum time). In general, there are three main sterilisation treatments with different conditions, according to Table 2.1. These methods—classic sterilisation or appertisation, indirect and direct ultra-high temperature (UHT) systems—are discussed in Sects. 2.1.2.1 and 2.1.2.2.

2.1.2.1 Appertisation

Historically, the ‘appertisation’ method—asccribed to Nicholas Francois Appert (1804)—has been considered a low sterilisation procedure because of the use of open baths with temperatures ≤ 100 °C (Borde 2006): foods were packaged in sealed glass containers. At present, appertisation is performed with autoclaves, pressurised chambers with boilers for the production of saturated steam (temperature >100 °C) and devices allowing the modification of temperature and pressure values. Moreover, the evolution of used containers—metal packages above all—and the enhanced thermal conductivity have to be taken into account. For these reasons, the classical sterilisation can be very useful when speaking of packaged solid foods; liquid foods are usually pasteurised (Larousse 1991).

2.1.2.2 UHT Pasteurisation

The UHT method differs from the classical sterilisation because it reaches rapidly higher temperatures with very short duration (direct UHT system); as a result, organoleptic and nutritional properties of treated foods should suffer a few variations only. The UHT method also applies to bulk products (heat penetration is extremely rapid) and is associated with aseptic packaging techniques.

Two different UHT systems can be used (Andreini et al. 1990; Clare et al. 2005; Morales et al. 2000; Rerkrai et al. 1987):

- (a) Indirect method. In detail, thermal energy is supplied by means of an indirect steam injection system; fluids are forced to pass through close heat exchanger plates. It should be noted that thermal energy is diffused from the outside zone of UHT chambers to the inner 'core' (Corzo et al. 1994). As a consequence, external areas of the ideally considered fluid food may be superheated with possible variations (protein profiles, emersion of 'cooked' flavours because of the production of volatile sulphur components, etc.)
- (b) Direct system. The preheated food and the injected heating medium (steam) are in direct contact for a few seconds; subsequently, under-vacuum cooling is required, similarly to pasteurisation treatments, with the aim of eliminating extra water (a certain volumetric augment is observed in the process). The reduced time of contact explains apparently the absence of serious damages, from the sensorial viewpoint (Datta et al. 2002).

A brief comparison of indirect and direct UHT methods seems to highlight the importance and advantages of the direct injection: sensorial variations seem reduced; on the other hand, these techniques may cause localised burns. An interesting 'infusion' steam version—the fluid food is infused into steam—is reported to minimise this defect (Datta et al. 2002).

Nutritional variations may be observed; on the other side, it may be affirmed that digestibility is improved. For these reasons, different non-heat-based systems (freezing methods) are apparently preferred at present.

2.2 Cold Preservation

This method of food preservation is based on the ability of cold to interfere with the vital functions of microorganisms: essentially, cold systems can assure a reliable microbiostatic action.

The cold slows down until it stops degradative (chemical and enzymatic) reactions inside the food. This effect is mainly due to the inhibition of microbial enzymes: they are not denatured and therefore inactivated, but their action is inhibited (rendered useless) in relation to the low temperature of the food. Moreover, the inhibitory action of low temperatures occurs with the subtraction of free water from the system because of ice formation with the consequent increase in the concentration of extracellular solutions and the dehydration of microbial cells.

The minimum temperature accretion (MTA) of microorganisms is needed in order to effectively maintain cold conditions. MTA of pathogenic microorganisms is always >0 °C, while psychrophilic life forms remain viable at lower temperatures. The microbial growth can be considered negligible at -18 °C and lower thermal values. Moreover, cold has a partial microbicidal action: a number of

microbes suffer the irreversible alteration of cytoplasmic proteins and mechanic damages (formation of ice crystals) during the early stages of freezing treatments; however, surviving life forms may be ready to resume vital functions with spores and toxins, if thermal values are increased. For these reasons, foods should have excellent or good hygiene conditions before treatments.

2.2.1 Cold Storage: Different Preservation Techniques

2.2.1.1 Refrigeration

There are two different methods of food preservation by cold storage (Fig. 2.1): refrigeration and freezing systems. The first method, refrigeration preservation, concerns the storage of food commodities at temperatures low enough to slow down degradative (chemical and enzymatic) reactions inside stored products; in addition, water must remain liquid.

The choice of storage temperatures depends on the type of food and the required retention time: anyway, storage times are not excessive. Generally, refrigeration temperatures are between -1 and 8 °C. However, refrigerated storage shows good performances on condition that other factors are controlled; ventilation has to be




Fig. 2.1 A brief overview of cold preservation techniques. The first method, refrigeration, concerns the storage of food commodities at temperatures low enough to slow down degradative reactions inside stored products; in addition, water must remain liquid. Usually, temperatures remain between -1 and 8 °C. On the other side, freezing systems involve the exposure of foods and beverages to low or very low temperatures with the consequent water crystallisation and the solidification of the whole product. Four main systems—slow, rapid, flash and deep freezing—are used at present with different temperatures and processing times

assured into storage chills. Moreover, acceptable relative humidity (RH) values are between 80 and 95 %: reduced RH values may cause excessive water evaporation from food surfaces, while excessive RH may favour the development of moulds (visible mycelia).

Anyway, refrigerated storage can be used in different situations and for the following food categories at least:

- (a) Refrigerated products during transport and storage only (sold at room temperature): fruits, vegetables, eggs
- (b) Edible products distributed, stored and sold between -1 and 8 °C
- (c) Prepared and precooked foods: salads, sandwiches, fresh pastries, precooked and delicatessen products, meats and cold cuts.

Consequently, the diffusion of refrigeration systems has been progressively introduced everywhere in the last decades with excellent results (Bruhn and Schutz 1999), despite several problems may be observed when speaking of food safety practices in consumers' homes (Karabudak et al. 2008; Redmond and Griffith 2003).

Historically, the use of natural and artificially produced snow and ice was the first homemade application of food cold storage. With reference to industrial applications, refrigeration was mainly based on the liquefaction by compression of several allowed gases such as ammonia, sulphur dioxide, carbon dioxide and chloroform (Feist and Hadi 1965; Reif-Acherman 2009) and the concomitant removal of heat from a surrounding area. These 'static refrigerators' are not used at present: the current evolution (dynamic refrigerators) concerns the continuous circulation of cold air between different areas; moreover, frost formation is avoided by means of forced ventilation and thermal changes caused by frequent door openings are remarkably reduced.

In the industrial field, at the end of the preparation or the last heat treatment, foods or intermediate products should reach refrigeration temperatures very quickly because microbial spreading is favoured between 3 and 70 °C. Consequently, 'blast chillers'—discontinuous (cabinets) or continuous (tunnel) systems—are frequently used: refrigeration conditions are reached by means of the introduction of cold air (forced ventilation at about 1 °C) or cold water. Subsequently, treated foods have to be stored in 'permanent' chillers and/or dedicated trucks or rail cars capable of self-cooling. By the legal viewpoint, the transportation has to be carried out with the absolute warranty (digital or printed records) of thermal values below or equal to -18 °C (only for frozen products and long distances). Otherwise, simple small trucks with thermally insulated walls are required.

2.2.1.2 Freezing Techniques

Basically, freezing involves the exposure of foods and beverages to low or very low temperatures with the consequent water crystallisation and the solidification of the

whole product. A food is defined 'frozen' when 80–90 % of the total water amount is turned into ice.

It has to be noted that freezing methods affect only free water; a small fraction of aqueous amounts (2–5 %) is electrostatically bound to sugars, proteins and solutes; this water is also named 'non-freezing water' (Ali and Bettelheim 1985). Consequently, foods and beverages—liquid, solid or fluid water solutions—have to reach temperatures below 0 °C (the cryoscopic point or frost point of pure water at atmospheric pressure) for freezing purposes. Generally, the freezing point of free water is initially between –4 and –0.5 °C, but the freezing treatment should obtain a 'core' temperature between –10 and –25 °C (end of the process). In fact, a notable fraction of free water is constantly separated from the original food solution in form of ice. The end of the process is reached when 80–90 % of the total water content is turned into ice: the remaining non-freezing solvent and a certain amount of free water are also present. Interestingly, the residual free water (with a notable concentration of dissolved molecules) is located in very small channels and may be also able of free movement through the treated food.

Basically, the freezing process may be subdivided into two steps:

- (a) Nucleation and crystallisation. Temperature values: between 0 and –7 °C. Ice crystals are initially formed; this step implies the maximum separation of water in the solid state from the remaining food system, including non-freezing and residual free water
- (b) Crystal growth. Each ice crystal tends to augment its dimensions with the incorporation of other water molecules around the initial nucleus.

It has to be noted that the dimension of final ice crystals is inversely proportional to the number of initial ice nuclei. Moreover, the number of ice crystals depends on the freezing speed. Basically, two different situations can be observed (Zhao 2007):

- (1) Slow freezing. The food is subjected to temperature values between –8 and –20 °C with a low penetration speed. Should this be the situation, the initial number of ice crystals would be reduced (poor nucleation and crystallisation). Consequently, strong crystals of considerable size are easily observed. Unfortunately, the formation of a reduced number of high-scale ice crystals is the main cause for micro- and macro-damages in the food because macro-crystals easily damage cell walls (vegetable and animal tissues). Once defrosted, the food does not return to the original texture and fibrous-like products can be obtained; additionally, unpleasant flavours are often reported. For these reasons, the method is not recommended. Recently, new similar systems have been proposed: high-pressure freezing and dehydrofreezing (initial dehydration of foods and subsequent freezing), with the aim of avoiding volumetric augments and tissue damages (Li and Sun 2002)
- (2) Rapid freezing (required systems: air freezers; plate freezers; liquid-immersion freezers). The food is subjected to temperatures between –30 and –50 °C with notable penetration speed. Should this be the situation, the initial number of ice crystals would be remarkable (good or excellent nucleation and

crystallisation). Consequently, many microcrystals are easily observed (a few μm only). The formation of a notable number of small-scale ice crystals can be very useful because microcrystals do not easily damage cell walls (vegetable and animal tissues). Once defrosted, the food may return to the original texture with potentially small variations: fibrous-like products are not reported when speaking of rapid freezing. Sensorial features are substantially preserved: higher shelf life values have been also demonstrated. For these reasons, the method is recommended in the industrial ambit. The so-called ‘individual quick freezing’ (IQF) system is extremely useful when foods are not packaged. However, the main problem remains the superficial dehydration with related weight loss between 2 and 3 % (Pruthi 1999; Senesi 1984)

- (3) Flash or cryogenic freezing. The food is subjected to temperatures below $-50\text{ }^{\circ}\text{C}$ with high penetration speed. Should this situation be observed, the initial number of ice crystals would be higher than expected in comparison with rapid freezing. Consequently, the performance of this method is reported to be excellent in comparison with rapid freezing, despite some observation with concern to the real necessity (Farouk et al. 2004). The total treatment time is reduced (maximum time: some hours, depending on the dimensional features and composition of products). Once defrosted, the food may return to the original texture with potentially small variations: fibrous-like products are not reported when speaking of rapid freezing. Sensorial features are substantially preserved: higher shelf life values have been also demonstrated. For these reasons, this method is recommended in the industrial ambit. Generally, the use of liquid nitrogen or solid carbon dioxide is required in flash (cryogenic) systems because products are usually placed on a conveyor belt into dedicated tunnels and subsequently cooled with sprayed or dribbled cold gas. Subsequently, the product must reach the final temperature. Dehydration is substantially negligible
- (4) Deep freezing. This technique is discussed in Sect. 2.2.1.3.

The duration of treatments can vary, depending on the volume and thickness of the product, from a few minutes to a few hours.

2.2.1.3 Deep Freezing

Deep-frozen foods are products subjected to a special freezing process: the zone of maximum crystallisation has to be reached quickly and maintained constant in the whole product at temperatures $\leq -18\text{ }^{\circ}\text{C}$ until the end of durability periods. The mentioned thermal value is compulsory from the legal viewpoint. This process, also named ‘ultrafast’ system, can be performed with air or plate freezers: the only exception concerns the immersion in non-freezing liquids. Requisites are:

- The inner temperature (at the centre of the whole product) has to be minor or equal than $-18\text{ }^{\circ}\text{C}$
- The product has to be continually kept at temperatures $\leq -18\text{ }^{\circ}\text{C}$ until the sale to the final consumer.

Moreover, stricter conditions may be applied in several stages because of inevitable thermal changes in some point of the whole food chain. In particular, factory warehouses may be kept at $-30\text{ }^{\circ}\text{C}$ before the delivery of frozen foods, while intermediate storage chillers (in all possible points of the food chain except for the initial production plant) may remain at $-25\text{ }^{\circ}\text{C}$ value (subcooling).

It should be also noted that:

- (1) The use of peculiar packaging materials is strictly required when speaking of deep-frozen foods. These food packaging materials include aluminium (alone or combined with plastic films), tinplate, cardboard, coupled plastic films (nylon/polyethylene, polypropylene/polyethylene)
- (2) The label must contain all required information for the proper use of the food, in accordance with official Regulations.

2.3 Non-thermal Food Preservation Methods

Food products can be preserved by means of thermally based preservation methods or different (non-thermally conducted) systems. Actually, this subdivision is not so clear because a peculiar food product can be subjected to two different preservation systems. Basically, the main difference between above-mentioned procedures is the definition of ‘thermal processing method’. It may be inferred that all processes implying the use of heat (or heat removal) can be defined ‘thermal’ procedures.

Sterilisation procedures concern only preserved foods, which are packaged in a hermetically sealed container and have undergone a specific heat treatment or other authorised treatments with the aim of inactivating irreversibly dangerous enzymes and microorganisms. Two basic categories of sterilised foods (Sect. 2.1.2, Table 2.1) are known: acidic products with $\text{pH} \leq 4.5$ and non-acidic foods with $\text{pH} \geq 4.5$ (Soncini 2008).

On the other hand, semi-preserved foods are products whose stability over time depends on the preservation method and environmental conditions. In detail, the following products can be defined semi-preserved foods:

- (a) Pasteurised foods with $\text{pH} \geq 4.5$. These products are packaged in a hermetically sealed container, pasteurised (or subjected to analogous treatments with lower performances if compared with sterilisation) and their shelf life depends on environmental conditions (low temperatures, use of modified atmosphere, etc.). For these reasons, pasteurised semi-preserved foods have to be refrigerated

- (b) Non-thermally treated semi-preserved foods. These products are packaged in more or less hermetic containers and have undergone a specific technological treatment without heat addition. In addition, their performance and safety depend on storage conditions. Allowed technological systems are: smoking, drying, lyophilisation and addition of natural substances with inhibiting power. These products are normally stored as refrigerated or frozen foods.

The main objective of this chapter is the comparison of different preservation systems and techniques. Because of the existence of different foods and beverages belonging to the category of preserved (sterilised) products and the more complex world of semi-preserved foods, a critical comparison of different systems should be carried out. In addition, the association between several peculiar methods of technological preservation and selected thermal methods should be explained with reliable argumentation. Next sections are dedicated to a brief overview of non-thermal preservation systems (Fig. 2.2) and the explanation of synergic effects between these methods and thermal procedures.




Fig. 2.2 A brief overview of non-thermal preservation techniques. The main difference concerns the use of traditional preservation systems such as the immersion in salt solutions, sucrose, ethyl alcohol, vegetable oils and vinegar, or the use of ‘historical’ methods such as smoking, drying and fermentative processes. Actually, other systems may be considered in this category with the aim of discriminating them in comparison with high-technology procedures: dehydration, lyophilisation and blanching. High-technology preservation systems—MAP, CAS, intelligent packaging, superchilling, etc.—are often used in modern plants and distribution services

2.3.1 *Modified Atmosphere Packaging Technologies*

2.3.1.1 MAP Systems: General Features

Modified atmosphere packaging (MAP) techniques are one of the most recent and interesting applications in the field of food preservation. Basically, this technology can enhance shelf life values for certain products and reduce the addition of food preservatives.

Briefly, food degradation caused by chemical–physical reactions and microbiological spreading (enzymatic reactions are also included) can be delayed or inhibited to a certain extent by means of the simple modification of atmosphere compositions in the final food packaging (Micali et al. 2009; Parisi 2009, 2012). In this ambit, two possible MAP technologies can be used for food applications:

- (1) ‘Active’ MAP systems. The existing gas mixture into the container is removed and a different gaseous composition is inserted before packaging
- (2) ‘Passive’ MAP methods. The composition of food packaging materials is designed to allow the continuous modification of inner atmospheres caused by cellular ‘respiration’. This phenomenon is observed when speaking of fresh (minimally processed) fruits and vegetables, with special attention to cut products. Cellular respiration causes the continuous increase of carbon dioxide at the expense of oxygen; consequently, exceeding carbon dioxide has to be removed (Senesi 1984).

The effectiveness of MAP systems depends on the ‘empty’ volume occupied by the gaseous mixture (it should be equal to the space occupied by the packaged food). In addition:

- (a) The composition of modified atmospheres depends on the peculiar food. Generally, most used gases (in different proportions) are carbon dioxide, oxygen, nitrogen and/or argon; in particular, carbon dioxide may be extremely useful (Molin 2000). Possible mixtures have to be designed on the basis of the microbial ecology and approximate chemical compositions of packaged foods (Ercolini et al. 2006a). Some specific parameters such as water activity (a_w), pH and redox potential have to be taken into account; the latter of these measures, in particular, can be very useful to try to determine the probability of growth and survival of pathogenic and spoilage microorganisms (Delia et al. 2005; Parisi et al. 2004; Senesi et al. 2000)
- (b) The composition of packaged foods has to be taken into account, specifically with relation to fatty and moist foods and possible irregular agglomerations into MAP products (Sect. 2.3.1.2)
- (c) The augment of food contact surface is critical. As an example, diced cheeses (Sect. 2.3.1.2) can show enhanced shelf life values if compared with whole cheeses in the same MAP condition because the food/atmosphere interface is dimensionally increased. This interesting result is not expected normally

because cuts or different mechanical operations aiming to separate or subdivide solid foods cause generally lower durability values (Delia et al. 2005; Parisi 2002, 2009; Parisi et al. 2004).

2.3.1.2 MAP Systems: Technological Details

The ‘empty’ volume occupied by the gaseous mixture in MAP products has a remarkable importance. This space should be equal to the space occupied by the packaged food. However, this condition is not always possible because of the physical disposition of the edible mass into the package. In other terms, subdivided food products can physically occupy part of the theoretical empty volume into the MAP container. This situation has been reported clearly for MAP diced mozzarella cheeses (Parisi et al. 2004). This product tends to occupy large volumes of the inner package because of the irregular disposition of cheese cubes and the fatty composition of products, with consequent agglomeration and partial ‘fusion’ between single cheese parts.

MAP technology requires the combination of three interdependent elements: the packaging machine, the type of wrapping material and the composition of gaseous mixtures. In detail, the release of modified atmospheres (after the removal of pre-existing gases into the container) can be observed immediately before, during or immediately after the placement of whole or subdivided foods. With relation to gaseous mixtures, the so-called ‘reactivity’ or ‘inertness’ depends on the abundance of three main gases: carbon dioxide, nitrogen and oxygen (Micali et al. 2009). Should nitrogen be mainly present, the inner gas mixture would be defined ‘inert’; on the contrary, mixtures with carbon dioxide and/or oxygen (with the exclusion of inert nitrogen) are defined ‘reactive’. Intermediate compositions (carbon dioxide and nitrogen, with or without oxygen) are considered mixtures with a limited ‘reactivity’ by the technological viewpoint (Micali et al. 2009).

Substantially, it can be affirmed that MAP technology offers excellent results in terms of food durability, if compared with known traditional techniques such as vacuum packaging solutions. In addition, sensorial features remain acceptable. As a result, many different foods can be preserved in this way: baked products, pre-cooked foods, coffee, fruit juices, wines, chips, processed meats, etc.

MAP advantages are certainly referred to the commercial stability and the safety of products; at the same time, long shelf life values allow the optimisation of logistic operations and a more efficient management of transportation costs without the use of declared preservatives. Moreover, the ‘thorny’ matter of waste foods (expired products) may be partially solved.

However, the MAP technology cannot assure the complete safety and hygiene of treated food products without a solid and reliable basis: the use of good or excellent raw materials and packaging materials (and objects). In other words, the basic ‘support’ for MAP systems (and each preservation technique) is always the initial raw material (or the sum of used raw materials). As above-mentioned, some

chemical–physical data such as pH, a_w and redox potential have to be taken into account. In detail, redox potentials can be extremely useful because of the direct connection between these values and the probability of microbial spreading for selected microorganisms.

The determination of redox potentials in foods is recommended because of the connection between the increase towards higher values and the concomitant decline in the multiplication of life forms such as *Listeria monocytogenes*, *Escherichia coli* and *Salmonella* spp. Because of the influence of the edible ‘culture medium’ (the food itself) on the redox behaviour, several foods are not good enough when speaking of microbial spreading; as a consequence, these foods could be treated with MAP technologies with the aim of maintaining initial redox potentials in an unfavourable way for many anaerobic bacteria (Delia et al. 2005).

Generally, the first objective of MAP systems is the reduction of contacts between oxygen and food surfaces. In fact, oxygen is mainly responsible for the promotion of microbial spreading (aerobic or facultative anaerobic life forms). On the other hand, this gas could be useful in certain situations: as an example, the bright red colour of oxyhemoglobin in red meats and fish products can be notably enhanced in presence of oxygen; otherwise, the formation of meta-myoglobin would be easily expected with unacceptable modifications of colorimetric performances (Delia et al. 2005). However, oxygen can favour lipoxidation and fat rancidity in medium-fat and fat fish such as salmon and tuna fish. In addition, food degradation may be ascribed to the oxidant action of this gas, including the degradation of β -carotene.

On the other side, a real antimicrobial action is ascribed to carbon dioxide (this effect is maximum when the related concentration exceeds 30–40 % of the total gaseous mixture). Naturally, the efficacy of carbon dioxide depends also on the microbiological state of treated foods, in terms of initial microbial counts. On the other side, high concentrations of this gas may lead to fractures in the packaging material (collapse) with consequent severe alterations of sensorial features. The explanation of this strange phenomenon is mainly ascribed to the absorption of hydrated carbon dioxide (carbonic acid) in foods, with the concomitant pH lowering and the decrease of inner pressures into packages (Delia et al. 2005). pH low values are efficient enough in terms of microbial inhibition; in addition, the production of biogenic amines is delayed (Micali et al. 2009). The effect of this gas on vegetable products is substantially the inhibition of cellular respiration; moreover, maturation processes are delayed and possible damages to vegetable tissues because of cold temperatures are notably reduced (Micali et al. 2009). With reference to negative effects, it should also be remembered that excessive amounts of carbon dioxide may cause dehydration in meats.

Nitrogen is completely inert: as a consequence, there are no possible interaction with organic substances. Basically, this gas is used as a filler with the aim of preventing possible packaging collapses, if carbon dioxide is used and possibly dissolved in foods. Because of the complete inertness, pH is not affected.

Other possible gases are:

- (a) Carbon monoxide. This toxic gas has not inhibitory effect on microorganisms. However, its use has been often reported because of positive sensorial effects such as the formation of carboxymyoglobin with red colour enhancement in certain products (beef, fish). On the other hand, carbon monoxide is questioned because of possible effects on food operators. In addition, the above-mentioned formation of red-coloured carboxymyoglobin may be used for fraud purposes
- (b) Argon and other noble gases. These chemical elements may be a good alternative because of the substantial inertness, colourless, odourless and tasteless. In addition, gases such as argon can prevent crushing and deformation damages in MAP products.

2.3.1.3 MAP Systems: Synergic Applications with Thermal Processes and Possible Failures

The use of MAP systems can be very useful if these technologies are compared to more traditional (air or vacuum packaging) techniques. At present, many applications are possible when speaking of pasteurised and cooked foods: meat and sausages, baked foods, precooked foods, cheeses, milk powders, coffee, fruit juice, wine, chips, pretzels, etc. The additional removal of food preservatives has to be highlighted.

On the other side, possible failures of MAP systems can be caused by:

- Initial conditions of foods and beverages. In other words, incorrectly stored, manipulated or pasteurised foods cannot be well preserved with MAP (or vacuum) technologies. In particular, the possibility of certain greenish colours (Kröckel 1995) on the surface of processed meats may be favoured when using peculiar MAP gases
- Strange behaviours of packaged foods. In detail, certain sensorial variations have been often observed and correlated to the dissolution of MAP gases (carbon dioxide) in foods, where possible (cheeses)
- Inner or external package fractures caused by the permeability of food packaging materials, with consequent lower pressure into the MAP product (Parisi 2012; Sivertsvik et al. 2002). Other situations depend on pre-existing food packaging failures (Parisi 2013) or processing errors (during the replacement of the inner atmosphere)
- Incorrect storage temperatures (these products have to be stored in refrigerated chillers at least).

The above-discussed failures can be also observed in vacuum packaged foods: naturally, the frequency of possible defects, including food safety and hygiene situations, depends on the peculiar preservation system. In other terms, MAP

methods aim at modifying shelf life values of foods by means of the controlled variation of inner atmosphere; on the contrary, vacuum packaging techniques aim at inhibiting the development of microbial spreading and chemical–physical reactions based on the necessary presence of oxygen.

2.3.2 Controlled Atmosphere Storage Techniques

2.3.2.1 CAS Systems: General Features

Because of the effect of oxygen on processed and fresh foods (Sect. 2.3.2), another preservation system—controlled atmosphere storage (CAS)—can be considered into large-scale food warehouses in synergy with refrigeration procedures. In brief, the air inside refrigerated cells can be ejected and replaced with mixed gases, depending on the type of food product. The composition of ‘artificial’ gas mixtures is constantly monitored and kept constant by means of automatic control systems (analysers, absorbers and gas generators). Generally, atmospheres should have the following rough composition: nitrogen (92–95 %); carbon dioxide (2–4 %); oxygen (3–4 %); the temperature of chillers should not exceed 3–4 °C. Basically, the main modification concerns the substitution of oxygen with nitrogen, carbon dioxide or their mixtures (Sect. 2.3.1). CAS systems are mainly used in food warehouses and sea freight containers for apples, pears, citrus fruits, other fruits and vegetables, and flowers. Stored products are generally sensible to alterations because of cellular respiration phenomena and the consequent production of carbon dioxide, water vapour and several organic and aromatic compounds.

2.3.2.2 CAS Systems: Synergic Applications with Thermal Processes and Possible Failures

The CAS system may avoid the early degradation of fresh products if combined with cold-based technologies. In detail, the production of ethylene is reduced; the degradation of chlorophylls may be blocked; pectins may be preserved at some extent from hydrolysis. Stored products can be subsequently used for different productions and other preservation treatments, including thermal processing. With reference to possible defects, observed failures are generally correlated to the incorrect management of process parameters: chiller temperatures, adequate ventilation, surveillance and the continuous modification of inner atmospheres into chillers.

2.3.3 Vacuum Packaging and Hypobaric Packaging Systems

2.3.3.1 Vacuum and Hypobaric Packaging Systems: General Features

The amount of oxygen inside food packages may be also reduced if the inner pressure is lower than 2.66×10^3 Pa or between this value and 101,300 Pa (atmospheric pressure). Substantially, two different methods may be used in this ambit:

- (1) Vacuum packaging. The inner pressure is lower than 2.66×10^3 Pa. Recommended containers for these methods can be pouches, special rigid containers and thermoformed trays. The virtual absence of oxygen does not favour the aerobic activity of many microorganisms; consequently, deterioration processes are remarkably delayed, and sensorial features (appearance, colour, aroma, flavour and texture) suffer some little modification. The same thing can be affirmed for chemical and nutritional features (digestibility, protein amount, vitamins and starches). Moreover, environmental contaminants are removed with the originally contained air into the package
- (2) Hypobaric packaging. The inner pressure is between this value and 101,300 Pa. Performances can be interesting enough in terms of shelf life values, sensorial expectations and safety, although vacuum packaging is considered with more favour.

2.3.3.2 Vacuum and Hypobaric Packaging Systems: Synergic Applications with Thermal Processes and Possible Failures

Vacuum and hypobaric methods can be used in association with low temperatures (refrigeration) and thermal processes in general (including the simple steam cooking at 80 °C, for catering purposes) with excellent performances.

In particular, vacuum packaging meets fundamental food safety requirements. The main advantage is the extension of shelf life values: products are kept longer under refrigeration conditions and during the commercial transportation. Moreover, the possible addition of chemical preservatives, salt or strong spices can be advantageously replaced or reduced. Another important advantage, from the economic angle, is correlated with the virtual absence of weight loss because of evaporation and liquid release.

On the other side, vacuum techniques may be correlated with the possible modification of aroma and the limited evaporation of moisture and volatile compounds. However, the main failure for vacuum and hypobaric packaged foods is always correlated with the use of food packaging materials in terms of possible damages before use, incorrect design/choice and poor 'barrier' properties (Parisi 2012).

2.3.4 *Smart and Intelligent Packaging Systems*

2.3.4.1 **Smart and Intelligent Packaging: General Features**

The increasing demand for fresh products and the consequent necessity of reliable demonstrations—including also new and redeveloped analytical techniques (Parisi 2016)—has progressively stimulated the development of the so-called ‘functional’ packaging. Generally, these packaging materials are subdivided in two different categories: ‘active’ and ‘intelligent’ packaging systems (Parisi 2009), depending on the related function.

Briefly, ‘active’ or ‘smart’ packaging materials are designed with a very specific aim: the continuous (active) modification of food and atmosphere properties during the whole shelf life period (Parisi 2009) with the consequent enhancement of positive food properties, including safety. This objective can be achieved by means of the chemical interaction with the packaged food and the gas mixture inside the package. Smart systems can allow the possibility of an active control because of the direct and predictable interaction with packaged foods. As a result, food technologists may be able to influence the complex phenomena of chemical, microbiological, physical and mechanical modifications occurring during storage. Finally, it should be remembered that modern packaging materials have also a clear marketing function (Brunazzi et al. 2014); consequently, the use of smart systems should be evaluated in relation to preservation purposes and marketing strategies (Parisi 2012).

On the other side, ‘intelligent’ packages are designed with the aim of reporting product changes during the whole shelf life period, with the consequent possibility of diminished freshness and/or durability. The expression ‘intelligent packaging’ means a packaging technique that involves the use of an internal or external indicator, capable of actively representing the history of the product and the remaining shelf life (Parisi 2009).

The correlation of smart and intelligent systems with thermal processing is not apparently clear. Both solutions require the use of other preservation strategies, including thermal processing systems for unpackaged foods and cold-based techniques (refrigeration). Thermally processed foods can be packaged with modified atmospheres and interact with active compounds such as antimicrobials, antioxidants or other useful substances. The continuous modification of food composition can also be obtained by means of the variation of gaseous mixtures (head-space control): small accessories containing gas absorbers or scavengers (target gases: oxygen, ethylene, moisture) or generators of oxygen, carbon dioxide or ethanol can be used (Parisi 2009). In detail, the following systems are currently reported:

- Moisture absorbers (glycerol, silica gel, clays) for meats
- Humidity regulators (sodium and potassium chloride)
- Ethylene absorbers (activated carbon, zeolites) or emitters of carbon dioxide (ascorbic acid) for vegetables

- Carbon dioxide absorbers for packed coffee (calcium chloride and sodium or potassium hydroxide)
- Ethanol emitters for baked goods.

Moreover, the active packaging may induce natural antimicrobials such as essential oils or bacteriocins: these substances can inhibit microbial growth in foods (Ercolini et al. 2006b; Mauriello et al. 2004; Murray and Richard 1997; Ouwehand 1998; Skandamis and Nychas 2001; Tassou et al. 2000). However, strict management requirements are needed, and storage procedures (good manufacturing practices, high hygienic standards) are also required.

Anyway, active systems cannot fully give expected performances without reliable traditional preservation techniques. Cold storage is required: in fact, the design and the implementation of active approaches are based on the preventive application of refrigeration systems. Sometimes, these applications have not been recommended when speaking of frozen foods.

The intelligent packaging has to interact with the environment and the whole food/packaging system: recorded variations such as temperature abuses should be immediately visible by means of evident chromatic variations. Time temperature indicators (TTI) represent the most widespread forms of commercially available intelligent packaging. TTI are normally applied on the outer surface of packages: the thermal history of the whole food/packaging system is recorded and made available readily (Parisi 2009). In addition, some commercial application of oxygen and carbon dioxide indicators already exists: these applications—tablets or strips into containers—can vary their colour because of the effect of specific chemical and enzymatic reactions on the composition of inner atmospheres. Otherwise, a destructive analytical procedure would be required.

2.3.4.2 Smart and Intelligent Packaging: Synergic Applications with Thermal Processes and Possible Failures

Because of the peculiar design, smart and intelligent systems should be used if coupled with cold-based storage methods. For this basic reason, related advantages should be considered simply in terms of performance enhancement for cold-stored foods or refrigerated products packed with modified atmospheres. On the other hand, failures can be easily defined as the opposite of predictable (desired) results:

- (a) Insufficient modification of food properties
- (b) Insufficient modification of atmosphere properties
- (c) Inaccurate demonstration of thermal history
- (d) Inaccurate demonstration of qualitative variations (with relation to foods and/or gas mixtures).

Moreover, each mechanical damage caused by one or more of the above-mentioned defects can be ascribed (at first sight) to different reasons with the exclusion of smart devices. Actually, the simple insufficient modification of gaseous

compositions into packages can easily cause packaging defects such as fractures or container collapsing (Parisi 2012).

2.3.5 High-Pressure Treatment

An interesting non-thermal preservation process is based on the application of high pressures (Hugas et al. 2002; Patterson 2005; Simpson et al. 2012). This technique, generally named high-pressure processing (HPP), shows very good performances when speaking of food preservation because of the effect on several spoilage bacteria and pathogens (destruction) and some enzymes (inactivation). HHP may significantly reduce the presence of *Salmonella* spp and *L. monocytogenes* in raw meats and marinades.

HPP macroscopic effects on processed foods are different (pressures >200 MPa with maximum values >500 MPa):

- (1) Inactivation and death of microorganisms
- (2) Structural and functional modification of biopolymers (proteins, enzymes and polysaccharides). In particular, the tridimensional structure of enzymes protein molecules suffers important variations (Simpson et al. 2012)
- (3) Morphological changes (vacuolar gas compression, deformation/elongation)
- (4) Altered metabolism and biochemical reactions
- (5) Changes in the cell membrane (altered permeability).

The use of high pressures (400–700 MPa) on foods causes the permeabilisation of cell membranes and other phenomena such as the probable crystallisation of membrane phospholipids. Anyway, the microbial resistance to high pressures depends on several factors: the type of microorganism, its stage of development, process conditions [time, pressure, temperature, chemical parameters of treated foods (a_w , pH, etc.)].

HHP is recommended for the microbial stabilisation of acid and heat-sensible foods, the sanitisation of non-acidic products and the change in the technological functionality of ingredients. In addition, solid and liquid foods can be treated with HPP techniques; post-contamination is minimised; cleaning and sanitising procedures in food plants are easier.

With relation to potential resistances, it has to be highlighted that viruses and spores are very resistant to HHP; on the other side, HHP shows excellent performances with non-sporeforming pathogens (*L. monocytogenes*, *Staphylococcus aureus*, etc.).

Other potential disadvantages of the HPP system for packaged foods concern substantially the complex manipulation of foods, the limited choice for containers, long cycle times and the expensive cost of HPP lines. Actually, non-packaged foods can be HPP treated with interesting advantages: easy handling, possibility of many different containers and the absence of loading and unloading times; the process

may be defined ‘continuous’ (Parisi 2005). On the other side, only liquid and semi-liquid foods can be considered; in addition, aseptic packaging systems are required.

2.3.6 Irradiation

The so-called ‘cold pasteurisation’ has certainly interesting effects on enzymatic reactions and microbial activity. Basically, the exposure of foods to ionising radiations (X-rays, accelerated electron beams and γ -rays) enhances the shelf life expectation of certain products such as vegetables, fruits, meats and seafoods (Simpson et al. 2012). On the other hand, consumers appear to reject irradiation with relation to non-food accidents (Donati 2015); moreover, some decrease in nutrients has been reported (Simpson et al. 2012) and the use of this technique can be effective on condition that enzymatic inactivation is complete. For these reasons, the method is certainly powerful, but the current market does not seem to accept it at present.

2.3.7 Superchilling

Superchilling is an interesting method for the preservation of cod white fish on ships (Waterman and Taylor 2001). Substantially, the process concerns the partial freezing of water inside foods: the final temperature should be 1–2 °C below the initial freezing point. A notable extension of shelf life has been observed (Kaale et al. 2011; Magnussen et al. 2008; Schubring 2009), thereby extending the storage life of the fish. However, three main disadvantages have to be considered (Waterman and Taylor 2001):

- (1) Superchilled products cannot be processed or used immediately because they are partially frozen. A certain time loss is expected
- (2) Normal quality inspections cannot be performed on partially frozen fish. Once more, a certain time loss should be taken into account
- (3) Superchilling parameters (temperature) have to be strictly monitored and constant. This condition could not be always observed easily on trawlers.

2.4 Traditional Preservation Techniques

Basically, ‘traditional’ preservation methods involve the use of food additives with microbial actions (natural or artificial origin), peculiar historical systems such as smoking and fermentation, and the addition of natural food ingredients. This section

is dedicated to a brief description of these techniques, with possible advantages and risks if compared with thermal preservation.

2.4.1 Preservation with Traditional Additives and Pretreatments

2.4.1.1 Preservation with Sodium Chloride

The traditional use of salt for the preservation of meat, fish and vegetable products (Delgado et al. 2016) can also provide particular sensorial features. With exclusive relation to sodium chloride, the following positive actions have to be considered:

- (a) Increased osmotic pressure and consequent damages to microbial cells. Osmotic pressure depends exclusively on the number of dissolved solute particles in solution, while their size is not important. Consequently, small-size ions such as sodium cations and chloride anions can provide higher osmotic pressure values (and a better preservation of treated foods) if compared with other high-size solutes such as sucrose at the same concentration (g/l)
- (b) Decrease of a_w values with the consequent inhibition of microbial activities. Sodium and chloride ions tend to ‘block’ the surrounding water molecules: the amount of bioavailable free water has to decrease necessarily. In addition, the formation of hydroxide ions reduces a_w . Finally, sodium chloride and proteolytic enzymes may create new compounds with the consequent reduction of cellular decomposition.

The action of the salt is mainly bacteriostatic; however, the bactericidal action cannot be guaranteed because of the possible presence of osmophilic, halophilic or salt-tolerant life forms. For this reason, salting is often combined with other preservation methods such as low temperatures, the addition of nitrite and nitrate, cooking or smoking process.

Usually, commercially available salt is obtained from the evaporation of seawater, from natural brines and from underground warehouses. Different methods are available at present, including the addition of salt in direct contact (possibly mixed with spices, nitrates or nitrites), or by rubbing, or with stratification in alternate layers. Anyway, salt should be added at low temperatures: otherwise, certain defects could occur including dishomogeneity of water transfer to the external surface of foods and the formation of superficial crusts. Other well-known systems comprehend the addition of salt by means of the brine injection into meat tissues or by simple immersion (e.g. Grana and Parmigiano Reggiano cheeses). Anyway, cooking and smoking processes can be combined with salting.

2.4.1.2 Preservation with Sucrose

Sucrose can increase the osmotic pressure similarly to salt, but its preservative action is low if compared with sodium chloride; for this reason, high concentrations of added sucrose are needed (65–70 %). Consequently, treated foods are often jams, fruit jellies and candies.

It should be also noted that higher concentrations of added sucrose lead to crystallisation phenomena, while lower quantities cannot be effective enough (a synergic action with other preservation methods such as concentration, acidification and cold-temperature storage are required). Glucose and fructose may be used as possible sucrose substitutes because of their lower molecular weight and the consequent reduced amount (40 %).

Similarly to salt, sucrose has a prevailing bacteriostatic action without bactericidal effects; moreover, osmophilic life forms may remain unaffected in presence of sucrose. For these reasons, a synergic action with other methods is needed and recommended.

2.4.1.3 Preservation with Ethyl Alcohol

Ethanol—a non-toxic alcohol for food applications—is added to fruits for the production of peculiar products with amounts between 50 and 70 %. In these conditions, ethanol would be lethal against vegetative forms; naturally, treated foods would also obtain a pleasant and strong taste. On the other hand, initially present vitamins would be destroyed.

2.4.1.4 Preservation with Oils

Preserved foods in oil are one of the most ‘popular’ products at present (Delgado et al. 2016). Generally, two main food products—vegetables and fish—are preserved with vegetable oils. Intermediate foods are placed in dedicated containers (after cooking or other thermal treatments) and immersed in oil with the aim of covering completely the edible product. Substantially, the aim of food technologists is to create an oxygen-free environment, with the consequent inhibition of aerobic bacteria; vegetable oils do not show a direct preservative action.

For this reason, the use of oils is not effective and recommended when speaking of possible anaerobic activity by certain microorganisms (example: *Clostridium botulinum*). Moreover, moist foods cannot be immersed in oil and remain preserved because living microorganisms may be able to spread in the aqueous phase (example: undrained foods). Consequently, this traditional method may require the synergic action of different preservation systems such as pasteurisation and sterilisation (canned tuna is initially cooked before being salted, canned in oil and sterilised) or the addition of salt and vinegar (artichokes are boiled in water and vinegar before treatment in oil).

The storage in oil may determine a certain loss of nutrients (vitamins above all); in addition, the preservation of pickles is very different if compared to fresh products. Moreover, the composition of drained foods gives an excess of energy in comparison with the untreated product: sodium contents are rather high and lipids may be still 10 times greater if compared with fresh products.

2.4.1.5 Preservation with Vinegar

Vinegar is historically used to preserve vegetables and some fish products by means of its content in acetic acid (minimum quantity: 6 %). Substantially, the immersion in vinegar can determine:

- (1) A microbiostatic action (pH values are notably lowered)
- (2) A bactericidal action because of the known toxic effect of acetic acid.

The immersion in cold or heated vinegar (75–80 °C) has to be carried out after preliminary treatments: foods should be seared, salted or macerated in vinegar for a few days. Moreover, the acid solution can be ‘enriched’ with spices or herbs. These procedures allow to counteract the possible decrease of preservation effects because of the excessive dilution (intermediate solutions contain always more or less moist foods).

Almost all vegetables (peppers, eggplant, cucumbers, capers, etc.) can be preserved in vinegar. Initial foods can be either fresh or semi-preserved raw materials; in the latter case, these are kept in brine up to the time of the preservation process, when they are rinsed and put under vinegar.

The modern production of preserved foods in vinegar involves also the use of chemical additives (citric acid and ascorbic acid) or a pasteurisation treatment. For these reasons, radical modifications of organoleptic features (taste, colour and odour) have to be expected. Moreover, vitamins and other nutrients are likely lost. On the other time, pickles improve with time and can be stored for a very long time.

2.4.1.6 Blanching

Blanching is a thermal pretreatment with peculiar applications for vegetable raw materials. Actually, the process should be discussed in Sect. 2.1 because of the use of heat (water or steam at 70–105 °C). However, the importance of blanching is marginal when speaking of food preservation methods.

In fact, the main purpose of this technique is to inhibit only enzymatic activity in vegetables and some fruit types before subjecting them to ‘classical’ preservation systems. In other words, blanching has a limited importance when speaking of food safety, inhibition of spoilage microorganisms and elimination of pathogen agents; on the contrary, further preservation methods are required. By the chemical viewpoint, macroscopic effects of blanching are the modification of sensorial

features (colours above all) because of the transformation of chlorophylls and other organic substances such as β -carotene, ascorbic acid, etc. (Muftugil 1986; Negi and Roy 2000). Other effects include the elimination of unpleasant odours and gases enclosed into plant tissues (with reduced oxidation risks), the decrease of microbial counts and the coagulation of proteins with lower cooking times.

From the technological viewpoint, raw materials are rapidly heated to the desired temperature: after a certain time, they are rapidly cooled to room temperature.

2.4.2 Other Traditional and Modern Preservation Techniques

2.4.2.1 Dehydration

The term ‘dehydration’ means the progressive reduction of water content inside foods with the aim of inhibiting microbial spoilage; essentially, this process has microbiostatic effects. However, three different processes can be named ‘dehydration’:

- (a) Concentration
- (b) Drying
- (c) Freeze-drying.

Basically, these methods are performed with the aim of reducing the bioavailability of free water into foods. This availability is measured in terms of water activity; should a_w (Sects. 2.3.1 and 2.4.1.1) be low enough, the decrease in the growth rate of living microorganisms would be observed until the complete block of all metabolic activities. a_w can be reduced:

- (1) By means of the removal of water molecules from foods (dehydration), or
- (2) By means of the dissolution of peculiar substances (solutes) in the solid food solution (chemical preservation).

Dehydration has a certain microbiostatic action; in addition, water bioavailability is remarkably reduced. On the other side, toxins and spores remain substantially unaffected; for this reason, the use of dehydration cannot be recommended when using raw materials with appreciable microbial counts.

In general, the tripartite subdivision of dehydration systems depends on the technological solution for the elimination (separation) of the aqueous solvent from the food matrix.

Concentration techniques are generally performed on liquid foods or products with high water amounts; the objective is the reduction of volumes and weights (limitation of transportation costs). It has to be noted that preservation effects are rather limited and always associated with other treatments (pasteurisation, freezing, addition of chemicals, etc.).

Concentration by evaporation is the most used technique for the production of concentrated feed and food products. A basic requirement has to be satisfied: the food product has to be treated with temperatures close to the boiling point of the food itself. No substantial changes should be observed when speaking of sensorial and nutritional properties, on condition that the processing temperature is close but slightly lower than the maximum thermal value (boiling point of the solid or liquid mixture). For these reasons, the continuous monitoring of rheological properties is required because of the necessity of eliminating possible foams and the consequent local heating (imperfect heat circulation and steam removal). The control of pressure is also recommended: under partial vacuum, boiling points are lowered and the removal of steam is easy at low temperatures. Generally, this process (increase of thermal values) may damage thermally sensible products such as milks, alcoholic beverages and juices.

The second dehydration process, cryoconcentration, is similar to classical freezing systems: liquid foods (solid products cannot be discussed here) are treated with the aim of removing water in form of ice crystals (final concentration of solutes: up to 60 %, while the normal evaporation process should assure only 30 %). The crystallisation of water molecules is carried out by treating the food between -3 and -7 °C: this procedure aims to favour the growth of ice crystals instead of the simple nucleation (Sect. 2.2.1.2). Consequently, food damages (in liquid foods) are limited and the removal of ice crystals is more efficient. Generally, cryoconcentration procedures are used for tea and spices (before drying), coffee extracts and similar products, vinegar, fruit juices, milk and wines. Clearly, the following advantages are determined by the notable removal of volumes (and weights):

- Excellent aroma retention
- Different logistic solutions during transportation and storage steps in the food chain
- Virtual absence of sensory variations and damages to certain organic substances (vitamins).

On the other side, the incorrect management of processing parameters and the insufficient water removal could cause different defects, including microbial spreading and possible phase separations in the liquid product.

The third concentration procedure allows the partial water removal by means of the use of artificial membranes (concentration of solutes may reach 50 %). Interestingly, this removal can easily be performed at room temperature, without classical thermal damages (nutritional loss, variation of sensorial features, etc.). One of the most known versions of concentration by membrane processes, the direct osmosis, is based on the contact (13.5 ± 1.5 h) between whole, cut or sliced foods on the one side and a (possibly heated) concentrated solution (65–70 % of sucrose). In this situation, cellular membranes in foods provide the physical interface between foods and the concentrated solution. The addition of food additives such as ascorbic acid may be recommended because of possible browning effects.

Naturally, this method is completely 'natural'; on the other hand, sweet tastes should be notably enhanced in the final product. A possible countermeasure is the concomitant addition of salt (it can also reduce browning effects and increase osmotic pressures).

Anyway, direct osmosis is not a standalone preservation treatment: on the contrary, it should be recommended as a preliminary process before other drastic preservation procedures (drying, freezing and lyophilisation). In addition, recent modifications of the original system imply the use of relatively short times (up to 20 min) and high temperatures (75 ± 10 °C) with the declared aim of combining synergically dehydration and pasteurisation. At present, the new frontier is represented by the use of artificial membranes (organic polymers): new and improved processes are named 'reverse osmosis' systems.

2.4.2.2 Drying

Drying is historically well known: this method implies the heat treatment of solid or liquid products with the aim of removing large water amounts: the final aqueous content should not exceed 15 %. Differences between historical and modern systems include the following points:

- (a) Heat sources. Ancient systems imply the exposure of foods to sun and air for weeks or months. Industrial systems, surely more manageable than historical procedures, are based on the development of artificially heated rooms and the subdivision of the whole process in three steps: food preparation before drying, drying and final packaging
- (b) Drying procedures. Historical systems do not require peculiar standards when speaking of monitoring and controls. On the other hand, modern systems—hot air drying; infrared, ultrasound or microwave drying; direct food contact drying—require the critical control of heat transfer from emitting sources to the product and the concomitant water removal until a new thermohygro-metric equilibrium with the surrounding environment. Other critical parameters are the gradient heating of foods (core temperatures should not exceed 70 °C), the application of very low pressures and the management of relative humidity (desired values between 10 and 20 %). Insufficient controls could cause serious damages to foods, similarly to failures observed in certain thermal processes (pasteurisation, sterilisation).

2.4.2.3 Lyophilisation

Lyophilisation consists in the under-vacuum dehydration of previously frozen products by sublimation (four steps are required). Temperature has to be <0 °C. Treated foods have a reduced weight (final values can reach 92 %): rehydrated products can show the virtual absence of nutritional losses. In detail, the following

products—high-protein and high-calorie soups, granular and orange juice, vegetable soups, meat, etc.—can be preserved for very long times. On the other hand, the main disadvantage of these systems is correlated with the incorrect management of these processes and the hygroscopic behaviour of treated foods: packaging processes should be carried out packaged under vacuum or under nitrogen in triple-layer envelopes.

2.4.2.4 Smoking

Smoking treatments mean the introduction of peculiar substances in foods by means of the simple immersion in a saturated atmosphere or in a liquid bath. Basically, these compounds can:

- (a) Penetrate the inner structure of treated foods towards the centre of the product (depending on immersion time) by food contact on edible surfaces with the ‘smoking’ source
- (b) Remain into the food after immersion
- (c) Inhibit microbial growth during immersion and after the treatment (prolonged effects).

Smoking procedures are generally performed on already salted foods: the synergic action of salting and smoking can increase dehydration and enhance peculiar aroma and taste features.

From the technological viewpoint, smoking is recognised to show antiseptic, antioxidant and antimicrobial properties. Historically, foods were smoked by hanging over burning wood: as a result, the action of smoking was associated with heat penetration and superficial drying. At present, modern operations have different processing parameters. In detail, two main industrial applications are available:

- (1) ‘Cold’ smoking. foods are heated between 20 and 45 °C; total processing times are measurable as days or weeks
- (2) ‘Hot’ smoking. Intermediate products are heated between 50 and 90 °C for a few hours only.

Because of the nature of smoking compounds, sensorial features of treated foods are completely different from original raw materials or intermediates. As a result, smoked products are extremely appreciated as a new food category, both for hedonistic reasons and historical features. The following list shows the most known types of smoked food products:

- (a) Cheeses such as palmero, provola, ewe cheese, etc. (Guillén et al. 2007; Majcher et al. 2011; Naccari et al. 2008)
- (b) Cured, cooked and raw meats, such as beef muscle cooked in brine, wurstel, sausages, Prague ham, Speck, etc. (Budig 2012; Lucarini et al. 2013)

- (c) Fish (salmon, herring, sturgeon, swordfish, tuna, mackerel, trout and eel). Interestingly, smoking treatments are associated with a certain increase in sodium amounts.

From the technological viewpoint, it should be considered that smoked products are certainly dry versions of the original foods; for this reason, durabilities are notably enhanced (the influence of salting has to be taken into account).

On the other hand, smoking is questioned enough at present because of certain safety concerns (presence of potentially carcinogenic substances such as benzopyrene). As a result, the food industry is currently evaluating smoke flavourings or 'liquid smoke' solutions. In particular, the liquid solution corresponds to the result of the condensation of volatile smoke products (from wood combustion) in water; after a filtration process, the liquid smoke is free from heavy oils and ready for use.

New smoking systems may assure other advantages if compared with traditional methods, including a notable uniformity with relation to flavours, aroma and colours on food surfaces. Moreover, processing times are remarkably faster. Disadvantages of these methods, typically caused by the improper management of times and/or the incorrect dilution of the initial smoke in the final water solution, might be worsened by salting and thermal preservation systems. In particular, excessive salting may enhance crust formation in certain cheeses, and smoking treatments could have more visible effects (brownish colours) than expected; the same thing may occur if preheating treatments give more or less concentrated products, with concern to moisture amounts.

2.4.2.5 Fermentation

Fermentation means the process—or the sum of concomitant processes—that allows certain microorganisms, including degradative life forms such as moulds and yeasts, to break down carbohydrates with the production of alcohol and organic acids. These substances can efficiently inhibit the growth of putrefactive microorganisms and increase shelf life periods. In addition, fermentative processes can modify partially chemical compositions and sensorial features, the digestibility of preserved foods, and other properties. At present, the role of biotechnology is relevant when speaking of industrial fermentation processes: more than 3500 different types of fermented foods are known and produced worldwide, including yoghurts, cheeses, wines, beers and sauerkrauts (Baglio 2014).

Apparently, fermentation can give a number of advantages when speaking of food safety and product categorisation. However, there are some risks because fermentative processes can assure good performances on condition that 'food supports' are good enough. In other words, heat-based thermal treatments have to destroy all degradative microorganisms with the aim of eliminating all possible causes of microbial competition between fermentative life forms and other microbial agents; should not this condition be respected, raw materials could give

unexpected results. At the same time, strongly pasteurised raw materials could have a reduced amount of fermentative life forms, and the normal fermentation process could be seriously impeded. With relation to cold-based storage treatments, several microorganisms might suffer important damages and be inhibited in terms of fermentation. Finally, drastic heat-based preservation treatments could decrease the bioavailability of certain nutrients, with comprehensible effects on the normal fermentation yield.

References

- Ahmed SH, Luksas AJ (1988) Spreadable butter-like composition and method for production thereof. US Patent 4,769,255, 6 Sept 1988
- Alais C (1984) *Science du Lait. Principes des techniques laitières*, 4th edn. S.E.P.A.I.C., Paris
- Ali S, Bettelheim FA (1985) Non-freezing water in protein solutions. *Colloid Polym Sci* 263 (5):396–398. doi:10.1007/BF01410387
- Andersen HJ, Bertelsen G, Ohlen A, Skibsted LH (1990) Modified packaging as protection against photodegradation of the colour of pasteurized, sliced ham. *Meat Sci* 28(1):77–83. doi:10.1016/0309-1740(90)90021-W
- Andreini R, Chiodi J, Noni ID, Resmini P, Battelli G, Cecchi L, Todesco R, Cattaneo TMP, Rampilli M, Foschino R (1990) Evaluation of heat damage in UHT and in-bottle sterilized milk samples traded in Italy. *Sci Tec Latt Casearia* 41(6):472–492
- Baglio E (2014) The modern yoghurt: introduction to fermentative processes. In: Baglio E (ed) *Chemistry and technology of Yoghurt Fermentation*. SpringerBriefs in Chemistry of Foods, Springer International Publishing, Cham. doi:10.1007/978-3-319-07377-4_1
- Borde C (2006) La Revolution sardiniere: Pecheurs et conserveurs en Bretagne sud au XIXe siecle (review). *Technol Cult* 47(1):200–201. doi:10.1353/tech.2006.0058
- Brant AW, Patterson GW, Walters RE (1968) Batch pasteurization of liquid whole egg. 1. Bacteriological and functional property evaluation. *Poult Sci* 47(3):878–885. doi:10.3382/ps.0470878
- Brunazzi G, Parisi S, Pereno A (2014) Packaging and quality. In: Brunazzi G, Parisi S, Pereno A (eds) *The importance of packaging design for the chemistry of food products*. Springer International Publishing, Cham. doi:10.1007/978-3-319-08452-7_5
- Bruhn CM, Schutz HG (1999) Consumer food safety knowledge and practices. *J Food Saf* 19 (1):73–87. doi:10.1111/j.1745-4565.1999.tb00235.x
- Budig J (2012) Prague ham: the past and the present. *Maso Int J Food Sci Technol* 1:77–80. <http://www.maso-international.cz/category/volume-012012/>. Accessed 28 Apr 2016
- Chandan RC (2015) Dairy processing and quality assurance: an overview. In: Chandan RC, Kilara A, Shah NP (eds) *Dairy processing and quality assurance*. Wiley, Chichester. doi:10.1002/9781118810279.ch01
- Cinquanta L, Albanese D, Curcurullo G, Di Matteo M (2010) Effect on orange juice of batch pasteurization in an improved pilot-scale microwave oven. *J Food Sci* 75(1):E46–E50. doi:10.1111/j.1750-3841.2009.01412.x
- Clare DA, Bang WS, Cartwright G, Drake MA, Coronel P, Simunovic J (2005) Comparison of sensory, microbiological, and biochemical parameters of microwave versus indirect UHT fluid skim milk during storage. *J Dairy Sci* 88(12):4172–4182. doi:10.3168/jds.S0022-0302(05)73103-9
- Corzo N, López-Fandiño R, Delgado T, Ramos M, Olano A (1994) Changes in furosine and proteins of UHT-treated milks stored at high ambient temperatures. *Z Lebensm Unters Forsch* 198(4):302–306

- Cunningham FE (1986) Egg-product pasteurization. In: Stadelman WJ, Cotteril OJ (eds) *Egg science and technology*, 3rd edn. P. Macmillan, London, pp 243–272
- Datta N, Elliott AJ, Perkins ML, Deeth HC (2002) Ultra-high-temperature (UHT) treatment of milk: comparison of direct and indirect modes of heating. *Aust J Dairy Technol* 57(3):211–227
- Degraeve P, Saurel R, Coutel Y (2003) Vacuum impregnation pretreatment with pectin-methylesterase to improve firmness of pasteurized fruits. *J Food Sci* 68(2):716–721. doi:10.1111/j.1365-2621.2003.tb05738.x
- Delgado A, Vaz de Almeida MD, Parisi S (2016) *Chemistry of the Mediterranean diet*. Springer International Publishing, Cham. doi:10.1007/978-3-319-29370-7
- Delia S, Laganà P, Parisi S (2005) *Materiali e metodi di confezionamento nella conservazione dei prodotti alimentari refrigerati*. Proceedings of the National Conference “Microbiologia degli alimentari conservati in stato di refrigerazione”. Oxoid S.p.A., Garbagnate M.se, pp 93–111
- Donati C (2015) Risk analysis, contaminants and impact on health in imports of non-animal origin: the EU context. In: Montanari F, Jezso V, Donati C (eds) *Risk regulation in non-animal food imports*. SpringerBriefs in Chemistry of Foods, Springer International Publishing, Cham. doi:10.1007/978-3-319-14014-8_1
- Engelman MS, Sani RL (1983) Finite-element simulation of an in-package pasteurization process. *Numer Heat Transf* 6(1):41–54. doi:10.1080/01495728308963073
- Ercolini D, Russo F, Torrieri E, Masi P, Villani F (2006a) Changes in the spoilage-related microbiota of beef during refrigerated storage under different packaging conditions. *Appl Environ Microbiol* 72(7):4663–4671. doi:10.1128/AEM.00468-06
- Ercolini D, La Storia A, Villani F, Mauriello G (2006b) Effect of a bacteriocin-activated polyethylene film on *Listeriamonocytogenes* as evaluated by viable staining and epifluorescence microscopy. *J Appl Microbiol* 100(4):765–772. doi:10.1111/j.1365-2672.2006.02825.x
- Farouk MM, Wieliczko KJ, Merts I (2004) Ultra-fast freezing and low storage temperatures are not necessary to maintain the functional properties of manufacturing beef. *Meat Sci* 66(1):171–179. doi:10.1016/S0309-1740(03)00081-0
- Feist EM, Hadi HT (1965) Liquefaction of gases. US Patent 3,224,207, 21 Dic 1965
- Fry DE (2001) In vino veritas. *Surg Infect* 2(3):185–191. doi:10.1089/109629601317202669
- Guillén MD, Palencia G, Sopelana P, Ibargoitia ML (2007) Occurrence of polycyclic aromatic hydrocarbons in artisanal Palermo cheese smoked with two types of vegetable matter. *J Dairy Sci* 90(6):2717–2725. doi:10.3168/jds.2006-452
- Samani BH, Khoshtaghaza MH, Minaei S, Zareifouros H, Eshtiaghi MN, Rostami S (2016) Design, development and evaluation of an automatic fruit-juice pasteurization system using microwave-ultrasonic waves. *J Food Sci Technol* 53(1):88–103. doi:10.1007/s13197-015-2026-6
- Hugas M, Garriga M, Monfort JM (2002) New mild technologies in meat processing: high pressure as a model technology. *Meat Sci* 62(3):359–371. doi:10.1016/S0309-1740(02)00122-5
- Kaale LD, Eikevik TM, Rustad T, Kolsaker K (2011) Superchilling of food: a review. *J Food Eng* 107(2):141–146. doi:10.1016/j.jfoodeng.2011.06.004
- Karabudak E, Bas M, Kiziltan G (2008) Food safety in the home consumption of meat in Turkey. *Food Control* 19(3):320–327. doi:10.1016/j.foodcont.2007.04.018
- Kröckel L (1995) Bacterial fermentation of meats. In: Campbell-Platt G, Cook PE (eds) *Fermented meats*. Springer US, New York, pp 69–109. doi:10.1007/978-1-4615-2163-1_4
- International Dairy Foods Association (2006) *Pasteurization*. <http://www.idfa.org/news-views/media-kits/milk/pasteurization>. Accessed 27 Apr 2016
- Larousse J (1991) Evolution technologique de l'appertisation et de la pasteurisation des fruits et légumes. *Ind Aliment Agric* 108(6):475–478
- Li B, Sun DW (2002) Novel methods for rapid freezing and thawing of foods—a review. *J Food Eng* 54(3):175–182. doi:10.1016/S0260-8774(01)00209-6
- Lucarini M, Sacconi G, D'Evoli L, Tufi S, Aguzzi A, Gabrielli P, Marletta L, Lombardi-Boccia G (2013) Micronutrients in Italian ham: a survey of traditional products. *Food Chem* 140(4):837–842. doi:10.1016/j.foodchem.2012.10.020

- Magnussen OM, Haugland A, Hemmingsen AKT, Johansen S, Nordtvedt TS (2008) Advances in superchilling of food—Process characteristics and product quality. *Trends Food Sci Technol* 19 (8):418–424. doi:[10.1016/j.tifs.2008.04.005](https://doi.org/10.1016/j.tifs.2008.04.005)
- Mangalassary S, Dawson PL, Rieck J, Han IY (2004) Thickness and compositional effects on surface heating rate of bologna during in-package pasteurization. *Poult Sci* 83(8):1456–1461. doi:[10.1093/ps/83.8.1456](https://doi.org/10.1093/ps/83.8.1456)
- Mangalassary S, Han I, Rieck J, Acton J, Dawson P (2008) Effect of combining nisin and/or lysozyme with in-package pasteurization for control of *Listeria monocytogenes* in ready-to-eat turkey bologna during refrigerated storage. *Food Microbiol* 25(7):866–870. doi:[10.1016/j.fm.2008.05.00](https://doi.org/10.1016/j.fm.2008.05.00)
- Majcher MA, Goderska K, Pikul J, Jeleń HH (2011) Changes in volatile, sensory and microbial profiles during preparation of smoked ewe cheese. *J Sci Food Agric* 91(8):1416–1423. doi:[10.1002/jsfa.4326](https://doi.org/10.1002/jsfa.4326)
- Mauriello G, Ercolini D, La Storia A, Casaburi A, Villani F (2004) Development of polyethylene films for food packaging activated with an antilisterial bacteriocin from *Lactobacillus curvatus* 32Y. *J Appl Microbiol* 97(2):314–322. doi:[10.1111/j.1365-2672.2004.02299.x](https://doi.org/10.1111/j.1365-2672.2004.02299.x)
- Micali M, Parisi S, Minutoli E, Delia S, Laganà P (2009) Alimenti confezionati e atmosfera modificata. Caratteristiche basilari, nuove procedure, applicazioni pratiche. *Ind Aliment* 48 (489):35–43
- Molin G (2000) Modified atmospheres. In: Lund BM, Baird-Parker TC, Gould GW (eds) *The microbiological safety and quality of food*, vol 1. Aspen Publishers Inc, Gaithersburg
- Morales FJ, Romero C, Jiménez-Pérez S (2000) Characterization of industrial processed milk by analysis of heat-induced changes. *Int J Food Sci Technol* 35(2):193–200. doi:[10.1046/j.1365-2621.2000.00334.x](https://doi.org/10.1046/j.1365-2621.2000.00334.x)
- Muftugil N (1986) Effect of different types of blanching on the color and the ascorbic acid and chlorophyll contents of green beans. *J Food Proc Preserv* 10(1):69–76. doi:[10.1111/j.1745-4549.1986.tb00006.x](https://doi.org/10.1111/j.1745-4549.1986.tb00006.x)
- Murphy RY, Duncan LK, Driscoll KH, Marcy JA (2003) Lethality of *Salmonella* and *Listeria innocua* in fully cooked chicken breast meat products during postcook in-package pasteurization. *J Food Prot* 66(2):242–248
- Murphy RY, Hanson RE, Johnson NR, Chappa K, Berrang ME (2006) Combining organic acid treatment with steam pasteurization to eliminate *Listeria monocytogenes* on fully cooked frankfurters. *J Food Prot* 69(1):47–52
- Murray M, Richard JA (1997) Comparative study of the antilisterial activity of nisin A and pediocin AcH in fresh ground pork stored aerobically at 5°C. *J Food Prot* 60:1534–1540
- Naccari C, Cristani M, Licata P, Giofrè F, Trombetta D (2008) Levels of benzo [a] pyrene and benzo [a] anthracene in smoked “Provola” cheese from Calabria (Italy). *Food Addit Contam I* (1):78–84. doi:[10.1080/19393210802236968](https://doi.org/10.1080/19393210802236968)
- Negi PS, Roy SK (2000) Effect of blanching and drying methods on β -carotene, ascorbic acid and chlorophyll retention of leafy vegetables. *LWT-Food Sci Technol* 33(4):295–298. doi:[10.1006/fstl.2000.0659](https://doi.org/10.1006/fstl.2000.0659)
- Ouwehand AC (1998) Antimicrobial components from lactic acid bacteria. In: Salminen S, von Wright A (eds) *Lactic acid bacteria: microbiology and functional aspects*, 2nd edn. Marcel Dekker Inc., New York, pp 139–159
- Parisi S (2002) I fondamenti del calcolo della data di scadenza degli alimenti: principi ed applicazioni. *Ind Aliment* 41(417):905–919
- Parisi S (2005) La produzione “continua” è anche “costante”? Confutazione di alcuni luoghi comuni nel settore industriale/manifatturiero. *Chim Ital* 16(3–4):10–18
- Parisi S (2009) Intelligent packaging for the food industry. In: Carter EJ (ed) *Polymer electronics —a flexible technology*. Smithers Rapra Technology Ltd, Shawbury
- Parisi S (2012) *Food packaging and food alterations*. Smithers Rapra Technology Ltd, Shawbury
- Parisi S (2013) *Food Industry and packaging materials: performance-oriented guidelines for users*. Smithers Rapra Technology Ltd, Shawbury

- Parisi S (2016) Alberto Escarpa, María Cristina González, and Miguel Angel López: Agricultural and food electroanalysis. *Anal Bioanal Chem* 408(9):2185–2186. doi:[10.1007/s00216-016-9347-9](https://doi.org/10.1007/s00216-016-9347-9)
- Parisi S, Laganà P, Delia S (2004) Cubed Mozzarella Cheese in modified atmosphere packages: evolutive profiles of chemical and microbiological parameters during Shelf Life. In: Programme & Abstracts Book of the 3rd ILSI Europe International Symposium on Food Packaging, Barcelona, 17–19 Nov 2004
- Patterson MF (2005) Microbiology of pressure-treated foods. *J Appl Microbiol* 98(6):1400–1409. doi:[10.1111/j.1365-2672.2005.02564.x](https://doi.org/10.1111/j.1365-2672.2005.02564.x)
- Potter NN, Hotchkiss JH (1995) Milk and milk products. Food Science. Springer, US, New York, pp 279–315
- Pruthi JS (1999) Quick freezing preservation of foods: foods of plant origin, vol 2. Allied Publishers Ltd, New Delhi
- Rahman MS (ed) (2007) Handbook of food preservation, 2nd edn. CRC Press LLC, Boca Raton
- Redmond EC, Griffith CJ (2003) Consumer food handling in the home: a review of food safety studies. *J Food Prot* 66(1):130–161
- Reif-Acherman S (2009) Several motivations, improved procedures, and different contexts: the first liquefactions of helium around the world. *Int J Refrig* 32(5):738–762. doi:[10.1016/j.ijrefrig.2009.02.019](https://doi.org/10.1016/j.ijrefrig.2009.02.019)
- Rerkrai S, Jeon IJ, Bassette R (1987) Effect of various direct ultra-high temperature heat treatments on flavor of commercially prepared milks. *J Dairy Sci* 70(10):2046–2054. doi:[10.3168/jds.S0022-0302\(87\)80252-7](https://doi.org/10.3168/jds.S0022-0302(87)80252-7)
- Rigo M (2010) Sistemi di pastorizzazione a radiofrequenza. Dissertation, University of Padua
- Rogers WR (2013) Background to sterilisation—an historical introduction. In: Rogers WR (ed) Healthcare sterilisation: introduction and standard practices, vol 1. Smithers Rapra Technology, Shawbury
- Schubring GR (2009) “Superchilling”—an “old” variant to prolong shelf life of fresh fish and meat requicked. *Fleischwirtsch* 89(9):S.104–113
- Senesi E (1984) Surgelazione dei prodotti vegetali. In: Ottaviani F (ed) Microbiologia dei prodotti di origine vegetale – ecologia ed analisi microbiologica. Chiriotti Editori, Pinerolo
- Senesi E, Bignardi B, Gennari M (2000) Influence of pre-treatment and protective atmosphere packaging on quality indices of fresh-cut musk melon. *Fruttic* 11:69–73
- Simpson BK, Rui X, Klomkloa S (2012) Enzymes in food processing. In: Simpson BK (ed) Food biochemistry and food processing, Second Edition. Wiley-Blackwell, Oxford. doi:[10.1002/9781118308035.ch9](https://doi.org/10.1002/9781118308035.ch9)
- Sivertsvik M, Jeksrud WK, Rosnes JT (2002) A review of modified atmosphere packaging of fish and fishery products—significance of microbial growth, activities and safety. *Int J Food Sci Technol* 37(2):107–127
- Skandamis PN, Nychas GJE (2001) Effect of oregano essential oil on microbiological and physico-chemical attributes of minced meat stored in air and modified atmospheres. *J Appl Microbiol* 91(6):1011–1022. doi: [10.1046/j.1365-2672.2001.01467.x](https://doi.org/10.1046/j.1365-2672.2001.01467.x)
- Soncini G (2008) Frodi e alterazioni delle conserve. Proceedings of the Fourth ‘Meeting Nazionale sulla Sicurezza Alimentare: le Frodi Alimentari e le Alterazioni degli Alimenti’, Rome, 23–24 May 2008
- Stabel JR (2001) On-farm batch pasteurization destroys *Mycobacterium paratuberculosis* in waste milk. *J Dairy Sci* 84(2):524–527. doi:[10.3168/jds.S0022-0302\(01\)74503-1](https://doi.org/10.3168/jds.S0022-0302(01)74503-1)
- Tassou C, Koutsoumanis K, Nychas GJE (2000) Inhibition of *Salmonella enteritidis* and *Staphylococcus aureus* in nutrient broth by mint essential oil. *Food Res Int* 33(3–4):273–280. doi:[10.1016/S0963-9969\(00\)00047-8](https://doi.org/10.1016/S0963-9969(00)00047-8)
- Waterman JJ, Taylor DH (2001) Superchilling. Torry Advisory Note No. 32. Torry Research Station, UK. FAO in partnership with Support Unit for International Fisheries and Aquatic Research, SIFAR
- Wheeler JL, Harrison MA, Koehler PE (1987) Presence and stability of patulin in pasteurized apple cider. *J Food Sci* 52(2):479–480. doi:[10.1111/j.1365-2621.1987.tb06643.x](https://doi.org/10.1111/j.1365-2621.1987.tb06643.x)
- Zhao Y (2007) Freezing process of berries. In: Zhao Y (ed) Berry fruit: value-added products for health promotion. CRC Press, Boca Raton



<http://www.springer.com/978-3-319-42461-3>

The Chemistry of Thermal Food Processing Procedures

Micali, M.; Fiorino, M.; Parisi, S.

2016, VI, 54 p. 3 illus., 2 illus. in color., Softcover

ISBN: 978-3-319-42461-3