

Contents

1	Introduction	1
	References	3
2	Theoretical Framework and Motivation	5
2.1	The Standard Model	5
2.1.1	Outstanding Issues in the Standard Model	6
2.2	Supersymmetry	9
2.2.1	Minimal Supersymmetric Standard Model	9
2.2.2	R-parity	10
2.2.3	SUSY Breaking Model	11
2.3	Wino LSP Scenarios and Their Phenomenology	11
2.3.1	Anomaly-Mediated Supersymmetry Breaking Model	11
2.3.2	Pure Gravity Mediation Model	13
2.3.3	Phenomenology	14
2.3.4	Theoretical and Experimental Indications	16
2.3.5	Previous Searches and Constraints	17
	References	21
3	LHC and ATLAS Detector	23
3.1	Large Hadron Collider	23
3.2	ATLAS Detector	25
3.2.1	ATLAS Coordinate	25
3.2.2	Magnet System	26
3.2.3	Inner Detector	27
3.2.4	Calorimeter System	33
3.2.5	The Electromagnetic Calorimeter	33
3.2.6	Hadronic Calorimeter	35
3.2.7	Muon Spectrometer	37
3.2.8	Luminosity Detector	40
3.2.9	Trigger System	42
	References	44

4	Data and Monte Carlo Simulation	47
4.1	Data Sample	47
4.1.1	Trigger	47
4.1.2	Luminosity and Pile-Up	48
4.2	Simulated Sample	49
4.2.1	Signal Monte Carlo Sample	49
4.2.2	Background Monte Carlo Sample	52
	References	54
5	Object Reconstruction	55
5.1	Track Reconstruction	55
5.1.1	Track Parametrization	55
5.1.2	Inside-Out Track Reconstruction	56
5.1.3	Outside-In Track Reconstruction	58
5.1.4	Re-tracking	58
5.2	Vetex Reconstruction	61
5.3	Jet	61
5.3.1	Reconstruction	61
5.3.2	Clustering of Calorimeter Cells	61
5.3.3	Jet Finding Algorithm	61
5.3.4	Calibration	62
5.3.5	Jet Energy Scale Uncertainty	62
5.4	Electron	64
5.4.1	Reconstruction	64
5.4.2	Identification Criteria	64
5.5	Muon	65
5.5.1	Reconstruction	65
5.5.2	Performance	67
5.6	Missing Transverse Energy	68
	References	69
6	Tracking Performance	71
6.1	Tracking Performance for Decaying Charginos	71
6.2	Tracking Performance in QCD Multi-jet Events	72
	Reference	77
7	Properties of Signal Events	79
7.1	Production Process	79
7.2	Finding Decaying Chargino Tracks	81
7.3	Properties of Chargino Tracks	83
	References	88
8	Event Selection	91
8.1	Online Selection	91
8.1.1	Trigger	91

8.2	Offline Selection	95
8.2.1	Primary Vertex Selection	95
8.2.2	Overlap Removal	95
8.2.3	Event Cleaning	96
8.2.4	Kinematic Selection Criteria	97
8.2.5	Disappearing Track Selection	98
	References	102
9	Background Estimation	103
9.1	Interacting-Hadron Tracks	104
9.1.1	Control Sample for Interacting-Hadron Tracks	104
9.1.2	Extraction of p_T Shape	106
9.2	Leptons Failing to Satisfy Identification Criteria	108
9.2.1	Probability of Identifying a Lepton as a Disappearing Track	109
9.2.2	p_T Spectrum of Lepton Background Tracks	112
9.2.3	Uncertainty of the Lepton Background Yield	113
9.3	Tracks with Mismeasured p_T	115
9.3.1	Control Data Sample of p_T -Mismeasured Tracks	117
9.3.2	Further Check on the Impact-Parameter Dependence	118
9.3.3	Systematic Uncertainty on the Track- p_T Shape	119
	References	120
10	Systematic Uncertainties	121
10.1	Uncertainty on the Signal Events	121
10.2	Uncertainty on the Background Events	122
	References	124
11	Extraction of Signal Yield	125
11.1	Unbinned Extended Maximum Likelihood Fit	125
11.2	Validation of the Signal Extraction Method	127
11.3	Null-Hypothesis Testing in the High- p_T Region	130
11.4	Full p_T -range Fit	131
11.5	Properties of the Excess in the Track- p_T Spectrum	132
11.5.1	Instrumental Background	132
11.5.2	Possibility of Underestimating the Muon Background	133
11.5.3	Signature of Decaying Charged Massive Particles	134
12	Interpretations and Discussion	139
12.1	Constraints on the AMSB Model	139
12.2	Constraints on the Pure Gravity Mediation Model	141
12.3	Constraints on Wino Dark Matter	142
12.4	Discussion	143
	References	146
13	Conclusions	147

Search for Charginos Nearly Mass-Degenerate with the
Lightest Neutralino

Based on a Disappearing-Track Signature in pp
Collisions at $\sqrt{s} = 8$ TeV

Kazama, S.

2016, XIII, 148 p., Hardcover

ISBN: 978-4-431-55656-5