

Chapter 2

Link of Zygotic Genome Activation and Cell Cycle Control

Boyang Liu and Jörg Grosshans

Abstract

The activation of the zygotic genome and onset of transcription in blastula embryos is linked to changes in cell behavior and remodeling of the cell cycle and constitutes a transition from exclusive maternal to zygotic control of development. This step in development is referred to as mid-blastula transition and has served as a paradigm for the link between developmental program and cell behavior and morphology. Here, we discuss the mechanism and functional relationships between the zygotic genome activation and cell cycle control during mid-blastula transition with a focus on *Drosophila* embryos.

Key words Cell cycle, Mid-blastula transition, Zygotic genome activation

1 Introduction

In most animals, from nematodes to chordates, embryogenesis starts with a series of rapid cleavage cell cycles after fertilization. These fast divisions lead to an exponentially increasing number of cells without an accompanied growth of the embryo. After a species-specific number of divisions, the cell cycle slows down and finally enters a pause. Subsequently, the embryo enters gastrulation with its characteristic morphogenetic movements, loss of symmetry, and cell type-specific differentiation. Mammalian embryogenesis is special in that it begins with differentiation of inner cell mass (ICM) and trophoblast, and the fast embryonic cleavage cycles eventually arise at late blastocyst stage [1–3]. Maternally supplied materials, including proteins, RNAs, and conceivably also metabolites contribute to the initial developmental processes. Maternal products exclusively control development during this first period, as the zygotic genome starts expression only with a delay after fertilization. Following zygotic genome activation (ZGA), both maternal and zygotic factors contribute to developmental control. The switch from maternal to zygotic control is especially prominent in species with large, externally deposited eggs. ZGA coincides with striking changes in cell behavior and molecular processes, including

cell cycle, DNA replication, maternal RNAs degradation, chromatin structure, metabolite composition, and status of DNA checkpoint. This morphologically visible switch in early development during the blastula stage was first described 120 years ago in sea urchin *Echinus microtuberculatus* and *Sphaerechinus granularis*, and later has been referred to as mid-blastula transition (MBT) [4, 5].

1.1 MBT in Model Organisms

Many model organisms are well studied in terms of MBT. Amphibian *Xenopus laevis*, for instance, undergoes 12 short and synchronized cleavage cycles with a lack of gap phases, 35 min each and proceeds with a series of progressively longer and less synchronized divisions from cycles 13 to 15. The transition period is defined as the MBT [5–8]. S phase progressively lengthens, and the cell cycle pauses in G1 or G2 phases during the MBT [9]. Concomitantly, maternal transcripts are deadenylated and degraded. The first zygotic transcripts are detected at cycle 7 and transcription rate increases up to and beyond MBT [10]. During the MBT, developmental control is handed over from maternal to zygotic factors (maternal-zygotic transition, MZT).

In zebrafish *Danio rerio* embryo, 9 rapid cycles with approximately 15 min each are followed by gradually longer cell cycles [11]. MBT begins at cycle 10, and the cell cycle loses synchrony with acquisition of a G1 phase in cycle 11 [12]. Similar to *Xenopus*, ZGA is regulated by the nuclear-cytoplasmic ratio, but DNA damage checkpoint acquisition is independent of zygotic transcription [13]. Maternal factors Nanog, Pou5f1, and SoxB1 are required for de novo zygotic transcription as well as inducing maternal clearance by activating the microRNA *miR-430* expression [14].

In the nematode *Caenorhabditis elegans* (*C. elegans*), zygotic transcription is already activated in the 4-cell stage. Multiple mechanisms and maternal factors, including OMA-1 and OMA-2, are involved and regulated by phosphorylation, nuclear shuttling, and protein destabilization [15, 16]. In contrast to the other species discussed above, cells divide asynchronously and asymmetrically following fertilization in *C. elegans* embryos [17, 18].

1.2 MBT in *Drosophila*

MBT is observed in embryos of *Drosophila melanogaster* at about 2 h post fertilization. Embryonic development starts with 13 rapid and meta-synchronized nuclear divisions, with extraordinary short S phases and no gap phases [19]. The extraordinary speed of about 10 min per pre-blastoderm cell cycle is achieved by fast replication of DNA and the absence of cytokinesis [20–22]. The syncytial mode of early development is a special feature of insect embryogenesis [23]. Due to the absence of cytokinesis, the early cell cycles are often referred to as nuclear cycles (NC). The onset of the embryonic cell cycle is regulated by *pan gu*, *plutonium*, and *giant nuclei* [24–27]. From NC8 to 9, the nuclei move from the interior

of the egg toward the periphery, forming the syncytial blastoderm. From NC10 to 13, nuclei undergo four more divisions at the egg cell cortex, until the nuclei number reaches approximately 6000. Some nuclei remain in the interior egg to differentiate into polyploid yolk nuclei. After mitosis 13, the cell cycle mode changes with the introduction of a long G2 phase, and the embryo enters into cellularization stage [19]. Following NC11, the cell cycle gradually slows down from 10 min in NC11 to 21 min in NC13 and an hour-long G2 pause in interphase 14 (25 °C) [19]. The S phase lengthens and by cycle 14 a difference between early and late replicating euchromatin and the satellite DNA becomes obvious. In addition, the usage of replication origins changes [28].

Interphase 14 corresponds to the MBT in *Drosophila*. Interphase 14 is the stage when the cell cycle pauses in a G2 phase, zygotic transcription strongly increases, and DNA replication switches to a slow replication mode. During interphase 14, visible morphology changes from the syncytial to cellular blastoderm, in a process called cellularization. Cellularization is the first morphological process that depends on zygotic gene products [29, 30].

However, the first signs of MBT are already visible earlier. As mentioned above, the extending interphases in NC11–14 depend on zygotic transcription. The first transcripts and activated RNA polymerase II (Pol II) can be already detected in pre-blastoderm stages. Transcription slowly increases until cycle 12. In cycle 13 many zygotic genes are clearly expressed [31]. Genome-wide analysis showed that gene expression is initiated at different time points throughout early development [32, 33], suggesting that rather than a sharp switch, MZT is likely regulated by multiple and diverse mechanisms [9, 34, 35]. The timing of these multiple and diverse mechanisms depends, to a certain degree, on the ratio of nuclear and cytoplasmic content (N:C ratio). This is further discussed in Subheading 5.

Approximately, two-thirds of all genes are contained in *Drosophila* eggs as maternal mRNAs [34, 36]. A third of all maternal transcripts are eliminated in stages leading to MBT in three ways [36]: First, maternally encoded factors activate mRNA degradation of over 20% of maternal transcripts after egg activation in a ZGA-independent manner [34, 37–39]. The RNA-binding protein Smaug is such a factor, acting together with the CCR4/POP2/NOT deadenylase complex [38, 40, 41]. Another RNA-binding protein, Brain Tumor, functions in a similar way [42]. Second, 15% of maternal mRNAs are eliminated depending on zygotic transcription during MBT [43, 44]. Third, microRNAs induce maternal RNA degradation. More than 100 maternal transcripts are degraded depending on zygotically expressed microRNAs from the *miR-309* cluster, which is activated by the early zygotic transcription factor Vielfältig/Zelda [45–47].

2 Mechanism of Zygotic Genome Activation

Transcription of the zygotic genome only begins shortly after fertilization [48]. The highly dynamic transcription profile was characterized by number of methods, including high-throughput strategies, global run-on sequencing (GRO-seq), and fluorescent labeling of nascent RNA [14, 49–52]. In general, the initiation of low-level zygotic transcription, mostly of signaling and patterning genes, already appears before NC10 ahead of large-scale ZGA [31, 53]. These include small and intron-less genes, as well as genes with TAGteam DNA motif in the control region [36]. A comparable profile is also observed in that of the zebrafish [54]. Full activation of zygotic transcription is observed during MBT, when thousands of genes are transcriptionally activated and transcribed in high levels. Taken together, the activation of the zygotic genome is a gradual process rather than a single sharp switch. This suggests that ZGA is triggered by multiple and diverse events [9, 34, 35].

A contribution to ZGA is intrinsically provided by the division of nuclei and doubling of DNA with every nuclear cycle. Even with a constant activity of the individual zygotic transcription units, the total number of transcripts would exponentially increase. In general, zygotic transcription is quantified in relation to the number of embryos, total mass of embryos (protein or total RNA content), or in comparison to an abundant maternal RNA, such as ribosomal RNA. Most of the older data are based on samples prepared from mixed stages comprising several nuclear division cycles. Alternatively, zygotic transcription may be normalized to the number of nuclei in an embryo. Given recent technological advances, transcription profiling can be conducted with few or even single *Drosophila* embryos, allowing highly accurate staging according to the nuclear division cycle [33, 55]. Such normalization is important to reveal the actual transcriptional activity of a locus.

This hypothesis was tested with normalized transcriptional profiles of selected early zygotic genes (Fig. 1) based on a data set from manually staged embryos [56]. Normalization to the number of nuclei was performed with the assumption of a doubling with every cell cycle. In case of a doubling transcript number from one cycle to the next, this results in a zero value. An increase in transcript number higher than a factor two results in a positive number, whereas an increase less than a factor two, in a negative number (Fig. 1). This simple and exemplary calculation indicates that both the increasing number of nuclei and an increased activity of the transcription units contribute to the overall increase in zygotic transcripts per embryo. There is, however, also transcript-dependent variation. A similar finding was reported recently for dorsoventrally patterning genes [57]. This indicates that depending on the zygotic gene, both an increased activity of individual transcription units and an increased number of transcription units/nuclei contribute to ZGA.




Fig. 1 Zygotic transcription and number of nuclei. (a) Number of selected zygotic transcripts based on NanoString analysis with extracts from manually staged embryos plotted on a logarithmic scale. (b) The number of transcripts was normalized to the number of nuclei that double with every cycle. Plotted is the difference of \log_2 of the number of transcripts from one cycle to the previous cycle minus 1. The number of transcripts in pre-blastoderm stages is not included. Transcripts for the ribosomal protein L32 serve as a reference. Staging by the nuclear cycle, pre-blastoderm stage (Pre) and late cellularization (14-I). Data are from Sung et al. [56]

2.1 *Vielfältig/Zelda* Functions in ZGA Regulation

The zinc-finger protein *Vielfältig/Zelda* (*Vfl/Zld*) plays a major role in ZGA. *Vfl/Zld* specifically binds to TAGteam elements in the early *Drosophila* embryo. The TAGteam CAGGTAG sequence was identified by genome-wide studies as a general *cis*-regulatory element and as the most highly enriched regulatory motif in genes involved in anterior-posterior patterning [36, 58, 59]. *Vfl/Zld* is an essential transcriptional activator during early zygotic gene expression, as demonstrated by the strongly reduced (but not absent) expression of many early zygotic genes in embryos from females with *Vfl/Zld* mutant germline [60]. *Vfl/Zld* is maternally deposited and uniformly distributed throughout the egg and early embryo. The *Vfl/Zld* protein levels increase coincidentally with the activation of zygotic genome during pre-blastoderm stage, prior to large-scale transcription [49, 61].

Vfl/Zld consists of a cluster of four zinc fingers and a low-complexity activation domain, both of which are required for promoting DNA binding and mediating transcriptional activation [62]. *Vfl/Zld* binding to promoters is detected already in NC8 for particular genes and roughly a thousand genes during NC10 [63, 64]. The DNA binding is maintained at least until NC14 [49]. During ZGA, *Vfl/Zld*-binding sites are highly enriched specifically in regions of accessible chromatin, allowing transcription factors to subsequently bind and drive zygotic transcription [63, 64]. Thus, *Vfl/Zld* acts as a co-activator during MZT. *Vfl/Zld* also controls the accurate temporal and spatial expression of microRNAs [46].

2.2 RNA Polymerase II Pausing

The binding of Pol II to promotor sequences is the key to transcriptional activation and elongation. Pol II regulates ZGA by three distinct binding statuses: active, no binding, and stalled/paused [65]. Among them, paused Pol II is critical in *Drosophila* ZGA, because approximately 100 genes are bound by active Pol II from NC8 to 12, yet in NC14, over 4000 promoters are occupied by Pol II at the transcription start site (TSS) [55, 66]. Furthermore, compared with NC12, loci with paused Pol II near the TSS show a significant increase in NC13 [67].

2.3 Epigenetics and ZGA

Epigenetic marks, including histone modifications and chromatin remodeling, dramatically change in early embryogenesis and MBT. Formation of heterochromatin correlates with the emergence of late replication. Heterochromatin Protein 1 (HP1) together with histone modifications on H3K9 and H3K4 is involved in establishing of tightly packed chromatin structure [68, 69]. Modifications of lysine acetylation and methylation in histones H3 and H4 appear during MZT. In zebrafish, a striking change in histone modification correlates with ZGA [70]. An increase in histone methylation during MZT matches high level of zygotic transcription [70, 71]. In *Xenopus* embryo, maternally provided histones H3/H4 and their modification states control the regulation of transcriptional activation and cell cycle lengthening [72, 73]. Similarly, during *Drosophila* early development, genome-wide studies showed that domains of histone methylation H3K4me1, H3K4me3, H3K27me3, and H3K36me3 increased from undetectable to widespread level at NC14 [48, 55, 74]. Levels of acetylation on H3K9 appear correspondingly to methylation marks, whereas H3K18ac, H3K27ac, and H4K8ac levels are evidently precocious at NC12 [48]. These early appearing acetylation marks are strongly correlated with maternal DNA-binding protein Vfl/Zld, demonstrating that Vfl/Zld may regulate transcriptional activation by recruiting histone acetylation, thus allowing opening of genome state [34, 48]. In contrast, the mark H4K5ac, whose level was previously shown to bookmark active transcription in mammalian cells, decreases from NC8 with the slowdown of the cleavage cycles [48, 75]. In addition to histone modifications, remodeling of nucleosomes and linker histones with histone variants may contribute to ZGA. *Drosophila* maternal-specific linker histone H1 dBigH1 is replaced by somatic H1 in early development [76]. dBigH1 seems to suppress ZGA, since increased levels of activated Pol II and expression of zygotic genes are observed in embryos with reduced dBigH1 levels [76].

Both histone modification and Vfl/Zld DNA binding ultimately affect transcriptional activation by altering chromatin accessibility. Highly accessible chromatin regions are locally and globally marked by H3/H4 acetylation and Vfl/Zld enrichment from NC8 to 12 in *Drosophila* [77]. In NC13, however, thousands of

enhancers and promoters with nucleosome-free regions accumulate additional transcription factors in a cascade way [48, 78]. This phenomenon has also been observed in zebrafish [79].

2.4 Other Regulators

Drosophila zygotic transcription is modulated by multiple factors including *cis*-regulatory elements. For instance, TATA-dependent promoters, as well as enhancers, are central in transcriptional regulation [80, 81]. Distinct enhancer-core-promoter specificities ensure that developmental and housekeeping genes are activated precisely across the entire genome [81]. Likewise, the post-transcriptional regulation of TATA-binding protein (TBP) affects transcription pattern together with the earliest transcribed genes during the MZT [55]. Smaug may involve ZGA regulation through maternal clearance of transcription factor *tramtrack* mRNA, which is involved in triggering transcription of transcripts depending on the N:C ratio [38, 53].

3 Switch in Cell Cycle Mode During the MBT

The cell cycle switch from a fast syncytial mode to a mode with slow replication and extended G2 phase is the most obvious aspect of MBT in morphological terms. A long-standing question is the functional relationship of the cell cycle switch with ZGA. According to one model, the cell cycle switch allows for the strong increase in zygotic transcription (Fig. 2) [82]. In the opposing model, zygotic




Fig. 2 Models for the control of cell cycle remodeling during MBT. **(a)** The onset of zygotic transcription leads to the activation of the DNA checkpoint due to interference of transcription and replication as well as expression of mitotic inhibitors. These two processes lead to the cell cycle remodeling. **(b)** Activation of the DNA checkpoint, caused by limiting amounts of replication factors, for example, triggers a slowdown and subsequent pause of the cell cycle. The longer interphase promotes zygotic transcription

transcription triggers the remodeling of the cell cycle [56, 67]. Depending on the experimental system, strong experimental evidence speaks in favor of the first or the second model. A synthesis has not been achieved, yet.

3.1 Cell Cycle Regulation in *Drosophila* Early Embryogenesis

Cyclin and its partner cyclin-dependent kinase (Cdk) are essential for cell cycle control. In *Drosophila*, cyclin A/B/B3:Cdk1 complexes regulate entry into M phase [20, 83]. The rapid S phases in pre-MBT cycles are maternally controlled, and the catalytic activity level of cyclin:Cdk1 complexes determines the timing for mitotic entry [21, 84]. Distinct mechanisms regulate cyclin:Cdk1 complexes in pre-MBT: First, during each nuclear division, Cyclin A, B and B3 proteins are synthesized in S phase by maternally supplied mRNA [85, 86], and degraded in mitosis by the ubiquitin pathway [87, 88]. Cyclin A, B, and B3 fulfill a redundant but essential function, as RNAi-mediated depletion stops the syncytial cycles [20, 89]. Cyclin B levels also contribute to the cell cycle switch as changes in *cyclin B* gene dose affect the number of nuclear divisions [90]. Second, the inhibitory phosphorylation of T14Y15 sites of Cdk1 are pairwise regulated by maternally supplied kinases Wee1/Myt1 and phosphatase Cdc25/Twine [85, 91–95]. Therefore, Cdk1 is timely activated and inactivated by controlling T14Y15 inhibitory phosphorylation sites [96].

3.2 Cdc25/Twine Degradation at the MBT

In NC14 and to a certain degree already in NC12 and 13, S phase lengthens and a G2 phase is introduced. Central to these changes is the induced inactivation and final degradation of the phosphatase Cdc25/Twine [97, 98] (Fig. 3). *Drosophila* Cdc25/Twine is a dual specificity phosphatase that activates cyclin:Cdk1 complexes by removing inhibitory phosphates from the ATP-binding sites T14 and Y15 [22, 87, 99, 100]. Twine protein is present in high levels during the pre-MBT cycles. Twine protein localization is dynamic with a nuclear accumulation during interphases and uniform dispersal during mitosis [98]. The half-life of Twine was estimated to about 20 min during pre-MBT cycles [98]. Yet with the beginning of NC14, Twine becomes destabilized as indicated by the shortening of its half-life to only about 5 min [98]. Degradation of Twine is required for the cell cycle switch because embryos expressing a more stable version of Twine protein (Twine¹⁰⁶⁻¹⁸⁰) undergo an extra mitotic division [98]. The rapid destabilization is the key to the cell cycle switch during MBT, as it depends on the N:C ratio and on zygotic transcription [98].

Prior to MBT, the steady-state level of Twine is relatively stable due to balanced synthesis and degradation. The link of zygotic transcription and the switch-like decrease in the half-life of Twine suggests that zygotic factors may be involved. One of these factors is the pseudokinase Tribbles [101–103], as RNAi-mediated depletion of *tribbles* accelerates Twine degradation [97].




Fig. 3 Model of cell cycle remodeling in *Drosophila*. Cyclin:Cdk1 is activated by the phosphatase Cdc25 and inactivated by the kinases Myt1/Wee1. In pre-MBT Cyclin:Cdk1 activity is high and promotes fast cell cycles. During MBT the balance of Cyclin:Cdk1 control is shifted toward low activity. Cdc25 is inhibited by the DNA checkpoint, which is activated by DNA stress caused by interference of DNA replication and zygotic transcription. In addition, the zygotic mitotic inhibitors, Tribbles and Frühstart, promote Cdc25 degradation and inhibition of the Cyclin:Cdk1 complexes, respectively

However, *tribbles* is not essential for the cell cycle switch, since embryos deficient for maternal and zygotic *tribbles* do not undergo an extra nuclear cycle [101, 102]. The mechanism for how *tribbles* induces Twine degradation remains unknown, but in other organisms such as yeast, *Xenopus*, and human cells, Cdc25 (or Cdc25C) degradation is induced by phosphorylation due to multiple pathways [56, 104, 105]. In addition to induced destabilization of Cdc25/Twine at NC14, additional mechanisms control pre-MBT levels and activity of Twine. The number of pre-MBT cell cycles is rather insensitive to changes in *twine* gene dose. A tripling of *twine* gene dose to $6 \times \text{twine}[+]$ induces an extra nuclear division in only a few embryos [106], suggesting that mechanisms exist that make Twine protein levels independent of gene dose.

The second *Drosophila* homologue of Cdc25, String, has distinct developmental functions in cell cycle control [84, 107]. String but not Twine is required for mitotic entry in zygotically controlled cycles 14–16. In contrast to these later stages, *string* is not required for progression of the syncytial cell cycles [84]. Premature expression of *string* is sufficient to trigger mitotic entry during later stages

of embryonic development but not in pre-MBT stages [84]. Although both *string* and *twine* mRNAs are destructed in interphase 14 [106], String protein stability gradually decreases during syncytial cycle without a sharp switch before MBT [97]. String protein turnover is due to increased checkpoint activity [98].

3.3 DNA Replication Checkpoint at NC13

Before the switch in cell cycle mode in NC14 in *Drosophila*, S phases show a progressive lengthening from 3.4 min in NC8 to 14 min in NC13 [21, 108]. A critical regulator of the slowdown of replication is the *Drosophila* homologue of checkpoint kinase Chk1, Grapes [109]. Grapes starts to inhibit cyclin:Cdk1 activity by promoting the activity of kinases Wee1/Myt1 and suppressing the activity of phosphatase Cdc25, thereby shifting the balance to T14Y15 inhibitory phosphorylation of Cdk1 from NC11 onward [109, 110]. Grapes mediates the DNA replication checkpoint and ensures that cells do not enter mitosis while replication is ongoing. *grapes* mutants prematurely enter mitosis during syncytial divisions, which leads to mitotic catastrophe, as incompletely replicated chromosomes cannot be segregated in anaphase [109, 110]. The checkpoint kinase, ataxia telangiectasia and Rad3-related (ATR, Mei-41 in *Drosophila*), acts upstream and activates Chk1/Grapes similar as in *Xenopus* [111, 112]. *mei-41* mutants show a similar phenotype during syncytial divisions as *grapes*, indicating a functional replication checkpoint is required at the MBT [67].

In *Drosophila* the DNA checkpoint is triggered by ZGA. Blocking transcription by α -amanitin in *Drosophila* pre-MBT embryos does not suppress lethality of *mei-41* mutant [67]. Nonetheless, embryos from *mei-41 Vfl/Zld* double mutant mothers could partially suppress the mitotic catastrophe, indicating that replication has been finished in time [67]. These observations are consistent with the model that zygotic transcription reduces replication speed and induces DNA stress, leading to DNA checkpoint activation at ZGA [56, 67].

3.4 Other Regulators

In *Drosophila*, cyclin-dependent kinase inhibitor (CKI) Frühstart is another zygotic regulator, which functions to inhibit cyclin:Cdk1 activity by binding the hydrophobic patch of cyclins, thereby interfering with Cdk1 substrate recognition [101, 113, 114]. Together with large-scale ZGA, *frühstart* starts transcription immediately after mitosis 13, and generates a uniform cell cycle pause in cycle 14 [114]. In the absence of Frühstart, embryos enter an extra round of nuclear division especially in embryos with extra copies of *twine*[+] [114]. The expression of Frühstart depends on the N:C ratio, suggesting that Frühstart is involved in the link of N:C with cell cycle regulation [115]. Wee1 and Myt1 kinases are Cdk1 inhibitors that oppose functions to Cdc25 phosphatases [91–93, 116, 117] (Fig. 3). Wee1 can be activated by Grapes, and inhibits Cdk1 activity by adding inhibitory phosphorylation at T14 and Y15 sites [9, 118, 119]. Cyclin:Cdk1 activity is also influenced by

some other factors such as mitotic kinase Aurora-A and acquisition of late-replicating heterochromatin domains [95, 120].

In summary, the switch of the cell cycle from a fast syncytial mode to a slow embryonic mode is controlled on two levels of inhibition: (1) indirectly by interference of zygotic transcription with DNA replication and subsequent activation of the DNA checkpoint, (2) directly by expression of zygotic genes encoding mitosis inhibitors.

4 What Is the Trigger for MBT?

The MBT cell cycle switch depends on ZGA (Fig. 2). First, injection of α -amanitin, a Pol II inhibitor, before MBT induces an extra synchronized mitotic division, indicating that widespread zygotic transcription is required for the cell cycle switch in *Drosophila* [106]. Second, ZGA correlates with DNA stress. About 80% of the RpA-70-GFP-binding sites in early MBT cycles also have RNA Pol II bound [67]. RpA70-GFP marks sites of DNA stress [121]. This indicates that ZGA causes DNA stress and activates the DNA checkpoint [67]. Third, a precocious onset of zygotic transcription is sufficient for an earlier MBT [56]. Fourth, dependent on ZGA, Tribbles and other factors trigger Twine destruction in NC14, resulting in inhibition of Cdk1 activation, thereby pausing the cell cycle [101, 102].

The essential role of the DNA checkpoint for triggering MBT was initially shown by the analysis of the checkpoint mutants, *grapes/Chk1* and *mei-41/ATR*, in *Drosophila* [109, 111]. Embryos from *grapes* females do not switch the cell cycle mode and do not enter MBT, indicating that the DNA checkpoint is required for MBT in *Drosophila* [67, 109]. Based on the observation that *grapes* embryos would not express zygotic genes, the authors concluded that the checkpoint would be upstream of ZGA [109]. Recent data clearly show, however, that ZGA is normal in checkpoint-deficient embryos and that the initial observation was probably due to technical difficulties in detecting expression of early zygotic genes [67].

An alternative source for checkpoint activation beside interference of replication and transcription are limiting amounts of replication factors. Experiments from mostly *Xenopus* support this model (Fig. 2). In *Xenopus* embryos slowdown of DNA replication has been proposed to be upstream of ZGA [82]. The replication factors Cut5, RecQ4, Treslin, and Drf1 become limiting in MBT, which leads to an activation of the DNA checkpoint, slowdown of the cell cycle, and ZGA [82].

In summary, in vivo and genetic experiments provide strong evidence for the model that ZGA is the trigger for MBT in *Drosophila*. ZGA acts upstream of cell cycle control, including the DNA checkpoint and degradation of Cdc25/Twine. First, ZGA is required for MBT and timely cell cycle pause; second, ZGA is

associated with induction of replication stress in time and space (on the chromosome); third, precocious ZGA leads to precocious MBT. In other organisms experimental evidence mainly in *Xenopus* speaks in favor of the alternative model, i.e., that cell cycle control acts upstream ZGA. However not all three criteria are fulfilled in vivo: the mechanism should be necessary, sufficient, and temporally and spatially associated with MBT.

5 What Is the Timer for MBT?

A central unresolved question concerning MBT is the timing mechanism for the associated processes including ZGA and number of pre-MBT cell cycles. Tight control of the cell cycle is important for further embryonic development, since the number of divisions determines the cell number and size. Too few cells may be incompatible with the formation of stripes of pair-rule gene expression, for example, as stripes should be at least one cell wide.

5.1 Molecular Clocks

With the onset of embryonic development, fertilization may trigger a molecular clock, on which MBT and its associated processes may depend. A conceivable mechanism is translation of certain maternal mRNAs, which would lead to a time-dependent accumulation of the product following onset after fertilization. Translational regulators such as FMRP are required for MBT regulation in *Drosophila*, through dynamically regulating RNA metabolism and controlling the availability of specific transcripts, as well as mediating the *frühstart* mRNA activation level [122, 123]. A target for translational regulation may be Vfl/Zld, whose protein level increases during blastoderm concomitantly with activation of zygotic transcription [34, 124].

Maternal RNA degradation may represent a second such a mechanism constituting a molecular clock. A large fraction of these maternal RNAs is degraded following egg activation and independent of zygotic transcription. For some RNAs at least, the degradation proceeds with a constant speed [38, 56], and may in this manner constitute a molecular clock. It has been proposed that the speed of RNA degradation affects the number of nuclear divisions, as expression levels of *smaug* affect the timing of MBT [40, 125]. Distinct from Vfl/Zld, Smaug reaches its peak expression level at NC10, and performs downregulation at the MBT [38, 125]. Smaug is functional to mRNA clearance, and times the ZGA through inducing the destruction of maternal transcriptional inhibitor [27].

5.2 N:C Ratio as a Clock

In contrast to a molecular clock as an absolute timer, more evidence speaks in favor of a regulatory process. The morphologically visible MBT depends on genome ploidy, because haploid embryos undergo one more division and tetraploid embryos, one less

division [11]. It has been proposed that the N:C ratio represents the timer for MBT. Nuclear content is determined by the amount of DNA or chromatin, which doubles with every cell cycle, whereas cytoplasmic content remains constant during cleavage divisions. The embryo may measure the N:C in that the increasing amount of chromatin titrates a constant cytoplasmic factor until this becomes rate-limiting [6, 53]. Potential cytoplasmic factors are repressors of transcription, replication, or the cell cycle, for example. In *Xenopus* embryos, DNA content is important for MBT [5, 7]. Injection of purified DNA leads to precocious onset of zygotic transcription, as measured by total transcription rate [7]. However, the amount of DNA seems not to be the only determinant, since an increased or decreased nuclear volume, while keeping the DNA content unchanged, leads to a precocious or delayed MBT including zygotic activation and corresponding cell cycle remodeling [126]. Similar findings come from zebrafish that the timing of ZGA is governed by the N:C ratio [13].

It is unclear what is titrated by the exponentially increasing amount of DNA and chromatin, but maternal histones proteins H3/H4 may be a central factor [72]. Depletion and overexpression of H3/H4 delay the cell cycle switch, and also induce premature transcriptional activation [72]. In *Drosophila* embryos, the maternal form of the linker histone H1 dBigH1 has been implicated in the timing of MBT [76]. Maternal dBigH1 is replaced by the somatic form in early embryogenesis. Embryos with half of the maternal contribution and lacking zygotic expression show increased levels of activated Pol II and zygotic gene expression. However, the link of dBigH1 to MBT remains unclear as mutant defects and embryonic genotypes were not analyzed with sufficiently high temporal resolution and with respect to MBT and ZGA.

The replication factors Cut5, RecQ4, Treslin, and Drf1 have been found to be limiting for replication initiation during MBT in *Xenopus* embryos [82]. Titration of the maternal pool of these replication factors by the exponentially increasing chromatin leads to slower replication initiation, ZGA, longer interphases, and DNA checkpoint activation.

Other cytoplasmic factors may also be titrated, such as metabolites. It has been proposed that deoxynucleotides may serve as a marker for the cytoplasm [127]. The maternal pool may be incorporated in the exponentially increasing amounts of DNA. The existence of such a maternal pool is well known, as inhibition of zygotic synthesis by hydroxyurea (HU), which inhibits the NDP reductase, causes a cell cycle arrest only briefly before MBT [127].

Although it is clear that ploidy determines the number of pre-MBT cell cycles in model organisms, it is much less clear whether all of the MBT-associated processes, including ZGA, cell cycle, RNA degradation, are controlled by the N:C ratio. Haploid *Drosophila* embryos switch the cell cycle mode only after an extra

division 14 in NC15 [115, 128]. In contrast, ZGA does not depend on the N:C ratio in *Drosophila*. Although older data indicated a link of ploidy and ZGA in *Drosophila* [53], genome-wide analysis of embryonic transcripts with carefully staged *Drosophila* embryos revealed that the majority of zygotic transcripts (127 out of 215 genes) show an expression profile comparable between haploid and diploid embryos [115]. These data suggest that ZGA timing is controlled by a molecular clock in *Drosophila*. However, a small set of zygotic transcripts (88 out of 215 genes) shows clearly delayed expression in haploid embryos [115]. This small gene set includes genes encoding mitotic inhibitors such as *Frühstart* [114], which are involved in the MBT-associated remodeling of the cell cycle.

6 Conclusions

Recent years brought striking advances in our understanding of zygotic genome activation and its relation to MBT. This is mainly due to improved technology now allowing to analyze transcriptional activity and chromosome status with high resolution and importantly with very little material, down to single embryos. In this way, the variation and limited temporal resolution of mixtures of many embryos can be overcome. Despite this progress, there is no unifying model for zygotic genome activation, MBT, and cell cycle control. Conclusion on central questions and favored models depend on the experimental system. Strong evidence supports the model that DNA replication onset triggers MBT and ZGA in *Xenopus*. However, the alternative model is supported by convincing experiments from *Drosophila*, where ZGA triggers MBT and cell cycle remodeling. It will be the task for future work to reconcile these opposing views. Having the new technologies available and standardized, we can expect new and surprising findings to come.

Acknowledgment

BL was supported by China Scholarship Council. The work in JG's laboratory was in part supported by the German Research Council (Deutsche Forschungsgemeinschaft (DFG) GR1945/3-1, SFB937/TP10).

References

1. Hiiragi T, Solter D (2004) First cleavage plane of the mouse egg is not predetermined but defined by the topology of the two apposing pronuclei. *Nature* 430(6997):360–364. doi:10.1038/nature02595
2. O'Farrell PH, Stumpff J, Su TT (2004) Embryonic cleavage cycles: how is a mouse like a fly? *Curr Biol* 14(1):R35–R45
3. O'Farrell PH (2015) Growing an embryo from a single cell: a hurdle in animal life. *Cold*

- Spring Harb Perspect Biol 7(11):a019042. doi:10.1101/cshperspect.a019042
4. Boveri T (1893) An organism produced sexually without characteristics of the mother. *Am Soc Nat* 27(315):222–232
 5. Gerhart JC (1980) Mechanisms regulating pattern formation in the amphibian egg and early embryo. In: Goldberger R (ed) *Biological regulation and development*, vol 2. Springer, Boston, MA, pp 133–316
 6. Newport J, Kirschner M (1982) A major developmental transition in early *Xenopus* embryos: I. Characterization and timing of cellular changes at the midblastula stage. *Cell* 30(3):675–686
 7. Newport J, Kirschner M (1982) A major developmental transition in early *Xenopus* embryos: II. Control of the onset of transcription. *Cell* 30(3):687–696
 8. Newport JW, Kirschner MW (1984) Regulation of the cell cycle during early *Xenopus* development. *Cell* 37(3):731–742
 9. Farrell JA, O'Farrell PH (2014) From egg to gastrula: how the cell cycle is remodeled during the *Drosophila* mid-blastula transition. *Annu Rev Genet* 48:269–294. doi:10.1146/annurev-genet-111212-133531
 10. Collart C, Owens ND, Bhaw-Rosun L, Cooper B, De Domenico E, Patrushev I, Sesay AK, Smith JN, Smith JC, Gilchrist MJ (2014) High-resolution analysis of gene activity during the *Xenopus* mid-blastula transition. *Development* 141(9):1927–1939. doi:10.1242/dev.102012
 11. Kane DA, Kimmel CB (1993) The zebrafish midblastula transition. *Development* 119(2):447–456
 12. Zamir E, Kam Z, Yarden A (1997) Transcription-dependent induction of G1 phase during the zebra fish midblastula transition. *Mol Cell Biol* 17(2):529–536
 13. Zhang M, Kothari P, Mullins M, Lampson MA (2014) Regulation of zygotic genome activation and DNA damage checkpoint acquisition at the mid-blastula transition. *Cell Cycle* 13(24):3828–3838. doi:10.4161/15384101.2014.967066
 14. Lee MT, Bonneau AR, Takacs CM, Bazzini AA, DiVito KR, Fleming ES, Giraldez AJ (2013) Nanog, Pou5f1 and SoxB1 activate zygotic gene expression during the maternal-to-zygotic transition. *Nature* 503(7476):360–364. doi:10.1038/nature12632
 15. Robertson S, Lin R (2015) The maternal-to-zygotic transition in *C. elegans*. *Curr Top Dev Biol* 113:1–42. doi:10.1016/bs.ctdb.2015.06.001
 16. Guven-Ozkan T, Nishi Y, Robertson SM, Lin R (2008) Global transcriptional repression in *C. elegans* germline precursors by regulated sequestration of TAF-4. *Cell* 135(1):149–160. doi:10.1016/j.cell.2008.07.040
 17. Sulston JE, Schierenberg E, White JG, Thomson JN (1983) The embryonic cell lineage of the nematode *Caenorhabditis elegans*. *Dev Biol* 100(1):64–119
 18. Rose L, Gonczy P (2014) Polarity establishment, asymmetric division and segregation of fate determinants in early *C. elegans* embryos. *WormBook*:1–43. doi:10.1895/wormbook.1.30.2
 19. Foe VE, Alberts BM (1983) Studies of nuclear and cytoplasmic behaviour during the five mitotic cycles that precede gastrulation in *Drosophila* embryogenesis. *J Cell Sci* 61:31–70
 20. McClelland ML, O'Farrell PH (2008) RNAi of mitotic cyclins in *Drosophila* uncouples the nuclear and centrosome cycle. *Curr Biol* 18(4):245–254. doi:10.1016/j.cub.2008.01.041
 21. Shermoen AW, McClelland ML, O'Farrell PH (2010) Developmental control of late replication and S phase length. *Curr Biol* 20(23):2067–2077. doi:10.1016/j.cub.2010.10.021
 22. Farrell JA, Shermoen AW, Yuan K, O'Farrell PH (2012) Embryonic onset of late replication requires Cdc25 down-regulation. *Genes Dev* 26(7):714–725. doi:10.1101/gad.186429.111
 23. Rabinowitz M (1941) Studies on the cytology and early embryology of the egg of *Drosophila melanogaster*. *J Morphol* 69(1):1–49
 24. Axton JM, Shamanski FL, Young LM, Henderson DS, Boyd JB, Orr-Weaver TL (1994) The inhibitor of DNA replication encoded by the *Drosophila* gene plutonium is a small, ankyrin repeat protein. *EMBO J* 13(2):462–470
 25. Fenger DD, Carminati JL, Burney-Sigman DL, Kashevsky H, Dines JL, Elfring LK, Orr-Weaver TL (2000) PAN GU: a protein kinase that inhibits S phase and promotes mitosis in early *Drosophila* development. *Development* 127(22):4763–4774
 26. Lee LA, Van Hoewyk D, Orr-Weaver TL (2003) The *Drosophila* cell cycle kinase PAN GU forms an active complex with PLUTONIUM and GNU to regulate embryonic divisions. *Genes Dev* 17(23):2979–2991. doi:10.1101/gad.1132603
 27. Laver JD, Marsolais AJ, Smibert CA, Lipshitz HD (2015) Regulation and function of maternal gene products during the maternal-to-zygotic transition in *Drosophila*. *Curr Top Dev Biol* 113:43–84. doi:10.1016/bs.ctdb.2015.06.007

28. Schubeler D, Scalzo D, Kooperberg C, van Steensel B, Delrow J, Groudine M (2002) Genome-wide DNA replication profile for *Drosophila melanogaster*: a link between transcription and replication timing. *Nat Genet* 32(3):438–442. doi:10.1038/ng1005
29. Edgar BA, Schubiger G (1986) Parameters controlling transcriptional activation during early *Drosophila* development. *Cell* 44(6): 871–877
30. Merrill PT, Sweeton D, Wieschaus E (1988) Requirements for autosomal gene activity during precellular stages of *Drosophila melanogaster*. *Development* 104(3):495–509
31. Ali-Murthy Z, Lott SE, Eisen MB, Kornberg TB (2013) An essential role for zygotic expression in the pre-cellular *Drosophila* embryo. *PLoS Genet* 9(4):e1003428. doi:10.1371/journal.pgen.1003428
32. Graveley BR, Brooks AN, Carlson JW, Duff MO, Landolin JM, Yang L, Artieri CG, van Baren MJ, Boley N, Booth BW, Brown JB, Cherbas L, Davis CA, Dobin A, Li R, Lin W, Malone JH, Mattiuzzo NR, Miller D, Sturgill D, Tuch BB, Zaleski C, Zhang D, Blanchette M, Dudoit S, Eads B, Green RE, Hammonds A, Jiang L, Kapranov P, Langton L, Perrimon N, Sandler JE, Wan KH, Willingham A, Zhang Y, Zou Y, Andrews J, Bickel PJ, Brenner SE, Brent MR, Cherbas P, Gingeras TR, Hoskins RA, Kaufman TC, Oliver B, Celniker SE (2011) The developmental transcriptome of *Drosophila melanogaster*. *Nature* 471(7339):473–479. doi:10.1038/nature09715
33. Lott SE, Villalta JE, Schroth GP, Luo S, Tonkin LA, Eisen MB (2011) Noncanonical compensation of zygotic X transcription in early *Drosophila melanogaster* development revealed through single-embryo RNA-seq. *PLoS Biol* 9(2):e1000590. doi:10.1371/journal.pbio.1000590
34. Harrison MM, Eisen MB (2015) Transcriptional activation of the zygotic genome in *Drosophila*. *Curr Top Dev Biol* 113:85–112. doi:10.1016/bs.ctdb.2015.07.028
35. Lee MT, Bonneau AR, Giraldez AJ (2014) Zygotic genome activation during the maternal-to-zygotic transition. *Annu Rev Cell Dev Biol* 30:581–613. doi:10.1146/annurev-cellbio-100913-013027
36. De Renzis S, Elemento O, Tavazoie S, Wieschaus EF (2007) Unmasking activation of the zygotic genome using chromosomal deletions in the *Drosophila* embryo. *PLoS Biol* 5(5):e117. doi:10.1371/journal.pbio.0050117
37. Tadros W, Houston SA, Bashirullah A, Cooperstock RL, Semotok JL, Reed BH, Lipshitz HD (2003) Regulation of maternal transcript destabilization during egg activation in *Drosophila*. *Genetics* 164(3):989–1001
38. Tadros W, Goldman AL, Babak T, Menzies F, Vardy L, Orr-Weaver T, Hughes TR, Westwood JT, Smibert CA, Lipshitz HD (2007) SMAUG is a major regulator of maternal mRNA destabilization in *Drosophila* and its translation is activated by the PAN GU kinase. *Dev Cell* 12(1):143–155. doi:10.1016/j.devcel.2006.10.005
39. Tadros W, Lipshitz HD (2009) The maternal-to-zygotic transition: a play in two acts. *Development* 136(18):3033–3042. doi:10.1242/dev.033183
40. Semotok JL, Cooperstock RL, Pinder BD, Vari HK, Lipshitz HD, Smibert CA (2005) Smaug recruits the CCR4/POP2/NOT deadenylase complex to trigger maternal transcript localization in the early *Drosophila* embryo. *Curr Biol* 15(4):284–294. doi:10.1016/j.cub.2005.01.048
41. Chen L, Dumelie JG, Li X, Cheng MH, Yang Z, Laver JD, Siddiqui NU, Westwood JT, Morris Q, Lipshitz HD, Smibert CA (2014) Global regulation of mRNA translation and stability in the early *Drosophila* embryo by the Smaug RNA-binding protein. *Genome Biol* 15(1):R4. doi:10.1186/gb-2014-15-1-r4
42. Laver JD, Li X, Ray D, Cook KB, Hahn NA, Nabeel-Shah S, Kekis M, Luo H, Marsolais AJ, Fung KY, Hughes TR, Westwood JT, Sidhu SS, Morris Q, Lipshitz HD, Smibert CA (2015) Brain tumor is a sequence-specific RNA-binding protein that directs maternal mRNA clearance during the *Drosophila* maternal-to-zygotic transition. *Genome Biol* 16:94. doi:10.1186/s13059-015-0659-4
43. Walser CB, Lipshitz HD (2011) Transcript clearance during the maternal-to-zygotic transition. *Curr Opin Genet Dev* 21(4):431–443. doi:10.1016/j.gde.2011.03.003
44. Bashirullah A, Halsell SR, Cooperstock RL, Kloc M, Karaïskakis A, Fisher WW, Fu W, Hamilton JK, Etkin LD, Lipshitz HD (1999) Joint action of two RNA degradation pathways controls the timing of maternal transcript elimination at the midblastula transition in *Drosophila melanogaster*. *EMBO J* 18(9): 2610–2620. doi:10.1093/emboj/18.9.2610
45. Bushati N, Stark A, Brennecke J, Cohen SM (2008) Temporal reciprocity of miRNAs and their targets during the maternal-to-zygotic transition in *Drosophila*. *Curr Biol* 18(7): 501–506. doi:10.1016/j.cub.2008.02.081
46. Fu S, Nien CY, Liang HL, Rushlow C (2014) Co-activation of microRNAs by Zelda is essential for early *Drosophila* development. *Development* 141(10):2108–2118. doi:10.1242/dev.108118

47. Huntzinger E, Izaurralde E (2011) Gene silencing by microRNAs: contributions of translational repression and mRNA decay. *Nat Rev Genet* 12(2):99–110. doi:10.1038/nrg2936
48. Li XY, Harrison MM, Villalta JE, Kaplan T, Eisen MB (2014) Establishment of regions of genomic activity during the *Drosophila* maternal to zygotic transition. *Elife* 3. doi:10.7554/eLife.03737
49. Harrison MM, Li XY, Kaplan T, Botchan MR, Eisen MB (2011) Zelda binding in the early *Drosophila* melanogaster embryo marks regions subsequently activated at the maternal-to-zygotic transition. *PLoS Genet* 7(10):e1002266. doi:10.1371/journal.pgen.1002266
50. Lecuyer E, Yoshida H, Parthasarathy N, Alm C, Babak T, Cerovina T, Hughes TR, Tomancak P, Krause HM (2007) Global analysis of mRNA localization reveals a prominent role in organizing cellular architecture and function. *Cell* 131(1):174–187. doi:10.1016/j.cell.2007.08.003
51. Saunders A, Core LJ, Sutcliffe C, Lis JT, Ashe HL (2013) Extensive polymerase pausing during *Drosophila* axis patterning enables high-level and pliable transcription. *Genes Dev* 27(10):1146–1158. doi:10.1101/gad.215459.113
52. Ferraro T, Lucas T, Clemot M, De Las Heras Chanes J, Desponds J, Coppey M, Walczak AM, Dostatni N (2016) New methods to image transcription in living fly embryos: the insights so far, and the prospects. *Wiley Interdiscip Rev Dev Biol* 5(3):296–310. doi:10.1002/wdev.221
53. Pritchard DK, Schubiger G (1996) Activation of transcription in *Drosophila* embryos is a gradual process mediated by the nucleocytoplasmic ratio. *Genes Dev* 10(9):1131–1142
54. Heyn P, Kircher M, Dahl A, Kelso J, Tomancak P, Kalinka AT, Neugebauer KM (2014) The earliest transcribed zygotic genes are short, newly evolved, and different across species. *Cell Rep* 6(2):285–292. doi:10.1016/j.celrep.2013.12.030
55. Chen K, Johnston J, Shao W, Meier S, Staber C, Zeitlinger J (2013) A global change in RNA polymerase II pausing during the *Drosophila* midblastula transition. *Elife* 2:e00861. doi:10.7554/eLife.00861
56. Sung HW, Spangenberg S, Vogt N, Grosshans J (2013) Number of nuclear divisions in the *Drosophila* blastoderm controlled by onset of zygotic transcription. *Curr Biol* 23(2):133–138. doi:10.1016/j.cub.2012.12.013
57. Sandler JE, Stathopoulos A (2016) Quantitative single-embryo profile of *Drosophila* genome activation and the dorsal-ventral patterning network. *Genetics* 202(4):1575–1584. doi:10.1534/genetics.116.186783
58. ten Bosch JR, Benavides JA, Cline TW (2006) The TAGteam DNA motif controls the timing of *Drosophila* pre-blastoderm transcription. *Development* 133(10):1967–1977. doi:10.1242/dev.02373
59. Li XY, MacArthur S, Bourgon R, Nix D, Pollard DA, Iyer VN, Hechmer A, Simirenko L, Stapleton M, Luengo Hendriks CL, Chu HC, Ogawa N, Inwood W, Sementchenko V, Beaton A, Weiszmänn R, Celniker SE, Knowles DW, Gingeras T, Speed TP, Eisen MB, Biggin MD (2008) Transcription factors bind thousands of active and inactive regions in the *Drosophila* blastoderm. *PLoS Biol* 6(2):e27. doi:10.1371/journal.pbio.0060027
60. Liang HL, Nien CY, Liu HY, Metzstein MM, Kirov N, Rushlow C (2008) The zinc-finger protein Zelda is a key activator of the early zygotic genome in *Drosophila*. *Nature* 456(7220):400–403. doi:10.1038/nature07388
61. Staudt N, Fellert S, Chung HR, Jackle H, Vorbruggen G (2006) Mutations of the *Drosophila* zinc finger-encoding gene *vielfaltig* impair mitotic cell divisions and cause improper chromosome segregation. *Mol Biol Cell* 17(5):2356–2365. doi:10.1091/mbc.E05-11-1056
62. Hamm DC, Bondra ER, Harrison MM (2015) Transcriptional activation is a conserved feature of the early embryonic factor Zelda that requires a cluster of four zinc fingers for DNA binding and a low-complexity activation domain. *J Biol Chem* 290(6):3508–3518. doi:10.1074/jbc.M114.602292
63. Foo SM, Sun Y, Lim B, Ziukaite R, O'Brien K, Nien CY, Kirov N, Shvartsman SY, Rushlow CA (2014) Zelda potentiates morphogen activity by increasing chromatin accessibility. *Curr Biol* 24(12):1341–1346. doi:10.1016/j.cub.2014.04.032
64. Schulz KN, Bondra ER, Moshe A, Villalta JE, Lieb JD, Kaplan T, McKay DJ, Harrison MM (2015) Zelda is differentially required for chromatin accessibility, transcription factor binding, and gene expression in the early *Drosophila* embryo. *Genome Res* 25(11):1715–1726. doi:10.1101/gr.192682.115
65. Zeitlinger J, Stark A, Kellis M, Hong JW, Nechaev S, Adelman K, Levine M, Young RA (2007) RNA polymerase stalling at developmental control genes in the *Drosophila* melanogaster embryo. *Nat Genet* 39(12):1512–1516. doi:10.1038/ng.2007.26

66. Boettiger AN, Levine M (2009) Synchronous and stochastic patterns of gene activation in the *Drosophila* embryo. *Science* 325(5939): 471–473. doi:10.1126/science.1173976
67. Blythe SA, Wieschaus EF (2015) Zygotic genome activation triggers the DNA replication checkpoint at the midblastula transition. *Cell* 160(6):1169–1181. doi:10.1016/j.cell.2015.01.050
68. Rudolph T, Yonezawa M, Lein S, Heidrich K, Kubicek S, Schafer C, Phalke S, Walther M, Schmidt A, Jenuwein T, Reuter G (2007) Heterochromatin formation in *Drosophila* is initiated through active removal of H3K4 methylation by the LSD1 homolog SU(VAR)3-3. *Mol Cell* 26(1):103–115. doi:10.1016/j.molcel.2007.02.025
69. Yuan K, O'Farrell PH (2016) TALE-light imaging reveals maternally guided, H3K9me2/3-independent emergence of functional heterochromatin in *Drosophila* embryos. *Genes Dev*. doi:10.1101/gad.272237.115
70. Lindeman LC, Andersen IS, Reiner AH, Li N, Aanes H, Ostrup O, Winata C, Mathavan S, Muller F, Alestrom P, Collas P (2011) Prepatterning of developmental gene expression by modified histones before zygotic genome activation. *Dev Cell* 21(6):993–1004. doi:10.1016/j.devcel.2011.10.008
71. Vastenhouw NL, Zhang Y, Woods IG, Imam F, Regev A, Liu XS, Rinn J, Schier AF (2010) Chromatin signature of embryonic pluripotency is established during genome activation. *Nature* 464(7290):922–926. doi:10.1038/nature08866
72. Amodeo AA, Jukam D, Straight AF, Skotheim JM (2015) Histone titration against the genome sets the DNA-to-cytoplasm threshold for the *Xenopus* midblastula transition. *Proc Natl Acad Sci U S A* 112(10):E1086–E1095. doi:10.1073/pnas.1413990112
73. Hontelez S, van Kruijsbergen I, Georgiou G, van Heeringen SJ, Bogdanovic O, Lister R, Veenstra GJ (2015) Embryonic transcription is controlled by maternally defined chromatin state. *Nat Commun* 6:10148. doi:10.1038/ncomms10148
74. Boettiger AN, Bintu B, Moffitt JR, Wang S, Beliveau BJ, Fudenberg G, Imakaev M, Mirny LA, Wu CT, Zhuang X (2016) Super-resolution imaging reveals distinct chromatin folding for different epigenetic states. *Nature* 529(7586):418–422. doi:10.1038/nature16496
75. Zhao R, Nakamura T, Fu Y, Lazar Z, Spector DL (2011) Gene bookmarking accelerates the kinetics of post-mitotic transcriptional reactivation. *Nat Cell Biol* 13(11):1295–1304. doi:10.1038/ncb2341
76. Perez-Montero S, Carbonell A, Moran T, Vaquero A, Azorin F (2013) The embryonic linker histone H1 variant of *Drosophila*, dBigH1, regulates zygotic genome activation. *Dev Cell* 26(6):578–590. doi:10.1016/j.devcel.2013.08.011
77. Li XY, Thomas S, Sabo PJ, Eisen MB, Stamatoyannopoulos JA, Biggin MD (2011) The role of chromatin accessibility in directing the widespread, overlapping patterns of *Drosophila* transcription factor binding. *Genome Biol* 12(4):R34. doi:10.1186/gb-2011-12-4-r34
78. Thomas S, Li XY, Sabo PJ, Sandstrom R, Thurman RE, Canfield TK, Giste E, Fisher W, Hammonds A, Celniker SE, Biggin MD, Stamatoyannopoulos JA (2011) Dynamic reprogramming of chromatin accessibility during *Drosophila* embryo development. *Genome Biol* 12(5):R43. doi:10.1186/gb-2011-12-5-r43
79. Zhang Y, Vastenhouw NL, Feng J, Fu K, Wang C, Ge Y, Pauli A, van Hummelen P, Schier AF, Liu XS (2014) Canonical nucleosome organization at promoters forms during genome activation. *Genome Res* 24(2):260–266. doi:10.1101/gr.157750.113
80. Juven-Gershon T, Kadonaga JT (2010) Regulation of gene expression via the core promoter and the basal transcriptional machinery. *Dev Biol* 339(2):225–229. doi:10.1016/j.ydbio.2009.08.009
81. Zabidi MA, Arnold CD, Schernhuber K, Pagani M, Rath M, Frank O, Stark A (2015) Enhancer-core-promoter specificity separates developmental and housekeeping gene regulation. *Nature* 518(7540):556–559. doi:10.1038/nature13994
82. Collart C, Allen GE, Bradshaw CR, Smith JC, Zegerman P (2013) Titration of four replication factors is essential for the *Xenopus laevis* midblastula transition. *Science* 341(6148): 893–896. doi:10.1126/science.1241530
83. Lehner CF, O'Farrell PH (1990) *Drosophila* cdc2 homologs: a functional homolog is coexpressed with a cognate variant. *EMBO J* 9(11):3573–3581
84. Edgar BA, O'Farrell PH (1990) The three postblastoderm cell cycles of *Drosophila* embryogenesis are regulated in G2 by string. *Cell* 62(3):469–480
85. Edgar BA, Sprenger F, Duronio RJ, Leopold P, O'Farrell PH (1994) Distinct molecular mechanisms regulate cell cycle timing at successive stages of *Drosophila* embryogenesis. *Genes Dev* 8(4):440–452
86. Yuan K, O'Farrell PH (2015) Cyclin B3 is a mitotic cyclin that promotes the metaphase-anaphase transition. *Curr Biol* 25(6):811–816. doi:10.1016/j.cub.2015.01.053

87. Sigrist S, Ried G, Lehner CF (1995) Dmcdc2 kinase is required for both meiotic divisions during *Drosophila* spermatogenesis and is activated by the Twine/cdc25 phosphatase. *Mech Dev* 53(2):247–260
88. Glotzer M, Murray AW, Kirschner MW (1991) Cyclin is degraded by the ubiquitin pathway. *Nature* 349(6305):132–138. doi:10.1038/349132a0
89. Yuan K, Farrell JA, O'Farrell PH (2012) Different cyclin types collaborate to reverse the S-phase checkpoint and permit prompt mitosis. *J Cell Biol* 198(6):973–980. doi:10.1083/jcb.201205007
90. Ji JY, Squirrell JM, Schubiger G (2004) Both cyclin B levels and DNA-replication checkpoint control the early embryonic mitoses in *Drosophila*. *Development* 131(2):401–411. doi:10.1242/dev.00944
91. Jin Z, Homola EM, Goldbach P, Choi Y, Brill JA, Campbell SD (2005) *Drosophila* Myt1 is a Cdk1 inhibitory kinase that regulates multiple aspects of cell cycle behavior during gametogenesis. *Development* 132(18):4075–4085. doi:10.1242/dev.01965
92. Price D, Rabinovitch S, O'Farrell PH, Campbell SD (2000) *Drosophila* weel has an essential role in the nuclear divisions of early embryogenesis. *Genetics* 155(1):159–166
93. Stumpff J, Duncan T, Homola E, Campbell SD, Su TT (2004) *Drosophila* Wee1 kinase regulates Cdk1 and mitotic entry during embryogenesis. *Curr Biol* 14(23):2143–2148. doi:10.1016/j.cub.2004.11.050
94. Edgar BA, O'Farrell PH (1989) Genetic control of cell division patterns in the *Drosophila* embryo. *Cell* 57(1):177–187
95. Blythe SA, Wieschaus EF (2015) Coordinating cell cycle remodeling with transcriptional activation at the *Drosophila* MBT. *Curr Top Dev Biol* 113:113–148. doi:10.1016/bs.ctdb.2015.06.002
96. Ayeni JO, Varadarajan R, Mukherjee O, Stuart DT, Sprenger F, Srayko M, Campbell SD (2014) Dual phosphorylation of cdk1 coordinates cell proliferation with key developmental processes in *Drosophila*. *Genetics* 196(1):197–210. doi:10.1534/genetics.113.156281
97. Farrell JA, O'Farrell PH (2013) Mechanism and regulation of Cdc25/Twine protein destruction in embryonic cell-cycle remodeling. *Curr Biol* 23(2):118–126. doi:10.1016/j.cub.2012.11.036
98. Di Talia S, She R, Blythe SA, Lu X, Zhang QF, Wieschaus EF (2013) Posttranslational control of Cdc25 degradation terminates *Drosophila*'s early cell-cycle program. *Curr Biol* 23(2):127–132. doi:10.1016/j.cub.2012.11.029
99. Edgar BA, Lehner CF (1996) Developmental control of cell cycle regulators: a fly's perspective. *Science* 274(5293):1646–1652
100. Alphey L, Jimenez J, White-Cooper H, Dawson I, Nurse P, Glover DM (1992) twine, a cdc25 homolog that functions in the male and female germline of *Drosophila*. *Cell* 69(6):977–988
101. Grosshans J, Wieschaus E (2000) A genetic link between morphogenesis and cell division during formation of the ventral furrow in *Drosophila*. *Cell* 101(5):523–531
102. Mata J, Curado S, Ephrussi A, Rorth P (2000) Tribbles coordinates mitosis and morphogenesis in *Drosophila* by regulating string/CDC25 proteolysis. *Cell* 101(5):511–522
103. Rorth P, Szabo K, Texido G (2000) The level of C/EBP protein is critical for cell migration during *Drosophila* oogenesis and is tightly controlled by regulated degradation. *Mol Cell* 6(1):23–30
104. Frazer C, Young PG (2012) Phosphorylation mediated regulation of Cdc25 activity, localization and stability. In: Huang C (ed) *Protein phosphorylation in human health, Biochemistry, genetics and molecular biology*. InTech, Rijeka, Croatia, pp 395–436. doi:10.5772/48315
105. Murphy JM, Nakatani Y, Jamieson SA, Dai W, Lucet IS, Mace PD (2015) Molecular mechanism of CCAAT-enhancer binding protein recruitment by the TRIB1 pseudokinase. *Structure* 23(11):2111–2121. doi:10.1016/j.str.2015.08.017
106. Edgar BA, Datar SA (1996) Zygotic degradation of two maternal Cdc25 mRNAs terminates *Drosophila*'s early cell cycle program. *Genes Dev* 10(15):1966–1977
107. Chen F, Archambault V, Kar A, Lio P, D'Avino PP, Sinka R, Lilley K, Laue ED, Deak P, Capalbo L, Glover DM (2007) Multiple protein phosphatases are required for mitosis in *Drosophila*. *Curr Biol* 17(4):293–303. doi:10.1016/j.cub.2007.01.068
108. Blumenthal AB, Kriegstein HJ, Hogness DS (1974) The units of DNA replication in *Drosophila melanogaster* chromosomes. *Cold Spring Harb Symp Quant Biol* 38:205–223
109. Sibon OC, Stevenson VA, Theurkauf WE (1997) DNA-replication checkpoint control at the *Drosophila* midblastula transition. *Nature* 388(6637):93–97. doi:10.1038/40439
110. Fogarty P, Campbell SD, Abu-Shumays R, Phalle BS, Yu KR, Uy GL, Goldberg ML, Sullivan W (1997) The *Drosophila* grapes gene is related to checkpoint gene chk1/

- rad27 and is required for late syncytial division fidelity. *Curr Biol* 7(6):418–426
111. Sibon OC, Laurencon A, Hawley R, Theurkauf WE (1999) The *Drosophila* ATM homologue Mei-41 has an essential checkpoint function at the midblastula transition. *Curr Biol* 9(6):302–312
 112. Shimuta K, Nakajo N, Uto K, Hayano Y, Okazaki K, Sagata N (2002) Chk1 is activated transiently and targets Cdc25A for degradation at the *Xenopus* midblastula transition. *EMBO J* 21(14):3694–3703. doi:10.1093/emboj/cdf357
 113. Gawlinski P, Nikolay R, Goursot C, Lawo S, Chaurasia B, Herz HM, Kussler-Schneider Y, Ruppert T, Mayer M, Grosshans J (2007) The *Drosophila* mitotic inhibitor Fruhstart specifically binds to the hydrophobic patch of cyclins. *EMBO Rep* 8(5):490–496. doi:10.1038/sj.embor.7400948
 114. Grosshans J, Muller HA, Wieschaus E (2003) Control of cleavage cycles in *Drosophila* embryos by fruhstart. *Dev Cell* 5(2):285–294
 115. Lu X, Li JM, Elemento O, Tavazoie S, Wieschaus EF (2009) Coupling of zygotic transcription to mitotic control at the *Drosophila* mid-blastula transition. *Development* 136(12):2101–2110. doi:10.1242/dev.034421
 116. Campbell SD, Sprenger F, Edgar BA, O'Farrell PH (1995) *Drosophila* Wee1 kinase rescues fission yeast from mitotic catastrophe and phosphorylates *Drosophila* Cdc2 in vitro. *Mol Biol Cell* 6(10):1333–1347
 117. Bettencourt-Dias M, Giet R, Sinka R, Mazumdar A, Lock WG, Balloux F, Zafiropoulos PJ, Yamaguchi S, Winter S, Carthew RW, Cooper M, Jones D, Frenz L, Glover DM (2004) Genome-wide survey of protein kinases required for cell cycle progression. *Nature* 432(7020):980–987. doi:10.1038/nature03160
 118. Fasulo B, Koyama C, Yu KR, Homola EM, Hsieh TS, Campbell SD, Sullivan W (2012) Chk1 and Wee1 kinases coordinate DNA replication, chromosome condensation, and anaphase entry. *Mol Biol Cell* 23(6):1047–1057. doi:10.1091/mbc.E11-10-0832
 119. Royou A, McCusker D, Kellogg DR, Sullivan W (2008) Grapes(Chk1) prevents nuclear CDK1 activation by delaying cyclin B nuclear accumulation. *J Cell Biol* 183(1):63–75. doi:10.1083/jcb.200801153
 120. Kang Q, Srividhya J, Ipe J, Pomerening JR (2014) Evidence toward a dual phosphatase mechanism that restricts Aurora A (Thr-295) phosphorylation during the early embryonic cell cycle. *J Biol Chem* 289(25):17480–17496. doi:10.1074/jbc.M113.527622
 121. Zou L, Elledge SJ (2003) Sensing DNA damage through ATRIP recognition of RPA-ssDNA complexes. *Science* 300(5625):1542–1548. doi:10.1126/science.1083430
 122. Papoulas O, Monzo KF, Cantin GT, Ruse C, Yates JR 3rd, Ryu YH, Sisson JC (2010) dFMRP and Caprin, translational regulators of synaptic plasticity, control the cell cycle at the *Drosophila* mid-blastula transition. *Development* 137(24):4201–4209. doi:10.1242/dev.055046
 123. Monzo K, Papoulas O, Cantin GT, Wang Y, Yates JR 3rd, Sisson JC (2006) Fragile X mental retardation protein controls trailer hitch expression and cleavage furrow formation in *Drosophila* embryos. *Proc Natl Acad Sci U S A* 103(48):18160–18165. doi:10.1073/pnas.0606508103
 124. Nien CY, Liang HL, Butcher S, Sun Y, Fu S, Gocha T, Kirov N, Manak JR, Rushlow C (2011) Temporal coordination of gene networks by Zelda in the early *Drosophila* embryo. *PLoS Genet* 7(10):e1002339. doi:10.1371/journal.pgen.1002339
 125. Benoit B, He CH, Zhang F, Votruba SM, Tadros W, Westwood JT, Smibert CA, Lipshitz HD, Theurkauf WE (2009) An essential role for the RNA-binding protein Smaug during the *Drosophila* maternal-to-zygotic transition. *Development* 136(6):923–932. doi:10.1242/dev.031815
 126. Jevtic P, Levy DL (2015) Nuclear size scaling during *Xenopus* early development contributes to midblastula transition timing. *Curr Biol* 25(1):45–52. doi:10.1016/j.cub.2014.10.051
 127. Vastag L, Jorgensen P, Peshkin L, Wei R, Rabinowitz JD, Kirschner MW (2011) Remodeling of the metabolome during early frog development. *PLoS One* 6(2):e16881. doi:10.1371/journal.pone.0016881
 128. Edgar BA, Kiehle CP, Schubiger G (1986) Cell cycle control by the nucleo-cytoplasmic ratio in early *Drosophila* development. *Cell* 44(2):365–372

Zygotic Genome Activation

Methods and Protocols

Lee, K. (Ed.)

2017, XI, 272 p. 51 illus., 41 illus. in color., Hardcover

ISBN: 978-1-4939-6986-9

A product of Humana Press