

Chapter 2

Late Triassic Global Plate Tectonics

Jan Golonka, Ashton Embry, and Michal Krobicki

Abstract The Late Triassic was the time of the Early Cimmerian and Indosinian orogenies that closed the Paleotethys Ocean, which occurred earlier in the Alpine-Carpathian-Mediterranean area, later in the Eastern Europe-Central Asia and latest in the South-East Asia. The Indochina Southeastern Asian and Qiangtang plates were sutured to South China. The new, large Chinese-SE Asian plate, including North and South China, Mongolia and eastern Cimmerian plates, was consolidated by the end Triassic, leaving open a large embayment of Panthalassa, known as Mongol-Okhotsk Ocean, between Mongolia and Laurasia. The Uralian Orogeny, which sutured Siberia and Europe continued during Late Triassic times and was recorded in Novaya Zemlya. The onset of Pangaea break-up constitutes the main Late Triassic extensional event. Continental rifts originating then were filled with clastic deposits comprising mainly red beds. The pulling force of the north-dipping subduction along the northern margin of Neotethys caused drifting of a new set of plates from the passive Gondwana margin, dividing the Neotethys Ocean. Carbonate sedimentation dominated platforms on the Neotethys and Paleotethys margins as well as the Cimmerian microplates. Synorogenic turbidites and postorogenic molasses were associated with the Indosinian orogeny. The late stages of the Uralian orogeny in Timan-Pechora, Novaya Zemlya and eastern Barents regions filled the foreland basin with fine-grained, molasse sediments. Siliciclastics were common in the Siberia and Arctic regions. The widespread, large magnitude, base-level changes of the Late Triassic are interpreted as an expression of relatively rapid and substantial changes in the horizontal and vertical stress fields that affected the Pangaea

J. Golonka (✉)

Faculty of Geology, Geophysics and Environmental Protection, AGH University of Science and Technology, Al. Mickiewicza 30, 30-059 Kraków, Poland
e-mail: jgonlonka@agh.edu.pl

A. Embry

Geological Survey of Canada, Calgary, AB, Canada
e-mail: ashton.embry@canada.ca

M. Krobicki

Faculty of Geology, Geophysics and Environmental Protection, AGH University of Science and Technology, Al. Mickiewicza 30, 30-059 Kraków, Poland

Carpathian Branch, Polish Geological Institute – National Research Institute, Kraków, Poland
e-mail: krobicki@geol.agh.edu.pl

supercontinent. Such stress changes may be due to abrupt changes in the speed and/or direction of plate movements, which episodically affected Pangaea.

Keywords Paleogeography • Plate tectonics • Paleoenvironment • Paleolithofacies • Paleoclimate • Sea level changes

2.1 Introduction

The Triassic maps used here (Figs. 2.1, 2.2, 2.3, 2.4, 2.5, 2.6, 2.7 and 2.8) were derived from a series of global and regional Phanerozoic paleogeographic and plate tectonic maps which depicted present day coastlines, plate boundaries (sutures), selected transform faults, spreading centers, rifts, normal and thrust faults as well as paleoenvironment and lithofacies (Golonka 2000, 2002, 2007a, b, 2011; Golonka et al. 2003a, 2006a,b). Also included is a corrected and improved version of the Triassic maps previously presented (Golonka 2007a, b). The base maps, (past position of present day coastlines and plate boundaries) were generated by PLATES, PALEOMAP and GPLATES computer software (see Sect. 1.2). The definitions of mapped time slices were presented by Golonka and Kiessling (2002), however, recently the simple stratigraphic “Late Triassic” slice was used (Golonka 2007a, b). The name “Triassic” was derived from the German Trias defined by von Alberti (1834), referring to the division of the period into three stages: the Buntsandstein, Muschelkalk, and Keuper (see Köppen and Carter 2000; Feist-Burkhardt et al. 2008; Scheck-Wenderoth et al. 2008; McKie and Williams 2009 and references therein). This sequence is valid for Central Europe (Germany, Poland), but causes many problems when applied to other regions. The global Late Triassic (Ogg et al. 2016) is now divided into the Carnian, Norian and Rhaetian ages (Fig. 2.9). For the environment and facies assembly we used two units, applying the methods used for the Phanerozoic reefs map (Kiessling and Flügel 1999) and also presented by Golonka (2007a, b). The base maps (Figs. 2.1 and 2.2) depict the configuration of land masses, rifts, spreading centers and subduction and the beginning (Fig. 2.1) and end (Fig. 2.2) of the Late Triassic. The paleoenvironments and lithofacies (Figs. 2.3, 2.4, 2.5, 2.6, 2.7 and 2.8) represent the whole of the Late Triassic Epoch. They are posted on the 224 Ma base maps.

2.2 Methods

The Phanerozoic maps were constructed using a plate tectonic model that describes the relative motions between approximately 300 plates and terranes (Golonka 2000). This model was originally constructed using PLATES and PALEOMAP software, later the GPLATES program was used (see the detailed reconstruction




Fig. 2.1 Global plate tectonic map of Late Triassic at 224 Ma ago. Molweide Projection. (1) oceanic spreading center and transform faults, (2) subduction zone, (3) thrust fault, (4) normal fault, (5) transform fault




Fig. 2.2 Global plate tectonic map of Late Triassic at 200 Ma ago. Molweide Projection. (1) oceanic spreading center and transform faults, (2) subduction zone, (3) thrust fault, (4) normal fault, (5) transform fault

methodology in Golonka et al. 2003b. The rotation file was presented in Golonka (2007a), and is shown in the appendix of that paper.

We modified this model using new paleomagnetic data, especially in the Tethys and Arctic areas (Kravchinsky et al. 2002; Hounslow and Nawrocki 2008; Kovalenko 2010; Metelkin et al. 2011, 2012; Uno et al. 2011; Domeier et al. 2012; Choulet et al. 2013; Vernikovskiy et al. 2013; Wang et al. 2013; Song et al. 2015; Huang and Opdyke 2016; Li et al. 2016a, b; Zhou et al. 2016). We left the position of the major




Fig. 2.3 Plate tectonic, paleoenvironment and lithofacies map of the western Tethys, future Central Atlantic and adjacent areas during Late Triassic time. Molweide Projection. Modified from Golonka (2007b)

continent unchanged due to the absence of important new data. For example, according to Metelkin et al. (2011) there is an absence of authentic data for the Middle and Late Triassic from Siberia.

The facies were reconstructed using established sedimentological concepts for reefs and other sedimentary environments (Kiessling and Flügel 1999; Kiessling et al. 2003) and also presented by Golonka et al. (2006b) and Golonka (2007a, b). The calculated paleolatitudes and paleolongitudes were used to generate computer maps in Microstation design (.dgn format) converted later into Corel Draw (.cdr format). Facies and paleoenvironment information were posted after reviewing database files, regional paleogeographic maps and relevant papers. Information from several general and regional paleogeographic papers were filtered and utilized (Vinogradov 1968; Ziegler 1982, 1988; Hongzen 1985; Ronov et al. 1989; Cook 1990; Zonenshain et al. 1990; Doré 1991; Dercourt et al. 1993, 2000; Golonka et al. 1994, 2006a; Metcalfe 1994, 2011, 2013a, b; Veevers 1994, 2006, 2013; Nikishin et al. 1996; Sengör and Natalin 1996; Puchkov 1997; Kiessling and Flügel 1999; Golonka 2000, 2002, 2007a, b, 2011; Golonka and Ford 2000; Ford and Golonka 2003; Scotese 2004; Miller et al. 2006; Robertson 2007; Feist-Burkhardt et al. 2008; Heydari 2008; Maurer et al. 2008; Miall and Blakey 2008; Miall et al. 2008; Pčelina and Korčinskaja 2008; Scheck-Wenderoth et al. 2008; Schmid et al. 2008; Peng et al. 2009; McKie and Williams 2009; Glørstad-Clark et al. 2010; Metelkin et al.




Fig. 2.4 Explanations to Figs. 2.3, 2.4, 2.5, 2.7. Qualifiers: *B* bauxites/laterites, *C* coals, *E* evaporites, *F* flysch, *Fe* Iron, *G* glauconite, *M* marls, *O* oolites, *P* phosphates, *R* red beds, *Si* silica, *T* tillites, *V* volcanics

2011, 2012; Schettino and Turco 2011; Sibuet et al. 2012; Li and Huang 2013; Luo et al. 2014; Ershova et al. 2015a, b; Pease et al. 2015; Lane and Stephenson 2016; Müller et al. 2016; Toro et al. 2016; Cai et al. 2017; Centeno-García 2017).

2.3 Convergent Tectonics

The Late Paleozoic supercontinent Pangaea included North America, South America, Africa, Australia, Europe and Siberia and was surrounded by the Panthalassa Ocean (Figs. 2.1 and 2.2). The collision between Siberia and Europe




Fig. 2.5 Plate tectonic, paleoenvironment and lithofacies map of eastern Tethys and adjacent areas during Late Triassic time. Molweide Projection




Fig. 2.6 Plate tectonic, paleoenvironment and lithofacies map of the Paleotethys, Chinese plates and adjacent areas during Late Triassic time. Molweide Projection. Modified from Golonka (2007b)




Fig. 2.7 Plate tectonic, paleoenvironment and lithofacies map of the western Gondwana and adjacent areas during Late Triassic time. Molweide Projection. Modified from Golonka (2007b)

formed the Ural Mountains during the Uralian Orogeny (Zonenshain et al. 1990; Nikishin et al. 1996; Puchkov 1997). The last episode of this orogeny occurred at the end of the Triassic in Novaya Zemlya (Toro et al. 2016; Zhang et al. 2017a). Deformation also affected the Taimyr Peninsula (Torsvik and Anderson 2002; Golonka 2007a, b). According to Zhang et al. (2017b) the Taimyr Permo-Triassic magmatic rocks were locally folded and faulted as a result of Late Triassic to Early Jurassic dextral transpression. According to Vernikovsky (1995) and Vernikovsky et al. (2003) the formation of the Taimyr structures is connected with the collision of the Kara microcontinent with Siberia. The uplift of the adjacent areas of Europe and Siberia was related to these orogenic events (Figs. 2.2 and 2.8). The Crockerland uplifted area of the Alaska-Chukotka micro-plate supplied sediments to the adjacent Sverdrup Basin in North America and was linked with Siberia at this time (Fig. 2.8; Anfinson et al. 2016). The subduction zones, known as the Late Paleozoic Pangaeian Rim of Fire, were still active during the Triassic (Golonka and Ford 2000; Golonka 2002, 2004, 2007a, b; Matthews et al. 2016). This Rim of Fire was especially active along the western coast of Pangaea (Figs. 2.1 and 2.2). Active volcanism, terrane accretion, and back-arc basin development accompanied the subduction zones (Golonka 2007a, b). The subduction accompanied by magmatism was active in Central and North America (Goodge 1989, 1990; Dorsey and LaMaskin 2007; Centeno-García et al. 2008; Arvizu and Iriondo 2015) as well as




Fig. 2.8 Plate tectonic, paleoenvironment and lithofacies map of the Arctic during Late Triassic time. Stereographic polar. Projection Modified from Golonka (2011)

in South America (Bustamante and Juliani 2011; del Rey et al. 2016). The movement of terranes within Panthalassa was related to the activity of this subduction (Figs. 2.1 and 2.2). According to Dorsey and LaMaskin (2007), the collision of terranes in North America happened during Late Triassic times in the Blue Mountains of Oregon. The position of these terranes is a subject of controversy, however (e.g. Engebretson et al. 1985; Panuska 1985; Debiche et al. 1987; Sengör and Natalin 1996; Keppie and Dostal 2001; Belasky et al. 2002; Trop et al. 2002; Piercey et al. 2006; Golonka 2007a, b; Colpron and Nelson 2011; Roniewicz 2013; Matthews et al. 2016). The relationship between Panthalassa terranes and Cimmerian plates was previously postulated and mapped (Golonka 2007a, b). The Panthalassa terranes bearing reef complexes were also mentioned by Flügel




Fig. 2.9 Late Triassic stratigraphy of various basins contains sequence boundaries of basal Carnian, mid-Carnian, basal Norian, mid-Norian, basal Rhaetian, and latest Rhaetian age. The features of these boundaries indicate they represent relatively short-lived, tectonic episodes. Each tectonic episode was characterized by a rapid base level fall followed by rapid rise which punctuated the relatively slow, long term subsidence of the basins

(2002). According to Peyberness et al. (2016, see also Stanley and Onoue 2015) the Western Panthalassa reefs from Japan corresponds with those of the Tethys Ocean during the Late Triassic. The Late Triassic was the time of the collisions now known as the Early Cimmerian and Indosinian orogenies. Blocks of the Cimmerian provenance and Eurasia (Sengör 1984; Sengör et al. 1984; Sengör and Natalin 1996) were involved in these collisions with the southern margin of Eurasia (Golonka 2000, 2002, 2007a, b; Golonka et al. 2003a, 2006a,b; Robertson 2007; Richards 2015). This series of collisions closed the Paleotethys Ocean. The closure happened earlier in the Alpine-Carpathian-Mediterranean area, later in the Eastern Europe-Central Asia and latest in the South-East Asia (Figs. 2.1 and 2.2). Microplates now included in the Alpine-Carpathian systems formed the marginal part of Europe. Subduction developed south of this zone. Late Triassic collisional events occurred also in the Moesia-Rhodopes areas (Tari et al. 1997; Golonka 2004, 2007a, b; Okay and Nikishin 2015; Petrík et al. 2016). The Alborz and the

South Caspian Microcontinent collided with the Scythian platform in Eastern Europe, and the other Iranian plates, including the large Lut block, collided with the Turan platform (Zonenshain et al. 1990; Kazmin 1991; Nikishin et al. 1996, 1998a; Golonka 2004, 2007a, b; Heydari 2008; Wilmsen et al. 2009; Masoodi et al. 2013; Okay and Nikishin 2015; Zanchi et al. 2009, 2016). Compressional deformations were recorded in the Caucasus, and Kopet Dagh areas, accompanied by the general uplift of the Fore-Caucasus, Caucasus and Middle Asia regions (Golonka 2004). According to Okay and Nikishin (2015), the accretion of an oceanic plateau was recorded by Late Triassic eclogites in the Pontides. Collisional events were also noted in Afghanistan and Pamir areas (Sengör 1984; Zonenshain et al. 1990; Golonka 2004, 2007a, b; Montenat 2009; Robinson 2015).

The Paleotethys between Qiangtang and Eurasia was closed during Late Triassic times (Figs. 2.1 and 2.2; Metcalfe 2013a; Zhai et al. 2013; Zhu et al. 2013; Luo et al. 2014; Song et al. 2015; Wu et al. 2016). The eastern Cimmerian plates were involved in the Indosinian orogeny. This name was derived from Indochina, the region where the orogeny was noted over one hundred years ago (Deprat 1913, 1914; Fromaget 1927, 1934, 1941, 1952). A major unconformity was observed in Northwest Vietnam. The deformed Lower – lowermost Upper Triassic (up to Carnian) marine metamorphosed rocks arranged into nappes and thrusts are covered by Upper Triassic continental red conglomerates (“terrains rouges”, see Deprat 1913, 1914, also Golonka et al. 2006b). According to Lepvrier et al. (2004 see also Maluski et al. 2001, 2005; Lepvrier and Maluski 2008 and references therein), the main metamorphic event occurred during the Early Triassic, 250–240 Ma. The Late Triassic unconformity and 225–205 Ma postorogenic plutonism was noted by Faure et al. (2014). Hung (2010) describes magmatism in northeastern Vietnam related to Triassic Indosinian orogeny. According to Faure et al. (2014) the Jinshajiang and Ailaoshan belts in China and their geodynamic evolution, with Vietnam orogeny marking the same Indosinian Orogeny. It was related to the closure of Paleotethys Ocean along Raub-Bentong, Sra Kaeo and Nan-Uttaradit suture between Sibumasu and Indochina and Ailaoshan suture between Sibumasu and South China (Metcalfe 1994, 1996, 2000, 2011, 2013a, b; Golonka et al. 2006b and references therein).

One of the best examples of the Late Triassic orogenic event occurs in the Thailand/Myanmar trans-border zone. The Triassic-Jurassic succession in the Mae Sot area (northern Thailand), belongs to the Shan-Thai terrane. This block is subdivided into several zones from the west to east, including the Mae Sariang zone, where the Mae Sot area is located. This zone contains rocks of Triassic cherts (radiolarites), carbonates and flysch (turbiditic) facies, which indicate both pelagic condition and synorogenic deposits. From a paleogeographic point of view, the Shan-Thai block was a remnant of Paleotethys Ocean (Meesook and Sha 2010), which occupied a wide realm between Cimmerian Continent and Eurasian plate during Late Paleozoic-Early Mesozoic times. On the other hand, the Late Triassic Indosinian orogenic event has been associated with the docking and amalgamation of the Indoburma, Shan-Thai (Sibumasu) and Indochina terranes, which recently

constituted the main part of Southeast Asia. Therefore, entire Jurassic units of these regions are represented by post-orogenic continental-shelf deposits, which are underlain discontinuously by older rocks. The oldest Jurassic bed, or the youngest Triassic bed, is the so-called “base-conglomerate”, in local nomenclature, and is characterized by limestone and chert pebbles-bearing conglomerate, which is significant for the understanding of the tectonic evolution of the Shanthai terrane (Ishida et al. 2006; Meesook and Sha 2010). The underlying cherts are dated biostratigraphically (based on radiolarians) as Middle-Late Triassic. Limestone and chert pebbles from the “base-conglomerate” are dated as Early-Late Triassic by conodonts and as Middle-Late Triassic by radiolarians, respectively. These microfossils from pebbles constrain the age of the Indosinian (ShanThai = Mae Sariang) orogeny. Additionally, the youngest clasts, both limestones and siliceous rocks, indicate a strictly pelagic character of sedimentation up to Late Triassic time (see Ishida et al. 2006). A full open ocean condition must have existed at least before the end of the Triassic. The “base-conglomerate” is characterized by poorly-sorted, chaotically organized, pebble/fragment-bearing sedimentary breccia with no evidence of bivalve borings on their surfaces. The multicolored clasts are subrounded and subangular, and occur within reddish silt matrix. Chert clasts are red, green and grey and carbonate pebbles are represented both by micritic, pelagic limestones and the entire spectrum of packstones and grainstones, including extremely shallow-water bioclastic limestones (with bivalve fragments, crinoids, fragments of corals, etc.) with ooids and coated grains. The “base-conglomerate” is overlain by limestones and marls with mudstone intercalations of the Khun Huai Formation of the Hua Fai Group, dated by ammonites and bivalves as Early Toarcian. These facts indicate, by superposition, that the “base-conglomerate” is the latest Triassic or earliest Jurassic in age, according to the latest Triassic age of the chert and limestone pebbles within it. Sedimentological features indicate, on the other hand, a very rapid sedimentation event during its origin, such as erosion of steep, submarine “cliffs” that formed proximal aprons of debris flows. Additionally, the composition of this conglomerate, which has both deep-marine clasts and shallow-water ones, without any evidence of their long-distance transport, suggests erosion of different type of source material, which most probably originally took place in a different part of the primary Paleotethys Ocean. Then, they were removed, folded (forming nappes?) and overthrust to another location where they were destroyed and eroded, and produced marine molasse-type deposits unconformably overlying Indosinian deformed rocks. In fact, these data indicate both time and space reorganization of this orogenic system, which took place possibly during latest Triassic to earliest Jurassic time. The examination of the main orogenic events in the Southeast Asia regions indicates diachronous, multi-stages movements of the Indosinian orogeny. These include Early Triassic and Carnian/Norian orogenic pulses in Vietnam (Lepvrier et al. 2004), late Middle Triassic–early Late Triassic activity, the so-called second Indosinian event (Hahn 1984; Lepvrier and Maluski 2008, see also Cai et al. 2017) and close to the Triassic/Jurassic boundary in Thailand, as the Asian plate docked first on the East and later on the West (in modern coordinates).

Additionally, the Late Triassic volcanogenic-sedimentary event in Myanmar correlates presumably with synorogenic processes, which are represented by the Late Triassic flysch deposits with basaltic pillow lavas of the Shweminbon Group (Upper Triassic – Lower Jurassic turbidites), formerly part of Loi-an Group, the Bawgyo Group (Upper Triassic) and their equivalents, and with Upper Triassic turbidites represented by the Thanbaya/Pane Chaung Group/formations (Bannert et al. 2011; Win Swe 2012; Cai et al. 2017).

The collision between the South Chinese plate and the North Chinese block began during the Late Permian and continued during the Triassic (Yin and Nie 1996; Golonka et al. 2006b; Golonka 2007a, b). The Qinling orogenic belt records this collision. According to Dong et al. (2011) the Shangdan zone between the North and South Qinling belts is the suture separating the convergence and collision between North South Chinese plates. The post-suturing plutons were emplaced along the suture zone and on the adjacent plates (Bao et al. 2015; Liang et al. 2015; Lu et al. 2016). Consolidation of North China and Mongolia occurred mainly earlier but continued during the Triassic between North China and Mongolia. The newly formed, larger plate contains volcanics and collisional granites (Fig. 2.6; Chen et al. 2000; Wu et al. 2002; Shi et al. 2016). This consolidation left open a large embayment between Mongolia and Laurasia, the so-called Mongol-Okhotsk Ocean (Zonenshain et al. 1990; Golonka 2000, 2007a; Zeng et al. 2014). Active subduction existed along the margin of this ocean (Figs. 2.6 and 2.8), dipping cratonwards towards East Siberia (Zonenshain et al. 1990; Golonka 2007a, b), and granitic intrusions occurred along the Siberian margin (Zonenshain et al. 1990; Donskaya et al. 2013, 2016). The new, large Chinese-Southeast Asian plate including North and South China, Mongolia and eastern Cimmerian plates was consolidated at the Triassic-Jurassic Boundary (Fig. 2.2).

2.4 Extensional Tectonics

The onset of Pangaeian break-up constitutes the main Late Triassic extensional tectonic event (Golonka 2007a, b). The rift basins originated between North America and Africa. The extensional rifting was accompanied by strike-slip faulting and block rotation (Ford and Golonka 2003; Laville et al. 2004; Golonka 2007a, b). Incipient continental rifting occurred also between northern Europe and North America (Fig. 2.2), reactivating the Late Paleozoic fracture system (Ziegler 1982; Doré 1991; Nikishin et al. 2002; Golonka 2011), and activating the North Sea rifts. The Central European Permian rift system known as the Polish/Danish Aulacogene was still active during Late Triassic times. The Upper Permian (Zechstein) salt went into salt tectonic phase with incipient salt diapirism and extrusion (Kutek 2001; Krzywiec 2012). Continental extension also began in isolated areas in South America during the Late Triassic (Macdonald et al. 2003; Ford and Golonka 2003; Golonka 2007a, b). Additionally, rift basins developed behind the subduction zone along the western Pangaeian margin (Goode 1989, 1990; Golonka and Ford 2000; Golonka 2007a, b;

Centeno-García et al. 2008; Dickinson 2009; Bustamante and Juliani 2011; Giambiagi et al. 2011; Baby et al. 2013; Helbig et al. 2013; Spikings et al. 2016; Centeno-García 2017).

The Pangaea rift systems extended also to the Barents shelf, Arctic, and Siberia (Golonka 2011; Golonka et al. 2003a, 2006b). Rifting in Siberia was associated with the subduction zone at the Mongol-Okhotsk Ocean margin (Figs. 2.1, 2.2 and 2.8). Late Triassic sea-floor spreading in Siberia constituted an extension of the Anyui Ocean, which existed between the Alaska-Chukotka and Verkhoyansk terranes (Fig. 2.8; Zonenshain et al. 1990; Sengör and Natalin 1996; Golonka et al. 2003a; Golonka 2011). The opening of the Amerasia Basin appears to have begun near the Norian/Rhaetian boundary resulting in the rotational separation of the Alaska-Chukotka terrane from northern Laurasia (Embry and Anfinson 2014).

The volcanics (flows and intrusions) of the Central Atlantic Magmatic Province (CAMP), were emplaced at the end of Triassic and beginning of the Early Jurassic (e.g., Olsen 1997; Withjack et al. 1998; Marzoli et al. 1999, 2004, 2011; Knight et al. 2004; Golonka 2007a; Cirilli et al. 2009). CAMP constitutes one of the largest known Phanerozoic flood basalt provinces. It triggered climate changes and the end-Triassic extinction event (Wignall 2001; Lucas and Tanner 2008; Preto et al. 2010; Bond and Wignall 2014; Müller et al. 2016). The Late Triassic northward drift of the Cimmerian continent was accompanied by active seafloor spreading within the Neotethys Ocean. The spreading was driven by trench-pulling forces related to the north-dipping subduction, as well as the ridge-pushing forces related to mantle upwelling, expressed by hot spot activity (Golonka and Bocharova 2000; Golonka 2004, 2007a, b). Rifting and the opening of oceanic type basins could have occurred in the Alpine, Carpathian, Balkans and future Mediterranean area (Figs. 2.1, 2.2 and 2.3; Golonka et al. 2006a). The opening of the incipient Pindos–Maliac Ocean was related to the establishment of the Pelagonian, Sakariya and Kirsehir blocks as separate microplates within the Western (Robertson et al. 1991, 1996; Ferriere et al. 2016). The proto-Transylvanian and Vardar oceans originated within Carpathian-Balkan. The Tisa block was perhaps fully separated from the European margin by the Meliata-Halstatt Ocean. The positions of the Vardar, Meliata-Halstatt, Transylvanian, Pindos, Maliac oceans and their embayments within the Western Tethys remain quite speculative and are subjects of the debate (e.g., Kozur and Krahel 1987; Săndulescu 1988; Kozur 1991; Channell and Kozur 1997; Mock et al. 1998; Ivan 2002; Golonka 2004; Haas and Pero 2004; Golonka et al. 2006a; Dallmeyer et al. 2008; Schmid et al. 2008; Hoeck et al. 2009; Gawlick and Missoni 2015; Meinhold and Kostopoulos 2013). The Eurasian platform east of the Carpathians and Meliata Ocean was dissected by rifts that extended from the Dobrogea, through the proto-Black Sea area and along the margins of Scythianturan platform and probably were connected with Polish/Danish Aulacogene (Fig. 2.3; Zonenshain et al. 1990; Kazmin 1990, 1991; Nikishin et al. 1998a, b; Golonka 2004). The Tauric basin, which belonged to this rift system, was located between Pontides and the Dobrogea-Crimea segment of the Scythian platform (Golonka et al. 2006a). The North Dobrogea part of the rift zone separated Moesia and Eastern European platform (Muttoni et al. 2000; Golonka 2004; Golonka et al. 2006a). Several blocks

were located between the rifted zone and the Neotethys (Golonka 2004; Golonka et al. 2006a; Okay and Nikishin 2015). This rifted zone can be interpreted as a back-arc basin resulting from the northward subduction of the Neotethys Ocean (Figs. 2.1, 2.2 and 2.3). The deep-water basin was located between Apulia, the Taurus platform and the African continent (Fig. 2.1; Catalano et al. 1991; Kozur 1991; Marsella et al. 1993; Golonka 2004, Golonka et al. 2006b). It was connected eastwards with an oceanic-type basin recorded by the Mamonia ophiolites complex in Cyprus (Robertson and Woodcock 1979; Morris 1996; Robertson 1998). The rifts cutting Apulia were connected with the western part of Neotethys.

The whole Paleotethys was closed in the western part of the Tethyan realm in the Early Jurassic (Fig. 2.2). The pulling force of the north-dipping subduction along the northern margin of Neotethys caused the drift of a new set of plates from the passive Gondwanian margin. These plates divided the Neotethys Ocean into northern and southern branches (Golonka 2004). Metcalfe (2013a) distinguished Cenotethys as the southern branch. The Lhasa block was the most prominent plate which drifted away from Gondwana (Sengör 1984; Dercourt et al. 1993; Metcalfe 1994, Metcalfe 2013a, b; Sengör and Natalin 1996; Yin and Nie 1996; Golonka 2004; Cai et al. 2016; Li et al. 2016a, b; Lu et al. 2016; Meng et al. 2016; Zhou et al. 2016). According to Li et al. (Li et al. 2016a, b), the Kirsehir, Sakarya (Robertson et al. 1991, 1996), and perhaps the Lesser Caucasus-Sanandaj-Sirjan, Biston-Avoraman plates drifted in the central Neotethys area (Adamia 1991; Robertson et al. 1991, 1996, 2004; Arfania and Shahriari 2009; Mehdi-pour Ghazi and Moazzen 2015; Nouri et al. 2016). According to Metcalfe (2013a, b) South West Borneo and East Java-West Sulawesi were separated from Northwest Australia in the Late Triassic in the easternmost Tethys area. The consolidation of the Chinese and south-eastern Asian blocks was followed by extensional tectonics caused by the pulling force if the new Neotethys subduction. Consequently, rift basins developed in China and adjacent areas (Golonka et al. 2006b; Luo et al. 2014). This process was enhanced by the Panthalassa (Paleo-Pacific) plate sliding beneath the Eurasian plate (Luo et al. 2014; Li et al. 2016a, b).

2.5 Sedimentation and Paleolithofacies

Continental rifts, which originated during Triassic times, were filled with clastic deposits, particularly abundant red beds consisting of fluvial deposits and accompanied by evaporites (Ziegler 1988; Withjack et al. 1998; Golonka and Ford 2000; Kutek 2001; Feist-Burkhardt et al. 2008). Mixed siliciclastics, carbonates and evaporates were deposited in Central Europe (Figs. 2.3 and 2.9) as to the upper part (Keuper) of the Central European tripartite facies sequence that gave the Triassic its name (Köppen and Carter 2000). The Keuper Formation encompasses the Carnian, Norian and Rhaetian stages (Fig. 2.9). The accumulation of sediments in this area reached up to 4000 m due to significant subsidence (Köppen and Carter 2000; Kutek 2001; Golonka 2007a, b; Feist-Burkhardt et al. 2008). Meanwhile, continental red

beds were deposited in the eastern United States while a marine shelf existed on the western North America margin. Continental rifting occurred between northern Europe and Greenland (Fig. 2.8). The Pangaea rift systems extended from the Newark and Central Europe basins through the North Atlantic, to the Barents shelf and Arctic Alaska (Figs. 2.3, 2.8 and 2.9). These rifts were filled primarily with red continental clastics reflecting arid climate (Ronov et al. 1989; Olsen 1997; Golonka et al. 2003a, 2006a, b; Golonka 2007a, b; Dickinson 2004, 2008, 2009; Miall et al. 2008; Miall and Blakey 2008). Carbonate sedimentation dominated in the Alps and Carpathians (Golonka 2004, 2007a, b; Feist-Burkhardt et al. 2008). This sedimentation was associated with existence of platforms on the Neotethys and Paleotethys margins as well as on Cimmerian microplates. Shallow-water limestones and dolomites with algal/coral-dominated reefs were deposited on these platforms (Golonka 2007a, b). They were accompanied by fine grained clastics (Figs. 2.3 and 2.9). Many of the western Tethyan reefs were located on these platforms. Triassic carbonate platforms and reefs were formed not only in the Tethys, but also in the western and eastern parts of the Panthalassa (Paleo-Pacific) Ocean (Golonka 2007a, b). A large carbonate platform that spread from Apulia to the Taurus zone provides an example (Dercourt et al. 1993, 2000; Golonka 2004, 2007a, b; Feist-Burkhardt et al. 2008) in that it contains significant numbers of reefs (Kiessling and Flügel 1999; Flügel 2002) and was connected with the Alpine-Inner Carpathian carbonate platforms, which also contained abundant reefs (Kiessling and Flügel 1999; Flügel 2002). Dolomitization of the platform limestones was common and dolomites are widespread in Southern Europe and Central Asia. The Dolomia Principale (Fig. 2.9) represents a classic example of the Tethyan dolomites. The Dolomites range in the Italian Southern Alps took their name from the mineral and rock dolomite, which in turn were named after the French geologist Dieudonné Sylvain Guy Tancrède de Gratet de Dolomieu by de Saussure (1792). Dolomites were also widespread on the southern margin of Eurasia in the Caspian area and in Central Asia (Figs. 2.5 and 2.7). Continental and marginal marine sediments with evaporites and volcanics were also deposited in this part of Eurasia (Zonenshain et al. 1990; Dercourt et al. 1993, 2000; Nikishin et al. 1996, 1998a, b; Brunet et al. 2002; Zharkov and Chumakov 2001; Golonka 2004, 2007a, b). The neritic and lagoonal sediments of so-called Carpathian Keuper were deposited in the Northern Carpathians during the latest Triassic, marking the uplift of the Inner Carpathian plate (Kotański 1961; Golonka 2004; Feist-Burkhardt et al. 2008; Rychliński 2008). The Neotethyan margins of Greater India, Arabia and Australia (Figs. 2.5 and 2.7) were occupied by mixed carbonate-clastic facies (Cook 1990; Alsharhan and Magara 1994; Golonka and Ford 2000; Golonka 2007a, b). Basins containing Triassic continental red bed deposits were located in Gondwana (Fig. 2.7), in South America, Africa, Antarctica, Madagascar and India (Golonka 2007a, b). The deposition of synorogenic flysch sequences in South-East Asia was linked to the Indosinian orogenic collisional events (Hahn 1984; Golonka et al. 2006b; Lepvrier and Maluski 2008; Cai et al. 2017). They were accompanied by pelagic cherts, cherty limestones and fine-grained clastics as well as by volcanoclastics and pillow lavas (Ishida et al. 2006; Bannert et al. 2011; Win Swe 2012; Cai et al. 2017) and followed by post-orogenic

molasses. Post-orogenic Upper Triassic continental red conglomerates are known as “terrains rouges” of Deprat (1913, 1914) in Vietnam (Golonka et al. 2006a, b). These red-bed postorogenic facies that follow synorogenic turbidites and pelagic cherts are also known from the Malaysian Peninsula (Oliver and Prave 2013; Ridd 2013). Flysch sequences and volcanoclastic deposits occur on the Lhasa plate, South Tibet (Liu et al. 2012). Shallow marine, carbonate and clastic sedimentation dominated on the Qiantang plate (Zhu et al. 2013; Wu et al. 2016). Various stratigraphic sequences representing different paleogeographic facies existed in South China where paralic clastics, shallow marine clastics, shelf carbonate platform facies and deep water turbidites can be distinguished. Siliciclastic sedimentation prevailed in North China, including shallow marine clastics, marginal marine deposits such as deltas, as well as turbidites accompanied by volcanoclastics (Fig. 2.6; Hongzen 1985; Tong and Yin 2002; Golonka 2007a, b; Cao et al. 2010; Luo et al. 2014; Li et al. 2014). The sediments consisting mainly of fine-grained molasse-type filled the foreland basins following the Uralian orogeny in Timan-Pechora, Novaya Zemlya and eastern Barents regions. Siliciclastics were common in the Siberia and Arctic regions (Figs. 2.8 and 2.9; Embry 1988, 1993, 1997; Nikishin et al. 1996; Golonka and Ford 2000; Golonka et al. 2003a; Golonka 2007a, b, 2011; Toro et al. 2016); the Sverdrup Basin of Arctic Canada was a main depocenter with the Late Triassic succession of fluvial to marine slope deposits being over 2500 m thick (Embry 1997). Triassic, restricted-marine shelf basins contain black shales that have source rock potential (Leith et al. 1993; Golonka et al. 2003a; Golonka 2007b). Upper Triassic source rocks, important for hydrocarbon exploration in the North Atlantic, were identified in the Jameson Land Basin, East Greenland (Andrews et al. 2014).

2.6 Global Base-Level Changes

In this section, we briefly review postulated base-level changes that have been interpreted to affected numerous basins throughout Pangaea during the Late Triassic. We first look at small scale changes with frequencies of less than 500,000 years. Then we address large scale base-level changes with frequencies of greater than 2 million years.

Tanner (2010) comprehensively reviewed the literature for high frequency, small scale cycles for the entire Triassic. Such cycles have been recorded in various Late Triassic successions with the best documentation being from the rift valley deposits of the Newark Group of the northeastern USA (Olsen and Kent 1996) and the carbonate platforms of the Italian Alps (Cozzi et al. 2005; Schwarzhacher 2006). The Late Triassic small scale cycles of the Italian Alps are characterized by the presence of exposure surfaces and paleosols. This leaves little doubt as to such cycles being generated by base-level changes caused by either eustasy or tectonics. It must be noted that any high frequency, small scale cycles which do not include exposure surfaces may well have an auto-cyclic explanation for their generation.

Given the occurrence of such high-frequency base level changes in the Late Triassic of the Italian Alps and the apparent coincidence of the calculated frequencies with those of the Milankovitch spectrum (Cozzi et al. 2005; Schwarzscher 2006), it seems reasonable to assume that small-scale, global sea-level changes driven by climate changes characterize the Late Triassic. However, as cautioned by Tanner (2010), this interpretation cannot be considered as unassailable for two main reasons. Given the greenhouse climate of the Late Triassic (Preto et al. 2010) and the consequent unlikelihood that substantial amounts of water could have been stored as ice during cold periods, there are no obvious mechanisms for climate changes to drive eustatic sea level change of the magnitude seemingly recorded by the cycles. The other problem is the general lack of precise radiometric age dates to constrain the interpreted cycle periods.

More studies are needed for Late Triassic, very shallow water carbonate and siliciclastic strata in a number of basins of Pangaea to see if they are characterized by high-frequency cycles that are capped by exposure surfaces. If Milankovitch climate change cycles were operating during the Late Triassic, then such cycles should be present in the successions of most, if not, every basin. In summary, it is quite possible that Milankovitch climate cycles were operating during the Late Triassic but further studies are needed to confirm or deny such a phenomenon.

Large scale, base-level changes are recorded in most Late Triassic successions and are expressed as large-magnitude, sequence boundaries. Such boundaries are characterized by an extensive unconformable portion on the basin margins and are the product of base-level changes that can exceed 100 m. Both eustatic and tectonic explanations have been offered for the generation of these boundaries.

Late Triassic, large-magnitude, sequence boundaries, which have been recorded in different basins throughout Pangaea, have been biostratigraphically dated as near the base Carnian, mid-Carnian, near the base Norian, mid-Norian, near the base Rhaetian and latest Rhaetian. Initially, these boundaries were interpreted to be the product of eustasy, including a significant sea level fall followed by sea level rise (Haq et al. 1987, 1988; Embry 1988; De Zanche et al. 1993; Gianolla and Jacquin 1998). Given a climate change/continental glaciation explanation was not possible, the authors appealed to changes in the volume of the world ocean (tectono-eustasy) as the main driver of such large scale eustatic changes.

Embry (1989, 1997) reversed his earlier interpretation and postulated that the large-magnitude sequence boundaries, which punctuated the entire Mesozoic succession of the Sverdrup Basin of Arctic Canada, were of tectonic origin. This interpretation was based on various characteristics of such boundaries which strongly favor a tectonic origin. Such characteristics included:

- A widespread, often angular, unconformity on the basin margins and positive elements
- A major change in depositional regime
- A notable change in tectonic regime and subsidence pattern
- A change in provenance for siliciclastic sediments
- A widespread transgression with significant deepening directly following the boundary.

Furthermore Embry (1997) demonstrated that the five, Late Triassic large-magnitude sequence boundaries present in the Sverdrup Basin are also present in basins in western Canada, southwestern USA, Barents Sea, Germany, Italian Alps, western Siberia, and northern Himalayas (Fig. 2.9). Notably, the unconformities in all these areas exhibit characteristics which favor a tectonic origin.

To explain the occurrence of simultaneous tectonic episodes in multiple and widely separated basins of Pangaea, Embry (1997) invoked the tectonic model of Cloetingh et al. (1985). The widespread, large magnitude base-level changes of the Late Triassic were interpreted to be an expression of relatively rapid and substantial changes in the horizontal and vertical stress fields that affected the Pangaea super-continent (Fig. 2.9). Such stress changes would be possibly due to somewhat abrupt changes in the speed and/or direction of the plate movements that episodically affected Pangaea. Notably, it is possible that secondary tectono-eustatic effects were associated with such plate tectonic reorganizations (Embry 1997).

2.7 Climate Change and Episodic Tectonism

The climate of the Triassic has been reviewed by Preto et al. (2010) and they have interpreted that it “was characterized by a non-zonal pattern, dictated by a strong global monsoon system with effects that are most evident in the Tethys realm”. For the Late Triassic, Preto et al. (2010) postulated that the monsoonal climate had its maximum expression and that there were three climatic zones which did not have a clear latitudinal distribution. These three zones included a dry climate for the western margin of Tethys and the central part of Pangaea, a wet and dry climate for the coasts of eastern Laurasia and Gondwana and the western coasts of Pangaea, and a wet climate in the high latitudes.

Although, in general, there was not much variability in climate throughout the Late Triassic, significant climate changes seem to be associated with the five tectonic episodes discussed in the last section. The most well-known of these is the “Carnian Pluvial Episode” (Ruffell et al. 2015) which corresponds with the mid-Carnian tectonic episode. This event was marked by warmer, more humid conditions in various parts of Pangaea and a notable increase of siliciclastic supply to numerous basins (Ruffell et al. 2015). Climate changes seem to have occurred associated with the other four tectonic episodes as shown by the marked changes in spore/pollen ratios associated with these boundaries (Hochuli and Vigran 2010). Climate change associated with the latest Rhaetian has been documented by various workers as summarized by Preto et al. (2010). The CAMP flood basalts, which were associated with the extensional phase of the latest Rhaetian tectonic episode, produced enormous amounts of CO₂, triggered global warming, and increased ocean acidification. These factors caused the end of Triassic extinction event (Wignall 2001; Lucas and Tanner 2008; Preto et al. 2010; Bond and Wignall 2014; Müller et al. 2016).

2.8 Concluding Summary

Herein, we present a new set of global and regional paleogeographic maps for the Late Triassic (Carnian-Rhaetian) time interval. The global maps depict the plate tectonic configuration, present day coastlines, subduction zones, selected transform faults, spreading centers and rifts during the beginning (224 Ma) and end (200 Ma) of Late Triassic. The regional maps illustrate the Late Triassic paleoenvironment and paleolithofacies distribution for most important regions. The stratigraphic chart shows Late Triassic stratigraphy of various basins and sequence boundaries of basal Carnian, mid-Carnian, basal Norian, mid-Norian, basal Rhaetian, and latest Rhaetian age.

The Late Triassic was a time of collisional events, now known as Early Cimmerian and Indosinian orogenies. This series of collisions closed the Paleotethys Ocean. The closure happened earlier in the Alpine-Carpathian-Mediterranean area, later in the Eastern Europe-Central Asia and latest in the South-East Asia. The Indochina, Southeastern Asian and Qiangtang plates were sutured to South China. The new, large Chinese-Southeast Asian plate, including the North and South China, Mongolia and eastern Cimmerian plates, was consolidated at the Triassic-Jurassic Boundary. This consolidation left open a large embayment of Panthalassa, between Mongolia and Laurasia, known as Mongol-Okhotsk Ocean. The Uralian Orogeny, which sutured Siberia and Europe continued during Late Triassic times and was recorded in Novaya Zemlya.

The onset of the break-up of Pangaea constitutes the main Late Triassic extensional tectonics event. Continental rifts, which originated during this event, were filled with clastic deposits. Abundant red beds, accompanied by fluvial deposits and evaporites, were deposited in classic sedimentary systems. The pulling force of the north-dipping subduction along the northern margin of Neotethys caused the drift of a new set of plates from the passive Gondwana margin. These plates divided the Neotethys Ocean. Carbonate sedimentation was associated with existence of platforms on the Neotethys and Paleotethys margins as well as on Cimmerian microplates. Synorogenic turbidites and postorogenic molasses were associated with the Indosinian orogeny. The late stages of the Uralian orogeny in Timan-Pechora, Novaya Zemlya and eastern Barents regions included the filling of the foreland basin with fine-grained, molasse sediments. Siliciclastics were common in the Siberia and Arctic regions.

The widespread, large magnitude, base level changes of the Late Triassic are interpreted to be an expression of relatively rapid and substantial changes in the horizontal and vertical stress fields that affected the Pangaea supercontinent. Such stress changes would be possibly due to somewhat abrupt changes in the speed and/or direction of plate movements, which episodically affected Pangaea. The Late Triassic climate changes seem to be associated with the main tectonic episodes. The most well-known of these is the “Carnian Pluvial Episode” which corresponds with the mid-Carnian tectonic episode. The Central Atlantic Magmatic Province flood basalts, which were associated with the extensional phase of the latest Rhaetian

tectonic episode, produced enormous amounts of CO₂, triggering global warming, increasing ocean acidification, and causing the latest Triassic extinction event.

Acknowledgements The research was partially supported by Faculty of Geology, Geophysics and Environmental Protection, AGH University of Science and Technology, Kraków, Poland Statutory grant no. 11.11.140.005. Embry thanks the Geological Survey of Canada for supporting his research on the Triassic of the Canadian Arctic for the past 40 years.

References

- Adamia SA (1991) The Caucasus oil and gas province, Occasional Publications. ESRI, New Series No. 7(I-II), Part I: 53–74
- Alsharhan AS, Magara K (1994) The Jurassic of the Arabian Gulf Basin: facies, depositional setting and hydrocarbon habitat. In: Embry AF, Beauchamp B, Glass DJ (eds) *Pangea; global environments and resources*. Can Soc Petr Geol Mem 17:397–412
- Andrews SD, Kelly SRA, Braham W, Kaye M (2014) Climatic and Eustatic controls on the development of a Late Triassic source rock in the Jameson Land Basin, East Greenland. *J Geol Soc* 171(5):609–619
- Anfinson OA, Embry AF, Stockli DF (2016) Geochronologic Constraints on the Permian-Triassic Northern Source Region of the Sverdrup Basin, Canadian Arctic Islands. *Tectonophysics* 691:206–219
- Arfania R, Shahriari S (2009) Role of southeastern Sanandaj-Sirjan Zone in the tectonic evolution of Zagros Orogenic Belt, Iran. *Island Arc* 18(4):555–576
- Arvizu HE, Iriondo A (2015) Temporal control and geology of the Permo-Triassic magmatism in Sierra Los Tanques, NW Sonora, Mexico: Evidence for the beginning of cordilleran arc magmatism in SW of Laurentia [Control temporal y geología del magmatismo Permo-Triásico en Sierra Los Tanques, NW Sonora, México: Evidencia del inicio del arco magmático cordillerano en el SW de Laurentia]. *Boletín de la Sociedad Geológica Mexicana* 67(3):545–586
- Baby P, Rivadenreira M, Barragàn R, Christophoul F (2013) Thick-skinned tectonics in the Oriente foreland basin of Ecuador. *Geol Soc London Spec Publ* 377(1):59–76
- Bannert D, Sang Lyen A, Htay T (2011) The geology of the Indoburman ranges in Myanmar. *Geol Jb B* 101:5–101
- Belasky P, Stevens CH, Hanger RA (2002) Early Permian location of western North American terranes based on brachiopod, fusulinid, and coral biogeography. *Palaeogeogr Palaeoclimatol Palaeoecol* 179:245–266
- Bao Z, Wang CY, Zeng L, Sun W, Yao J (2015) Slab break-off model for the Triassic syn-collisional granites in the Qinling orogenic belt, Central China: Zircon U-Pb age and Hf isotope constraints. *Internat Geol Rev* 57(4):492–507
- Bond DPG, Wignall PB (2014) Large igneous provinces and mass extinctions: an update. In: Keller G, Kerr AC (eds) *Volcanism, impacts, and mass extinctions: causes and effects*. *Geol Soc Am Spec Pap* 505: 29–55
- Brunet M-F, Korotaev MV, Ershov AV, Nikishin AM (2002) The South Caspian basin: a review of the evolution with the approach of the subsidence modelling. In: Brunet M-F, Cloetingh S (eds) *Integrated Peri-Tethyan Basins Studies*. *Sed Geol*, vol 156, pp 119–148
- Bustamante A, Juliani C (2011) Unraveling an antique subduction process from metamorphic basement around Medellín city, Central Cordillera of Colombian Andes. *J S Am Earth Sci* 32(3):210–221
- Cai F, Ding L, Laskowski AK, Kapp P, Wang H, Xu Q, Zhang L (2016) Late Triassic paleogeographic reconstruction along the Neo-Tethyan Ocean margins, southern Tibet. *Earth Planet Sci Lett* 435:105–114

- Cai F, Ding L, Yao W, Laskowski AK, Xu Q, Zhang J, Sein K (2017) Provenance and tectonic evolution of Lower Paleozoic–Upper Mesozoic strata from Sibumasu terrane, Myanmar. *Gond Res* 41:325–336
- Cao J, Wang X, Wei D, Sun P, Hu W, Jia D, Zha Y (2010) Complex petroleum migration and accumulation in central region of southern Junggar basin, Northwest China. *J Earth Sci* 21(1):83–93
- Catalano R, Di Stefano P, Kozur H (1991) Permian circum-pacific deep-water faunas from the western Tethys (Sicily, Italy): new evidence for the position of the Permian Tethys. *Palaeogeogr Palaeoclimatol Palaeoecol* 87:75–108
- Centeno-García E (2017) Mesozoic tectono-magmatic evolution of Mexico: An overview. *Ore Geol Rev* 81:1035–1052
- Centeno-García E, Guerrero-Suastegui M, Talavera-Mendoza O (2008) The Guerrero composite terrane of western Mexico: collision and subsequent rifting in a supra-subduction zone. In: Drau A, Clift PD, Scholl DW (eds) *Formation and applications of the sedimentary record in arc collision zones*. *Geol Soc Am Spec Pap* 436: 279–308
- Channell JET, Kozur H (1997) How many oceans? Meliata, Vardar, and Pindos oceans in Mesozoic Alpine paleogeography. *Geology* 25:183–186
- Chen B, Jahn B-M, Wilde S, Xu B (2000) Two contrasting paleozoic magmatic belts in northern Inner Mongolia, China: petrogenesis and tectonic implications. *Tectonophysics* 328:157–182
- Choulet F, Chen Y, Cogné J-P, Rabillard A, Wang B, Lin W, Faure M, Cluzel D (2013) First Triassic palaeomagnetic constraints from Junggar (NW China) and their implications for the Mesozoic tectonics in Central Asia. *J Asian Earth Sci* 78:371–394
- Cirilli S, Marzoli A, Tanner L, Bertrand H, Buratti N, Jourdan F, Bellieni G, Kontak D, Renne PR (2009) Latest Triassic onset of the Central Atlantic Magmatic Province (CAMP) volcanism in the Fundy Basin (Nova Scotia): New stratigraphic constraints. *Earth Planet Sci Lett* 286(3–4):514–525
- Cloetingh S, McQueen H, Lambeck K (1985) On a plate tectonic mechanism for regional sea level variations. *Earth Planet Sci Lett* 75:157–166
- Colpron M, Nelson JL (2011) A Palaeozoic NW passage and the Timanian, Caledonian and Uralian connections of some exotic terranes in the North American Cordillera. In: Spencer AM, Embry AF, Gautier DL, Stoupakova AV, Sørensen K (eds) *Arctic petroleum geology*. *Geol Soc London Mem* 35: 463–484
- Cook PI (1990) *Australia—evolution of a continent*. Australian Government Publishing Services, Canberra
- Cozzi A, Hinnov LA, Hardie LA (2005) Orbitally forced Lofer cycles in the Dachstein Limestone of the Julian Alps (northeastern Italy). *Geol* 33:789–792
- Dallmeyer RD, Neubauer F, Fritz H (2008) The Meliata suture in the Carpathians: regional significance and implications for the evolution of high-pressure wedges within collisional orogens. *Geol Soc London Spec Publ* 298:101–115
- De Zanche V, Gianolla P, Mietto P, Siorpaes C, Vail PR (1993) Triassic sequence stratigraphy in the Dolomites (Italy). *Memorie di Scienze Geologiche* 45:1–27
- Debiche MG, Cox A, Engebretson D (1987) The motion of allochthonous terranes across the North Pacific basin. *Geol Soc Am Spec Pap* 207:1–49
- del Rey A, Deckart K, Arriagada C, Martínez F (2016) Resolving the paradigm of the late Paleozoic–Triassic Chilean magmatism: Isotopic approach. *Gond Res* 37:172–181
- Deprat J (1913) Les chantages de la région de la Rivière Noire sur les feuilles de Thanh-ba et Van-Yên. *Mémoires de Service Géologique d'Indochine* 2:47–65
- Deprat J (1914) Etude des plissements et des zones d'écrasement de la moyenne et de la basse Rivière Noire. *Mémoires de Service Géologique d'Indochine* 3:1–59
- Dercourt J, Ricou LE, Vrielynck B (eds) (1993) *Atlas Tethys Paleoenvironmental maps*. Gauthier-Villars, Paris
- Dercourt J, Gaetani M, Vrielynck B, Barrier E, Biju-Duval B, Brunet M-F, Cadet JP, Crasquin S, Sandulescu M (eds) (2000) *Atlas Peri-Tethys Paleogeographical maps*. CCGM/CGMW, Paris
- Dickinson WR (2004) Evolution of the North American Cordillera. *Annu Rev Earth Planet Sci* 32:13–45

- Dickinson WR (2008) Accretionary Mesozoic-Cenozoic expansion of the Cordilleran continental margin in California and adjacent Oregon. *Geosphere* 4(2):329–353
- Dickinson WR (2009) Anatomy and global context of the North American Cordillera. In: Kay SM, Ramos VA, Dickinson WR (eds) *Backbone of the Americas: shallow subduction, plateau uplift, and ridge and terrane collision*. *Geol Soc Am Mem* 204: 1–29
- Domeier M, Van Der Voo R, Torsvik TH (2012) Paleomagnetism and Pangaea: The road to reconciliation. *Tectonophysics* 514–517:14–43
- Dong Y, Zhang G, Neubauer F, Liu X, Genser J, Hauzenberger C (2011) Tectonic evolution of the Qinling orogen, China: Review and synthesis. *J Asian Earth Sci* 41(3):213–237
- Donskaya TV, Gladkochub DP, Mazukabzov AM, Ivanov AV (2013) Late Paleozoic - Mesozoic subduction-related magmatism at the southern margin of the Siberian continent and the 150 million-year history of the Mongol-Okhotsk Ocean. *J Asian Earth Sci* 62:79–97
- Donskaya TV, Gladkochub DP, Mazukabzov AM, Wang T, Guo L, Rodionov NV, Demanterova EI (2016) Mesozoic granitoids in the structure of the Bezmyannyi metamorphic-core complex (western Transbaikalia). *Russ Geol Geophys* 57(11):1591–1605
- Doré AG (1991) The structural foundation and evolution of Mesozoic seaways between Europe and the Arctic. *Palaeogeogr Palaeoclimatol Palaeoecol* 87:441–492
- Dorsey RJ, LaMaskin TA (2007) Stratigraphic record of Triassic-Jurassic collisional tectonics in the Blue Mountains province, northern Oregon. *Am J Sci* 307(10):1167–1193
- Embry AF (1988) Triassic sea-level changes: evidence from the Canadian Arctic Archipelago. In: Wilgus CK, Posamentier H, Ross CA, Kendall, CGStC (eds) *Sea-level changes: an integrated approach*. *Soc Econ Paleontol Mineralog Spec Publ* 42: 249–259
- Embry A (1989) A tectonic origin for third-order depositional sequences in extensional basins implications for basin modelling. In: Cross T (ed) *Quantitative dynamic stratigraphy*. Prentice Hall, Upper Saddle River, NJ, p. 491–502
- Embry AF, (1993) Crockerland—the northern source area for the Sverdrup Basin, Canadian Arctic Archipelago. In: Vorren T, Bergsager E, Dahl-Stamnes O, Holter E, Johansen B, Lie E, Lund T (eds) *Arctic geology and petroleum potential: Norweg Petrol Soc Spec Publ* 2: 205–216
- Embry AF (1997) Global sequence boundaries of the Triassic and their recognition in the Western Canada Sedimentary Basin. *Bull Can Petrol Geol* 45:415–433
- Embry AF, Anfinson OA (2014) The initiation of the rift phase of the Amerasia Basin (Arctic Ocean). *CSPG Geoconvention 2014, Abstr. Calgary, AB, Canada* p 1–4
- Engelbreton DC, Cox A, Gordon RG (1985) Relative motions between oceanic and continental plate in the Pacific basin. *Geol Soc Am Spec Pap* 206:1–59
- Ershova VB, Prokoviev AV, Khudoley AK, Sobolev NN, Petrov EO (2015a) U/Pb dating of detrital zircons from late Palaeozoic deposits of Belkovsky Island (New Siberian Islands): Critical testing of Arctic tectonic models. *Internat Geol Rev* 57(2):199–210
- Ershova VB, Prokoviev AV, Khudoley AK, Sobolev NN, Petrov EO (2015b) Detrital zircon ages and provenance of the Upper Paleozoic successions of Kotel'ny Island (New Siberian Islands archipelago). *Lithosph* 7(1):40–45
- Faure M, Lepvrier C, Nguyen VV, TV V, Lin W, Chen Z (2014) The South China block-Indochina collision: Where, when, and how? *J Asian Earth Sci* 79:260–274
- Feist-Burkhardt S, Götz AE, Szulc J, Borkhataria R, Geluk M, Haas J, Hornung J, Jordan P, Kempf O, Michalik J, Nawrocki J, Reinhardt L, Ricken W, Röhling H-G, Rüffer T, Török Á, Zühlke R (2008). Triassic. In: McCann T (ed) *The geology of Central Europe. Volume 2: Mesozoic and Cenozoic*, *Geol Soc, London*, p 749–821
- Ferriere J, Baumgartner PO, Chancier F (2016) The Maliac Ocean: the origin of the Tethyan Hellenic ophiolites. *Internat J Earth Sci* 105(7):1941–1963
- Flügel E (2002) Triassic Reef Patterns. In: Kiessling W, Flügel E, Golonka J (eds) *Phanerozoic reef patterns*. *Soc Econ Paleontol Mineral Spec Publ* 72: 391–464
- Ford D, Golonka J (2003) Phanerozoic paleogeography, paleoenvironment and lithofacies maps of the circum-Atlantic margins. In: Golonka J (ed) *Thematic set on paleogeographic reconstruction and hydrocarbon basins: Atlantic, Caribbean, South America, Middle East, Russian Far East, Arctic*. *Mar Petrol Geol* 20: 249–285

- Fromaget J (1927) Etudes géologiques sur le Nord de l'Indochine centrale. *Bull Serv Géol Indoch* 16(2):1–368
- Fromaget J (1934) Observations et reflexions sur la geologic stratigraphique et structurale de l'Indochine. *Bull Soc Geol Fr* 5(4):101–164
- Fromaget J (1941) L'Indochine Française, sa structure geologique, ses roches, ses mines et leurs relations possibles avec la tectonique. *Bull Serv Géol Indoch* 26(2):1–140
- Fromaget J (1952) Etudes géologiques sur le nord-Ouest du Tonkin et le Nord du Haut-Laos. 2ème et 3ème parties. *Bull Serv Géol Indoch* 29(6):1–198
- Gawlick H-J, Missoni S (2015) Middle Triassic radiolarite pebbles in the Middle Jurassic Hallstatt Mélange of the Eastern Alps: implications for Triassic–Jurassic geodynamic and paleogeographic reconstructions of the western Tethyan realm. *Facies* 61(3):1–19
- Giambiagi L, Mescua J, Bechis F, Martínez A, Folguera A (2011) Pre-Andean deformation of the Precordillera southern sector, southern Central Andes. *Geosphere* 7(1):219–239
- Gianolla P, Jacquin T (1998) Triassic sequence stratigraphic framework of Western European basins. In: de Graciansky PC, Hardenbol J, Jacquin T, Vail PR (eds) *Mesozoic and Cenozoic sequence stratigraphy of European basins*. Soc Econ Paleontol Mineral Spec Publ 60: 643–650
- Glørstad-Clark E, Faleide JJ, Lundschieen BA, Nystuen JP (2010) Triassic seismic sequence stratigraphy and paleogeography of the western Barents Sea area. *Marine and Petroleum Geol* 27(7):1448–1475
- Golonka J (2000) Cambrian-Neogene Plate Tectonic Maps. Wydawnictwa Uniwersytetu Jagiellońskiego, Kraków
- Golonka J (2002) Plate-tectonic maps of the Phanerozoic. In: Kiessling W, Flügel E, Golonka J (eds) *Phanerozoic reef patterns*. Soc Econ Paleontol Mineral Spec Publ 72: 21–75
- Golonka J (2004) Plate tectonic evolution of the southern margin of Eurasia in the Mesozoic and Cenozoic. *Tectonophysics* 381:235–273
- Golonka J (2007a) Late Triassic and Early Jurassic paleogeography of the world. *Palaeogeogr Palaeoclimatol Palaeoecol* 244:297–307
- Golonka J (2007b) Phanerozoic Palaeoenvironment and Paleolithofacies Maps. Mesozoic. Mapy paleośrodowiska i paleolitofacje fanerozoiku. Mezozoik. Kwartalnik AGH. *Geologia* 33(2):211–264
- Golonka J (2011) Phanerozoic palaeoenvironment and palaeolithofacies maps of the Arctic region. In: Spencer, AM, Embry AF, Gautier DL, Stoupakova A, Sørensen K (eds) *Arctic petroleum geology*. *Geol Soc London Mem* 35: 79–129
- Golonka J, Bocharova NY (2000) Hot spots activity and the break-up of Pangaea. *Palaeogeogr Palaeoclimatol Palaeoecol* 161:49–69
- Golonka J, Ford DW (2000) Pangaeian (Late Carboniferous–Middle Jurassic) palaeoenvironment and lithofacies. *Palaeogeogr Palaeoclimatol Palaeoecol* 161:1–34
- Golonka J, Kiessling W (2002) Phanerozoic time scale and definition of time slices. In: Kiessling W, Flügel E, Golonka J (eds) *Phanerozoic reef patterns*. Soc Econ Paleontol Mineral Spec Publ 72:11–20
- Golonka J, Ross MI, Scotese C R (1994) Phanerozoic paleogeographic and paleoclimatic modeling maps. In: Embry AF, Beauchamp B, Glass DJ (eds) *Pangaea; global environments and resources*. *Can Soc Petrol Geol Mem* 17: 1–47
- Golonka J, Bocharova NY, Ford D, Edrich ME, Bednarczyk J, Wildharber J (2003a) Paleogeographic reconstructions and basins development of the Arctic. In: Golonka J (ed) *Thematic set on paleogeographic reconstruction and hydrocarbon basins: Atlantic, Caribbean, South America, Middle East, Russian Far East, Arctic*. *Mar Petrol Geol* 20: 211–248
- Golonka J, Krobicki M, Oszczypko N, Ślaczka A, Słomka T (2003b) Geodynamic evolution and palaeogeography of the Polish Carpathians and adjacent areas during Neo-Cimmerian and preceding events (latest Triassic–earliest Cretaceous). In: McCann T, Saintot A (eds) *Tracing tectonic deformation using the sedimentary record*. *Geol Soc London Spec Publ* 208: 138–158
- Golonka J, Krobicki M, Paják J, Van Giang N, Zuchiewicz W (2006a) Global plate tectonics and paleogeography of Southeast Asia. *Arkadia*, Kraków

- Golonka J, Gahagan L, Krobicki M, Marko F, Oszczytko N, Ślaczka A (2006b) Plate Tectonic Evolution and Paleogeography of the Circum-Carpathian Region. In: Golonka J. & Picha F (eds) *The Carpathians and their foreland: Geology and hydrocarbon resources*: Am Assoc Petrol Geol Mem 84: 11–46
- Goodge JW (1989) Polyphase metamorphic evolution of a Late Triassic subduction complex, Klamath Mountains, northern California. *Am J Sci* 289(7):874–943
- Goodge JW (1990) Tectonic evolution of a coherent Late Triassic subduction complex, Stuart Fork terrane, Klamath Mountains, northern California. *Geol Soc Am Bull* 102(1):86–101
- Haas J, Pero S (2004) Mesozoic evolution of the Tisza Mega-unit. *Internat J Earth Sciences* 93:297–313
- Hahn L (1984) The Indosinian Orogeny in Thailand and adjacent areas. *Mémoires de la Société géologique de France* 147:71–82
- Haq BU, Hardenbol J, Vail PR (1987) Chronology of fluctuating sea levels since the Triassic. *Science* 235:1156–1167
- Haq BU, Hardenbol J, Vail PR (1988) Mesozoic and Cenozoic chronostratigraphy and cycles of sea-level change In: Wilgus, CK, Posamentier H, Ross CA, Kendall, CGStC (eds) *Sea-level Changes: an Integrated Approach*. Soc Econ Paleontol Min Spec Publ 42: 71–108
- Helbig M, Keppie JD, Murphy JB, Solari LA (2013) Exotic rifted passive margin of a back-arc basin off western Pangaea: Geochemical evidence from the Early Mesozoic Ayú Complex, southern Mexico. *Internat Geol Rev* 55(7):863–881
- Heydari E (2008) Tectonics versus eustatic control on supersequences of the Zagros Mountains of Iran. *Tectonophysics* 451(1–4):56–70
- Hochuli PA, Vigran JO (2010) Climate variations in the Boreal Triassic—inferred from palynological records from the Barents Sea. *Palaeogeogr Palaeoclimatol Palaeoecol* 290:20–42
- Hoecck V, Ionescu C, Balintoni I, Koller F (2009) The Eastern Carpathians “ophiolites” (Romania): Remnants of a Triassic ocean. *Lithos* 108(1–4):151–171
- Hongzen W (1985) *Atlas of Paleogeography of China*. Cartographic Publishing House, Beijing
- Hounslow MW, Nawrocki J (2008) Palaeomagnetism and magnetostratigraphy of the Permian and Triassic of Spitsbergen: a review of progress and challenges. *Pol Res* 27(3):502–522
- Huang K, Opdyke ND (2016) Paleomagnetism of the Upper Triassic rocks from south of the Ailaoshan Suture and the timing of the amalgamation between the South China and the Indochina Blocks. *J Asian Earth Sci* 119:118–127
- Hung KT (2010) Overview of magmatism in northwestern Vietnam. *Ann Soc Geol Pol* 80(2):185–226
- Ishida K, Nanba A, Hirsch F, Kozai T, Meesook A (2006) New micropaleontological evidence for a Late Triassic Shan–Thai orogeny. *Geosci J* 10(3):181–194
- Ivan P (2002) Relics of the Meliata Ocean crust: Geodynamic implications of mineralogical, petrological and geochemical proxies. *Geol Carpath* 53(4):245–256
- Kazmin VG (1990) Early Mesozoic reconstruction of the Black Sea–Caucasus region. Evolution of the northern margin of the Tethys. *Mémoires de la Société géologique de France. Nouvelle Series* 54:147–158
- Kazmin VG (1991) Collision and rifting in the Tethys Ocean: geodynamic implications. *Tectonophysics* 123:371–384
- Keppie JD, Dostal J (2001) Evaluation of the Baja controversy using paleomagnetic and faunal data, plume magmatism, and piercing points. *Tectonophysics* 339:427–442
- Kiessling W, Flügel E, Golonka J (1999) Paleo Reef Maps: Evaluation of a comprehensive database on Phanerozoic reefs: *Am Assoc Petrol Geol Bull* 83: 1552–1587
- Kiessling W, Flügel E, Golonka J (2003) Patterns of Phanerozoic carbonate platform sedimentation. *Lethaia* 36:195–226
- Knight KB, Nomade S, Renne PR, Marzoli A, Bertrand H, Youbi N (2004) The Central Atlantic magmatic province at the Triassic–Jurassic boundary: paleomagnetic and $^{40}\text{Ar}/^{39}\text{Ar}$ evidence from Morocco for brief, episodic volcanism. *Earth Planet Sci Lett* 228:143–160

- Köppen A, Carter A (2000) Constraints on provenance of the central European Triassic using detrital zircon fission track data. *Palaeogeogr Palaeoclimatol Palaeoecol* 161:193–204
- Kotański Z (1961) Tectogenesis and paleogeographic reconstruction of the Hg-Tatra chain in Tatra Mountains (In Polish, English Summary). *Acta Geologica Polonica* 11: 401–415
- Kovalenko DV (2010) Paleomagnetism of Late Paleozoic, Mesozoic, and Cenozoic rocks in Mongolia. *Russ Geol Geophys* 51(4):387–403
- Kozur H (1991) The evolution of the Meliata-Halstatt ocean and its significance for the early evolution of the Eastern Alps and Western Carpathians. *Palaeogeogr Palaeoclimatol Palaeoecol* 87:109–130
- Kozur H, Krahel W (1987) Erster Nachweis von Radiolarien in tethyalen Perm Europas. *N Jahrb Geol Paleontol Abh* 174:357–372
- Kravchinsky VA, Cogne J-P, Harbert W, Kuzmin MI (2002) Evolution of the Mongol-Okhotsk Ocean as constrained by new paleomagnetic data from the Mongol-Okhotsk suture zone, Siberia. *Geophys J Internat* 148:34–57
- Krzywiec P (2012) Mesozoic and Cenozoic evolution of salt structures within: the Polish Basin: an overview. *Geol Soc Spec Publ* 363(1):381–394
- Kutek J (2001) The Polish Permo-Mesozoic Rift Basin. In: Ziegler PA, Cavazza W, Robertson AHF, Crasquin-Soleau S (eds) *Peri-Tethys Memoir 6: Peri-Tethyan Rift/Wrench Basins and Passive margins*. Mémoires du Muséum national d'Histoire naturelle 186: 213–236
- Lane LS, Stephenson RA (2016) Circum-Arctic lithosphere-basin evolution: An overview. *Tectonophysics* 691:1–7
- Laville E, Pique A, Amrhar M, Charroud M (2004) A restatement of the Mesozoic Atlantic Rifting (Morocco). *J Afric Earth Sci* 38:145–153
- Leith TL, Weiss HM, Mørk A, Arhus N, Elvebakk N, Embry A F, Brooks P W, Stewart KR, Pchelina TM, Bro EG, Verba M L, Danushevskaya A, Borisov AV (1993) Mesozoic hydrocarbon source rocks of the Arctic region. In: Vorren TO, Bergsær E, Dahl-Stammes OA, Holter E, Johansen B, Lie E, Lund TB (eds) *Arctic geology and petroleum potential*. *Norw Petrol Geol Spec Publ* 2: 1–25
- Lepvrier C, Maluski H (2008) The Triassic Indosinian orogeny in East Asia. *Compt Rendus Geosci* 340:75–82
- Lepvrier C, Maluski H, Layreloup A, Van Tich V, Thi PT, Vuong NV (2004) The Early Triassic Indosinian orogeny in Vietnam (Truong Son Belt and Kontum Massif): implications for the geodynamic evolution of Indochina. *Tectonophysics* 393:87–118
- Li H-Y, Huang X-L (2013) Constraints on the paleogeographic evolution of the North China Craton during the Late Triassic-Jurassic. *J Asian Earth Sci* 70-71(1):308–320
- Li S, Wilde SA, He Z, Jiang X, Liu R, Zhao L (2014) Triassic sedimentation and postaccretionary crustal evolution along the Solonker suture zone in Inner Mongolia, China. *Tectonics* 33(6):960–981
- Li X, Zheng J, Li S, Liu B, Xiang L, Wang Y, Liu X (2016a) Late Triassic orogenic collapse and Palaeo-Pacific slab roll-back beneath central South China: constraints from mafic granulite xenoliths and structural features. *Geol J* 51:123–136
- Li Z, Ding L, Lippert PC, Song P, Yue Y, van Hinsbergen DJJ (2016b) Paleomagnetic constraints on the Mesozoic drift of the Lhasa terrane (Tibet) from Gondwana to Eurasia. *Geol* 44(9):727–730
- Liang W, Zhang G, Bai Y, Jin C, Nantasin P (2015) New insights into the emplacement mechanism of the Late Triassic granite plutons in the Qinling orogen: A structural study of the Mishuling pluton. *Geol Soc Am Bull* 127(11–12):1583–1603
- Liu X, Hsu KJ, Ju Y, Li G, Liu X, Wei L, Zhou X, Zhang X (2012) New interpretation of tectonic model in south Tibet. *J Asian Earth Sci* 56:147–159
- Lu YH, Zhao Z-F, Zheng Y-F (2016) Geochemical constraints on the source nature and melting conditions of Triassic granites from South Qinling in central China. *Lithos* 264:141–157
- Lucas SG, Tanner LH (2008) Reexamination of the end-Triassic mass extinction. In: Elewa AMT (ed) *Mass Extinction*. Springer, p 66–103

- Luo M, Lu L, Jia J, Wang S, Xu Y, He W (2014) Evolution of sedimentary basins in China during Mesozoic. *Earth Sci J China Univ Geosci* 39(8):954–976
- Macdonald D, Gomez-Perez I, Francese J, Spalletti L, Lawver L, Gahagan L, Dalziel I, Thomas C, Trewin N, Hole M, Paton D (2003). Mesozoic break-up of SW Gondwana: implications for regional hydrocarbon potential of the southern South Atlantic. In: Golonka J (ed) Thematic set on paleogeographic reconstruction and hydrocarbon basins: Atlantic, Caribbean, South America, Middle East, Russian Far East, Arctic. *Mar Petrol Geol* 20: 287–308
- Maluski H, Lepvrier C, Jolivet L, Carter A, Roques D, Beyssac O, Ta TT, Thang ND, Avigad D (2001) Ar–Ar and fission-track ages in the Song Chay Massif: Early Triassic and Cenozoic tectonics in Northern Vietnam. *J Asian Earth Sci* 19:233–248
- Maluski H, Lepvrier C, Layrelop A, Van Tich V, Thi PT (2005) ^{40}Ar – ^{39}Ar geochronology of the charnokites and granulites of the Kan Nack Complex, Kon Tun Massif, Vietnam. *J Asian Earth Sci* 25:653–677
- Marsella E, Kozur H, D'Argenio B (1993) Monte Facito Formation (Scythian - Middle Carnian. A deposit of the ancestral Lagonegro Basin in Southern Apennines. *Bolletino de Servizio Geologico Italia* 119: 225–248
- Marzoli A, Bertrand H, Knight KB, Cirilli S, Buratti N, Verati C, Nomade S, Renne PR, Youbi N, Martini R, Allenbach K, Neuwerth R, Rapaille C, Zaninetti L, Bellieni G (2004) Synchrony of the Central Atlantic magmatic province and the Triassic-Jurassic boundary climatic and biotic crisis. *Geol* 32:973–976
- Marzoli A, Jourdan F, Puffer JH, Cuppone T, Tanner LH, Weems RE, Bertrand H, Cirilli S, Bellieni G, De Min A (2011) Timing and duration of the Central Atlantic magmatic province in the Newark and Culpeper basins, eastern U.S.A. *Lithos* 122(3–4):175–188
- Marzoli A, Renne PR, Piccirillo EM, Ernesto M, Bellieni G, De Min A (1999) Extensive 200-million-year-old continental flood basalts of the Central Atlantic Magmatic Province. *Science* 284:616–618
- Masoodi M, Yassaghi A, Nogole Sadat MAA, Neubauer F, Bernroider M, Friedl G, Genser J, Houshmandzadeh A (2013) Cimmerian evolution of the Central Iranian basement: Evidence from metamorphic units of the Kashmar-Kerman Tectonic Zone. *Tectonophysics* 588:189–208
- Matthews KJ, Maloney KT, Zahirovic S, Williams SE, Seton M, Müller RD (2016) Global plate boundary evolution and kinematics since the late Paleozoic. *Glob Planet Ch* 46:226–250
- McKie T, Williams B (2009) Triassic palaeogeography and fluvial dispersal across the northwest European basins. *Geol J* 44(6):711–741
- Meesook A, Sha J (2010) The Jurassic system of Thailand. Univ Sci Technol China Press, Hefe
- Mehdipour Ghazi J, Moazzen M (2015) Geodynamic evolution of the Sanandaj-Sirjan Zone, Zagros Orogen, Iran. *Turk J Earth Sci* 24(5):513–528
- Meinhold G, Kostopoulos DK (2013) The Circum-Rhodope Belt, northern Greece: Age, provenance, and tectonic setting. *Tectonophysics* 595-596:55–68
- Meng Y, Xu Z, Santosh M, Ma X, Chen X, Guo G, Liu F (2016) Late Triassic crustal growth in southern Tibet: Evidence from the Gangdese magmatic belt. *Gond Res* 37:449–464
- Metcalfe I (1994) Late Paleozoic and Mesozoic Paleogeography of Eastern Pangaea and Tethys. In: Embry AF, Beauchamp B, Glass DJ (eds) Pangaea; global environments and resources. *Can Soc Petrol Geol Mem* 17: 97–111
- Metcalfe I (1996) Gondwanaland dispersion, Asian accretion and evolution of eastern Tethys, In: Li Z X., Metcalfe I, Powell CM (eds) Breakup of Rodinia and Gondwanaland and assembly of Asia. *Austral J Earth Sci* 43: 605–623
- Metcalfe I (2000) The Bentong-Raub Suture zone. *J Asian Earth Sci* 18:691–712
- Metcalfe I (2011) Palaeozoic-Mesozoic history of SE Asia. In: Hall R, Cottam MA, Wilson MEJ (eds) The SE Asian Gateway: history and tectonics of the Australia–Asia Collision. *Geol Soc Spec Publ* 355: 7–35
- Metcalfe I (2013a) Gondwana dispersion and Asian accretion: Tectonic and palaeogeographic evolution of eastern Tethys. *J Asian Earth Sci* 66:1–33
- Metcalfe I (2013b) Tectonic evolution of the Malay Peninsula. *J Asian Earth Sci* 76:195–213

- Maurer F, Rettori R, Martini R (2008) Triassic stratigraphy, facies and evolution of the Arabian shelf in the northern United Arab Emirates. *Internat J Earth Sci* 97(4):765–784
- Metelkin DV, Kazansky AY, Vernikovskiy VA (2011) Paleomagnetic evidence for Siberian plate tectonics from Rodinia through Pangaea to Eurasia. In: Closson D (ed) *Tectonics*. InTech, Zagreb, pp 159–236
- Metelkin DV, Vernikovskiy VA, Kazansky AY (2012) Tectonic evolution of the Siberian paleocontinent from the Neoproterozoic to the Late Mesozoic: Paleomagnetic record and reconstructions. *Russ Geol Geophys* 53(7):675–668
- Miall AD, Blakey RC (2008) The Phanerozoic Tectonic and Sedimentary Evolution of North America. *Sedimentary Basins of the World* 5(C):1–29
- Miall AD, Balkwill HR, McCracken J (2008) The Atlantic Margin Basins of North America. *Sedimentary Basins of the World* 5(C):473–504
- Miller EL, Toro J, Gehrels G, Amato JM, Prokopiev A, Tuchkova MI, Akinin VV, Dumitru TA, Moore TE, Cecile MP (2006) New insights into Arctic paleogeography and tectonics from U-Pb detrital zircon geochronology. *Tecton* 25(3.) TC3013
- Mock R, Sýkora M, Aubrecht R, Ožvoldová L, Kronome B, Reichwalder P, Jablonský J (1998) Petrology and stratigraphy of the Meliaticum near the Meliata and Jaklovce Villages, Slovakia. *Slovak Geol Mag* 4:223–260
- Montenat C (2009) The Mesozoic of Afghanistan. *GeoArabia* 14(1):147–210
- Morris A (1996) A review of paleomagnetic research in the Troodos ophiolite, Cyprus. In: Morris A, Tarling DH (eds) *Paleomagnetism and Tectonics of the Mediterranean Region*. Geol Soc London Spec Publ 105:311–324
- Müller RD, Seton M, Zahirovic S, Williams SE, Matthews KJ, Wright NM, Shephard GE, Maloney KT, Barnett-Moore N, Hosseinpour M, Bower DJ, Cannon J (2016) Ocean basin evolution and global-scale plate reorganization events since Pangaea breakup. *Annu Rev Earth Planet Sci* 44:107–138
- Muttoni G, Gaetani M, Budurov K, Zagorchev I, Trifonova E, Ivanova D, Petrounova L, Lowrie W (2000) Middle Triassic paleomagnetic data from northern Bulgaria: constraints on Tethyan magnetostratigraphy and paleogeography. *Palaeogeogr Palaeoclimatol Palaeoecol* 160:223–237
- Nikishin AM, Ziegler PA, Cloething S, Stephenson RA, Furne AV, Fokin PA, Ershov AV, Bolotov SN, Koraeve MV, Alekseev AS, Gorbachev I, Shipilov EV, Lankrejer A, Shalimov IV (1996) Late Precambrian to Triassic history of the East European Craton: dynamics of sedimentary basin evolution. *Tectonophysics* 268:23–63
- Nikishin AM, Cloething S, Bolotov SN, Baraboshkin EY, Kopaevich LF, Nazarevich BP, Panov DI, Brunet M-F, Ershov AV, Il'ina VV, Kosova SS., Stephenson RA (1998a) Scythian platform: chronostratigraphy and polyphase stages of tectonic history. In: Crasquin-Soleau S, Barrier E (eds) *Peri-Tethys Memoir 3: Stratigraphy and Evolution of Peri-Tethyan Platforms*. Mémoires du Muséum national d'Histoire naturelle 177: 151–162
- Nikishin AM, Cloething S, Brunet M-F, Stephenson RA, Bolotov SN, Ershov AV (1998b) Scythian Platform and Black Sea region: Mesozoic-Cenozoic tectonic and dynamics. In: Crasquin-Soleau S, Barrier E (eds) *Peri-Tethys Memoir 3: Stratigraphy and Evolution of Peri-Tethyan Platforms*. Mémoires du Muséum national d'Histoire naturelle 177: 163–176
- Nikishin AM, Ziegler PA, Abbott D, Brunet M-F, Cloething S (2002) Permo-Triassic intraplate magmatism and rifting in Eurasia: implications for mantle plumes and mantle dynamics. *Tectonophysics* 351:3–39
- Nouri F, Azizi H, Golonka J, Asahara Y, Orihashi Y, Yamamoto K, Tsuboi M, Anma R (2016) Age and petrogenesis of Na-rich felsic rocks in western Iran: evidence for closure of the southern branch of the Neo-Tethys in the Late Cretaceous. *Tectonophysics* 671:151–172
- Ogg JG, Ogg GM, Gradstein FM (2016) *A concise geologic time scale 2016*. Elsevier, Amsterdam
- Okay AI, Nikishin AM (2015) Tectonic evolution of the southern margin of Laurasia in the Black Sea region. *Internat Geol Rev* 57(5–8):1051–1076
- Oliver G, Prave A (2013) Palaeogeography of Late Triassic red-beds in Singapore and the Indosinian Orogeny. *J Asian Earth Sci* 76:214–224

- Olsen PE, Kent DV (1996) Milankovitch climate forcing in the tropics of Pangaea during the Late Triassic. *Palaeogeogr Palaeoclimatol Palaeoecol* 122:1–26
- Olsen PE (1997) Stratigraphic record of the early Mesozoic breakup of Pangaea in the Laurasia-Gondwana rift system. *Annu Rev Earth Planet Sci* 25:337–401
- Panushka BC (1985) Paleomagnetic evidence for a post-Cretaceous accretion of Wrangellia. *Geol* 13:880–883
- Pčelina TM, Korčinskaja MV (2008) Palaeogeographic reconstructions of the Russian Boreal areas and Svalbard during the Triassic. *Polar Res* 27(3):491–494
- Pease VL, Kuzmichev AB, Danukalova MK (2015) The new Siberian Islands and evidence for the continuation of the Uralides, Arctic Russia. *J Geol Soc* 172(1):1–4
- Peng Z-M, Peng S-M, Wu Z-P, Li W, Kong X (2009) Basin pattern and evolution of Triassic in North China. *J Xi'an Shiyu Univ, Nat Sci Ed* 24(2):34–38
- Petrík I, Janák M, Froitzheim N, Georgiev N, Yoshida K, Sasinková V, Konečný P, Milovská S (2016) Triassic to Early Jurassic (c. 200 Ma) UHP metamorphism in the Central Rhodopes: Evidence from U-Pb-Th dating of monazite in diamond-bearing gneiss from Chepelare (Bulgaria). *J Met Geol* 34(3):265–291
- Peybernes C, Chablais J, Onoue T, Escarguel G, Martini R (2016) Paleocology, biogeography, and evolution of reef ecosystems in the Panthalassa Ocean during the Late Triassic: Insights from reef limestone of the Sambosan Accretionary Complex, Shikoku, Japan. *Palaeogeogr Palaeoclimatol Palaeoecol* 457:31–51
- Piercey SJ, Nelson JL, Colpron M, Dusel-Bacon C, Simard R-L, Roots CF (2006) Paleozoic magmatism and crustal recycling along the ancient Pacific margin of North America, northern Cordillera. In: Colpron M, Nelson JL (eds) *Paleozoic Evolution and Metallogeny of Pericratonic Terranes at the Ancient Pacific Margin of North America*. Canadian and Alaskan Cordillera *Geol Ass Can Spec Pap* 45: 281–322
- Preto N, Kustatscher E, Wignall PB (2010) Triassic climates - State of the art and perspectives. *Palaeogeogr Palaeoclimatol Palaeoecol* 290(1–4):1–10
- Puchkov N (1997) Structure and geodynamics of the Uralian Orogen. In: Burg J-P, Ford M (eds) *Orogeny through time*. *Geol Soc London Spec Publ* 121: 201–236
- Richards JP (2015) Tectonic, magmatic, and metallogenic evolution of the Tethyan orogen: From subduction to collision. *Ore Geol Rev* 70:323–345
- Ridd MF (2013) A Middle Permian-Middle Triassic accretionary complex and a Late Triassic foredeep basin: Forerunners of an Indosinian (Late Triassic) thrust complex in the Thailand-Malaysia border area. *J Asian Earth Sci* 76:99–114
- Robertson AHF (1998) Mesozoic-tertiary tectonic evolution of the Easternmost Mediterranean area: integration of the Marine and land evidence. In: Robertson AHF, Richter C, Camerlenghi CA (eds) *Proceedings of the Ocean Drilling Program Scientific Results* 60: 723–782
- Robertson AHF (2007) Overview of tectonic settings related to the rifting and opening of Mesozoic ocean basins in the Eastern Tethys: Oman, Himalayas and Eastern Mediterranean regions. *Geol Soc Spec Publ* 282:325–388
- Robertson AHF, Woodcock NH (1979) Mamonia complex, southwest Cyprus: evolution and emplacement of a Mesozoic continental margin. *Geol Soc Am Bull* 90:651–665
- Robertson AHF, Clift PD, Degnan P, Jones G (1991) Paleogeographic and paleotectonic evolution of Eastern Mediterranean Neotethys. *Palaeogeogr Palaeoclimatol Palaeoecol* 87:289–344
- Robertson AHF, Dixon JE, Brown S, Collins A, Morris A, Pickett EA, Sharp I, Ustaömer T (1996) Alternative tectonic models for the Late Palaeozoic-Early Tertiary development of Tethys in the Eastern Mediterranean region. In: Morris A, Tarling DH, (eds) *Palaeomagnetism and tectonics of the Mediterranean region*. Geological Society London Special Publication 105: 239–263
- Robertson AHF, Ustaömer T, Pickett EA, Collins AS, Andrew T, Dixon JE (2004) Testing models of Late Palaeozoic-Early Mesozoic orogeny in Western Turkey: Support for an evolving open-Tethys model. *J Geol Soc* 161(3):501–511
- Robinson AC (2015) Mesozoic tectonics of the Gondwanan terranes of the Pamir plateau. *J Asian Earth Sci* 102:170–179

- Roniewicz E, Stanley Jr GD (2013) Upper Triassic corals from Nevada, Western North America, and the implications for paleoecology and paleogeography. *J Paleontol* 87(5): 934–964
- Ronov A, Khain V, Balukhovskiy A (1989) Atlas of Lithological Paleogeographical Maps of the World: Mesozoic and Cenozoic of the Continents. USSR Acad Sci, Leningrad
- Ruffell A, Simms M, Wignall P (2015) The Carnian Humid Episode of the late Triassic: a review. *Geol Mag* 153:271–284
- Rychliński T (2008) Facies development and sedimentary environments of the Carpathian Keuper deposits from the Tatra mountains. Poland and Slovakia *Annales Societatis Geologorum Poloniae* 78(1):1–18
- Săndulescu M (1988) Cenozoic tectonic history of the Carpathians. In: Royden L, Horváth F (eds) *The Pannonian Basin: a study in basin evolution*. *Am Assoc Petrol Geol Mem* 45: 17–25
- de Saussure le fils M (1792) Analyse de la dolomie. *J Physique* 40:161–173
- Scheck-Wenderoth M, Krzywiec P, Zühlke R, Maystrenko Y, Froitzheim N (2008) Permian to Cretaceous tectonics. *Geol Centr Eur* 6(2):999–1030
- Schettino A, Turco E (2011) Tectonic history of the Western Tethys since the Late Triassic. *Geol Soc Am Bull* 123(1–2):89–105
- Schmid SM, Bernoulli D, Fügenschuh B, Matenco L, Schefer S, Schuster R, Tischler M, Ustaszewski K (2008) The Alpine-Carpathian-Dinaridic orogenic system: Correlation and evolution of tectonic units. *Swiss J Geosci* 101(1):139–183
- Schwarzacher W (2006) The stratification and cyclicity of the Dachstein Limestone in Lofer, Leogang and Steinernes Meer (Northern Calcareous Alps, Austria). *Sed Geol* 181:93–106
- Scotese CR (2004) A Continental Drift Flipbook. *J Geol* 112:729–741
- Sengör AMC (1984) The Cimmeride orogenic system and the tectonics of Eurasia. *Geol Soc Am Spec Pap* 195:1–82
- Sengör AMC, Natalin BA (1996) Paleotectonics of Asia: fragment of a synthesis. In: An Y, Harrison TM (eds) *The Tectonic Evolution of Asia*. Cambridge Univ Press, Cambridge
- Sengör AMC, Yilmaz Y, Sungurlu O (1984) Tectonics of the Mediterranean Cimmerides: Nature and evolution of the western termination of Palaeo-Tethys. *Geol Soc London Spec Publ* 17:77–112
- Shi Y, Anderson JL, Li L, Ding J, Liu C, Zhang W, Shen C (2016) Zircon ages and Hf isotopic compositions of Permian and Triassic A-type granites from central Inner Mongolia and their significance for late Palaeozoic and early Mesozoic evolution of the Central Asian Orogenic Belt. *Intern Geol Rev* 58(8):967–982
- Sibuet J-C, Rouzo S, Srivastava S (2012) Plate tectonic reconstructions and paleogeographic maps of the central and North Atlantic oceans. *Can J Earth Sci* 49(12):1395–1415
- Song P, Ding L, Li Z, Lippert PC, Yang T, Zhao X, Fu J, Yue Y (2015) Late Triassic paleolatitude of the Qiangtang block: Implications for the closure of the Paleo-Tethys Ocean. *Earth Planet Sci Lett* 424:69–83
- Spikings R, Reitsma MJ, Boekhout F, Mišković A, Ulianov A, Chiaradia M, Gerdes A, Schaltegger U (2016) Characterisation of Triassic rifting in Peru and implications for the early disassembly of western Pangaea. *Gond Res* 35:124–143
- Stanley GD, Onoue T (2015) Upper Triassic reef corals from the Sambosan Accretionary Complex, Kyushu, Japan. *Facies* 61(2):1–27
- Tanner LH (2010) Cyclostratigraphic record of the Triassic: A critical examination. In: Lucas SG (ed) *The Triassic Timescale*. *Geol Soc London Spec Publ* 334: 119–137
- Tari G, Dicea O, Faulkerson J, Georgiev G, Popov S, Stefanescu M, Weir G (1997) Cimmerian and Alpine stratigraphy and structural evolution of the Moesian Platform (Romania/Bulgaria). In: Robinson A G (ed) *Regional and petroleum geology of the Black Sea and surrounding regions*. *Am Assoc Petrol Geol Mem* 68: 63–90
- Tong J, Yin H (2002) The Lower Triassic of South China. *J Asian Earth Sci* 20:803–815
- Toro J, Miller EL, Prokopiev AV, Zhang X, Veselovskiy R (2016) Mesozoic orogens of the Arctic from Novaya Zemlya to Alaska. *J Geol Soc* 173:989–1006
- Torsvik TH, Anderson TB (2002) The Taimyr fold belt, Arctic Siberia: timing of prefold remagnetisation and regional tectonics. *Tectonophysics* 352:335–348

- Trop JM, Ridgway KD, Manuszak JD, Layer P (2002) Mesozoic sedimentary-basin development on the allochthonous Wrangellia composite terrane, Wrangell Mountains basin, Alaska: a long-term record of terrane migration and arc construction. *Geol Soc Am Bull* 114:693–717
- Uno K, Furukawa K, Hada S (2011) Margin-parallel translation in the western Pacific: Paleomagnetic evidence from an allochthonous terrane in Japan. *Earth Planet Sci Lett* 303(1–2):153–165
- Veevers JJ (2006) Updated Gondwana (Permian–Cretaceous) earth history of Australia. *Gond Res* 9(3):231–260
- Veevers JJ (2013) Pangaea: Geochronological correlation of successive environmental and stratigraphic phases in Europe and Australia. *Earth-Sci Rev* 127:48–95
- Veevers JJ (1994) Pangaea: Evolution of a supercontinent and its consequences for Earth's paleoclimate and sedimentary environments. In: Klein GD (ed) *Pangaea: Paleoclimate, Tectonics and Sedimentation during Accretion, Zenith and Breakup of a Supercontinent*. *Geol Soc Am Spec Pap* 228: 13–23
- Vernikovsky VA (1995) The geodynamic evolution of the Taimyr folded area. In: Simakov KV, Thurston DK (eds) *Proceedings of the International Conference on Arctic Margins*, Magadan, September, p. 186–193
- Vernikovsky VA, Pease VL, Vernikovskaya AE, Romanov AP, Gee DG, Travin AV (2003) First report of early Triassic A-type granite and syenite intrusions from Taimyr: Product of the northern Eurasian superplume. *Lithos* 66(1–2):23–36
- Vernikovsky VA, Metelkin DV, Tolmacheva TY, Malyshev NA, Petrov OV, Sobolev NN, Matushkin NY (2013) Concerning the issue of paleotectonic reconstructions in the Arctic and of the tectonic unity of the New Siberian Islands Terrane: New paleomagnetic and paleontological data. *Doklady Earth Sci* 451(2):791–797
- Vinogradov AP (ed) (1968) *Atlas of the Lithological-Paleogeographical Maps of the USSR*. Vol. III: Triassic, Jurassic and Cretaceous. *Ministr Geol USSR, Acad Sci USSR*
- von Alberti F (1834) *Beitrag zu einer Monographie des bunten Sandsteins, Muschelkalks und Keupers, und die Verbindung dieser Gebilde zu einer Formation*. Mit 2 Taf. lith. Stuttgart & Tübingen
- Wang B, Zhang G, Yang Z (2013) New Mesozoic paleomagnetic results from the northeastern Sichuan basin and their implication. *Tectonophysics* 608:418–427
- Wignall PB (2001) Large igneous provinces and mass extinctions. *Earth Sci Rev* 53(1–2):1–33
- Wilmsen M, Fürsich FT, Seyed-Emami K, Majidifard MR, Taheri J (2009) The Cimmerian Orogeny in northern Iran: Tectono-stratigraphic evidence from the foreland. *Terra Nova* 21(3):211–218
- Swe W (2012) Outline geology and economic mineral occurrences of the Union of Myanmar. *J Myanmar Geosci Soc* 1:1–215
- Withjack MO, Schlische RW, Olsen PO (1998) Diachronous rifting, drifting, and inversion on the passive margin of central eastern North America: an analog for other passive margins. *Am Assoc Petrol Geol Bull* 82:817–835
- Wu F, Sun D, Li H, Bor M, Wilde S (2002) A-type granites in northeastern China: age and geochemical constraints on their petrogenesis. *Chem Geol* 187:143–173
- Wu H, Li C, Chen J, Xie C (2016) Late Triassic tectonic framework and evolution of Central Qiangtang, Tibet, SW China. *Lithosphere* 8(2):141–149
- Yin A, Nie S (1996) A Phanerozoic palinspastic reconstruction of China and its neighboring regions. In: Yin AN, Harrison TM (eds) *The Tectonic Evolution of Asia*. Cambridge Univ Press, Cambridge, p. 442–485
- Zanchi A, Zanchetta S, Balini M, Ghassemi MR (2016) Oblique convergence during the Cimmerian collision: Evidence from the Triassic Aghdarband Basin, NE Iran. *Gond Res* 38:149–170
- Zanchi A, Zanchetta S, Berra F, Mattei M, Garzanti E, Molyneux S, Nawab A, Sabouri J (2009) The Eo-Cimmerian (Late? Triassic) orogeny in North Iran. *Geol Soc London Spec Publ* 312:31–55
- Zeng WS, Zhou JB, Dong C, Cao JL, Wang B (2014) Mongolia - Okhotsk oceanic subduction record: Evidence Ergun metamorphic complex area of the mall. *Acta Petrol Sin* 30(7):1948–1960
- Zhai Q-G, Jahn B-M, Su L, Wang J, Mo X-X, Lee H-Y, Wang K-L, Tang S (2013) Triassic arc magmatism in the Qiangtang area, northern Tibet: Zircon U-Pb ages, geochemical and Sr-Nd-Hf isotopic characteristics, and tectonic implications. *J Asian Earth Sci* 63:162–178

- Zhang X, Pease V, Carter A, Scott R (2017b). Reconstructing Palaeozoic and Mesozoic tectonic evolution of Novaya Zemlya: combining geochronology and thermochronology. In: Pease V and Coakley B (eds) Circum-Arctic Lithosphere Evolution. Geol Soc London Spec Publ 460. doi.org/10.1144/SP460.13
- Zhang X, Pease V, Carter A, Kostuychenko S, Suleymanov A, Scott R (2017a) Timing of exhumation and deformation across the Taimyr fold–thrust belt: insights from apatite fission track dating and balanced cross-sections In: Pease V, Coakley B (eds) Circum-Arctic Lithosphere Evolution. Geol Soc London Spec Publ 460, doi.org/10.1144/SP460.3
- Zharkov MA, Chumakov NM (2001) Paleogeography and sedimentation settings during Permian–Triassic reorganization in biosphere. *Stratigr Geol Correl* 9:340–363
- Zhou Y, Cheng X, Yu L, Yang X, Su H, Peng X, Xue Y, Li Y, Ye Y, Zhang J, Li Y, Wu H (2016) Paleomagnetic study on the Triassic rocks from the Lhasa Terrane, Tibet, and its paleogeographic implications. *J Asian Earth Sci* 121:108–119
- Zhu T, Feng X, Wang X, Zhou M (2013) Reconstruction of the Triassic tectonic lithofacies paleogeography in Qiangtang region, Northern Qinghai-Tibet Plateau, China. *Acta Geol Sin* 87(2):378–394
- Ziegler PA (1982) Geological atlas of western and central Europe. Shell Internationale Petroleum Mij, B.V., The Hague
- Ziegler PA (1988) Evolution of the Arctic North Atlantic and the Western Tethys. *Am Assoc Petrol Geol Mem* 43:1–198
- Zonenshain LP, Kuzmin ML, Natapov LN (1990) Geology of the USSR: A Plate-Tectonic Synthesis. *Am Geophys Un, Geodynam. Ser* 21:1–242

The Late Triassic World

Earth in a Time of Transition

Tanner, L. (Ed.)

2018, XI, 805 p. 190 illus., 115 illus. in color., Hardcover

ISBN: 978-3-319-68008-8