

Table 3. F, Shortened constancy table of the meadow birch forest (Geranio-Betuletum)

	Geranio-Betuletum		total/ mean
	subass. of Trollius eur.	typical SA	
no. of relevés	22	24	46
no. of species (total)	36.8	32.8	34.8
height of birch (m)	7.8	6.9	7.4
height of <i>Melanelia olivacea</i> (m)	0.8	0.7	0.8
coverage total (%)	94.6	95.0	94.8
tree layer (%)	55.7	58.6	57.2
shrub layer (%)	16.3	21.2	18.7
herb layer (%)	86.8	83.3	85.1
moss layer (%)	23.6	28.5	26.1
lichen layer (%)	0.3	0.6	0.4
tree layer			
<i>Betula pubescens</i> ssp. <i>czerepanovii</i>	V	V	V
<i>Salix phylicifolia</i>	+	I	+
shrub layer			
<i>Betula pubescens</i> ssp. <i>czerepanovii</i>	IV	III	III
<i>Juniperus communis</i>	IV	III	III
<i>Salix glauca</i>	II	III	III
field layer			
<i>Geranium sylvaticum</i>	V	IV	V
<i>Solidago virgaurea</i>	V	V	V
<i>Rubus saxatilis</i>	IV	IV	IV
<i>Trollius europaeus</i>	V	II	IV
<i>Saussurea alpina</i>	IV	IV	IV
<i>Vaccinium vitis-idaea</i>	IV	III	III
<i>Trientalis europaea</i>	IV	III	III
<i>Vaccinium uliginosum</i>	III	III	III
<i>Barbilophozia lycopodioides</i>	III	III	III
<i>Equisetum pratense</i>	III	IV	III
<i>Hylocomium splendens</i>	IV	III	III
<i>Equisetum arvense</i>	III	III	III
<i>Pleurozium schreberi</i>	III	III	III
<i>Festuca ovina</i>	II	IV	III
<i>Deschampsia flexuosa</i>	III	II	III
<i>Filipendula ulmaria</i>	III	II	III
<i>Sanionia uncinata</i>	III	II	III
degree of presence: + = 1-10%, I = 11-20%, II = 21-40%, III = 41-60%, IV = 61-80%, V = 81-100%			