

Fig. 33B-11-001. $\text{Cs}_{1-x}(\text{NH}_4)_x\text{H}_2\text{AsO}_4$ (CDA-ADA). Θ vs. x [94Son]. PG: proton glass region. Open square: Θ_f ; open circle: freezing temperature; full circle: Θ_a ; open triangle: glass transition temperature. Hatched area: mixed region.




Fig. 33B-11-002. $\text{Cs}_{1-x}(\text{NH}_4)_x\text{H}_2\text{AsO}_4$ (CDA-ADA, polycrystal). κ' vs. T [94Son]. Parameter: x, f .




Fig. 33B-11-003. $\text{Cs}_{1-x}(\text{NH}_4)_x\text{H}_2\text{AsO}_4$ (CDA-ADA, $x = 0.22$, polycrystal). κ' , κ'' vs. T [94Son]. Parameter: f .




Fig. 33B-11-004. $\text{Cs}_{1-x}(\text{NH}_4)_x\text{H}_2\text{AsO}_4$ (CDA-ADA, $x = 0.20$). κ'_a , κ''_a vs. T [94Son]. Parameter: f .




Fig. 33B-11-005. $\text{Cs}_{1-x}(\text{NH}_4)_x\text{H}_2\text{AsO}_4$ (CDA-ADA, $x = 0.20$). κ'_c , κ''_c vs. T [94Son]. Parameter: f .




Fig. 33B-11-006. $\text{Cs}_{1-x}(\text{NH}_4)_x\text{H}_2\text{AsO}_4$ (CDA-ADA, $x = 0.22$, polycrystal). f_c vs. $(T - \Theta_0)^{-1}$ [94Son]. f_c : cutoff frequency. Θ_0 : characteristic temperature in Vogel-Fulcher law: $f_c = f_0 \exp[-E_c/(T - \Theta_0)]$. $\Theta_0 = 10.1$ K.