




**Fig. 46A-9-001.** (CH<sub>3</sub>)<sub>2</sub>NH<sub>2</sub>Ga(SO<sub>4</sub>)<sub>2</sub> · 6H<sub>2</sub>O. Crystal structure projected along [010] in phase II [95Pie]. *T* = 125 K.




**Fig. 46A-9-002.** (CH<sub>3</sub>)<sub>2</sub>NH<sub>2</sub>Ga(SO<sub>4</sub>)<sub>2</sub> · 6H<sub>2</sub>O. Crystal structure projected along [010] in phase III [95Pie]. *T* = 100 K. Unit cell is chosen with 2<sub>1</sub> axis at 1/4, *y*, 1/4.




**Fig. 46A-9-003.** (CH<sub>3</sub>)<sub>2</sub>NH<sub>2</sub>Ga(SO<sub>4</sub>)<sub>2</sub> · 6H<sub>2</sub>O.  $a$ ,  $b$ ,  $c$ ,  $\beta$ ,  $V$  vs.  $T$  [95Pie].




**Fig. 46A-9-004.** (CH<sub>3</sub>)<sub>2</sub>NH<sub>2</sub>Ga(SO<sub>4</sub>)<sub>2</sub> · 6H<sub>2</sub>O.  $\kappa_X$ ,  $\kappa_Y$ ,  $\kappa_Z$ ,  $1/\kappa_X$  vs.  $T$  [92Sob]. Relation between  $X$ ,  $Y$ ,  $Z$  directions and the crystallographic axes: see Fig. 46A-8-004.  $X \parallel P_s$ .




**Fig. 46A-9-005.** (CH<sub>3</sub>)<sub>2</sub>NH<sub>2</sub>Ga(SO<sub>4</sub>)<sub>2</sub> · 6H<sub>2</sub>O.  $\kappa'_X$  vs.  $T$  [92Sob]. Parameter:  $f$ .



**Fig. 46A-9-006.** (CH<sub>3</sub>)<sub>2</sub>NH<sub>2</sub>Ga(SO<sub>4</sub>)<sub>2</sub> · 6H<sub>2</sub>O. Cole-Cole diagram of complex dielectric constant [92Sob]. Parameter:  $T$ .



**Fig. 46A-9-007.** (CH<sub>3</sub>)<sub>2</sub>NH<sub>2</sub>Ga(SO<sub>4</sub>)<sub>2</sub> · 6H<sub>2</sub>O.  $\tau$  vs.  $T$  [92Sob].  $\tau$ : dielectric relaxation time. 1: relaxation of dipoles, 2: relaxation due to domain motion.



**Fig. 46A-9-008.**  $(\text{CH}_3)_2\text{NH}_2\text{Ga}(\text{SO}_4)_2 \cdot 6\text{H}_2\text{O}$ .  $P_s$  vs.  $T$  [92Sob].