

M6FRINTN2S

Fresnel's formulas as function of angle of incidence for $n_1 > n_2$

for $R_p = r_p^2$, $R_s = r_s^2$, and $T_p = 1 - R_p$, $T_s = 1 - R_s$

1. Amplitude reflection coefficients.

$$r_p(\theta) := \frac{\left(\frac{n_2}{n_1}\right) \cdot \cos\left(2 \cdot \frac{\pi}{360} \cdot \theta\right) - \sqrt{1 - \left[\left(\frac{n_1}{n_2}\right) \cdot \sin\left(2 \cdot \frac{\pi}{360} \cdot \theta\right)\right]^2}}{\frac{n_2}{n_1} \cdot \cos\left(2 \cdot \frac{\pi}{360} \cdot \theta\right) + \sqrt{1 - \left[\left(\frac{n_1}{n_2}\right) \cdot \sin\left(2 \cdot \frac{\pi}{360} \cdot \theta\right)\right]^2}}$$

$$r_s(\theta) := \frac{\cos\left(2 \cdot \frac{\pi}{360} \cdot \theta\right) - \left(\frac{n_2}{n_1}\right) \cdot \sqrt{1 - \left[\left(\frac{n_1}{n_2}\right) \cdot \sin\left(2 \cdot \frac{\pi}{360} \cdot \theta\right)\right]^2}}{\cos\left(2 \cdot \frac{\pi}{360} \cdot \theta\right) + \left(\frac{n_2}{n_1}\right) \cdot \sqrt{1 - \left[\left(\frac{n_1}{n_2}\right) \cdot \sin\left(2 \cdot \frac{\pi}{360} \cdot \theta\right)\right]^2}}$$


$$n_1 \equiv 1.5 \quad n_2 \equiv 1 \quad \theta \equiv 0..90$$

2. Reflection: Intensities


$$R_p(\theta) := r_p(\theta) \cdot \overline{r_p(\theta)}$$

$$R_s(\theta) := r_s(\theta) \cdot \overline{r_s(\theta)}$$

p - case




s - case



3. Transmission: Intensities

p - case

$$T_p(\theta) := 1 - R_p(\theta)$$



s - case

$$T_s(\theta) := 1 - R_s(\theta)$$

