

<b>Child-Turcotte stage and Pugh modification</b>	<b>A 1 point</b>	<b>B 2 points</b>	<b>C 3 points</b>
<b>1. Bilirubin (mg/dl, <math>\mu</math>mol/l)</b> <b>2. Albumin (g/dl)</b> <b>3. Ascites</b> <b>4. Encephalopathy</b> <b>5. Quick's value (%)*) (or INR)</b>	<b>&lt; 2.0 &lt; 35</b> <b>&gt; 3.5</b> <b>none</b> <b>none</b> <b>&gt; 70 (&lt; 1.7)</b>	<b>2–3.0 35–51</b> <b>2.8–3.5</b> <b>easy to treat</b> <b>stages I, II</b> <b>40–70 (1.8–2.3)</b>	<b>&gt; 3.0 &gt; 51</b> <b>&lt; 2.8</b> <b>difficult to treat</b> <b>stages III, IV</b> <b>&lt; 40 (&gt; 2.3)</b>
<b><i>Points score (used by Pugh)</i></b>	<b>5–6</b>	<b>7–9</b>	<b>10–15</b>
<b>Survival rate: 1 and 2 year(s)</b>	<b>100%; 85%</b>	<b>80%; 60%</b>	<b>45%; 35%</b>