

Patient's initials.: | | |
First name Second name

Date of birth: | : | : | : |
Day Month Year

Extravasation of cytotoxic agents – Documentation (I)

Cannula used: ☐ Butterfly® ☐ Venflon® ☐ Other.
Diameter G

Cannula fixated with:

Site of puncture: ☐ Left Arm ☐ Right Arm ☐ Port-a-cath system
☐ Forearm ☐ Antecubital fossa ☐ Central venous catheter
☐ Wrist ☐ Dorsum of hand
☐ Other:



Was it necessary to puncture the same limb more than once?

☐ Yes ☐ No

Where – in relation to the original puncture site – was the vein punctured?

☐ Proximal ☐ Distal ☐ Medial/lateral

Has the patient any of the following symptoms:

→ Upper blockage to inflow: ☐ Yes ☐ No
→ Lymphoedema (same arm): ☐ Yes ☐ No
→ Haematoma (same arm): ☐ Yes ☐ No

| Sequence of application | Amount | Substance or trade name | Volume | |
|-------------------------|--------|-------------------------|--------|-------------------------------------|
| 1. | mg | | in ml | <input type="radio"/> extravascular |
| 2. | mg | | in ml | <input type="radio"/> extravascular |
| 3. | mg | | in ml | <input type="radio"/> extravascular |
| 4. | mg | | in ml | <input type="radio"/> extravascular |
| 5. | mg | | in ml | <input type="radio"/> extravascular |

Estimated volume of extravasated drug: ml

Type of administration: ☐ i.v. ☐ i.a.
☐ Bolus ☐ Infusion ☐ Infusion pump

Date of birth: |__:__|__:__|__:__|
Day Month Year

First name Second name

Day Month Year

Extravasation of cytotoxic agents – Documentation (III)

| | ✓ = Applies | ↑ Deterioration | | | = No change | | | ↓ Improvement | | |
|-------------------------------|-------------------------------|-----------------|------------|------------|-------------|------------|------------|---------------|--|--|
| | Status post paravasationem | 1. Control | 2. Control | 3. Control | 4. Control | 5. Control | 6. Control | | | |
| Date | | | | | | | | | | |
| Doctors's paraph | | | | | | | | | | |
| Symptoms after extravasation: | | | | | | | | | | |
| Pain (burning, stinging) | | | | | | | | | | |
| Oedema | | | | | | | | | | |
| Erythema | | | | | | | | | | |
| Blistering | | | | | | | | | | |
| Discolouration | | | | | | | | | | |
| Induration | | | | | | | | | | |
| Functional impairment | | | | | | | | | | |
| Ulceration | | | | | | | | | | |
| Necrosis | | | | | | | | | | |
| Demarcation | | | | | | | | | | |
| Formation of eschar | | | | | | | | | | |
| Infection | | | | | | | | | | |
| Complete healing | | | | | | | | | | |

Extent of extravasation:

| Two biggest diameters \leftrightarrow in cm | | | | | | |
|---|--|--|--|--|--|--|
| | | | | | | |

Measures:[illegible]

Notes: